

Emotion dysregulation and emotional impulsivity among adults with attention-deficit/hyperactivity disorder: Results of a preliminary study.

By: John T. Mitchell, Christopher D. Robertson, Arthur D. Anastopolous, Rosemary O. Nelson-Gray, and Scott H. Kollins

Mitchell, J.T., [Robertson, C.D.](#), [Anastopoulos, A.D.](#), [Nelson-Gray, R.O.](#) & Kollins, S.H. (2012). Emotion dysregulation and emotional impulsivity among adults with attention-deficit/hyperactivity disorder: Results of a preliminary study. *Journal of Psychopathology and Behavioral Assessment*, 34(4): 510-519.

Made available courtesy of Springer. The original publication is available at:

<http://link.springer.com/article/10.1007%2Fs10862-012-9297-2>

*****Reprinted with permission. No further reproduction is authorized without written permission from Springer. This version of the document is not the version of record. Figures and/or pictures may be missing from this format of the document. *****

Abstract:

Recent reviews argue that emotion dysregulation is an important feature of attention-deficit/hyperactivity disorder (ADHD) and involves a failure to inhibit negative emotions that leads to negative affectively-driven impulsive behavior (i.e., emotional impulsivity). The goal of the current study was to assess (a) whether emotion dysregulation and emotional impulsivity was higher in a group of adults diagnosed with ADHD and (b) if the relationship between core ADHD symptoms (i.e., inattention and hyperactivity-impulsivity) and emotional impulsivity is mediated by emotion dysregulation symptoms. A group of adults with ($n = 18$) and without ($n = 23$) ADHD completed measures of core ADHD symptoms, emotion dysregulation, and emotional impulsivity. A series of one-way analyses of covariance indicated significant between-group differences in emotion dysregulation and emotional impulsivity when current depression and oppositional defiant disorder ratings were covaried. In addition, the relationship between ADHD symptoms and emotional impulsivity was mediated by emotion dysregulation symptoms. These findings suggest that emotion dysregulation and emotional impulsivity are higher in adults diagnosed with ADHD and that emotion dysregulation symptoms have predictive value beyond core ADHD symptoms.

Keywords: ADHD | emotion dysregulation | emotional impulsivity | clinical psychology | Attention-Deficit/Hyperactivity Disorder | behavioral assessment

Article:

Attention-deficit/hyperactivity disorder (ADHD) is a developmental disorder that persists into adulthood for the majority of cases (Barkley et al. 2008; Mannuzza et al. 1993; G. Weiss and Hechtman 1993) and occurs among approximately 4 % of adults (Kessler et al. 2006). Though developmentally inappropriate symptoms of inattention and hyperactivity-impulsivity are the two core symptom sets of the disorder (American Psychiatric Association 2000), recent reviews argue that emotion dysregulation is an additional feature of ADHD that should be incorporated into theoretical conceptualizations and diagnostic criteria (Barkley 2010; Martel 2009; Skirrow et al. 2009).

Even though there is no consistently agreed upon definition of emotion dysregulation (Putnam and Silk 2005), most accounts describe it as a multidimensional construct (Gratz and Roemer 2004) that includes (a) the lack of inhibition of behavior associated with strong negative or positive emotion and (b) subsequent failure to engage in self-regulatory actions, including self-soothing, refocusing attention, moderating the initial emotion, and organizing for coordinated action in the service of goal-directed behavior (Barkley 2010; Gottman and Katz 1989; Hinshaw 2003; Martel 2009; Melnick and Hinshaw 2000). According to one account of emotion dysregulation in ADHD (Barkley 2010), those with the disorder are less likely to inhibit their emotions, particularly those pertaining to frustration, impatience, and anger, as a result of deficient cognitive control. The outcome of this poor inhibitory process is emotional impulsivity, which “refers to the quickness or speed with which and the greater likelihood that an individual with ADHD will react with negative emotions in response to events relative to others of the same developmental level or age without ADHD” (p. 5, Barkley 2010). In other words, deficient emotion regulation involves a failure to inhibit negative emotions that leads to negative affectively-driven impulsive behavior in ADHD. Thus, emotions are less moderated by top-down, effortful executive control (Barkley 2010).

Empirical support for the inclusion of emotion dysregulation as a core feature of the disorder comes from several sources. First, neuroanatomical networks associated with ADHD also underlie emotional functioning (see Barkley 2010, for a review). For example, the anterior cingulate cortex and the lateral prefrontal cortex are identified in studies of ADHD (Bush et al. 2005; Paloyelis et al. 2007; Valera et al. 2007) and emotion regulation (Ochsner and Gross 2005). Emotion dysregulation may emerge in ADHD because of structural and functional deficits in top-down emotion regulation in these regions that support goal-directed behavior (Bush et al. 2000).

Second, several studies involving behavioral observations, rating scales, and clinician ratings have supported overlap between emotion dysregulation and ADHD among children and adults (Anastopoulos et al. 2011; Barkley and Fischer 2010; Barkley and Murphy 2010; Maedgen and Carlson 2000; Melnick and Hinshaw 2000; Reimherr et al. 2010; Reimherr et al. 2005; Walcott and Landau 2004). For instance, a recent study reported that emotion dysregulation is higher in

children with ADHD and that it plays an important role in determining functional impairment and comorbidity outcomes (Anastopoulos et al. 2011). Another study in adults with ADHD found that poor emotional self-control (a) was higher in a group of adults with ADHD than in clinical and community control groups and (b) added explanatory and predictive power to various forms of impairment (e.g., occupational, educational, criminal history, driving outcomes, marital satisfaction) above and beyond core inattentive and hyperactive-impulsive symptoms (Barkley and Murphy 2010). In a longitudinal sample of hyperactive children followed into adulthood, these findings were replicated in similar domains of impairment among those meeting ADHD criteria into adulthood (Barkley and Fischer 2010). In addition, clinician-rated emotion dysregulation symptoms of ADHD (i.e., items assessing emotional overreactivity, temper, and affective lability) were higher among adults with ADHD than non-diagnosed individuals (Reimherr et al. 2010; Reimherr et al. 2005). These emotion dysregulation symptoms were responsive to both stimulant and non-stimulant ADHD pharmacological treatments. Finally, evidence from studies indicating that ADHD is associated with other emotion-related abilities (Miller et al. 2011) and emotion-related constructs (Martel et al. 2009; Mitchell and Nelson-Gray 2006; Nigg et al. 2002) indirectly suggest a relationship between ADHD and emotion dysregulation. For example, in one study neuroticism, a construct that includes components of emotion dysregulation (Kokkonen and Pulkkinen 2001), was associated with ADHD symptoms (Mitchell and Nelson-Gray 2006).

To our knowledge, no studies to date have assessed how emotion dysregulation and emotional impulsivity (i.e., impulsive behavior accompanied by negative emotions) are related to one another in those with and without ADHD. Further, limited empirical work on emotion dysregulation and ADHD has been conducted, particularly among adults. The overall goal of this cross-sectional pilot study, therefore, was to assess how emotion dysregulation symptoms, emotional impulsivity, and ADHD are related. The first and central aim of this study was to assess whether emotion dysregulation and emotional impulsivity was higher in a group of adults diagnosed with ADHD than controls. The ADHD group was hypothesized to score higher than the control group on measures of both constructs. Given that co-occurring symptoms of other forms of psychopathology have overlapping emotion dysregulation features (Barkley 2010) and can also account for any relationship between ADHD and emotion dysregulation, current symptoms of depression and oppositional defiant disorder were assessed as covariates in the analyses to test our hypotheses.¹ The second aim was to assess if the relationship between core ADHD symptoms (i.e., inattention and hyperactivity-impulsivity) and emotionally impulsive behavior is mediated by emotion dysregulation. Emotion dysregulation scores were predicted to mediate this relationship and therefore demonstrate explanatory value beyond core DSM-IV defined ADHD symptoms.

Method

Participants

Participants included a group of adults meeting criteria for ADHD, Combined Type (ADHD, $n = 18$) and a group of psychiatrically healthy adults who did not meet criteria for ADHD or other disorders ($n = 23$). ADHD participants were recruited from an ADHD specialty clinic ($n = 11$) and from an undergraduate participant pool within a large public university ($n = 7$). All control group participants were recruited from an undergraduate participant pool at the same university. See Table 1 for a summary of the demographic and clinical characteristics of both groups. Groups did not differ in age, sex, ethnicity, or education.

Table 1

Sample demographic and clinical characteristics

	ADHD Group ($n = 18$)			Control Group ($n = 23$)			
	<i>n</i>	<i>M</i>	<i>SD</i>	<i>n</i>	<i>M</i>	<i>SD</i>	<i>p</i>
Sex							0.99
Male	7			9			
Female	11			14			
Ethnicity							0.73
Caucasian	15			16			
African American	2			4			
Hispanic	0			1			
Other	1			1			
Age		24.83	4.93		22.61	5.60	0.19

	ADHD Group (<i>n</i> = 18)			Control Group (<i>n</i> = 23)			
	<i>n</i>	<i>M</i>	<i>SD</i>	<i>n</i>	<i>M</i>	<i>SD</i>	<i>p</i>
Education level		13.88	1.75		13.17	0.83	0.10
CAARS HI Symptoms (raw scores)		16.72	5.49		5.87	3.02	<0.001
CAARS IA Symptoms (raw scores)		19.67	4.55		5.00	3.53	<0.001
CAARS Total Symptoms (raw scores)		36.39	7.91		10.87	5.51	<0.001
CAARS ADHD Index (raw scores)		23.11	6.64		8.04	5.33	<0.001
No. of Current HI Symptoms		5.67	2.03		1.22	1.24	<0.001
No. of Current IA Symptoms		6.44	1.89		0.43	0.99	<0.001
No. of Childhood HI Symptoms		6.83	2.48		2.43	2.11	<0.001
No. of Childhood IA Symptoms		7.59	1.94		2.26	2.75	<0.001
Functional Impairment from Current ADHD Symptoms		17.28	4.27		5.87	6.08	<0.001
Oppositional Defiant Disorder Symptom Severity		7.83	5.01		3.74	3.57	0.004

	ADHD Group (<i>n</i> = 18)			Control Group (<i>n</i> = 23)			
	<i>n</i>	<i>M</i>	<i>SD</i>	<i>n</i>	<i>M</i>	<i>SD</i>	<i>p</i>
(current)							
Beck Depression Inventory		14.56	9.23		8.57	7.46	0.027

Ethnicity was not reported for one control group participant, *IA* inattentive, *HI* hyperactive-impulsive, the number of current and previous ADHD symptom counts are based on ADHD-RS scores.

Diagnostic criteria for the ADHD group included clinician-rated endorsement of six or more clinician-rated inattentive symptoms and four or more hyperactive-impulsive symptoms (see Murphy and Barkley 1996, regarding the age-appropriateness of this cut-off to capture developmental deviance for adults) on a modified version of the ADHD module in the Diagnostic Interview Schedule for Children (DISC-IV) (NIMH 1997) administered by a doctoral student working under the supervision of a senior project psychologist. Aside from the modified symptom count criteria for hyperactive-impulsive symptoms, full DSM diagnostic criteria were assessed via the DISC-IV. In addition, evidence of functional impairment related to reported symptoms on a functional impairment scale (see below), a T score greater than 60 on either the DSM-IV Inattentive, DSM-IV Hyperactive-Impulsive, or ADHD Index subscales of the Conners Adult ADHD Rating Scale (CAARS) (Conners et al. 1999), and a T score greater than 55 on the DSM-IV Total ADHD Symptoms subscale was required. Current and childhood ADHD symptoms were also assessed on a modified ADHD Rating Scale (ADHD-RS; DuPaul et al. 1998). All but one of the ADHD group participants reported a previous diagnosis of ADHD by a mental health care professional.

Control group participants were excluded if they were previously diagnosed with ADHD, met symptom count criteria based on the ADHD-RS, or yielded a T score greater than 55 on the Total Symptoms CAARS scale. Participants from both groups were excluded if they had a history of psychosis, neurological disability, traumatic brain injury, mental retardation, major sensory or motor impairment, or a pervasive developmental disorder. Forty-four percent of the ADHD group (*n* = 8) reported currently taking medication for ADHD.2 None of the participants in the control group were taking medication for ADHD.

Procedure

Participants (adults age 18 and above) were recruited from an ADHD specialty clinic and from a local public university. Consent was obtained in accordance with local institutional review board guidelines. Following completion of a consent form, the measures listed above were administered in randomized order along with other measures not included in this study. Sessions ranged from 60 to 90 min.

Materials

ADHD Symptoms

The CAARS (Conners et al. 1999) is a 66-item self-report measure of current ADHD behaviors that has considerable normative data. Response options are on a four-point Likert scale. The CAARS yields the following DSM-based subscale symptom severity scores: Hyperactivity-Impulsivity symptoms, Inattentive symptoms, and Total DSM symptoms. A derived ADHD Index subscale score can also be calculated which provides an estimate of the likelihood that the respondent would meet full criteria for the disorder. Internal consistency for these subscales in the current study was very good (α 's ranged from 0.91 to 0.96 for the scales listed above). The CAARS has demonstrated adequate reliability and validity (Erhardt et al. 1999). The ADHD-RS (DuPaul et al. 1998) lists DSM-IV inattentive and hyperactive-impulsive symptoms. Participants rate the frequency of each behavior on a four-point Likert scale. A modified adult self-report version of this scale was administered similar to that used by Barkley and Murphy (2006) to establish current and retrospective symptom counts. Internal consistency for the ADHD-RS items in this sample was very good (α 's ranged from 0.90 to 0.95 for the current and retrospective scales). In general, self-reported ADHD symptoms on questionnaires are correlated with collateral reports and diagnostic interviews (Belendiuk et al. 2007; Van Voorhees et al. 2011). Finally, a modified version of the DISC-IV ADHD module (NIMH 1997) was administered to the clinical group. This ADHD module assesses DSM-IV symptoms of inattention and hyperactivity-impulsivity, as well as onset, course, and functional impairment. Modification of this interview involved revised wording of some items to match developmental appropriateness of ADHD symptoms more commonly seen in adults and noted in the DSM-IV (e.g., subjective restlessness), and is similar to other interviews that assess ADHD in adulthood (Epstein et al. 2000).

Functional Impairment

Functional impairment resulting from ADHD symptoms in ten different domains was assessed from a subset of items from the Current Symptom Scale (Barkley and Murphy 2006). Participants were asked to indicate the extent to which problems indicated on the ADHD-RS for current ADHD symptoms interfered with their ability to function in several areas of life (e.g., relational functioning) on a four-point Likert scale. Internal consistency was very good for this scale ($\alpha = .92$).

Emotion Dysregulation

Emotion dysregulation was assessed by the Impulsivity/Emotional Lability scale from the CAARS (Conners et al. 1999). This 12-item subscale, which is administered in an identical format to the other CAARS scales described above, assesses temper, irritability, stress intolerance, and labile mood. High scorers on this scale are more likely to engage in impulsive acts, moods change more quickly and more frequently, and are more easily angered and irritable. Items includes, “Many things set me off easily” and “My moods are unpredictable.” Previous studies have administered the child version (Anastopoulos et al. 2011) and adult version (Mitchell et al. 2012) of this subscale to assess self-regulation of emotion. Internal consistency for this scale was very good ($\alpha = .94$). Although psychometric studies of the CAARS supports that the impulsivity and emotional functioning items from this scale (i.e., items separate from DSM symptoms) load onto one common factor (Conners et al. 1999), a potential confound in the current study was that the impulsivity items included in this subscale would artificially inflate the relationship between (a) ADHD and emotion dysregulation, (b) and emotional impulsivity and emotion dysregulation. Therefore, we also calculated a total score from the six emotion dysregulation items (e.g., “I am easily frustrated” and “Many things set me off easily”) on this subscale by summing them consistent with other applications of this scale (Mitchell et al. 2012). Internal consistency for the six items in this revised subscale that only included emotion dysregulation items and excluded any impulsivity items was good ($\alpha = .88$). For the analyses, raw scores are reported for the full Impulsivity/Emotional Lability subscale and the modified subscale assessing only emotion dysregulation items. Descriptively, this scale measures the failure to engage in self-regulatory actions component of emotion dysregulation.

Emotional Impulsivity

The UPPS (Urgency, Premeditation, Perseverance, and Sensation Seeking) Impulsive Behavior scale (Whiteside and Lynam 2001) is a 45-item self-report measure that proposes a conceptual framework of four different types of impulsive behavior. Response options are on a four-point Likert scale. One of these subscales, Urgency, is composed of 12-items and was used for this study. Items were summed to yield a raw score. Urgency refers to a tendency to engage in rash, impulsive behavior in the context of experiencing negative emotions. Items include “Sometimes when I feel bad, I can’t seem to stop what I am doing even though it is making me feel worse” and “When I am upset I often act without thinking.” Urgency is associated with the impulsiveness facet of neuroticism within the five factor model of personality and can be accompanied by negative emotions such as anger, anxiety, or depression. Psychometric studies support the reliability and validity of this measure (Miller et al. 2003; Whiteside and Lynam 2001; Whiteside et al. 2005). Internal consistency for this scale in the current sample was good ($\alpha = .88$).

Oppositional Defiant Disorder and Depressive Symptoms

Current DSM-IV oppositional defiant disorder symptoms were assessed with the Current Symptoms Scale (Barkley and Murphy 2006); depressive symptoms were assessed with the Beck Depression Inventory (Beck and Steer 1984). Respondents rated symptom severity on a four-point Likert scale for both measures. Internal consistency for both depressive and oppositional defiant disorder scales in the current sample was good (α 's = .88 and .85, respectively).

Data Analysis

All analyses were conducted using SPSS version 19.0. The sample was dichotomized into two groups, ADHD and non-ADHD, to assess the first hypothesis that emotion dysregulation and emotional impulsivity would be higher in the ADHD group. A series of one-way analyses of covariance (ANCOVA) were conducted. Current oppositional defiant disorder and depressive symptoms were covaried in these analyses. Eta-square was used as an effect size estimate and was interpreted in accordance with standard guidelines in which small effect sizes are ≥ 0.0099 , medium effect sizes are ≥ 0.0588 , and large effect sizes are ≥ 0.1379 (Cohen 1988).

To test the second hypothesis that emotion dysregulation scores would mediate the relationship between core ADHD symptoms and emotional impulsivity, a series of hierarchical regressions was conducted following traditional mediational analysis guidelines (Baron and Kenny 1986). As a more rigorous test of the mediational model, we utilized a bootstrapping (resampling) strategy to test the significance of the indirect path from core ADHD symptoms \rightarrow emotion dysregulation \rightarrow emotional impulsivity based on recommendations by Preacher and Hayes (2008). Bootstrapping is a nonparametric statistic using repeated sampling with replacement of the available data to generate an approximation of the sampling distribution. Bootstrapping provides relatively more accurate type I error rates and greater statistical power in detecting indirect effects than traditional mediational analysis. Moreover, it is a better estimate for samples that are small in size or not normally distributed (Preacher and Hayes 2008). An SPSS macro for bootstrapping using 10,000 samples was applied to conduct this analysis (Preacher and Hayes 2008). The significance and magnitude of indirect effects estimated through the bootstrap technique was evaluated with 95 % confidence intervals (CIs). Similar to previous studies assessing the relationship between ADHD and emotional functioning (e.g., Barkley and Fischer 2010; Barkley and Murphy 2010), both groups were collapsed together for the dimensional analysis.

Results

Table 1 indicates that the ADHD and control groups did not differ on demographic or clinical characteristics (non-significant t s < 1.69 and χ^2 s < 1.32), other than ratings of ADHD symptoms

(current and past), ADHD functional impairment, and depression and oppositional defiant disorder symptoms (significant *t*s ranged from 2.30 to 13.16).

Consistent with the hypothesis that the ADHD group would score higher on a measure of emotion dysregulation than the control group, a one-way ANCOVA indicated significant between-group differences while statistically controlling for current oppositional defiant disorder and depressive symptoms (ADHD group: *M* = 19.44, adjusted *M* = 16.80, *SD* = 7.06, Control group: *M* = 7.22, adjusted *M* = 9.29, *SD* = 5.21), *F* (1, 37) = 25.14, $\eta^2 = .41$, *p* < .001. This group difference remained when the modified, six-item emotion dysregulation scale was considered (ADHD group: *M* = 8.67, adjusted *M* = 7.12, *SD* = 3.85, Control group: *M* = 3.74, adjusted *M* = 4.95, *SD* = 3.11), *F* (1, 37) = 7.37, $\eta^2 = .17$, *p* = .01. Consistent with the hypothesis that the ADHD group would report higher emotional impulsivity than the control group, a one-way ANCOVA indicated significant between-group differences while statistically controlling for current oppositional defiant disorder and depressive symptoms (ADHD group: *M* = 34.11, adjusted *M* = 31.76, *SD* = 7.03, Control group: *M* = 25.17, adjusted *M* = 27.01, *SD* = 6.31), *F* (1, 37) = 6.02, $\eta^2 = .14$, *p* = .019.

To test the hypothesis that the relationship between ADHD symptoms and emotional impulsivity was mediated by emotion dysregulation, a series of regression analyses were conducted first. Values for skewness (ranging from 0.03 to 0.71) and kurtosis (ranging from -0.22 to -1.18) indicated that the variables were normally distributed when the groups were collapsed together (Kline 2005). Table 2 summarizes the correlational relationship between these variables. Tolerance (ranging from 0.28 to 0.46) and variance inflation factor (ranging from 2.17 to 3.53) were considered and minimized concerns about multicollinearity following standard guidelines (Menard 1995).

Table 2

Correlational relationship among emotional impulsivity, emotion dysregulation, and ADHD symptoms

	Emotional Impulsivity	Emotion Dysregulation (12 item)	Emotion Dysregulation (6 item)	ADHD Symptoms
Emotional Impulsivity	–	0.75*	0.80*	0.60*
Emotion Dysregulation (12 item)		–	0.93*	0.85*
Emotion Dysregulation (6 item)			–	0.73*

	Emotional Impulsivity	Emotion Dysregulation (12 item)	Emotion Dysregulation (6 item)	ADHD Symptoms
ADHD Symptoms				–

Emotional Impulsivity = Urgency subscale; Emotion Dysregulation = CAARS Impulsivity Emotional Lability subscale (12 item version and 6 item version including only emotion dysregulation items, respectively); ADHD Symptoms = CAARS DSM-IV Inattentive and Hyperactive-Impulsive subscales.

* $p < .001$

A series of separate regression analyses indicated that both ADHD symptoms ($\beta = .60, p < .001$) and emotion dysregulation ($\beta = .75, p < .001$) predicted emotional impulsivity, and that ADHD symptoms predicted emotion dysregulation ($\beta = .85, p < .001$). These findings indicated that a test of a mediated model was appropriate. Once emotion dysregulation scores were first entered in a regression model, ADHD symptoms were no longer predictive of emotional impulsivity ($\beta = -.11, p > .05$). See Table 3 for a summary of these regressions. An analysis was then conducted using 10,000 bootstrap samples following recommendations by Preacher and Hayes (2008) to assess the indirect path from ADHD symptoms → emotion dysregulation → emotional impulsivity. Using bias corrected confidence intervals, the results were significant (indirect effect = 1.21, 95 % CI [0.89, 1.67], $z = 6.19, SE = .20, p < .001$) and indicated that emotion dysregulation symptoms mediate the relationship between ADHD symptoms and emotional impulsivity.

Table 3

Summary of hierarchical regression analyses testing mediation between symptoms of ADHD and emotional impulsivity

	Emotional Impulsivity	
	ΔR^2	β
Step1		
ADHD Symptoms	0.36	0.60*
Step 1		

	Emotional Impulsivity	
	ΔR^2	β
Emotion Dysregulation (12 item)	0.56	0.75*
Step2		
ADHD Symptoms	<0.01	-0.11

Emotional Impulsivity = Urgency subscale; Emotion Dysregulation = CAARS Impulsivity Emotional Lability subscale (12 item version); ADHD Symptoms = CAARS DSM-IV Inattentive and Hyperactive-Impulsive subscales.

* $p < .001$

The mediation analysis was repeated with the modified, six-item emotion dysregulation scale that only included emotion regulation items. Both ADHD symptoms ($\beta = .60, p < .001$) and modified emotion dysregulation ($\beta = .80, p < .001$) predicted emotional impulsivity in separate regression models. In addition, ADHD symptoms predicted the modified emotion dysregulation scale ($\beta = .73, p < .001$) in a separate regression model. Again, once modified emotion dysregulation scores were considered in the regression model, ADHD symptoms were no longer predictive of emotional impulsivity ($\beta = .04, p > .05$). The indirect path from ADHD symptoms \rightarrow emotion dysregulation (six-item scale version) \rightarrow emotional impulsivity following bootstrapping methods resulted in similar findings (indirect effect = 0.99, 95 % CI [0.43, 1.53], $z = 3.52, SE = .28, p < .001$).

Discussion

The current study assessed the relationship among ADHD, emotion dysregulation, and emotional impulsivity (i.e., negative affectively-driven impulsive behavior). Scores of emotion dysregulation and emotional impulsivity were higher in the ADHD group after covarying depressive and oppositional defiant disorder symptoms. In addition, a series of regression analyses indicated that emotion dysregulation scores mediated the relationship between core ADHD symptoms and emotional impulsivity. These findings are consistent with recent reviews and empirical findings that argue that emotion dysregulation is an important component of

ADHD, adds predictive value beyond traditional ADHD symptoms, and should be considered in diagnostic conceptualizations of ADHD in DSM-V (Barkley 2010; Martel 2009; Skirrow et al. 2009). To our knowledge, this is the first study to assess how both emotion dysregulation symptoms and emotional impulsivity, constructs emphasized in ADHD-emotion dysregulation research (Barkley 2010), are related to ADHD symptoms. Further, this pilot study adds to the small, but growing literature assessing the role of emotion dysregulation among adults diagnosed with ADHD.

These findings are consistent with neuropsychological accounts of ADHD. For instance, according to an executive functioning theory of ADHD (Barkley 1997), response inhibition allows for separable executive functions that serve to guide behavior across time delays toward future goals that serve to maximize delayed rather than immediate consequences. Emotional inhibition is among these major executive functions and is argued to be disrupted in ADHD, therefore leading to emotion dysregulation (see Barkley 2010, for a summary). As another example, the delay aversion account of ADHD (Sonuga-Barke 2002, 2003) proposes that those with ADHD have a deficit in signaling for future rewards and therefore are characterized by a preference for smaller, immediate rewards over larger, delayed rewards. Accordingly, those who are delay averse learn to associate delay cues with failure due to the problematic functional outcomes of delay aversion behavioral patterns. Escape from or avoidance of situations associated with delay regulates negative affect (Sonuga-Barke et al. 2008). Thus, this pathway predicts that negative affect should be higher in ADHD, and that maladaptive ways of regulating negative emotions are likely to emerge as the result of negative reinforcement of escape or avoidant behaviors. Although these accounts predict how emotion dysregulation is related to these different phenotypes, these relationships have yet to be examined.

Emotion dysregulation deficits have implications for ADHD diagnosis and prognosis. Regarding diagnosis, some have argued that emotion dysregulation may comprise a separate subset of adults with ADHD with possibly more severe levels of functional impairment and comorbidities (Reimherr et al. 2010). In addition to diagnostic implications involving subtyping and differing developmental trajectories, future research should consider if emotion dysregulation is a mechanism that mediates higher rates of functional impairment and comorbidity in ADHD. Although some studies have reported that poor emotional self-control predicts functional impairment in multiple domains for adults diagnosed with ADHD (Barkley and Fischer 2010; Barkley and Murphy 2010), additional domains such as substance use have yet to be considered. In particular, one particular promising avenue of future research involves cigarette smoking. Adults with ADHD smoke at significantly higher rates, initiate smoking earlier, exhibit a higher level of nicotine dependence, and have greater difficulty quitting than non-ADHD smokers (McClernon and Kollins 2008). In addition, ADHD is an independent risk factor for smoking, even after accounting for comorbidity (Milberger et al. 1997; Molina and Pelham 2003). Smoking to regulate negative emotions is one proposed mechanism maintaining smoking behavior (Gehricke et al. 2007). Consistent with such proposals, smokers with ADHD endorse

higher rates of negative affect associated with withdrawal (McClernon et al. 2010). However, to date, no studies have explicitly addressed the role of emotion dysregulation in smoking and ADHD.

Another implication of the current findings involves ADHD treatment. Given that emotion dysregulation appears to be higher in ADHD samples, future pharmacological research should consider targeting these symptoms in addition to core ADHD symptoms. Indeed, clinician-rated emotion dysregulation symptoms are responsive to methylphenidate and atomoxetine (Reimherr et al. 2010; Reimherr et al. 2005). Also, cognitive-behavioral therapy has been demonstrated to be an efficacious treatment in recent randomized controlled trials (Safren et al. 2010; Solanto et al. 2010). Given the role of emotion dysregulation in ADHD, perhaps treatments should also target emotion dysregulation symptoms as well. Supplementary cognitive-behavioral treatments emphasizing mood regulation or treatments that can be incorporated within a cognitive-behavioral approach may be beneficial. For example, a feasibility study provides promising support for a mindfulness-based treatments in ADHD (Zylowska et al. 2008). Given that mindfulness has been incorporated into cognitive-behaviorally-based treatments for other forms of psychopathology (Segal et al. 2002) and emotion regulation is a proposed mechanism of change in mindfulness-based interventions (Gratz and Tull 2010), integrating mindfulness training with existing cognitive-behavioral interventions may be a promising avenue for future research. As an example, a modified group-based dialectical behavior therapy adapted by Philipsen and colleagues for adults with ADHD includes mindfulness and cognitive-behavioral skills training while also including a module that specifically targets emotion regulation (Philipsen et al. 2007).

Limitations & Future Directions

Although the current findings are novel and promising, they should be tempered by limitations. First, the sample size for this pilot study was small, which may have reduced statistical power. However, despite this limitation, the predicted relationships emerged and the estimated effect sizes were large. Future studies should also consider sample size to allow for an analysis of sex differences when considering emotion dysregulation among adults diagnosed with ADHD. A second limitation involved sample composition. For instance, the ADHD sample in the current study included some participants enrolled in college. Since fewer adults diagnosed with ADHD attempt or complete college (Biederman et al. 2006), the current study may have included a relatively higher functioning ADHD sample. Future studies should include a larger community sample to improve generalizability.

Third, the current study relied on self-report measures of emotion dysregulation as a unidimensional construct. Although unidimensional emotion dysregulation scores are not uncommon, self-report measures of emotion dysregulation also incorporate facets of emotion

dysregulation not assessed in the current study (Gratz and Roemer 2004). Future studies should consider the hypotheses of this study with laboratory tasks of emotion dysregulation (Gratz et al. 2006) or self-report measures specifically developed to operationalize emotion dysregulation, such as the Difficulties in Emotion Regulation Scale (Gratz and Roemer 2004). Fourth, the current study relied on self-report measures. This methodology leaves open the possibility of error caused by shared method variance. Future research should consider ADHD symptom severity and impairment from other reporters, in addition to laboratory tasks assessing emotion dysregulation.

Fifth, future research should explore the construct validity of emotional impulsivity and test additional features of the model proposed by Barkley (2010) involving core inhibitory control deficits in individuals with ADHD and emotionally dysregulated behavior. The current study represents only a preliminary test of features of a model in a small, but growing literature among adults diagnosed with ADHD. Although we administered an emotion dysregulation measure that has been administered in other studies assessing ADHD and ADHD symptoms (Anastopoulos et al. 2011; Mitchell et al. 2012), future studies should include other measures of emotion dysregulation (as noted above) to not just assess the role of emotion dysregulation in ADHD, but how it is associated with emotional impulsivity since these constructs are theoretically related but separable (Barkley 2010). These construct definitional issues require attention in future research to build upon existing models involving the role of emotion dysregulation in ADHD (e.g., Barkley 2010). Sixth, the current study did not fully assess different comorbidity patterns in ADHD. Those diagnosed with ADHD are heterogeneous in phenotypic expression, including comorbidity, and is indicative of possible variations in etiology which should be considered (Nigg, Goldsmith, & Sachek, 2004). Future research should more thoroughly evaluate the role of comorbidity in ADHD. Finally, a non-ADHD clinical comparison group was not included. This should be considered in future studies since it would allow for assessing how emotion dysregulation is specific to ADHD or perhaps is a transdiagnostic construct elevated in other psychiatric populations as well.

Conclusion

In conclusion, the role of emotion dysregulation in ADHD is an emerging field of research and the current findings add to previous studies supporting a relationship between ADHD and emotion dysregulation. In addition, the current study offers novel empirical insight into the role of emotional impulsivity and how emotion dysregulation symptoms mediate this relationship. Though replication with methodological and conceptual considerations listed above are warranted, the current study provides a foundation to generate hypotheses for future studies assessing the phenomenology of emotion regulation in this clinical population.

References

- Anastopoulos, A. D., Smith, T. F., Garrett, M. E., Morrissey-Kane, E., Schatz, N. K., Sommer, J. L., et al. (2011). Self-regulation of emotion, functional impairment, and comorbidity among children with ADHD. *Journal of Attention Disorders*, 15(7), 583–592. doi:10.1177/1087054710370567.
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders, Text Revision (DSM-IV-TR)* (4th ed.). Washington: American Psychiatric Association.
- Barkley, R. A. (1997). Behavioral inhibition, sustained attention, and executive functions: Constructing a unifying theory of ADHD. *Psychological Bulletin*, 121(1), 65–94. doi:10.1037/0033-2909.121.1.65.
- Barkley, R. A. (2010). Deficient emotional self-regulation is a core component of Attention-Deficit/Hyperactivity Disorder. *Journal of ADHD & Related Disorders*, 1(2), 5–37.
- Barkley, R. A., & Fischer, M. (2010). The unique contribution of emotional impulsiveness to impairment in major life activities in hyperactive children as adults. *Journal of the American Academy of Child and Adolescent Psychiatry*, 49(5), 503–513. doi:10.1016/j.jaac.2010.01.019.
- Barkley, R. A., & Murphy, K. R. (2006). *Attention-deficit/hyperactivity disorder: A clinical workbook* (Vol. 3rd). New York: Guilford Press.
- Barkley, R. A., & Murphy, K. R. (2010). Deficient emotional self-regulation in adults with attention-deficit/hyperactivity disorder (ADHD): The relative contributions of emotional impulsiveness and ADHD symptoms to adaptive impairments in major life activities. *Journal of ADHD & Related Disorders*, 1(4), 5–28.
- Barkley, R. A., Murphy, K. R., & Fischer, M. (2008). *ADHD in adults: What the science says*. New York: Guilford Press.
- Baron, R. M., & Kenny, D. A. (1986). The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*, 51(6), 1173–1182. doi:10.1037/0022-3514.51.6.1173.
- Beck, A. T., & Steer, R. A. (1984). Internal consistencies of the original and revised Beck Depression Inventory. *Journal of Clinical Psychology*, 40(6), 1365–1367.
- Belendiuk, K. A., Clarke, T. L., Chronis, A. M., & Raggi, V. L. (2007). Assessing the concordance of measures used to diagnose adult ADHD. *Journal of Attention Disorders*, 10(3), 276–287. doi:10.1177/1087054706289941.

Biederman, J., Faraone, S. V., Spencer, T. J., Mick, E., Monuteaux, M. C., & Aleardi, M. (2006). Functional impairments in adults with self-reports of diagnosed ADHD: A controlled study of 1001 adults in the community. *The Journal of Clinical Psychiatry*, *67*(4), 524–540.

Bush, G., Luu, P., & Posner, M. I. (2000). Cognitive and emotional influences in anterior cingulate cortex. *Trends in Cognitive Sciences*, *4*(6), 215–222. doi:10.1016/S1364-6613(00)01483-2.

Bush, G., Valera, E. M., & Seidman, L. J. (2005). Functional neuroimaging of attention-deficit/hyperactivity disorder: A review and suggested future directions. *Biological Psychiatry*, *57*(11), 1273–1284. doi:10.1016/j.biopsych.2005.01.034.

Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*. New York: Academic.

Conners, C. K., Erhardt, D., & Sparrow, E. (1999). *Conners' adult ADHD rating scales (CAARS) technical manual*. North Tonawanda: Multi-Health Systems, Inc.

DuPaul, G. J., Power, T. J., Anastopoulos, A. D., & Reid, R. (1998). *ADHD rating scale-IV: Checklists, norms, and clinical interpretation*. New York: Guilford Press.

Epstein, J. N., Johnson, D., & Conners, C. K. (2000). *Conners' adult ADHD diagnostic interview for DSM-IV*. North Tonawanda: Multi-Health Systems, Inc.

Erhardt, D., Epstein, J. N., Conners, C. K., Parker, J. D. A., & Sitarenios, G. (1999). Self ratings of ADHD symptoms in adults II: Reliability, validity, and diagnostic sensitivity. *Journal of Attention Disorders*, *3*, 153–158. doi:10.1177/108705479900300304.

Gehricke, J. G., Loughlin, S. E., Whalen, C. K., Potkin, S. G., Fallon, J. H., Jamner, L. D., et al. (2007). Smoking to self-medicate attentional and emotional dysfunctions. *Nicotine & Tobacco Research*, *9*(Suppl 4), S523–S536. doi:10.1080/14622200701685039.

Gottman, J. M., & Katz, L. F. (1989). Effects of marital discord on young children's peer interaction and health. *Developmental Psychology*, *25*, 373–381.

Gratz, K. L., & Roemer, L. (2004). Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure, and initial validation of the Difficulties in Emotion Regulation Scale. *Journal of Psychopathology and Behavioral Assessment*, *26*, 41–54. doi:10.1023/B:JOBA.0000007455.08539.94.

Gratz, K. L., Rosenthal, M. Z., Tull, M. T., Lejuez, C. W., & Gunderson, J. G. (2006). An experimental investigation of emotion dysregulation in borderline personality disorder. *Journal of Abnormal Psychology*, *115*(4), 850–855. doi:10.1037/0021-843X.115.4.850.

Gratz, K. L., & Tull, M. T. (2010). Emotion regulation as a mechanism of change in acceptance- and mindfulness-based treatments. In R. A. Baer (Ed.), *Assessing mindfulness and acceptance*

processes in clients: Illuminating the theory and practice of change (pp. 107–133). Oakland: Context Press/New Harbinger Publications.

Hinshaw, S. P. (2003). Impulsivity, emotion regulation, and developmental psychopathology: Specificity versus generality of linkages. *Annals of the New York Academy of Sciences*, 1008, 149–159. doi:10.1196/annals.1301.016.

Kessler, R. C., Adler, L., Barkley, R., Biederman, J., Conners, C. K., Demler, O., et al. (2006). The prevalence and correlates of adult ADHD in the United States: Results from the national comorbidity survey replication. *The American Journal of Psychiatry*, 163(4), 716–723. doi:163/4/716.10.1176/appi.ajp. 163.4.716.

Kline, R. B. (2005). *Principles and practice of structural equation modeling* (2nd ed.). New York: Guilford Press.

Kokkonen, M., & Pulkkinen, L. (2001). Extraversion and neuroticism as antecedents of emotion regulation and dysregulation in adulthood. *European Journal of Personality*, 15(6), 407–424. doi:10.1002/per.425.

Maedgen, J. W., & Carlson, C. L. (2000). Social functioning and emotional regulation in the attention deficit hyperactivity disorder subtypes. *Journal of Clinical Child Psychology*, 29(1), 30–42.

Mannuzza, S., Klein, R. G., Bessler, A., Malloy, P., & LaPadula, M. (1993). Adult outcome of hyperactive boys. Educational achievement, occupational rank, and psychiatric status. *Archives of General Psychiatry*, 50(7), 565–576.

Martel, M. M. (2009). Research review: A new perspective on attention-deficit/hyperactivity disorder: emotion dysregulation and trait models. *Journal of Child Psychology and Psychiatry*, 50(9), 1042–1051. doi:10.1111/j.1469-7610.2009.02105.x.

Martel, M. M., Nigg, J. T., & von Eye, A. (2009). How do trait dimensions map onto ADHD symptom domains? *Journal of Abnormal Child Psychology*, 37(3), 337–348. doi:10.1007/s10802-008-9255-3.

McClernon, F. J., English, J., Hallyburton, M., Van Voorhees, E. E., Holdaway, A., & Kollins, S. H. (2010). Smoking withdrawal symptoms and craving severity are worse among smokers with ADHD. Paper presented at the Poster presented at the annual meeting of the Society for Research on Nicotine and Tobacco, Baltimore, MD.

McClernon, F. J., & Kollins, S. H. (2008). ADHD and smoking: From genes to brain to behavior. *Annals of the New York Academy of Sciences*, 1141, 131–147. doi:NYAS1141016.10.1196/annals.1441.016.

- Melnick, S. M., & Hinshaw, S. P. (2000). Emotion regulation and parenting in AD/HD and comparison boys: Linkages with social behaviors and peer preference. *Journal of Abnormal Child Psychology*, 28(1), 73–86. doi:10.1023/A:1005174102794.
- Menard, S. (1995). *Applied logistic regression analysis: Sage university series on quantitative applications in the social sciences*. Thousand Oaks: Sage.
- Milberger, S., Biederman, J., Faraone, S. V., Chen, L., & Jones, J. (1997). ADHD is associated with early initiation of cigarette smoking in children and adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(1), 37–44. doi:10.1097/00004583-199701000-00015.
- Miller, J., Flory, K., Lynam, D., & Leukefeld, C. (2003). A test of the four-factor model of impulsivity-related traits. *Personality and Individual Differences*, 34(8), 1403–1418. doi:10.1016/S0191-8869(02)00122-8.
- Miller, M., Hanford, R. B., Fassbender, C., Duke, M., & Schweitzer, J. B. (2011). Affect recognition in adults with ADHD. *Journal of Attention Disorders*, 15(6), 452–460. doi:10.1177/1087054710368636.
- Mitchell, J. T., & Nelson-Gray, R. O. (2006). Attention-deficit/hyperactivity disorder symptoms in adults: Relationship to Gray's behavioral approach system. *Personality and Individual Differences*, 40, 749–760. doi:10.1016/j.paid.2005.08.011.
- Mitchell, J. T., Van Voorhees, E. E., Dennis, M. F., McClernon, F. J., Calhoun, P. S., Kollins, S. H., et al (2012). Assessing the role of attention-deficit/hyperactivity disorder symptoms in smokers with and without posttraumatic stress disorder. *Nicotine & Tobacco Research*. doi:10.1093/ntr/ntr245
- Molina, B. S., & Pelham, W. E., Jr. (2003). Childhood predictors of adolescent substance use in a longitudinal study of children with ADHD. *Journal of Abnormal Psychology*, 112(3), 497–507. doi:10.1037/0021-843X.112.3.497.
- Murphy, K. R., & Barkley, R. A. (1996). Prevalence of DSM-IV symptoms of ADHD in adult licensed drivers: Implications for clinical diagnosis. *Journal of Attention Disorders*, 1, 147–161.
- Nigg, J. T., Goldsmith, H. H., & Sachek, J. (2004). Temperament and attention deficit hyperactivity disorder: The development of a multiple pathway model. *Journal of Clinical Child and Adolescent Psychology*, 33(1), 42–53. doi:10.1207/S15374424JCCP3301_5.
- Nigg, J. T., John, O. P., Blaskey, L. G., Huang-Pollock, C. L., Willcutt, E. G., Hinshaw, S. P., et al. (2002). Big five dimensions and ADHD symptoms: Links between personality traits and clinical symptoms. *Journal of Personality and Social Psychology*, 83(2), 451–469. doi:10.1037/0022-3514.83.2.451.

- NIMH. (1997). *Diagnostic interview schedule for children – IV (DISC-IV)*. New York: NY Columbia University.
- Ochsner, K. N., & Gross, J. J. (2005). The cognitive control of emotion. *Trends in Cognitive Sciences*, 9(5), 242–249. doi:10.1016/j.tics.2005.03.010.
- Paloyelis, Y., Mehta, M. A., Kuntsi, J., & Asherson, P. (2007). Functional MRI in ADHD: A systematic literature review. *Expert Review of Neurotherapeutics*, 7(10), 1337–1356. doi:10.1586/14737175.7.10.1337.
- Philipsen, A., Richter, H., Peters, J., Alm, B., Sobanski, E., Colla, M., et al. (2007). Structured group psychotherapy in adults with attention deficit hyperactivity disorder: results of an open multicentre study. *The Journal of Nervous and Mental Disease*, 195(12), 1013–1019. doi:10.1097/NMD.0b013e31815c088b.
- Preacher, K. J., & Hayes, A. F. (2008). Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods*, 40(3), 879–891. doi:10.3758/BRM.40.3.879.
- Putnam, K. M., & Silk, K. R. (2005). Emotion dysregulation and the development of borderline personality disorder. *Development and Psychopathology*, 17(4), 899–925. doi:10.1017/S0954579405050431.
- Reimherr, F. W., Marchant, B. K., Olsen, J. L., Halls, C., Kondo, D. G., Williams, E. D., et al. (2010). Emotional dysregulation as a core feature of adult ADHD: Its relationship with clinical variables and treatment response in two methylphenidate trials. *Journal of ADHD & Related Disorders*, 1(4), 53–64.
- Reimherr, F. W., Marchant, B. K., Strong, R. E., Hedges, D. W., Adler, L., Spencer, T. J., et al. (2005). Emotional dysregulation in adult ADHD and response to atomoxetine. *Biological Psychiatry*, 58, 125–131. doi:10.1016/j.biopsych.2005.04.040.
- Safren, S. A., Sprich, S., Mimiaga, M. J., Surman, C., Knouse, L., Groves, M., et al. (2010). Cognitive behavioral therapy vs relaxation with educational support for medication-treated adults with ADHD and persistent symptoms: A randomized controlled trial. *Journal of the American Medical Association*, 304(8), 875–880. doi:10.1001/jama.2010.1192.
- Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002). *Mindfulness-based cognitive therapy for depression: A new approach to preventing relapse*. New York: Guilford Press.
- Skirrow, C., McLoughlin, G., Kuntsi, J., & Asherson, P. (2009). Behavioral, neurocognitive and treatment overlap between attention-deficit/hyperactivity disorder and mood instability. *Expert Review of Neurotherapeutics*, 9(4), 489–503. doi:10.1586/ern.09.2.

- Solanto, M. V., Marks, D. J., Wasserstein, J., Mitchell, K., Abikoff, H., Alvir, J. M., et al. (2010). Efficacy of meta-cognitive therapy for adult ADHD. *The American Journal of Psychiatry*, 167(8), 958–968. doi:10.1176/appi.ajp.2009.09081123.
- Sonuga-Barke, E. J. (2002). Psychological heterogeneity in AD/HD—A dual pathway model of behaviour and cognition. *Behavioural Brain Research*, 130(1–2), 29–36. doi:10.1016/S0166-4328(01)00432-6.
- Sonuga-Barke, E. J. (2003). The dual pathway model of AD/HD: An elaboration of neuro-developmental characteristics. *Neuroscience and Biobehavioral Reviews*, 27(7), 593–604. doi:10.1016/j.neubiorev.2003.08.005.
- Sonuga-Barke, E. J., Sergeant, J. A., Nigg, J., & Willcutt, E. (2008). Executive dysfunction and delay aversion in attention deficit hyperactivity disorder: Nosologic and diagnostic implications. *Child and Adolescent Psychiatric Clinics of North America*, 17(2), 367–384. doi:10.1016/j.chc.2007.11.008. ix.
- Valera, E. M., Faraone, S. V., Murray, K. E., & Seidman, L. J. (2007). Meta-analysis of structural imaging findings in attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 61(12), 1361–1369. doi:10.1016/j.biopsych.2006.06.011.
- Van Voorhees, E. E., Hardy, K. K., & Kollins, S. H. (2011). Reliability and validity of self- and other-ratings of symptoms of ADHD in adults. *Journal of Attention Disorders*, 15(3), 224–234. doi:10.1177/1087054709356163.
- Walcott, C. M., & Landau, S. (2004). The relation between disinhibition and emotion regulation in boys with attention deficit hyperactivity disorder. *Journal of Clinical Child and Adolescent Psychology*, 33(4), 772–782. doi:10.1207/s15374424jccp3304_12.
- Weiss, G., & Hechtman, L. T. (1993). *Hyperactive children grown up* (2nd ed.). New York: Guilford Press.
- Weiss, M. D., Wasdell, M., Gadow, K. D., Greenfield, B., Hechtman, L., & Gibbins, C. (2011). Clinical correlates of oppositional defiant disorder and attention-deficit/hyperactivity disorder in adults. *Postgraduate Medicine*, 123(2), 177–184. doi:10.3810/pgm.2011.03.2276.
- Whiteside, S. P., & Lynam, D. R. (2001). The five factor model and impulsivity: Using a structural model of personality to understand impulsivity. *Personality and Individual Differences*, 30(4), 669–689. doi:10.1016/S0191-8869(00)00064-7.
- Whiteside, S. P., Lynam, D. R., Miller, J. D., & Reynolds, S. K. (2005). Validation of the UPPS impulsive behaviour scale: A four-factor model of impulsivity. *European Journal of Personality*, 19(7), 559–574. doi:10.1002/per.556.

Zylowska, L., Ackerman, D. L., Yang, M. H., Futrell, J. L., Horton, N. L., Hale, T. S., et al. (2008). Mindfulness meditation training in adults and adolescents with ADHD: A feasibility study. *Journal of Attention Disorders*, 11(6), 737–746. doi:10.1177/1087054707308502.

Footnotes

1 We included a measure of oppositional defiant disorder in our adult sample given that emerging research suggests that this disorder can continue into adulthood for some and is functionally impairing (M. D. Weiss et al. 2011).

2 Of those taking medication for ADHD, 5 were from the student group.