

Cytotoxic Homoisoflavonoids from the Bulbs of *Bellevalia flexuosa*

By: [Tamam El-Elimat](#), José Rivera-Chávez, Joanna E. Burdette, Austin Czarnecki, Maram B. Alhawarri, Mohammad Al-Gharaibeh, Feras Alali, and [Nicholas H. Oberlies](#)

“Cytotoxic Homoisoflavonoids from the Bulbs of *Bellevalia flexuosa*.” Tamam El-Elimat, José Rivera-Chávez, Joanna E. Burdette, Austin Czarnecki, Maram B. Alhawarri, Mohammad Al-Gharaibeh, Feras Alali, and Nicholas H. Oberlies. *Fitoterapia*, 2018, 127, 201-206. PMID: 29471027; PMCID: PMC5984119; doi: 10.1016/j.fitote.2018.02.022

Made available courtesy of Elsevier: <https://doi.org/10.1016/j.fitote.2018.02.022>



This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](#).

***© Elsevier B.V. Reprinted with permission. This version of the document is not the version of record. ***

Abstract:

Four new homoisoflavonoids, 7-*O*-methyl-8-demethoxy-3'-hydroxy-3,9-dihydropunctatin (**4**), 6-hydroxy-8-demethoxy-4'-*O*-methyl-3,9-dihydropunctatin (**8**), 7,4'-*O*-dimethyl-8-demethoxy-3,3'-dihydroxy-3,9-dihydropunctatin (**13**), and 7-*O*-methyl-3-hydroxy-3,9-dihydropunctatin (**14**) were identified from a chloroform extract of the bulbs of *Bellevalia flexuosa*, along with 13 known analogues. The structures were determined by analysis of HRMS and NMR data, while ECD spectroscopy enabled the assignment of the absolute configurations of the new compounds **4**, **8**, **13** and **16**. The cytotoxic activities of the isolated compounds (**1–17**) were evaluated using a panel of human cancer cell lines. Compounds **2** and **7** were the most potent against the MDA-MB-435 (melanoma) cancer cell line with IC₅₀ values of 1.6 and 2.0 μM, respectively, and were essentially equipotent against the OVCAR3 (ovarian) cancer cell line with IC₅₀ values of 9.5 and 10.8 μM, respectively. However, compound **7**, with an IC₅₀ value of 3.6 μM, was the most potent against the MDA-MB-231 (breast) cancer cell line.

Keywords: Homoisoflavonoids | *Bellevalia* | Bulbs | Absolute configuration | Cytotoxicity | Human cancer cell lines

Article:

1. Introduction

Homoisoflavonoids are a rare subclass of flavonoids possessing an extra carbon atom in their skeleton [1, 2, 3, 4, 5]. Biosynthetically, chalcones are thought to be the precursors of homoisoflavonoids [3]. Currently, about 250 natural homoisoflavonoids have been reported in the literature, with the majority being isolated from several genera of Asparagaceae and Fabaceae [1, 2, 3, 4, 6, 7]. Based on their structures, homoisoflavonoids have been classified into five groups: 3-benzyl-4-chromanones, 3-benzyl-3-hydroxy-4-chromanones, 3-benzylidene-4-

chromanones (*E* or *Z*), 3-benzylchrom-2-en-4-ones, and the scillascillins [4]. Homoisoflavonoids have attracted attention because of their various biological activities, including antioxidant, anti-inflammatory, antimutagenic, antimicrobial, antiallergic and antihistaminic, anti-diabetic, cytotoxic, and anti-angiogenic effects, as well as protein tyrosine kinase inhibition activity [3,4].

Although Jordan is a small country of about 96,188 km², it has a remarkable diversity of wildlife, which could be attributed to its unique location at the intersection of three continents, encapsulating four bio-geographical zones: Mediterranean, Irano-Turanian, Saharo-Arabian, and Tropical (Sudanian penetration) [8, 9, 10]. Within these different zones, thirteen vegetation types are identified, reflecting Jordan's diverse landscape, climate, and geology [8,11]. >2600 vascular plant species belonging to 113 family and 810 genera are reported to grow in the wild, of which 100 are endemic and >70 species are considered extinct [8]. Hence, studies of Jordan's wild plants have been initiated [12, 13, 14] as a potential source of drug leads.

In the current study, the bulbs of *Bellevalia flexuosa* Boiss. (Asparagaceae) have been explored. *B. flexuosa*, which is known as “Common Roman Squill”, and in Jordan as “Drooping Onion” [15], is one of eleven species that are reported by Al-Eisawi to grow in Jordan in the wild [16]. Although no medical use is reported for this plant species by the local people of Jordan, a closely related species (*B. eigii*) was investigated recently, yielding a set of homoisoflavonoids [6]. *B. flexuosa* is a perennial herb with underground bulbs that is found flowering from February to March. It flourishes in mountains and waste grounds [16]. Chemical investigation of the chloroform extract of the bulbs of *B. flexuosa* resulted in the isolation and identification of seventeen homoisoflavonoids; of which thirteen were known (1–3, 5–7, 9–12, and 15–17) and four were new analogues: 7-*O*-methyl-8-demethoxy-3'-hydroxy-3,9-dihydropunctatin (4), 6-hydroxy-8-demethoxy-4'-*O*-methyl-3,9-dihydropunctatin (8), 7,4'-*O*-dimethyl-8-demethoxy-3,3'-dihydroxy-3,9-dihydropunctatin (13), and 7-*O*-methyl-3-hydroxy-3,9-dihydropunctatin (14). ECD spectroscopy was used to assign the absolute configurations of the new compounds 4, 8, 13, and 14. Homoisoflavonoids were reported to have a broad range of biological activities, including cytotoxic effects [3]. Therefore, the isolated compounds (1–17) were tested for their cytotoxicity using three human cancer cell lines, namely MDA-MB-435 (melanoma), MDA-MB-231 (breast), and OVCAR3 (ovarian).

2. Experimental

2.1. General experimental procedures

Optical rotations, UV data, and ECD spectra were obtained using a Rudolph Research Autopol III polarimeter (Rudolph Research Analytical), a Varian Cary 100 Bio UV-vis spectrophotometer (Varian Inc.), and an Olis DSM 17 ECD spectrophotometer (Olis, Inc.). NMR data were collected using either a JEOL ECA-500 NMR spectrometer operating at 500 MHz for ¹H and 125 MHz for ¹³C (JEOL Ltd.) or an Agilent 700 MHz NMR spectrometer (Agilent Technologies), equipped with a cryoprobe, operating at 700 MHz for ¹H and 175 MHz for ¹³C. Residual solvent signals were utilized for referencing. HRMS data were acquired using a Thermo QExactive Plus mass spectrometer equipped with an electrospray ionization source (Thermo Fisher Scientific). Gemini-NX C₁₈ analytical (5 μm; 250 × 4.6 mm) and preparative (5 μm; 250 × 21.2 mm) columns (both from Phenomenex) along with Atlantis T3 C₁₈ analytical (5 μm;

250 × 4.6 mm) and semipreparative (5 μm; 250 × 10.0 mm) columns (both from Waters Corp.) were used on a Varian Prostar HPLC system equipped with ProStar 210 pumps and a Prostar 335 photodiode array detector (PDA), with data collected and analyzed using Galaxie Chromatography Workstation software (version 1.9.3.2, Varian Inc.). Flash chromatography was performed on a Teledyne ISCO CombiFlash Rf 200 using Silica Gold columns (both from Teledyne Isco) and monitored by UV and evaporative light-scattering detectors. All other reagents and solvents were obtained from Fisher Scientific and were used without further purification.

2.2. Plant material

Bulbs of *B. flexuosa* were collected by Mohammad Bashabshah during flowering stage from the campus of the Jordan University of Science and Technology (JUST), Irbid, Jordan in March/April 2016. The plant material was identified by Mohammad Al-Gharaibeh, Plant Taxonomist, Faculty of Agriculture, JUST. A voucher specimen (PHS-122) was deposited in the herbarium of the Faculty of Pharmacy, JUST. The bulbs were cleaned of mud, sliced into small pieces, and air dried at rt. in a well-ventilated area.

2.3. Extraction and isolation

Air-dried bulbs of *B. flexuosa* were ground to a powder using a laboratory mill. The powdered bulbs (850 g) were extracted exhaustively with CHCl₃ by soaking at rt. The solvent was evaporated under reduced pressure to yield the CHCl₃ extract (3.1 g), which was reconstituted in a 500 mL mixture of 5:4:1 H₂O:CHCl₃:MeOH. The mixture was stirred for 30 min and left to separate in a separatory funnel. The organic layer was collected and evaporated to dryness under reduced pressure. The dried organic extract was reconstituted in 200 mL of 1:1 MeOH:CH₃CN and 200 mL of hexanes and transferred into a separatory funnel. The biphasic solution was shaken vigorously. The MeOH/MeCN layer was evaporated to dryness under vacuum. The dried MeOH/CH₃CN layer (2.1 g) was dissolved in CHCl₃ and mixed with Celite 545. Normal-phase flash chromatography was run using a gradient solvent system of hexanes-CHCl₃-MeOH, at a flow rate of 35 mL/min, and column volumes of 41.3 over a total run time of 30.7 min, to yield seven fractions. Fraction 3 (267.9 mg) was subjected to preparative HPLC over a Gemini column using a gradient system of 50:50 to 60:40 of MeOH-H₂O (0.1% formic acid) over 30 min at a flow rate of 20 mL/min to yield 8 subfractions, of which subfraction 3 was identified as compound **15** (2.1 mg). Subfraction 7 (1.9 mg) was subjected to semipreparative HPLC purification over Atlantis T3 column using a gradient solvent system of 40:60 to 50:50 CH₃CN-H₂O (0.1% formic acid) over 15 min at a flow rate of 4.6 mL/min to yield compounds **5** (0.7 mg) and **13** (0.4 mg). Subfraction 8 (8.3 mg) was subjected to semipreparative HPLC using Atlantis T3 column and a gradient solvent system of 40:60 to 60:40 CH₃CN-H₂O (0.1% formic acid) over 20 min at a flow rate of 4.6 mL/min to yield compounds **7** (5.4 mg) and **13** (0.3 mg).

Fraction 4 (544.7 mg) was subjected to preparative HPLC over a Gemini column using a gradient system of 60:40 to 70:30 of MeOH-H₂O (0.1% formic acid) over 20 min at a flow rate of 21.2 mL/min to yield compound **5** (335.6 mg).

Fraction 5 (116.0 mg) was subjected to preparative HPLC over a Gemini column using a gradient system of 50:50 to 70:30 of MeOH-H₂O (0.1% formic acid) over 20 min at a flow rate of 21.2 mL/min to yield seven sub-fractions. Further HPLC purification of subfractions 3 (3.4 mg) and 4 (3.2 mg) using semipreparative HPLC (Atlantis T3 column) and a gradient solvent systems of 40:60 to 50:50 CH₃CN-H₂O (0.1% formic acid) over 15 min at a flow rate of 4.6 mL/min yielded compounds **17** (1.3 mg) and **2** (2.1 mg) from subfractions 3 and 4, respectively. Subfractions 6 (2.2 mg) and 7 (14.3 mg) were purified using semipreparative HPLC (Atlantis T3 column) and a gradient solvent system of 40:60 to 60:40 CH₃CN-H₂O (0.1% formic acid) over 20 min at a flow rate of 4.6 mL/min to yield compounds **4** (0.9 mg) and **16** (10.8 mg) from subfractions 6 and 7, respectively.

Fraction 6 (379.3 mg) was purified using preparative HPLC (Gemini column) and a gradient solvent system of 50:50 to 55:45 of MeOH-H₂O (0.1% formic acid) over 20 min, hold for 5 min and then increasing to 60:40 over 10 min at a flow rate of 21.2 mL/min to yield 5 subfractions. Subfraction 1 (13.2 mg) was further purified using semipreparative (Atlantis T3 column) and a gradient solvent system of 30:70 to 40:60 CH₃CN-H₂O (0.1% formic acid) over 15 min at a flow rate of 4.6 mL/min to yield compound **8** (2.3 mg) and another subfraction that was further purified using semipreparative HPLC method over Atlantis T3 column and an isocratic solvent system of 28:72 CH₃CN-H₂O (0.1% formic acid) for 50 min at a flow rate of 4.6 mL/min to yield compounds **9** (2.3 mg) and **14** (1.9 mg). Subfractions 2 (15.6 mg) and 3 (39.0 mg) were purified using semipreparative HPLC (Atlantis T3 column) and a gradient solvent systems of 40:60 to 60:50 CH₃CN-H₂O (0.1% formic acid) over 15 min at a flow rate of 4.6 mL/min to yield compound **6** (9.1 mg) from subfraction 2 and compounds **3** (1.4 mg), **6** (1.9 mg), and **10** (19.2 mg) from subfraction 3. Subfractions 4 (3.4 mg) and 5 (3.8 mg) were purified using semipreparative HPLC (Atlantis T3 column) and gradient solvent systems of 40:60 to 50:50 CH₃CN-H₂O (0.1% formic acid) over 15 min at a flow rate of 4.6 mL/min to yield compounds **1** (0.5 mg) and **10** (0.5 mg) from fraction 4 and compounds **2** (0.9 mg), **5** (2.3 mg), **11** (0.9 mg), and **12** (1.5 mg) from subfraction 5.

2.3.1. 7-O-Methyl-8-demethoxy-3'-hydroxy-3,9-dihydropunctatin (4)

Yellowish oil; $[\alpha]_{\text{D}}^{28} - 27$ (*c* 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 334 (3.23), 288 (3.98), 220 (3.94) nm; ECD (*c* 0.9×10^{-4} M, MeOH) λ ($\Delta\epsilon$) 244 (+0.68) nm, 269 (+0.96) nm, 295 (-2.07), 320 (+0.18) nm (Fig. 3); HRESIMS *m/z* 317.1016 [M + H]⁺ (calcd for C₁₇H₁₇O₆, 317.1020).

2.3.2. 6-Hydroxy-8-demethoxy-4'-O-methyl-3,9-dihydropunctatin (8)

Light yellow amorphous powder; $[\alpha]_{\text{D}}^{28} - 36$ (*c* 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ) 360 (3.32), 291 (4.05), 242 (3.94), 229 (3.99) nm; ECD (*c* 0.9×10^{-4} M, MeOH) λ ($\Delta\epsilon$) 260 (+2.02) nm, 293 (-4.79), 323 (+0.50) nm (Fig. 3); HRESIMS *m/z* 317.1016 [M + H]⁺ (calcd for C₁₇H₁₇O₆, 317.1020).

2.3.3. 7,4'-O-dimethyl-8-demethoxy-3,3'-dihydroxy-3,9-dihydropunctatin (13)

White amorphous powder; $[\alpha]_D^{28} - 96$ (c 0.05, MeOH); UV (MeOH) λ_{\max} ($\log \epsilon$) 334 (3.35), 288 (4.08), 218 (4.10) nm; ECD (c 0.9×10^{-4} M, MeOH) λ ($\Delta\epsilon$) 253 (+1.80) nm, 295 (-3.52), 320 (-0.53) nm (Fig. 3); HRESIMS m/z 347.1122 $[M + H]^+$ (calcd for $C_{18}H_{19}O_7$, 347.1125).

2.3.4. 7-*O*-methyl-3-hydroxy-3,9-dihydropunctatin (14)

Light yellow amorphous powder; $[\alpha]_D^{28} - 170$ (c 0.1, MeOH); UV (MeOH) λ_{\max} ($\log \epsilon$) 348 (3.58), 292 (4.15), 224 (4.15) nm; ECD (c 0.9×10^{-4} M, MeOH) λ ($\Delta\epsilon$) 265 (+4.16) nm, 292 (-4.53), 313 (+0.42) nm (Fig. 3); HRESIMS m/z 347.1124 $[M + H]^+$ (calcd for $C_{18}H_{19}O_7$, 347.1125).

2.4. Cytotoxicity assay

Compounds (1–17) were tested for cytotoxicity against human melanoma cancer cells MDA-MB-435 [17], human breast cancer cells MDA-MB-231, and human ovarian cancer cells OVCAR3 as described previously [18,19]. Briefly, the cell lines were propagated at 37 °C in 5% CO₂ in RPMI 1640 medium, supplemented with fetal bovine serum (10%), penicillin (100 units/mL), and streptomycin (100 µg/mL). Cells in log phase growth were harvested by trypsinization followed by two washings to remove all traces of enzyme. A total of 5000 cells were seeded per well of a 96-well clear, flat-bottom plate (Microtest 96®, Falcon) and incubated overnight (37 °C in 5% CO₂). Samples dissolved in DMSO were then diluted and added to the appropriate wells. The cells were incubated in the presence of test substance for 72 h at 37 °C and evaluated for viability with a commercial absorbance assay (CellTiter 96® AQueous One Solution Cell Proliferation Assay, Promega Corp, Madison, WI) that measured viable cells. IC₅₀ values are expressed in µM relative to the solvent (DMSO) control. Taxol was used as a positive control.




Fig. 1. Structures of compounds 1–17.

3. Results and discussion

Air-dried bulbs of *B. flexuosa* were extracted with CHCl₃, and the resulting crude extract was reconstituted in a mixture of 5:4:1 H₂O:CHCl₃:MeOH. The dried organic extract was reconstituted in 1:1 MeOH/CH₃CN and then defatted using hexanes. The resulting dried MeOH/CH₃CN extract was fractionated using normal phase flash chromatography. HPLC methods, preparative and semipreparative, were used for purifications of the fractions leading to the isolation of seventeen homoisoflavonoids (**1–17**) (Fig. 1).

The structures of thirteen known homoisoflavonoids analogues (**1–3**, **5–7**, **9–12**, and **15–17**) were established by comparison of NMR (1D/2D), HRMS, and ECD data with literature values and were identified as: 4',5,7-trihydroxyhomoisoflavanone (**1**) [6], 3'-hydroxy-3,9-dihydroeucomin (**2**) [6], 3,9-dihdropunctatin (**3**) [20], 7-*O*-methyl-3,9-dihdropunctatin (**5**) [6], 8-*O*-demethyl-7-*O*-methyl-3,9-dihdropunctatin (**6**) [6], 7,4'-di-*O*-methyl-3'-hydroxy-3,9-dihdropunctatin (**7**) [6], 6-hydroxy-7-*O*-methyl-3,9-dihdropunctatin (**9**) [6], 7-*O*-methyl-3'-hydroxypunctatin (**10**), 3-(4-hydroxybenzyl)-5-hydroxy-6,7,8-trimethoxychroman-4-one (**11**) [21], 7-*O*-methyl-8-demethoxy-3-hydroxy-3,9-dihdropunctatin (**12**) [6], 5-hydroxy-7,8-dimethoxychroman-4-one (**15**) [6], 7-*O*-methylpunctatin (**16**) [6], and isomuscomosin (**17**) [6], (Figs. **S1–S3**, **S5–S7**, **S9–S12**, **S15–S17**, Supplementary Data).

Compound **4** (0.93 mg) was isolated as a yellowish oil with a molecular formula of C₁₇H₁₆O₆ (10 degrees of unsaturation) as determined by HRESIMS (m/z 317.1016 [M + H]⁺, calcd. 317.1020), which was further supported by ¹H, ¹³C, and edited HSQC NMR data (Table 1, Table 2, Fig. S4, Supplementary Data). Compound **4** showed characteristic NMR signals indicative of a 3-benzylchroman-4-one homoisoflavonoid analogue with very high structural similarity to **2**, both sharing the same molecular formula. HMBC data analysis indicated the structure of **4** to be different from that of **2** by the methylation position. HMBC correlations from H_{3–7'} (δ_{H} 3.88, s) to C-4' (δ_{C} 145.5) in **2** and from H_{3–10} (δ_{H} 3.81, s) to C-7 (δ_{C} 168.0) in **4** enabled the assignment of **2** as a 4'-*O*-methyl analogue and **4** as a 7-*O*-methyl analogue. In the latter, HMBC correlations from H-8 (δ_{H} 5.97, d, $J = 2.3$) to C-4a (δ_{C} 102.7), C-6 (δ_{C} 95.1), C-7 (δ_{C} 168.0), and C-8a (δ_{C} 163.0); from H-6 (δ_{H} 6.06, d, $J = 2.3$) to C-5 (δ_{C} 164.6), C-7, C-8 (δ_{C} 94.0), and C-4a, from 5-OH (δ_{H} 12.11, s) to C-4a, C-5, and C-6 established the substitution pattern of ring A. In addition, HMBC correlations from H-2' (δ_{H} 6.75, d, $J = 1.3$) to C-9 (δ_{C} 32.3), C-4' (δ_{C} 142.4), and C-6' (δ_{C} 121.8); from H-5' (δ_{H} 6.81, d, $J = 7.9$) to C-1' (δ_{C} 131.0) and C-3' (δ_{C} 143.8); from H-6' (δ_{H} 6.66, dd, $J = 7.9, 1.3$) to C-9, C-2' (δ_{C} 116.3), and C-4' established the substitution pattern of ring C. Further examination of the 2D NMR data established the structure of **4** (Fig. 2) as 7-*O*-methyl-8-demethoxy-3'-hydroxy-3,9-dihdropunctatin. The absolute configurations of 3-benzylchroman-4-one-type homoisoflavonoids are assigned using electronic circular dichroism (ECD) spectroscopy [22], in which a negative Cotton effect in the 287–295 nm region of the ECD curves is indicative of a 3*R* configuration [22]. Compound **4** showed a negative Cotton effect at 295 nm ($\Delta\epsilon = -2.07$) in the ECD spectrum, supporting an *R*-configuration at C-3 (Fig. 3).

Table 1. ¹H NMR data for compounds **4**, **8**, **13**, and **14** (500 MHz in CDCl₃)^a.

position	4	8	13	14
2	4.12, dd (11.4, 7.1)	4.08, dd (11.4, 7.1)	4.06, d (11.2)	4.09, d (11.2)
	4.28, dd (11.4, 4.1)	4.25, dd (11.4, 4.1)	4.23, d (11.2)	4.36, d (11.2)
3	2.80, m	2.78, m		
6	6.06, d (2.3)		6.11, d (2.3)	6.16, s
8	5.97, d (2.3)	6.05, s	6.05, d (2.3)	
9	2.65, dd (13.9, 10.4)	2.69, dd (13.8, 10.6)	2.89, d (13.8)	2.94, d (14.1)
	3.12, dd (13.9, 4.5)	3.16, dd (13.8, 4.5)	2.93, d (13.8)	2.98, d (14.1)
10	3.81, s		3.85, s	3.91, s
11			3.88, s	3.83, s
12		3.91, s		
2'	6.75, d (1.3)	7.08, d (8.5)	6.82, d (2.1)	7.04, d (8.5)
3'		6.78, d (8.5)		6.75, d (8.5)
5'	6.81, d (7.9)		6.76, d (8.1)	
6'	6.66, dd (7.9, 1.3)		6.65, dd (8.1, 2.1)	
3-OH			3.36, s	
5-OH	12.11, s	11.78, s	11.25, s	11.21, s
6-OH		5.03, br. s		
7-OH		4.81, br. s		
3'-OH	5.20 or 5.33 ^b , br. s		5.58, s	
4'-OH	5.20 or 5.33 ^b , br. s			4.83, br. s

^aδ in ppm, mult (*J* in Hz); ^bCould be swapped.

Table 2. ¹³C NMR data for **4**, **8**, **14** (125 MHz), and for **13** (175 MHz) in CDCl₃.

position	4	8	13	14
2	69.1	69.4	71.9	72.5
3	46.9	47.2	72.3	72.2
4	198.0	198.8	198.3	198.6
4a	102.7	102.5	100.6	100.4
5	164.6	148.2	164.1	160.0
6	95.1	156.1	95.5	93.7
7	168.0	127.4	168.8	162.3
8	94.0	91.1	94.6	129.9
8a	163.0	154.8	162.9	153.5
9	32.3	32.1	41.0	40.9
10	55.8		56.0	56.6
11			56.0	61.7
12		56.5		
1'	131.0	130.1	127.3	126.2
2'	116.3	130.5	116.8	131.9
3'	143.8	115.7	145.4	115.4
4'	142.4	154.5	145.9	155.0
5'	115.7		110.5	
6'	121.8		122.2	




Fig. 2. Key COSY and HMBC correlations of **4**, **8**, **13**, and **14**.




Fig. 3. ECD spectra for compounds A) **4**, B) **8**, C) **13**, and D) **14** [0.09 mM, MeOH, cell length 2 cm].

HRESIMS and NMR data of compound **8** (2.3 mg), which was obtained as a yellow amorphous powder, revealed its molecular formula as $C_{17}H_{16}O_6$ (m/z 317.1016 $[M+H]^+$, calcd 317.1020) (Table 1, Table 2, Fig. S8, Supplementary Data). As in compound **4**, the NMR data of **8** showed

characteristic signals indicative of a 3-benzylchroman-4-one-type homoisoflavonoid with structural similarity to **3** as they shared the same molecular formula. However, the 8-methoxy group of ring A ($\delta_{\text{H}}/\delta_{\text{C}}$ 3.95/61.7 for H₃-11/C-11) in **3** was replaced by an aromatic proton singlet ($\delta_{\text{H}}/\delta_{\text{C}}$ 6.05/91.1 for H-8/C-8) in **8**. In addition, the aromatic proton singlet ($\delta_{\text{H}}/\delta_{\text{C}}$ 6.13/96.0 for H-6/C-6) in **3** was replaced by an exchangeable proton singlet (δ_{H} 5.03 for 6-OH) in **8**. Moreover, the exchangeable proton singlet in **3** (δ_{H} 4.75 for 4'-OH) was replaced by a methyl group ($\delta_{\text{H}}/\delta_{\text{C}}$ 3.91/56.5 for H₃-7'/C-7'). The structure of **8** was inferred by further analysis of HMBC data. HMBC correlation from H-8 to C-4 (198.8), C-4a (102.5), C-6 (156.1), and C-8a (154.8) established the substitution pattern of ring A, while the HMBC correlation from H₃-7' (δ_{H} 3.91, s) to C-4' (δ_{C} 154.5) confirmed the methylation position in **8**. Further analysis of the 2D NMR data established the structure of **8** to which the trivial name 6-hydroxy-8-demethoxy-4'-*O*-methyl-3,9-dihydropunctatin was assigned (Fig. 2). A negative Cotton effect at 293 nm in the ECD spectrum of compound **8** ($\Delta\varepsilon = -4.79$) indicated an *R*-configuration at C-3 (Fig. 3).

Compound **13** (0.66 mg) was obtained as a white amorphous powder. HRESIMS (m/z 347.1122 [M + H]⁺, calcd 347.1125) NMR data indicated a molecular formula of C₁₈H₁₈O₇ (Table 1, Table 2, Fig. S13, Supplementary Data). The NMR data indicated compound **13** as a 3-benzyl-3-hydroxychroman-4-one homoisoflavonoid analogue. The compound showed high structural similarity to **12**. However, compound **13** showed a methoxy group, as indicated by ¹H and ¹³C NMR data ($\delta_{\text{H}}/\delta_{\text{C}}$ 3.88/56.0) and a hydroxyl group at δ_{H} 5.58 (3'-OH) consistent with the 30 amu difference in the HRMS data of **13** relative to **12**. The aromatic A₂B₂ system of ring B in **12** was replaced by an ABM spin system in **13** (δ_{H} 6.82, d, $J = 2.1$; 6.76, d, $J = 8.1$; and 6.65, dd, $J = 8.1, 2.1$, for H-2', H-5', and H-6', respectively), indicating the presence of a 1,3,4-trisubstituted benzene ring. An HMBC correlation from the 4'-OCH₃ protons to C-4' (δ_{C} 145.9) confirmed its connectivity. The structure of compound **13** was deduced by further inspection of the 2D NMR data, including COSY and HMBC spectra (Fig. 2). Compound **13** was given the trivial name 7,4'-*O*-dimethyl-8-demethoxy-3,3'-dihydroxy-3,9-dihydropunctatin. A negative Cotton effect ($\Delta\varepsilon = -3.52$) at 297 nm of the ECD spectrum of **13** indicated an *S*-configuration at C-3 (Fig. 3) [22].

The HRESIMS (m/z 347.1124 [M + H]⁺, calcd 347.1125) and NMR data of compound **14** (2.38 mg), which was obtained as a light yellow amorphous powder, indicated a molecular formula of C₁₈H₁₈O₇ (Table 1, Table 2, Fig. S14, Supplementary Data). The NMR data of **14** showed distinctive peaks indicative of a 3-benzyl-3-hydroxychroman-4-one homoisoflavonoid analogue. The compound showed high structural similarity to **12**. However, compound **14** had an extra methoxy group, as indicated by ¹H and ¹³C NMR data ($\delta_{\text{H}}/\delta_{\text{C}}$ 3.83/61.7), consistent with the 30 amu difference in the HRMS data of **14** relative to **12**, replacing the aromatic doublet (δ_{H} 6.05, d, $J = 2.1$ for H-8) in **12**. HMBC correlations from H-6 to C-4 (198.6), C-4a (100.4), C-5 (160.0), C-7 (162.3), and C-8 (129.9) confirmed the substitution pattern of ring A. The structure of compound **14** was deduced by further inspection of the 2D NMR data, including COSY and HMBC spectra (Fig. 2). Compound **14** was given the trivial name 7-*O*-methyl-3-hydroxy-3,9-dihydropunctatin. A negative cotton effect ($\Delta\varepsilon = -4.53$) at 292 nm in the ECD spectrum of **14** was indicative of an *S*-configuration at C-3 (Fig. 3) [22].

Compounds (**1–17**) were evaluated for their cytotoxic activities against the MDA-MB-435 (melanoma), MDA-MB-231 (breast), and OVCAR3 (ovarian) cancer cell lines. Compounds **2** and **7** were the most potent on the three cancer cell lines with IC₅₀ values of 1.6, 14.2, 9.5 and 2.0, 3.6, and 10.8 μM, respectively (Table 3). Compound **2** was reported to be active against colon cancer (HT-29 cell line ED₅₀ = 2.78 μM) and breast cancer (MDA-MB-435 cell line ED₅₀ = 1.33 μM). Compounds **2** and **7** were reported previously by our group to be active against breast cancer (MDA-MB-435 cell line) with IC₅₀ values of 1.0 and 1.1 μM, respectively [6]. Moreover, compound **7** showed moderate activity when tested against colon cancer (HT-29 cell line IC₅₀ = 17.3 μM) [6]. The cytotoxicity data of the new and known analogues isolated in the current work expanded our understanding of the structure-activity relationships of this unique class of flavonoids. The substitution pattern of ring B affects the activity significantly. Compound **1**, with a 4'-OH group, was inactive. However, compound **4**, with 3',4'-dihydroxy substituents, showed IC₅₀ values of 14.3, 17.5, and 24.8 μM against the MDA-MB-435, MDA-MB-231, and OVCAR3 cancer cell lines, respectively. Moreover, reducing the polarity of the ring B substituents results in improved cytotoxic activity. For example, the activity of compound **2**, with a 4'-methoxy substituent, increased by ~9- and 3-folds the activity against MDA-MB-435 and OVCAR3 cancer cell lines in comparison with compound **4** via IC₅₀ values of 1.6 and 9.5 μM, respectively. Similarly, although compound **10** was inactive, compound **7**, with a 4'-methoxy substituent, showed IC₅₀ values of 2.0, 3.6, and 10.8 μM against the MDA-MB-435, MDA-MB-231, and OVCAR3 cancer cell lines, respectively. The same pattern can be noticed when comparing the cytotoxic activities of compounds **12** and **13**. Compound **12** with a 4'-OH substituent was inactive, however compound **13**, with 3'-hydroxy and 4'-methoxy substituents, showed activity against the MDA-MB-435 cancer cell line with IC₅₀ value of 14.4 μM. In addition, introducing an OH group at C-3 increases the activity significantly, although compound **5** was inactive, compound **14**, with a 3-OH substituent, showed IC₅₀ value of 15.2 μM against the MDA-MB-435 cancer cell line.

Table 3. Cytotoxic activities of compounds **2**, **4**, **7**, **13**, and **14**.

compound ^a	IC ₅₀ values in μM ^b		
	MDA-MB-435	OvcAR3	MDA-MB-231
2	1.6	14.2	9.5
4	14.3	17.5	24.8
7	2.0	3.6	10.8
13	14.4	>25	>25
14	15.2	>25	>25
Taxol^c	0.0001	0.0015	0.17

^aCompounds **1**, **3**, **5**, **6**, **8–12**, **15–17** were inactive, IC₅₀ values >25 μM. ^bIC₅₀ is the concentration to inhibit 50% of growth with a 72 h incubation. ^cPositive control.

Conflict of interest

All the authors have no conflict of interest.

Acknowledgments

This research was supported, in part, by the Deanship of Research, Jordan University of Science and Technology, Irbid, Jordan (Grant No. 284/2017) and via program project grant P01

CA125066 from the National Cancer Institute/National Institutes of Health, Bethesda, MD, USA.

References

- [1] B.M. Abegaz, J. Mutanyatta-Comar, M. Nindi, Naturally occurring homoisoflavonoids: Phytochemistry, biological activities and synthesis, *Nat. Prod. Commun.* 2 (2007) 475-498.
- [2] W. Heller, C. Tamm, Homoisoflavanones and Biogenetically Related Compounds, in: P.A. Cadby, R.G. Cooke, J.M. Edwards, W. Heller, C.W. Jefford, E. Lederer, P. Lefrancier, S. Dev, C. Tamm, W. Herz, H. Grisebach, G.W. Kirby (Eds.), *Fortschritte Der Chemie Organischer Naturstoffe / Progress in the Chemistry of Organic Natural Products*, Springer Vienna, Vienna, 1981, pp. 105-152.
- [3] L.-G. Lin, Q.-Y. Liu, Y. Ye, Naturally occurring homoisoflavonoids and their pharmacological activities, *Planta Med.* 80 (2014) 1053-1066.
- [4] D.A. Mulholland, S.L. Schwikkard, N.R. Crouch, The chemistry and biological activity of the Hyacinthaceae, *Nat. Prod. Rep.* 30 (2013) 1165-1210.
- [5] M.I. Calvo, Three new homoisoflavanones from the bulbs of *Ledebouria floribunda*, *Fitoterapia* 80 (2009) 394-398.
- [6] F. Alali, T. El-Elimat, H. Albatineh, Q. Al-Balas, M. Al-Gharaibeh, J.O. Falkinham 3rd, W.L. Chen, S.M. Swanson, N.H. Oberlies, Cytotoxic homoisoflavones from the bulbs of *Bellevalia eigii*, *J. Nat. Prod.* 78 (2015) 1708-1715.
- [7] X. Zhou, Z. Yuping, H. Zhao, J. Liang, Y. Zhang, S. Shi, Antioxidant homoisoflavonoids from *Polygonatum odoratum*, *Food Chem.* 186 (2015) 63-68.
- [8] Jordanian Fifth National Report under the Convention on Biological Diversity, Ministry of Environment, Amman, Jordan, 2014.
- [9] N. Feinbrun-Dothan, *Flora Palestina*, The Israel Academy of Sciences and Humanities, Jerusalem, 1986.
- [10] D.M. Al-Eisawi, *Field Guide to Wild Flowers of Jordan and Neighbouring Countries*, Jordan, Jordan Press, Foundation Al-Rai, Amman, 1998, pp. 1-2.
- [11] D. Al-Eisawi, *Vegetation of Jordan*, UNESCO, Cairo office 1996.
- [12] F.Q. Alali, C.S.M. Amrine, T. El-Elimat, A. Alkofahi, K. Tawaha, M. Gharaibah, S.M. Swanson, J.O. Falkinham III, M. Cabeza, A. Sánchez, M. Figueroa, N.H. Oberlies, Bioactive withanolides from *Withania obtusifolia*, *Phytochem. Lett.* 9 (2014) 96-101.

- [13] F.Q. Alali, T. El-Elimat, C. Li, A. Qandil, A. Alkofahi, K. Tawaha, J.P. Burgess, Y. Nakanishi, D.J. Kroll, H.A. Navarro, J.O. Falkinham, M.C. Wani, N.H. Oberlies, New colchicinoids from a native Jordanian meadow saffron, *Colchicum brachyphyllum*: isolation of the first naturally occurring dextrorotatory colchicinoid, *J. Nat. Prod.* 68 (2005) 173-178.
- [14] F.Q. Alali, A. Gharaibeh, A. Ghawanmeh, K. Tawaha, N.H. Oberlies, Colchicinoids from *Colchicum crocifolium* Boiss.: a case study in dereplication strategies for (-)-colchicine and related analogues using LC-MS and LC-PDA techniques, *Phytochem. Anal.* 19 (2008) 385-394.
- [15] H. Taifour, A. El-Oqlah, Jordan Plant Red List, Royal Botanic Garden, Amman, Jordan, 2014.
- [16] D.M. Al-Eisawi, Field Guide to Wild Flowers of Jordan and Neighbouring Countries, Jordan Press Foundation Al Rai, Amman, 1998.
- [17] J. Rae, C. Creighton, J. Meck, B. Haddad, M. Johnson, MDA-MB-435 cells are derived from M14 melanoma cells-a loss for breast cancer, but a boon for melanoma research, *Breast Cancer Res. Treat.* 104 (2007) 13-19.
- [18] T. El-Elimat, M. Figueroa, H.A. Raja, S.M. Swanson, J.O. Falkinham, D.M. Lucas, M.R. Grever, M.C. Wani, C.J. Pearce, N.H. Oberlies, Sorbicillinoid analogues with cytotoxic and selective anti-*Aspergillus* activities from *Scytalidium album*, *J. Antibiot.* 68 (2015) 191-196.
- [19] T. El-Elimat, H.A. Raja, J.O. Falkinham, C.S. Day, N.H. Oberlies, Greensporones: Resorecylic acid lactones from an aquatic *Halenospora* sp, *J. Nat. Prod.* 77 (2014) 2088-2098.
- [20] M. Adinolfi, G. Barone, M. Belardini, R. Lanzetta, G. Laonigro, M. Parrilli, Homoisoflavanones from *Muscari comosum* bulbs, *Phytochemistry* 24 (1985) 2423-2426.
- [21] G. Amschler, A.W. Frahm, A. Hatzelmann, U. Kilian, D. Müller-Doblies, U. Müller-Doblies, Constituents of *Veltheimia viridifolia*; I. Homoisoflavanones of the bulbs, *Planta Med.* 62 (1996) 534-539.
- [22] M. Adinolfi, G. Barone, M.M. Corsaro, L. Mangoni, R. Lanzetta, M. Parrilli, Absolute configuration of homoisoflavanones from *Muscari* species, *Tetrahedron* 44 (1988) 4981-4988.