

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Faculty Publications: Department of  
Entomology

Entomology, Department of

---

2020

## Diabrotica undecimpunctata virus 2, a Novel Small RNA Virus Discovered from Southern Corn Rootworm, Diabrotica undecimpunctata howardi Barber (Coleoptera: Chrysomelidae)

Sijun Liu

Arnubio Valencia-Jiménez

Molly Darlington

Ana M. Vélez

Bryony C. Bonning

Follow this and additional works at: <https://digitalcommons.unl.edu/entomologyfacpub>



Part of the [Entomology Commons](#)

---

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



# *Diabrotica undecimpunctata virus 2*, a Novel Small RNA Virus Discovered from Southern Corn Rootworm, *Diabrotica undecimpunctata howardi* Barber (Coleoptera: Chrysomelidae)

Sijun Liu,<sup>a</sup> Arnubio Valencia-Jiménez,<sup>b,c</sup> Molly Darlington,<sup>c</sup> Ana M. Vélez,<sup>c</sup>  Bryony C. Bonning<sup>d</sup>

<sup>a</sup>Department of Entomology, Iowa State University, Ames, Iowa, USA

<sup>b</sup>Departamento de Producción Agropecuaria, Universidad de Caldas, Manizales, Colombia

<sup>c</sup>Department of Entomology, University of Nebraska—Lincoln, Lincoln, Nebraska, USA

<sup>d</sup>Department of Entomology and Nematology, University of Florida, Gainesville, Florida, USA

**ABSTRACT** The genome of *Diabrotica undecimpunctata virus 2* (DuV2), a putative positive-sense, single-stranded RNA virus identified from the southern corn rootworm transcriptome, comprises 5,313 nucleotides, including a short poly(A) tail. The two open reading frames encode a nonstructural polyprotein (p156) and a putative capsid protein (p25).

The southern corn rootworm (SCR), *Diabrotica undecimpunctata howardi* Barber, is one of three corn rootworm species that cause significant damage to agricultural and horticultural crops (1). Several small RNA viruses have been identified from the western corn rootworm, *Diabrotica virgifera virgifera* (2–4), but only one has been described from SCR to date (5). Here, we report on the genome of a second RNA virus identified from the transcriptome of SCR.

Eggs, larvae (first, second, and third instar), pupae, and adult (male and female) SCR were supplied by Crop Characteristics (Farmington, MN). The RNeasy minikit (Qiagen, Valencia, CA, USA) was used for isolation of total RNA from three biological replicates of 100 eggs, 50 first-instar larvae, 5 each of second- and third-instar larvae, 1 pupa, 1 adult female, and 1 adult male. cDNA libraries were constructed using the NEBNext Ultra RNA library prep kit for Illumina (NEB) for each stage-specific sample with poly(A) selection and paired-end sequencing with 150-nucleotide (nt) reads conducted on a HiSeq 2500 platform (Illumina, San Diego, CA). One or two replicate sequence data sets from each developmental stage were used for virus sequence discovery (5).

The Trinity assembler v2.6.6 (6) was used for assembly of high-quality reads. Default parameters were used for all software unless otherwise noted. A putative viral contig, with a short poly(A) tail, has 5,312 nt with a nucleotide composition of 37% G+C content. A total of 29,036 and 29,694 reads from two first-instar replicate data sets (comprising 50.7 million and 48.1 million reads and quality scores of 37.5 and 37.6, respectively) mapped to the viral contig with coverages of 819.8- and 838.3-fold, respectively. The virus is provisionally named *Diabrotica undecimpunctata virus 2* (DuV2). Sequences derived from DuV2 were observed in transcriptome data sets from all developmental stages of SCR. The near-full-length sequences were assembled from first-instar larvae, suggesting increased abundance and active replication of DuV2 in larvae relative to other developmental stages. The 5' coding sequence located between nt 156 and 4286 encodes a putative nonstructural protein of 1,376 amino acids (aa) (156.1 kDa, p156), while the more 3' sequence between nt 4366 and 5025 encodes a putative structural protein with 219 aa (25 kDa, p25). A conserved protein domain search revealed that p156 contains three conserved domains, two overlapping viral helicase domains (NCBI accession no. [pfam01443](https://.ncbi.nlm.nih.gov/nucl/pfam01443), aa 598 to 826; [cd17914](https://ncbi.nlm.nih.gov/nucl/cd17914), aa 593 to

**Citation** Liu S, Valencia-Jiménez A, Darlington M, Vélez AM, Bonning BC. 2020. *Diabrotica undecimpunctata virus 2*, a novel small RNA virus discovered from southern corn rootworm, *Diabrotica undecimpunctata howardi* Barber (Coleoptera: Chrysomelidae). *Microbiol Resour Announc* 9:e00380-20. <https://doi.org/10.1128/MRA.00380-20>.

**Editor** Kenneth M. Stedman, Portland State University


**Copyright** © 2020 Liu et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Bryony C. Bonning, [bbonning@ufl.edu](mailto:bbonning@ufl.edu).

**Received** 10 April 2020

**Accepted** 28 May 2020

**Published** 25 June 2020



**FIG 1** Phylogenetic analysis of DuV2 putative nonstructural protein p156 and putative structural protein p25. The replication-associated p156 is homologous to proteins from viruses with various hosts, including plants, fungi, insects, and hepatitis E-like viruses. Sequences with similarity to the capsid protein p25 were identified from viruses isolated from arthropods, although p25 also has a plant viral coat protein TMV domain. Viral proteins used for construction of the phylogenetic tree were selected from BLASTp analyses. Sequences were aligned with MAFFT v7.4.07 (using the AUTO strategy), and the maximum likelihood (ML) method was used for construction of the phylogenetic tree. ModelFinder implemented in IQ-TREE was used to identify the best partitioning scheme and models. ML analysis was performed using IQ-TREE with 10,000 ultrafast bootstrap replicates (10). Interactive Tree of Life (<http://itol.embl.de>) was used for visualization and annotation. The viruses and associated NCBI genome accession numbers are *Adelphocoris suturalis* virus, YP\_009336480.1; *Agaricus bisporus* virus 8, AQM49930.1; *Agaricus bisporus* virus 13, AQM49941.1; *Bactrocera oleae*, XP\_014085601.1; *Bat bastrovirus*, AWW67082.1; *Beet necrotic yellow vein virus*, ACA63016.1; *Beet soil-borne mosaic virus*, YP\_009513207.1; *Beihai charybdis crab virus 1*, YP\_009333243.1; *Burdock mottle virus*, YP\_008219063.1; *Chara australis* virus, AEJ33768.1 and AEJ33771.1; *Culex pipiens associated Tunisia virus*, YP\_009553258.1; *Goji berry chlorosis virus*, AYO99569.1; *Hepatitis E virus*, BAO47889.1; *Hubei beny-like virus 1*, APG77690.1; *Hubei virga-like virus 2*, YP\_009337413.1; *Hubei virga-like virus 11*, AVK59472.1 and YP\_009337245.1; *Lentinula edodes ssRNA mycovirus*, BAS04359.1; *Mangifera indica latent virus*, AMQ23297.1; *Megastigmus ssRNA virus*, QDZ71187.1; *Red clover RNA virus 1*, AVD98637.1; *Riboviria* sp. QDH86483.1; *Rice stripe necrosis virus*, QAR21011.1; *Rodent hepevirus*, AWW79978.1; *Sclerotium rolfsii beny-like virus 1*, AZF86092.1; *Wenling thamnaconus septentrionalis hepevirus*, AVM87557.1; *Wheat stripe mosaic virus*, AYD38100.1; and *Zeugodacus cucurbitae*, XP\_011177481.1.

726) (7), and one RNA-dependent RNA polymerase domain (RdRP\_2; [pfam00978](#)). A *Tomato mosaic virus* coat protein domain (TMV\_coat domain [pfam00721](#)) was identified in protein p25, supporting annotation as a capsid protein.

The highest BLASTp score for p156 (8) was *Hubei beny-like virus 1* (GenBank accession no. [KX883789.1](#)), an RNA virus isolated from dipteran insects (9) with 76% sequence coverage and 49% amino acid identity. Proteins similar to p156 included those from viruses of various host species (including fungi and plants) and soil samples (*Riboviria* sp.) (Fig. 1). BLASTp analysis for p25 identified the putative capsid protein of *Beihai charybdis crab virus 1* (GenBank accession no. [YP\\_009333243.1](#)), which has a genomic structure similar to that of DuV2.

**Data availability.** The genome sequence of DuV2 is available in GenBank under accession no. [MN646771](#). The raw reads for SRA accession no. [SRX8018653](#) to [SRX8018656](#) are available at BioProject no. [PRJNA615920](#).

## ACKNOWLEDGMENTS

This research was supported by the USDA National Institute of Food and Agriculture, Hatch project IOW05450; State of Iowa funds; University of Nebraska—Lincoln funds; the 2016 DuPont Young Professor Grant awarded to A. M. Vélez; and the University of Florida Institute of Food and Agricultural Sciences.

## REFERENCES

1. Krysan JL. 1986. Introduction: biology, distribution, and identification of pest *Diabrotica*, p 1–23. In Krysan JLM, Thomas A (ed), *Methods for the study of pest Diabrotica*. Springer, New York, NY.
2. Liu S, Chen Y, Sappington TW, Bonning BC. 2017. Genome sequence of a novel positive-sense, single-stranded RNA virus isolated from western corn rootworm, *Diabrotica virgifera virgifera* LeConte. *Genome Announc* 5:e00366-17. <https://doi.org/10.1128/genomeA.00366-17>.
3. Liu S, Chen Y, Sappington TW, Bonning BC. 2017. Genome sequence of *Diabrotica virgifera virgifera* virus 2, a novel small RNA virus of the western corn rootworm, *Diabrotica virgifera virgifera* LeConte. *Genome Announc* 5:e00365-17. <https://doi.org/10.1128/genomeA.00365-17>.
4. Liu S, Chen Y, Sappington TW, Bonning BC. 2017. Genome sequence of the first coleopteran iflavivirus isolated from western corn rootworm, *Diabrotica virgifera virgifera* LeConte. *Genome Announc* 5:e01530-16. <https://doi.org/10.1128/genomeA.01530-16>.
5. Liu S, Valencia-Jiménez A, Darlington M, Vélez AM, Bonning BC. 2020. Genome sequence of a small RNA virus of the southern corn rootworm, *Diabrotica undecimpunctata howardi* Barber (Coleoptera: Chrysomelidae). *Microbiol Resour Announc* 9:e00379-20. <https://doi.org/10.1128/MRA.00379-20>.
6. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber M, MacManes MD, Ott M, Orvis J, Pochet N, Strozzi F, Weeks N, Westerman R, William T, Dewey CN, Henschel R, Leduc RD, Friedman N, Regev A. 2013. *De novo* transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat Protoc* 8:1494–1512. <https://doi.org/10.1038/nprot.2013.084>.
7. Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, Gwadz M, Hurwitz DI, Lu F, Marchler GH, Song JS, Thanki N, Wang Z, Yamashita RA, Zhang D, Zheng C, Geer LY, Bryant SH. 2017. CDD/SPARCLE: functional classification of proteins via subfamily domain architectures. *Nucleic Acids Res* 45:D200–D203. <https://doi.org/10.1093/nar/gkw1129>.
8. Altschul SF, Wootton JC, Gertz EM, Agarwala R, Morgulis A, Schaffer AA, Yu YK. 2005. Protein database searches using compositionally adjusted substitution matrices. *FEBS J* 272:5101–5109. <https://doi.org/10.1111/j.1742-4658.2005.04945.x>.
9. Shi M, Lin XD, Tian JH, Chen LJ, Chen X, Li CX, Qin XC, Li J, Cao JP, Eden JS, Buchmann J, Wang W, Xu J, Holmes EC, Zhang YZ. 2016. Redefining the invertebrate RNA virosphere. *Nature* 540:539–543. <https://doi.org/10.1038/nature20167>.
10. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: improving the ultrafast bootstrap approximation. *Mol Biol Evol* 35:518–522. <https://doi.org/10.1093/molbev/msx281>.