

Association Between Bilirubin, Atazanavir, and Cardiovascular Disease Events Among People Living With HIV Across the United States

Heidi M. Crane, MD, MPH,^a Robin M. Nance, MPH,^a Susan R. Heckbert, MD, PhD,^a Corey Ritchings, PhD,^b Lisa Rosenblatt, PhD,^b Matthew Budoff, MD,^c Brian R. Wood, MD,^a David L. Tirschwell, MD,^a H. Nina Kim, MD,^a William C. Mathews, MD,^d Elvin Geng, MD,^e Richard D. Moore, MD,^f Peter W. Hunt, MD,^e Joseph J. Eron, MD,^g Greer A. Burkholder, MD,^h Daniel R. Drozd, MD,^a Felicia C. Chow, MD,^e Kyra J. Becker, MD,^a Joseph R. Zunt, MD,^a Emily L. Ho, MD,^a Rizwan Kalani, MD,^a Andrew Huffer, MD,^a Bridget M. Whitney, MS,^a Michael S. Saag, MD,^h Mari M. Kitahata, MD,^a and Joseph A. C. Delaney, PhDⁱ

Objective: Bilirubin is an antioxidant that may suppress lipid oxidation. Elevated bilirubin is associated with decreased cardiovascular events in HIV-uninfected populations. We examined these associations in people living with HIV (PLWH).

Methods: Potential myocardial infarctions (MIs) and strokes were centrally adjudicated. We examined MI types: type 1 MI (T1MI) from atherosclerotic plaque instability and type 2 MI (T2MI) in the setting of oxygen demand/supply mismatch such as sepsis. We used multivariable Cox regression analyses to determine associations between total bilirubin levels and outcomes adjusting for traditional and HIV-specific risk factors. To minimize confounding by hep-

atobiliary disease, we conducted analyses limited to bilirubin values <2.1 mg/dL; among those with fibrosis-4 values <3.25; and among everyone. We repeated analyses stratified by hepatitis C status and time-updated atazanavir use.

Results: Among 25,816 PLWH, there were 392 T1MI and 356 T2MI during follow-up. Adjusted hazard ratios for the association of higher bilirubin levels with T1MI were not significant. Higher bilirubin levels were associated with T2MI. By contrast, among PLWH on atazanavir, higher bilirubin levels were associated with fewer T2MI (hazard ratio 0.56:0.33–1.00). Higher bilirubin levels among those on atazanavir were associated with fewer T1MI combined with ischemic stroke.

Limitations: Analyses were conducted with total rather than unconjugated bilirubin.

Conclusions: Among PLWH, higher bilirubin levels were associated with T2MI among some subgroups. However, among those on atazanavir, there was a protective association between bilirubin and T2MI. These findings demonstrate different associations between outcomes and elevated bilirubin due to diverse causes and the importance of distinguishing MI types.

Key Words: HIV, myocardial infarction, atazanavir, bilirubin, stroke, cardiovascular disease

(*J Acquir Immune Defic Syndr* 2019;81:e141–e147)

Received for publication April 15, 2018; accepted March 11, 2019.

From the ^aDepartment of Medicine, University of Washington, Seattle, WA;

^bBristol Myers Squibb, Princeton, NJ; ^cDepartment of Medicine, University of California Los Angeles, Los Angeles, CA; ^dDepartment of Medicine, University of California San Diego, San Diego, CA; ^eDepartment of Medicine, University of California San Francisco, San Francisco, CA; ^fDepartment of Medicine, Johns Hopkins University, Baltimore, MD; ^gDepartment of Medicine, University of North Carolina, Chapel Hill, NC; ^hDepartment of Medicine, University of Alabama, Birmingham, AL; and ⁱDepartment of Epidemiology, University of Washington, Seattle, WA.

Supported by Bristol-Myers Squibb. Additional support came from the National Institute of Allergy and Infectious Diseases (CNICS R24 AI067039, UW CFAR NIAID Grant P30 AI027757; UNC CFAR Grant P30 AI50410, JHU CFAR Grant P30 AI094189, and UAB CFAR Grant P30 AI027767), the National Institute of Drug Abuse (U01DA036935) and the National Heart, Lung, and Blood Institute (R01HL126538) at the National Institutes of Health, and from the American Heart Association (13GRNT14560022). The following have received grant support or served as consultants or advisors: P.W.H. with Merck and Gilead, G.A.B. with DefiniCare, M.S.S. with BMS, Gilead, Merck, and ViiV, R.D.M. with Medscape, J.J.E. with Merck, ViiV, BMS, Abbvie, Gilead, and Tibotec/Janssen, D.R.D. with Gilead, H.M.C. with ViiV, and M.B. has consulted for General Electric. The following are affiliated with BMS: C.R. and L.R.

Presented in part at the 18th International Workshop on Comorbidities and Adverse Drug Reactions in HIV; September 2016; New York, NY.

The authors have no conflicts of interest to disclose.

Correspondence to: Heidi M. Crane, MD, MPH, University of Washington, Seattle, WA 98104 (e-mail: hcrane@uw.edu).

INTRODUCTION

Bilirubin is the end product of heme catabolism and an effective antioxidant that may suppress lipid oxidation.^{1–3} Bilirubin may also have anti-inflammatory, complement inhibitory, and possibly antithrombotic and lipid-lowering properties.^{4–6} Elevated bilirubin, particularly mildly elevated levels,⁴ has been associated with fewer myocardial infarctions (MIs) and strokes and with reduced carotid atherosclerosis, in the general population and individuals with Gilbert syndrome,^{2,3,7–11} although not always.^{12,13} The nature and

magnitude of the relationship between bilirubin levels and atherosclerotic events is undefined.^{9,10}

Relationships between bilirubin and atherosclerotic events are relevant to people living with HIV (PLWH), who have higher cardiovascular disease rates than the general population^{14–17} and may have abnormal bilirubin levels for several reasons. For example, the protease inhibitor atazanavir can cause hyperbilirubinemia due to inhibitor effects on UGT1A1 and has been a common component of HIV treatment regimens.^{18–20} A relationship has been reported between mild hyperbilirubinemia and slower progression of atherosclerosis in PLWH.²¹ It is unknown whether these associations translate into reduced risk of cardiovascular events such as MI with a recent study suggesting an association of higher bilirubin levels with lower cardiovascular disease risk among PLWH.²² However, this association was not statistically significant for MI.

These relationships may differ by the type of cardiovascular events. The Universal Definition of MI classifies MI into types according to the underlying mechanism of myocardial ischemia,^{23,24} with type 1 MI (T1MI) and type 2 MI (T2MI) being most common among PLWH. T1MI results spontaneously from atherosclerotic plaque instability. T2MI is related to oxygen demand–supply mismatch such as occurs with hypotension. Compared with the general population, where T2MIs comprise <10% of MIs,^{25–30} we found that half of MIs among PLWH were T2MI³¹. T2MI among PLWH is due to causes such as sepsis and cocaine-induced vasospasm.³¹ Similarly, there are multiple types of strokes including ischemic and hemorrhagic with multiple subtypes of ischemic stroke,³² which are also associated with different predisposing causes and risk factors.

We conducted this study to examine relationships between bilirubin and cardiovascular events among PLWH, whether these associations differ by MI type or among atazanavir users, and to evaluate the hypothesis that higher bilirubin levels may protect against T1MI.

METHODS

Study Setting

A longitudinal observational study conducted in the Centers for AIDS Research Network of Integrated Clinical Systems (CNICS), a multisite clinical cohort of PLWH receiving HIV care.³³

Study Participants

PLWH aged 18 years or older receiving HIV care at 6 CNICS sites after adjudication of cardiovascular events began (dates varied by site, ~2000–2017). We started follow-up at 6 months after enrollment and excluded individuals with events before the start of follow-up to ensure incident events.

Data Sources

The CNICS data repository integrates comprehensive clinical data from outpatient and inpatient encounters includ-

ing information on demographic characteristics, clinical and laboratory data, and medications. Laboratory data such as bilirubin values were collected as part of clinical care.

CNICS has an established state-of-the-art approach to cardiovascular outcome adjudication with MIs categorized by type.^{31,34} Potential MIs were identified in the centralized CNICS data repository using a comprehensive set of diagnostic and procedure codes and elevated cardiac biomarker values. For all potential events, sites assembled deidentified packets that included provider notes, electrocardiograms, and imaging results. Two physicians with expertise in cardiac disorders and adjudicating events reviewed each packet, followed by a third reviewer if discrepancies occurred. Reviewers categorized MIs by type, identified causes for T2MI, and also identified PLWH without an MI but with coronary interventions such as a coronary artery bypass graft surgery, which were grouped with the T1MI. MI types beyond T1MI and T2MI are not discussed further because of low numbers (eg, there were <10 type 4 and 5 MIs associated with cardiac procedures in CNICS to date). We used a similar ascertainment and adjudication approach for strokes using provider notes, procedure reports, and imaging results in deidentified packets for centralized review. Reviewers categorized each stroke as ischemic vs. hemorrhagic. Ischemic strokes were further classified by whether concurrent infection or illicit drug use/intoxication was a predisposing factor when the stroke occurred.

Outcomes

We examined MIs by type. We also conducted sensitivity analyses with a combined atherosclerotic outcome (T1MI plus ischemic strokes excluding strokes related to infection or illicit drug use).

Analyses

Multivariable Cox regression analyses were used to determine associations between time-updated total bilirubin and outcomes, adjusting for traditional and HIV-specific risk factors: time-varying viral load (VL) and CD4 count, and baseline age, sex, race/ethnicity, HIV transmission risk factor, smoking status, hepatitis C virus (HCV) infection, diabetes, treated hypertension, statin use, kidney function (estimated glomerular filtration rate <30, 30–60, ≥60), calendar time, and site. HCV status was based on positive laboratory values (antibody, genotype, or VL). Diabetes was based on any of the following: (1) hemoglobin A1c ≥ 6.5 OR (2) use of a diabetes-specific medication such as insulin OR (3) use of a diabetes-related medication not exclusively used to treat diabetes (eg, biguanides) in the setting of also having a diabetes diagnosis.³⁵

Bilirubin values ≤30 days before outcomes were excluded to ensure outcomes or concurrent predisposing factors for T2MI such as sepsis were not impacting bilirubin values. To minimize confounding by hepatobiliary disease, we conducted analyses limited to bilirubin values <2.1 mg/dL as has been performed in the general population¹¹ and repeated analyses limited to individuals who had fibrosis-4

(FIB-4) values <3.25 at baseline. The FIB-4 index, derived from age, liver aminotransferases, and platelet count, has been shown to predict liver-related events and death.^{36–38} We also conducted analyses where we examined everyone, regardless of the bilirubin or FIB-4 level. We examined associations in subgroups defined by HCV status. We examined the associations with total bilirubin as a continuous variable (\log_2 transformed), as a categorical variable (<0.6, 0.6–1.3, and ≥ 1.3 mg/dL, with cut points selected based on the normal adult range of up to 1.3 mg/dL), and using spline models to examine the association of a 0.1 mg/dL change in bilirubin within bilirubin categories.

We were particularly interested in any potential associations with atazanavir both to add clarity as to mechanisms and because of the potential clinical implications. We conducted stratified analyses examining person-time on or off atazanavir using time-updated models limited to those on antiretroviral therapy (ART) and repeated these analyses stratified by HCV status for those on non-atazanavir-containing regimens. All bilirubin values obtained while on ART were included. ART status varied over time based on whether individuals were currently receiving ART prescriptions.

We conducted sensitivity analyses excluding those with diabetes. We conducted sensitivity analyses stratified by sex. In models stratified by sex, we adjusted for HCV but did not stratify by HCV because of insufficient numbers among subgroups of stratified analyses. We conducted sensitivity analyses among the subset with indirect (unconjugated) bilirubin values.

RESULTS

We included 25,816 PLWH. The mean age at baseline was 40 years, 19% were female, 42% were white, and 18% had HCV (Table 1). There was a median of 9 total bilirubin values per person [interquartile range (IQR) 3–22]. The

TABLE 1. Demographic and Clinical Characteristics Overall (N = 25,816) and by Myocardial Infarction Status

Baseline	Overall	No MI	Type 1 MI	Type 2 MI
N	25,816	25,068	392	356
Age (mean)	40	40	47	45
Female, %	19	19	14	29
Race, %				
White	42	42	49	24
Black	40	40	38	65
Hispanic	13	13	10	8
Others/Missing	5	5	3	3
Hepatitis C, %	18	18	21	40
VL ≤ 400 , %	61	61	61	51
CD4 cell count cells/mm ³ (mean)	445	446	407	366
Bilirubin mg/dL (mean)	0.72	0.72	0.72	0.72

Percentages are the percent in each column with a particular characteristic. For example, overall 19% of the included participants are women. However, among those with a type 2 MI, 29% are women.

median baseline total bilirubin was 0.6 mg/dL (IQR 0.4–0.8) among men and 0.5 mg/dL (IQR 0.3–0.6) among women. During an average of 5.8 years of follow-up, 392 PLWH had a T1MI and 356 had a T2MI for an incidence rate of 2.6 per 1000 person-years [95% confidence interval (95% CI): 2.4 to 2.9] for T1MI and 2.4 per 1000 person-years (95% CI: 2.2 to 2.7) for T2MI. Of the T2MI, 127 (36%) were sepsis-related, 37 (10%) were related to cocaine-induced vasospasm, and 192 (54%) were related to a variety of other factors. There were 160 individuals with an ischemic stroke not related to infection or illicit drug use. There were 476 PLWH who had a combined outcome of a T1MI or ischemic stroke not associated with drug use or infection for an incidence rate of 3.2 per 1000 person-years (95% CI: 2.9 to 3.5). Those with both an MI and stroke were categorized by their first event.

Type 1 MI

In adjusted analyses, the hazard ratio (HR) for T1MI was <1.0 but not statistically significant (Table 2). For example, in analyses including bilirubin levels <2.1 mg/dL, the HR for the association of higher bilirubin levels with T1MI was 0.80 per 2-fold increase in bilirubin; 95% CI: 0.49 to 1.53 with similar findings among groups stratified by HCV status. Findings were not statistically different than 1.0 in analyses limited to those with baseline FIB-4 values <3.25 and in analyses that included everyone. Similar results were seen with categorical approaches (eg, bilirubin values between 1.3 and 2.1 vs. <0.6, HR 0.76; 95% CI: 0.33 to 1.72) and with spline models (data not shown). Similar patterns of nonsignificant findings were found in sensitivity analyses with an outcome of T1MI combined with ischemic strokes not associated with drug use or infection (Table 2).

Type 2 MI

In adjusted analyses limited to bilirubin levels <2.1 mg/dL, higher bilirubin levels were significantly associated with T2MI overall (HR 1.70 per 2-fold bilirubin increase; 95% CI: 1.04 to 2.78) and in analyses stratified by HCV status; higher bilirubin levels were significantly associated with T2MI among those without HCV (HR 2.52 per 2-fold bilirubin increase; 95% CI: 1.36 to 4.66) but not those with HCV (HR 1.01; 95% CI: 0.45 to 2.30) (Table 2; interaction *P* value, 0.01). Findings were similar in models limited to those with baseline FIB-4 values <3.25 but not in the models that included everyone (Table 2) where there was no difference by HCV status. In categorical analyses including bilirubin levels <2.1 mg/dL, higher categories of bilirubin were associated with T2MI, and this was driven by those without HCV. Higher bilirubin levels (compared with bilirubin levels <0.6) were associated with T2MI (bilirubin levels 0.6–1.3: HR 1.61; 95% CI: 1.15 to 2.27; bilirubin levels 1.3–2.1: HR 2.70; 95% CI: 1.34 to 5.46) among those without HCV. In spline models among everyone, a bilirubin increase of 0.1 in the range of 0.6–1.3 was associated with an increased risk of T2MI (1.09; 95% CI: 1.03 to 1.14), and this did not differ by

TABLE 2. Associations of Bilirubin Levels With Cardiovascular Outcomes Limited to Bilirubin Values <2.1 mg/dL, Limited to Those With FIB-4 Values <3.25, and Among Everyone Stratified by Hepatitis C Virus Coinfection Status in Adjusted Analyses*

Outcome	Hazard Ratio Per 2-Fold Increase in Bilirubin (95% CI)		
	HCV-Uninfected and Infected (Combined)	Stratified by HCV Status	
		HCV-Uninfected	HCV-Infected
N (limited to bilirubin values <2.1 mg/dL)	20,604	17,127	3477
Type 1 MI	0.80 (0.49 to 1.33)	0.78 (0.45 to 1.37)	0.88 (0.29 to 2.72)
Type 2 MI	1.70 (1.04 to 2.78)	2.52 (1.36 to 4.66)	1.01 (0.45 to 2.30)
Type I MI and ischemic stroke	0.87 (0.56 to 1.34)	0.98 (0.60 to 1.58)	0.53 (0.19 to 1.47)
N (limited to those with FIB-4 values <3.25 at baseline)	24,205	20,190	4015
Type 1 MI	1.01 (0.80 to 1.28)	1.04 (0.80 to 1.35)	0.90 (0.50 to 1.61)
Type 2 MI	1.35 (1.07 to 1.72)	1.56 (1.19 to 2.06)	1.00 (0.63 to 1.60)
Type I MI and ischemic stroke	0.99 (0.79 to 1.23)	0.99 (0.78 to 1.26)	0.97 (0.59 to 1.60)
N (all)	25,816	21,111	4705
Type 1 MI	0.98 (0.78 to 1.23)	0.99 (0.77 to 1.29)	0.93 (0.56 to 1.56)
Type 2 MI	1.47 (1.19 to 1.82)	1.53 (1.17 to 2.00)	1.42 (1.02 to 1.98)
Type I MI and ischemic stroke	0.98 (0.79 to 1.20)	0.94 (0.74 to 1.20)	1.06 (0.70 to 1.61)

*Analyses adjusted for time-varying viral load and CD4 count, and baseline age, sex, race/ethnicity, HIV transmission risk factor, smoking status, HCV status, diabetes, treated hypertension, statin use, kidney function (estimated GFR <30, 30–60, ≥60), calendar time, and site. Analyses stratified by HCV not adjusted for HCV. FIB-4, fibrosis-4 index; GFR, glomerular filtration rate; MI, myocardial infarction.

HCV status (without HCV, HR 1.09; 95% CI: 1.02 to 1.17; with HCV, HR 1.10; 95% CI: 1.01 to 1.19).

Atazanavir and ART

In models limited to those on ART, we stratified by atazanavir use (Table 3). During ART that did not include atazanavir, the pattern of findings was similar to the overall findings, showing a nonsignificant decreased risk of T1MI (HR 0.70; 95% CI: 0.43 to 1.15) and a significantly increased risk of T2MI associated with higher bilirubin levels (HR 2.00; 95% CI: 1.52 to 2.62). During ART that included atazanavir, the association with T1MI was similar to the main analyses with a HR <1.0 but not statistically significant. Higher

bilirubin levels were associated with significantly fewer of the combined outcome of T1MI and ischemic stroke (HR 0.64; 0.41 to 0.99). However, the association with T2MI was different than among those not on atazanavir: higher bilirubin levels were associated with significantly fewer T2MI events (HR 0.56; 95% CI: 0.33 to 1.00).

Sensitivity Analyses

For the analysis shown in Table 2, findings did not differ in analyses that excluded those with diabetes. In adjusted analyses adjusting for HCV status and stratified by sex limiting to those with bilirubin values <2.1, associations with T1MI were not significant, whereas for T2MI, higher

TABLE 3. Associations of Bilirubin Levels With Cardiovascular Outcomes Among Those on Antiretroviral Therapy Stratified by Atazanavir Use and by Hepatitis C Virus Coinfection Status in Adjusted Analyses*,†

Outcome	Hazard Ratio Per 2-Fold Increase in Bilirubin (95% CI)		
	HCV-Uninfected and Infected (Combined)	Stratified by HCV Status	
		HCV-Uninfected	HCV-Infected
N (limited to those on ART not including atazanavir)	20,305	16,706	3599
Type 1 MI	0.70 (0.43 to 1.15)	0.65 (0.36 to 1.17)	0.82 (0.33 to 2.04)
Type 2 MI	2.00 (1.52 to 2.62)	2.11 (1.37 to 3.25)	2.03 (1.40 to 2.95)
Type I MI and ischemic stroke	0.91 (0.61 to 1.35)	0.69 (0.41 to 1.16)	1.29 (0.75 to 2.22)
N (limited to those on ART including atazanavir)	5671		
Type 1 MI	0.68 (0.43 to 1.09)	‡	‡
Type 2 MI	0.56 (0.33 to 1.00)		
Type I MI and ischemic stroke	0.64 (0.41 to 0.99)		

*Analyses adjusted for time-varying viral load and CD4 count, and baseline age, sex, race/ethnicity, HIV transmission risk factor, smoking status, HCV status, diabetes, treated hypertension, statin use, kidney function (estimated GFR <30, 30–60, ≥60), calendar time, and site. Analyses stratified by HCV not adjusted for HCV.

†Note that ART use and atazanavir user are time-varying.

‡Atazanavir analyses not stratified because of small outcome numbers. GFR, glomerular filtration rate.

bilirubin levels were associated with increased risk for women only (4.58; 95% CI: 2.06 to 10.1); *P* value for bilirubin × sex interaction term, 0.02.

Findings were also similar but not significant in analyses among the subset of 4963 with indirect bilirubin values (data not shown).

DISCUSSION

In a large cohort of PLWH with carefully adjudicated cardiovascular outcomes, we found that overall, mildly elevated levels of total bilirubin (<2.1 mg/dL) were associated with increased risk of T2MI, particularly among those without HCV. Interestingly, among those on ART and including all bilirubin values, bilirubin levels were not associated with increased T2MI among those currently taking atazanavir; in fact, the opposite was the case. In the subset of individuals currently on atazanavir, elevated bilirubin levels were associated with reduced T2MI risk. Among those currently on atazanavir, elevated bilirubin levels were also associated with reduced risk of the combined outcome of T1MI and ischemic stroke. Mildly elevated bilirubin levels (<2.1 mg/dL) had a HR for the association with T1MI that was <1 consistent with protective findings seen in the general population; however, this association was not significant. To the best of our knowledge, these analyses are one of the largest to date evaluating the relationship between bilirubin and cardiovascular outcomes among PLWH and the first to do so distinguishing T1MI and T2MI.

PLWH with T2MI have previously been shown to have more severe HIV disease on average than those with T1MI.³¹ Elevated bilirubin in this setting possibly represents the greater burden of underlying comorbidity. However, we adjusted for HIV disease severity as measured by the time-varying CD4 count and VL and still found higher bilirubin levels were associated with T2MI. We hypothesized that liver disease might be a driver of this association; however, the association between higher bilirubin and T2MI was particularly notable in those without HCV. Heavy alcohol use and alcohol-related liver injury may also be a potential confounding factor, but our findings remained consistent in analyses that eliminated those with higher baseline FIB-4 values or with more elevated bilirubin levels, who likely comprise most patients with alcoholic hepatitis. We were concerned that higher bilirubin levels and T2MI may be surrogates for the same thing, such as sepsis that caused widespread organ ischemia; therefore, all analyses excluded bilirubin levels ≤30 days before the MI to minimize this.

These findings add to the limited literature on bilirubin, atazanavir, and cardiovascular disease outcomes in PLWH. A cross-sectional case-control study examining carotid intimal media thickness found little difference by the bilirubin level, but the small sample size (*N* = 78) limited the conclusions that could be drawn.³⁹ A study compared those on atazanavir (*N* = 11) vs. those not on atazanavir (*N* = 51) and found that both atazanavir use and higher baseline total serum bilirubin levels were associated with slower mean carotid intima media thickness progression.⁴⁰ Another study of 145 PLWH on atazanavir vs. efavirenz-based regimens found oxidative

stress measured by lipoprotein-associated phospholipase A2 and oxidized low-density lipoprotein was reduced in those on atazanavir and was correlated with elevated bilirubin levels.⁴¹ One of the more compelling studies (AIDS Clinical Trials Group 5260s) was a trial of ART-naïve individuals randomized to atazanavir vs. darunavir vs. raltegravir-based regimens. Those on atazanavir had slower carotid intima media thickness progression than those on darunavir, and in some analyses, raltegravir. This was associated in part with bilirubin levels >0.6 mg/dL,²¹ although bilirubin levels may also serve as a marker of good ART adherence for those on atazanavir.⁴² A recent large study conducted in the Veterans Affairs (VA) Health Care System suggested that higher bilirubin levels are associated with decreased cardiovascular events such as heart failure and ischemic stroke among PLWH, although associations with MI were not statistically significant and that study did not distinguish MI type.²² Another large study from the VA was conducted in 9500 PLWH.⁴³ A study strength was that it examined the association of atazanavir with MI and stroke rather than intermediate outcomes. However, MI and stroke were not adjudicated, raising concerns about accuracy, and surprisingly, there were higher rates of strokes than MI, raising questions about MI capture completeness. The VA study reported a lower incidence of MI and stroke among PLWH on atazanavir compared with other regimens; however, because this study was unable to distinguish T1MI from T2MI, it is not clear whether the findings were similar to ours.

We hypothesized that among PLWH on atazanavir, higher bilirubin levels would be associated with reduced MI risk, particularly T1MI. Instead, we found that higher bilirubin levels were associated with reduced risk of both types of MI, reaching statistical significance for T2MI risk or risk of the combined outcome of T1MI and ischemic stroke. Interestingly, this is in direct contrast to individuals on ART that did not include atazanavir, in whom higher bilirubin levels were significantly associated with increased T2MI risk. These findings are intriguing and suggest that the impact of bilirubin levels on cardiovascular disease in PLWH may differ compared with the general population. The difference in overall findings among those with and without HCV, and in particular, the protective association of elevated bilirubin levels among those on atazanavir, where presumably many of the higher bilirubin levels are due to atazanavir rather than illness, highlights the diverse causes of elevated bilirubin levels among PLWH. We suspect that bilirubin elevations due to benign causes such as atazanavir may be protective, similar to the protective associations seen with Gilbert syndrome in the general population. However, other factors such as liver disease including related to HIV disease itself, viral hepatitis, and hazardous alcohol use may all contribute to bilirubin elevation and may not necessarily have protective associations with either T1MI or T2MI.

We considered whether associations between atazanavir, bilirubin, and T2MI were due to channeling bias. There is always a concern about medication channeling, where the healthier participants are selectively prescribed a particular pharmacotherapy (eg, atazanavir). However, we suspect that if the protective association between bilirubin and T2MI among

patients on atazanavir vs. those on other regimens was due to health status differences and not direct effects of the atazanavir and bilirubin metabolism; we would have seen differences between those on atazanavir vs. those on other ART regimens with T1MI. The HRs for the association of bilirubin with T1MI are very similar among those on atazanavir vs. other regimens (HR 0.68 vs. 0.70). This difference in the bilirubin association by atazanavir use between MI types makes it less likely that the protective association between T2MI and bilirubin in the atazanavir group is due to channeling of healthier participants toward atazanavir.

A key finding is the importance of distinguishing T1MI and T2MI. The distinct associations by MI type add to previous studies demonstrating substantial differences in demographic and clinical characteristics and outcomes among PLWH who experienced a T1MI vs. T2MI.^{31,44} These differences in risk factors combined with the diverse pathophysiologies and concomitant comorbidities associated with different MI types add to the compelling argument that MI type in PLWH should be distinguished rather than combined.³¹

Strengths

The large sample size and well-defined outcomes are the strengths. The careful adjudication, including distinguishing T1MI and T2MI, is of value, particularly because we found that the association between bilirubin and atherosclerotic T1MI differs from the association with T2MI. We evaluated outcomes separately. We used several different bilirubin parameterizations based on the recommendations from studies not conducted in PLWH.^{9,10,45}

An additional strength is that we excluded bilirubin values ≥ 2.1 mg/dL from some models to minimize the impact of hepatic pathology. Although higher bilirubin levels have been associated with decreased atherosclerotic disease in the general population, this association has not always been found or different patterns such as u-shaped associations have been seen. This may be due to liver disease,⁹ which would likely have as big, if not bigger, impact among PLWH. However, excluding values ≥ 2.1 mg/dL may impact being able to evaluate a potential protective effect of hyperbilirubinemia associated with atazanavir because 2.1 mg/dL is not much higher than mean values on atazanavir.⁴⁶ Therefore, we repeated analyses using a FIB-4 cutoff and among everyone. We included all bilirubin values for the atazanavir analyses. We used time-updated models to examine atazanavir vs. other drugs and did not limit people to baseline regimens but allowed them to contribute whenever they were on ART.

Limitations

We analyzed total bilirubin, which combines conjugated and unconjugated bilirubin values, and does not distinguish between elevated bilirubin caused by hepatic pathology or by benign deficiency in excretory capacity.⁴ Some associations between bilirubin and reduced cardiovascular risk have been in the context of elevated unconjugated bilirubin levels such as with Gilbert syndrome.⁶ In our analysis, the overall results were similar in the subset with

unconjugated bilirubin levels available; it was a small subset and no associations were significant. In addition, we did not limit bilirubin levels to fasting values; associations may be stronger in studies using fasting bilirubin levels¹⁰ because fasting may impact bilirubin levels.⁴⁷ Our study is observational, so causality cannot be inferred. This limitation has also been noted in general population studies regarding whether relationships between bilirubin levels and cardiovascular disease may be causal or due to confounding and reverse causality.^{10,48} We stratified overall results by HCV status to better understand the influence of liver disease, but owing to low numbers, we were unable to stratify by hepatitis B.

More research seems warranted with carefully adjudicated T1MI vs. T2MI to better understand the mechanism of the associations between bilirubin and T2MI, the complicated relationships between bilirubin and cardiovascular outcomes in general, and whether there is a protective effect of mildly elevated bilirubin levels not due to liver disease and the role of specific antiretroviral medications, potentially done with unconjugated bilirubin levels rather than total bilirubin. Differences in associations between those with and without HCV and receiving or not receiving atazanavir suggest the need for a nuanced approach to evaluating bilirubin that considers causes of elevated levels among PLWH. In addition, the larger risk of T2MI associated with a 2-fold increase in bilirubin among women than men raises questions regarding reasons for differences between men and women worthy of future study.

Conclusions

In this large cohort, the HR for the association of higher bilirubin with T1MI was <1 , consistent with general population studies, but not statistically significant. If the antioxidant effect of bilirubin reduces cardiovascular risk among PLWH, the impact may be small and difficult to detect even with large samples. Higher bilirubin levels were associated with increased T2MI risk, particularly among those without HCV, raising questions about potential mechanisms. By contrast, among patients on atazanavir, higher bilirubin levels were associated with lower risk of T2MI. Among those on atazanavir, higher bilirubin levels were associated with lower risk of the combined outcome of T1MI and ischemic stroke. These findings demonstrate differences in impact of elevated bilirubin levels due to diverse causes on outcomes among PLWH and highlight the importance of carefully validated outcomes including MI types and looking at individual outcomes rather than composite outcomes where relationships differ.

ACKNOWLEDGMENTS

The authors acknowledge the patients, staff, and providers throughout CNICS.

REFERENCES

1. Stocker R, Yamamoto Y, McDonagh AF, et al. Bilirubin is an antioxidant of possible physiological importance. *Science*. 1987;235:1043–1046.
2. Vitek L. Bilirubin and atherosclerotic diseases. *Physiol Res*. 2017; 66(suppl 1):S11–S20.

3. Kang SJ, Lee C, Krusziak P. Effects of serum bilirubin on atherosclerotic processes. *Ann Med*. 2014;46:138–147.
4. Boon AC, Bulmer AC, Coombes JS, et al. Circulating bilirubin and defense against kidney disease and cardiovascular mortality: mechanisms contributing to protection in clinical investigations. *Am J Physiol Ren Physiol*. 2014;307:F123–F136.
5. Akboga MK, Canpolat U, Sahinarslan A, et al. Association of serum total bilirubin level with severity of coronary atherosclerosis is linked to systemic inflammation. *Atherosclerosis*. 2015;240:110–114.
6. Kundur AR, Singh I, Bulmer AC. Bilirubin, platelet activation and heart disease: a missing link to cardiovascular protection in Gilbert's syndrome? *Atherosclerosis*. 2015;239:73–84.
7. Vitek L, Jirsa M, Brodanová M, et al. Gilbert syndrome and ischemic heart disease: a protective effect of elevated bilirubin levels. *Atherosclerosis*. 2002;160:449–456.
8. Perlstein TS, Pande RL, Creager MA, et al. Serum total bilirubin level, prevalent stroke, and stroke outcomes: NHANES 1999–2004. *Am J Med*. 2008;121:781–788 e781.
9. Novotny L, Vitek L. Inverse relationship between serum bilirubin and atherosclerosis in men: a meta-analysis of published studies. *Exp Biol Med*. 2003;228:568–571.
10. Kunutsor SK, Bakker SJ, Gansevoort RT, et al. Circulating total bilirubin and risk of incident cardiovascular disease in the general population. *Arteriosclerosis, Thromb Vasc Biol*. 2015;35:716–724.
11. Kawamoto R, Ninomiya D, Hasegawa Y, et al. Mildly elevated serum total bilirubin levels are negatively associated with carotid atherosclerosis among elderly persons with type 2 diabetes. *Clin Exp Hypertens*. 2016;38:107–112.
12. Temme EH, Zhang J, Schouten EG, et al. Serum bilirubin and 10-year mortality risk in a Belgian population. *Cancer Causes Control*. 2001;12:887–894.
13. Kunutsor SK. Serum total bilirubin levels and coronary heart disease—Causal association or epiphenomenon? *Exp Gerontol*. 2015;72:63–66.
14. Freiberg MS, Chang CC, Kuller LH, et al. HIV infection and the risk of acute myocardial infarction. *JAMA Intern Med*. 2013;173:614–622.
15. Durand M, Sheehy O, Baril JG, et al. Association between HIV infection, antiretroviral therapy, and risk of acute myocardial infarction: a cohort and nested case-control study using Quebec's public health insurance database. *J Acquir Immune Defic Syndr*. 2011;57:245–253.
16. Triant VA, Lee H, Hadigan C, et al. Increased acute myocardial infarction rates and cardiovascular risk factors among patients with human immunodeficiency virus disease. *J Clin Endocrinol Metab*. 2007;92:2506–2512.
17. Currier JS, Lundgren JD, Carr A, et al. Epidemiological evidence for cardiovascular disease in HIV-infected patients and relationship to highly active antiretroviral therapy. *Circulation*. 2008;118:e29–35.
18. Petersen K, Riddle MS, Jones LE, et al. Use of bilirubin as a marker of adherence to atazanavir-based antiretroviral therapy. *AIDS*. 2005;19:1700–1702.
19. Zhang D, Chando TJ, Everett DW, et al. In vitro inhibition of UDP glucuronosyltransferases by atazanavir and other HIV protease inhibitors and the relationship of this property to in vivo bilirubin glucuronidation. *Drug Metab Dispos*. 2005;33:1729–1739.
20. Smith DE, Jegannathan S, Ray J. Atazanavir plasma concentrations vary significantly between patients and correlate with increased serum bilirubin concentrations. *HIV Clin Trials*. 2006;7:34–38.
21. Stein JH, Ribaldo HJ, Hodis HN, et al. A prospective, randomized clinical trial of antiretroviral therapies on carotid wall thickness. *AIDS*. 2015;29:1775–1783.
22. Marconi VC, Duncan MS, So-Armah K, et al. Bilirubin is inversely associated with cardiovascular disease among HIV-positive and HIV-negative individuals in VACS (Veterans Aging Cohort Study). *J Am Heart Assoc*. 2018;7:e007792.
23. Thygesen K, Alpert JS, White HD, et al. Universal definition of myocardial infarction. *Circulation*. 2007;116:2634–2653.
24. Thygesen K, Alpert JS, Jaffe AS, et al. Third universal definition of myocardial infarction. *Eur Heart J*. 2012;33:2551–2567.
25. Javed U, Aftab W, Ambrose JA, et al. Frequency of elevated troponin I and diagnosis of acute myocardial infarction. *Am J Cardiol*. 2009;104:9–13.
26. Morrow DA, Wiviott SD, White HD, et al. Effect of the novel thienopyridine prasugrel compared with clopidogrel on spontaneous and procedural myocardial infarction in the Trial to Assess Improvement in Therapeutic Outcomes by Optimizing Platelet Inhibition with Prasugrel-Thrombolysis in Myocardial Infarction 38: an application of the classification system from the universal definition of myocardial infarction. *Circulation*. 2009;119:2758–2764.
27. Melberg T, Burman R, Dickstein K. The impact of the 2007 ESC-ACC-AHA-WHF Universal definition on the incidence and classification of acute myocardial infarction: a retrospective cohort study. *Int J Cardiol*. 2010;139:228–233.
28. Szymanski FM, Karpinski G, Platek AE, et al. Clinical characteristics, aetiology and occurrence of type 2 acute myocardial infarction. *Kardiologia Pol*. 2014;72:339–344.
29. Stein GY, Herscovici G, Korenfeld R, et al. Type-II myocardial infarction—patient characteristics, management and outcomes. *PLoS One*. 2014;9:e84285.
30. Baron T, Hambraeus K, Sundström J, et al. Type 2 myocardial infarction in clinical practice. *Heart*. 2015;101:101–106.
31. Crane HM, Paramsothy P, Drozd DR, et al. Types of myocardial infarction among human immunodeficiency virus-infected individuals in the United States. *JAMA Cardiol*. 2017;2:260–267.
32. Adams HP Jr, Bendixen BH, Kappelle LJ, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke*. 1993;24:35–41.
33. Kitahata MM, Rodriguez B, Haubrich R, et al. Cohort profile: the Centers for AIDS Research Network of Integrated Clinical Systems. *Int J Epidemiol*. 2008;37:948–955.
34. Crane HM, Heckbert SR, Drozd DR, et al. Lessons learned from the design and implementation of myocardial infarction adjudication tailored for HIV clinical cohorts. *Am J Epidemiol*. 2014;179:996–1005.
35. Crane HM, Kadane JB, Crane PK, et al. Diabetes case identification methods applied to electronic medical record systems: their use in HIV-infected patients. *Curr HIV Res*. 2006;4:97–106.
36. Jain MK, Seremba E, Bhore R, et al. Change in fibrosis score as a predictor of mortality among HIV-infected patients with viral hepatitis. *AIDS Patient Care STDS*. 2012;26:73–80.
37. Bambha K, Pierce C, Cox C, et al. Assessing mortality in women with hepatitis C virus and HIV using indirect markers of fibrosis. *AIDS*. 2012;26:599–607.
38. Nunes D, Fleming C, Offner G, et al. Noninvasive markers of liver fibrosis are highly predictive of liver-related death in a cohort of HCV-infected individuals with and without HIV infection. *Am J Gastroenterol*. 2010;105:1346–1353.
39. Barber TJ, Moyle G, Hill A, et al. A cross-sectional study to evaluate the association of hyperbilirubinaemia on markers of cardiovascular disease, neurocognitive function, bone mineral density and renal markers in HIV-1 infected subjects on protease inhibitors. *HIV Clin Trials*. 2016;17:123–130.
40. Chow D, Kohorn L, Souza S, et al. Atazanavir use and carotid intima media thickness progression in HIV: potential influence of bilirubin. *AIDS*. 2016;30:672–674.
41. Estrada V, Monge S, Gómez-Garre MD, et al. Relationship between plasma bilirubin level and oxidative stress markers in HIV-infected patients on atazanavir- vs. efavirenz-based antiretroviral therapy. *HIV Med*. 2016;17:653–661.
42. Rekić D, Clewe O, Roshammam D, et al. Bilirubin—a potential marker of drug exposure in atazanavir-based antiretroviral therapy. *AAPS J*. 2011;13:598–605.
43. LaFleur J, Bress AP, Rosenblatt L, et al. Cardiovascular outcomes among HIV-infected veterans receiving atazanavir. *AIDS*. 2017;31:2095–2106.
44. Nance R, Delaney J, Heckbert S, et al. Death after myocardial infarction (MI): incidence and risk factors by MI type in HIV. 9th IAS Conference on HIV Science, July 2017; Paris, France, 2017.
45. Song YS, Koo BK, Cho NH, et al. Effect of low serum total bilirubin levels (≤ 0.32 mg/dl) on risk of coronary artery disease in patients with metabolic syndrome. *Am J Cardiol*. 2014;114:1695–1700.
46. Cotter AG, Brown A, Sheehan G, et al. Predictors of the change in bilirubin levels over twelve weeks of treatment with atazanavir. *AIDS Res Ther*. 2013;10:13.
47. Barrett PV. Hyperbilirubinemia of fasting. *JAMA*. 1971;217:1349–1353.
48. Stender S, Frikke-Schmidt R, Nordestgaard BG, et al. Genetically elevated bilirubin and risk of ischaemic heart disease: three Mendelian randomization studies and a meta-analysis. *J Intern Med*. 2013;273:59–68.