

RECEIVED
16 October 2017

REVISED
6 December 2017

ACCEPTED FOR PUBLICATION
15 December 2017

PUBLISHED
7 February 2018

Transition and post-transition metals in exhaled breath condensate

Andrew J Ghio^{1,3} , Michael C Madden¹ and Charles R Esther Jr²

¹ National Health and Environmental Effects Research Laboratory, Environmental Protection Agency, Chapel Hill NC, United States of America

² Pediatric Pulmonology, University of North Carolina, Chapel Hill NC, United States of America

³ Author to whom any correspondence should be addressed.

E-mail: ghio.andy@epa.gov

Keywords: lung, transition elements, iron, zinc, metals

Abstract

Water vapor in expired air, as well as dispersed non-volatile components, condense onto a cooler surface after exiting the respiratory tract. This exhaled breath condensate (EBC) provides a dilute sampling of the epithelial lining fluid. Accordingly, the collection of EBC imparts a capacity to provide biomarkers of injury preceding clinical disease. Concentrations of transition and post-transition metals in EBC are included among these endpoints. Iron and zinc are the metals with the highest concentration and are measurable in all EBC samples from healthy subjects; other metals are most frequently either at or below the level of detection in this group. Gender, age, and smoking can impact EBC metal concentrations in healthy subjects. EBC metal concentrations among patients diagnosed with particular lung diseases (e.g. asthma, chronic obstructive disease, and interstitial lung disease) have been of research interest but no definite pattern of involvement has been delineated. Studies of occupationally exposed workers confirm significant exposure to specific metals, but such EBC metal measurements frequently provide evidence redundant with environmental sampling. Measurements of metal concentrations in EBC remain a research tool into metal homeostasis in the respiratory tract and participation of metals in disease pathogenesis. The quantification of metal concentrations in EBC is currently not reliable for clinical use in either supporting or determining any diagnosis. Issues that must be addressed prior to the use of EBC metal measurements include the establishment of both standardized collection and measurement techniques.

1. Introduction

As a result of a large surface area (50–75 m²), a considerable volume of water is lost daily from the human body through its evaporation from the epithelial lining fluid of the respiratory tract [1]. While much of this volume is water vapor, a small fraction represents aerosol droplets generated through an action of turbulent flow on the epithelial lining fluid [2], or by the opening and closing of small airways [3]. Following this aerosolization, non-volatile molecules are dispersed into the water vapor. Water vapor in expired air, as well as the dispersed non-volatile components, condense onto a cooler surface after exiting the respiratory tract. Instrumentation can be added to lower the temperature of the surface and augment collection. This liquid sampling obtained from a subject, typically breathing at tidal breathing, is exhaled breath condensate (EBC).

The human respiratory tract represents the route of entry for many environmental and occupational exposures. Accordingly, volatile and non-volatile substances in EBC can function as biomarkers since concentrations can be impacted by these exposures. Such biomarkers may potentially precede evidence of injury and therefore predict clinical disease. Concentrations of transition and post-transition metals in EBC are included among these endpoints.

2. Metals in the respiratory tract

Transition and post-transition metals were selected in molecular evolution to carry out a wide range of biological functions and are required by all living organisms. They are utilized in almost every aspect of normal cell function and are particularly crucial for cellular metabolism. Consequently, almost all living

organisms require metals, including iron, zinc, copper, chromium, cobalt, manganese, molybdenum, nickel, tin, and vanadium.

Cells resident in the lung express proteins which participate in the import, export, storage, and transport of metals (e.g. divalent metal transporter I, ferroportin I, ferritin, and transferrin receptor respectively) [4]. Metal uptake by non-diseased, non-exposed lung with systemic translocation has been demonstrated [5]. However, the positioning of these proteins does not support metal uptake with systemic translocation to meet nutritional requirements as a normal function for which the lung was designed [6].

While all cells in the lower respiratory tract participate in metal homeostasis, the handling of metals is a primary responsibility of airway and alveolar macrophages [7, 8]. The macrophage has a capacity to mobilize, import, store, and release metals for transport to tissues of the reticuloendothelial system (e.g. the liver). The overwhelming majority of metals in the lung can be localized to macrophages [9]. Stains reflect this localization of metals in the respiratory tract to macrophages [10]. The macrophage, with its accumulated metals, can be removed from the lung via mucociliary clearance and lymphatics.

Metal homeostasis in the lung is a dynamic rather than a static process. Following exposure of the respiratory tract, metals are removed and some portion can be translocated outside of the lungs, reaching other tissues of the body [11]. This movement of metals from the respiratory tract into the systemic circulation with distribution to numerous different tissues is observed with particle-associated metals [12]. Such translocation from the respiratory tract can occur rapidly (e.g. over hours). Those metals with higher water solubility are more quickly transported systemically relative to those in an insoluble state. Clearance from the respiratory tract may also correlate with the availability in the lung of reductants and chelators which function to solubilize metals. Measurements of metals in samples of the lung will reflect the efficiency of the clearance of the metals and their systemic translocation. Accordingly, metal concentrations in the respiratory tract will be dependent on the duration of time since the initial exposure.

3. Collection of EBC

The American Thoracic Society and European Respiratory Society have provided guidelines for the collection of EBC [13]. Such collection necessitates that a subject breathe through a one-way valve into a tube cooled to condense exhaled water vapor. A collecting period of 10 min or more provides a volume of condensate suitable for metals analysis (2–3 ml). The device can be integrated with a pneumotachograph allowing a definition for the completion of collection as the total expired volume

rather than as a duration of time [14]. A device with audio and visual prompts can also be employed to control the respiratory rate, tidal volume, and expired volume and to terminate the effort [15].

At the low concentrations measured in EBC, metal contamination during the collection is a concern. The specific collection device can impact metal concentrations [16, 17]. To minimize metal contamination, the condensing surface should be a plastic polymer or fluoropolymer. The plasticware which the EBC is collected into should be disposable. Alternatively, components can be acid washed prior to use in an effort to diminish metal contamination.

Contamination of EBC due to metals included in particles in ambient air should be considered and eliminated if possible. All metals in the atmosphere, except for mercury, are associated with particles [18]. If one assumes that (1) a tidal volume of 0.5 L, (2) a respiratory rate of 20 min⁻¹, (3) a collection time of 10 min, (4) an ambient particulate level of 20 μg m⁻³, (5) a metal composition in the particles of 1%, and (6) lung deposition of particles which approximates 50%, it is calculated that with the collection of the exhaled particles, the metal concentrations could approximate levels of parts per billion [19]:

- (1) 0.5 L breath⁻¹ (tidal volume) × 20 breaths per minute × 10 min = 100 L or 0.1 m³ (inspired volume over 10 min of collection);
- (2) 0.1 m³ (inspired volume) × 20 μg m⁻³ (particle level) = 2 μg (particle mass in the inspired volume over 10 min of collection);
- (3) 2 μg (particle mass) × 50% (percentage not retained and expired) 1 ml⁻¹ (volume of EBC) = 1 μg particle 2 ml⁻¹ EBC;
- (4) 1 μg particle 2 ml⁻¹ EBC × 1% metals = 5 ppb metals (mass per volume).

With investigations providing measurements of metals in parts per trillion, particles in the ambient air potentially impact the results. This is particularly true with occupational studies where the ambient particle levels are elevated and there is a higher metal content (e.g. welders). An appropriately-sized filter attached to the inlet of the breathing train can be used to reduce the exposure to particles during EBC collection.

4. Measurements of metal concentrations in EBC

Metals in bronchoalveolar lavage (BAL) and EBC can be quantified using graphite furnace atomic absorption spectrometry (GFAAS) and inductively coupled plasma mass spectrometry (ICPMS). Detection limits for both GFAAS and ICPMS for metals are excellent. However, the ICPMS can measure more elements (greater than 75), has multi-element capability, and

Table 1. Concentrations of metal in lavage collected from healthy subjects.

Study	<i>n</i>	Subjects	Smoking status	Fe	Zn	Cu	Cr	Mn	Ni	Pb	Ti
Sabbioni <i>et al</i> 1987	4–22	Not provided	Not provided	508	510	215	10	2.7	145	34.6	<1
Romeo <i>et al</i> 1992	25	Patients	N	106			0.6	1.5		3.2	
Corhay <i>et al</i> 1995	45	Healthy	N and S	170	300	50	20		10		140
Nelson <i>et al</i> 1996	21	Healthy	N and S	2–9							
Harlyk <i>et al</i> 1997	157	Patients	Not provided	0–120	10–230	0–15					
Stites <i>et al</i> 1999	8	Healthy	N	0							
	8	Healthy	S	100							
Ghio <i>et al</i> 1998	22	Healthy	N	40–70							
Ghio <i>et al</i> 2003	28	Healthy	N	≈50							
Bargagli <i>et al</i> 2008	9	Healthy	N and S	32	8	3	5	10	3	10	
Ghio <i>et al</i> 2013	20	Healthy	N	<50							

N is non-smoker and S is smoker. Metal concentrations are provided in ppb ($\mu\text{g L}^{-1}$).

demonstrates better linearity of range and in-run precision [20]. The drawbacks of ICPMS include the fact that the volume of the sample required is much larger, the costs (operating and capital costs) are greater, and the training of the individual performing the assay must be exceptional. While the choice between GFAAS and ICPMS will frequently be determined by laboratory size, specialization, and resources, ICPMS must be considered the method of choice for multi-element screening in EBC.

Ultrapure water must be obtained and used for the dilution of the stock solutions and to prepare the blanks. Multi-element solutions can be purchased for external standards. An internal standard (e.g. yttrium) is recommended. Results can be reported as metals per volume condensate (i.e. microgram per ml and parts per billion (ppb)) rather than metals per expired volume as the latter is difficult to interpret and will reflect the variability of both the metal measurements and the spirometer. Both instrument detection limits and method detection limits should be calculated.

Metals in EBC should not be measured prior to acidification/digestion; the acid employed (HCl or HNO₃) should be very low in metals (e.g. Optima grade, Fisher Scientific). A direct assay of EBC for metals without prior homogenization/digestion will not provide an accurate analysis [21].

A significant increase in metal concentrations in samples was reported after their storage for several weeks [17]. This was presumed to be the result of leaching from the storage containers. Accordingly, it is currently recommended that the samples be analyzed as soon as possible.

Metal analysis of EBC can result in levels which are below the method detection limit in the majority of the samples [17]. Investigations have reported aluminum, cadmium, chromium, and tungsten levels to be below the limits of detection in the majority of samples [16, 22–25]. Subsequently, group effects can frequently indicate the number of samples with detectable concentrations [25]. Such findings must be interpreted with some caution.

Finally, as a result of a lack of standardized methodology for sampling and normalization for dilution and low absolute levels, some studies express results as concentrations of metallic elements relative to each other rather than as absolute concentrations. While this approach is not ideal, it can provide mechanistic insights into the pathophysiology underlying individual diseases.

5. Metals in EBC in normal populations

The epithelial lining fluid of the respiratory tract (i.e. the alveolar and airway lining fluids) is most commonly sampled using BAL and EBC. Regarding dilutions, it is estimated that approximately 1.0 +/- 0.1 ml of epithelial lining fluid was recovered per 100 ml BAL fluid [26]. In view of this, BAL fluid is considered an approximately 1/100 dilution of the epithelial lining fluid. The EBC is a more dilute sampling of the epithelial lining fluid [27]. Based on (1) the dilution of the BAL fluid approximating 1:100 and (2) the total protein concentration in BAL fluid and EBC approximating 100 and 1 $\mu\text{g ml}^{-1}$ respectively, an estimate of the dilution of epithelial lining fluid that EBC exhibits would be 1: 10 000 [28, 29]. Similar dilution estimates have been calculated using urea or electrolytes (sodium and potassium) as dilution markers [30, 31]. This dilution of EBC appears to be extremely variable both among individuals and within an individual over time [30–32]. This may correspond to changes in ventilation and condensation temperature as the main determinants of evaporation and efficiency of collection respectively [33], or to individual differences in the ability to generate aerosols of airway lining fluid [34].

Metals which have been quantified in BAL fluid collected from healthy subjects are few (table 1) [35–44]. Iron, zinc, copper, and chromium are the metals measured most frequently in BAL fluid from healthy subjects. The range of metal concentrations in these individuals is wide, indicating the different

Table 2. Concentrations of metal in EBC collected from healthy subjects.

Study	<i>n</i>	Subjects	Smoking status	Fe	Zn	Cu	Cr	Mn	Ni	Pb	Ti
Caglieri <i>et al</i> 2006	25	Healthy	N and EX				0.28				
Mutti <i>et al</i> 2006	50	Not provided	N and EX	≈10		≈1.25		≈0.1	<1	<0.10	
Goldoni <i>et al</i> 2008	20	Healthy	N				0.18				
Corradi <i>et al</i> 2009	33	Healthy	N	1.20	1.60	0.60	ND	0.10	0.02	0.02	
Vlasic <i>et al</i> 2009	22	Healthy	N	22							
Fox <i>et al</i> 2013	8	Not provided	Not provided				0.25	0.57	0.87		
Hulo <i>et al</i> 2014	16	Not provided	N and S	0.50				0.32	0.24		
Pelcova <i>et al</i> 2015	20	Not provided	N, S, and EX								ND
Leese <i>et al</i> 2017	22	Not provided	Not provided				0.01–0.09				

Metal concentrations are provided in ppb ($\mu\text{g L}^{-1}$). N is non-smoker, S is smoker, and EX is ex-smoker. ND is not detectable.

methodologies employed in performing both the procedure and the analysis. In addition, dissimilar populations of healthy subjects were used and could include non-smokers, smokers, and even patients. Despite the widely disparate approaches, it is accepted that iron and zinc are those metals in greatest concentrations in BAL fluid, with copper being less. It is uncertain that other metals are at quantifiable levels in BAL fluid collected from healthy subjects. This does not negate a significant role for other metals in the human lung of both healthy and diseased individuals, but defining metal participation in a biological effect may be problematic. While metals accumulate with age in numerous tissues, including the lung, an increase in BAL fluid metals among older, healthy subjects has not been shown [45, 46]. Smoking does increase levels of some metals in BAL fluids [41, 43]. A comparison of metal concentrations in BAL fluid with those in blood supports the possibility of blood being a source of iron, zinc, and copper in the epithelial lining fluid [47–50]. However, all other metals in BAL fluid are reported at approximately comparable levels as in blood, and it is proposed that environmental sources (e.g. air pollution particles, environmental tobacco smoke, and smoking) may impact these concentrations more than those in the vascular compartment.

EBC has numerous advantages over BAL fluid as a sampling of epithelial lining fluid in that its collection (1) does not demand a difficult preparation, (2) is non-invasive, (3) requires a short duration of time, (4) can be repeated multiple times, and (5) can be based outside of a medical clinic or hospital [27, 51]. Accordingly, metals have also been measured in EBC collected from healthy subjects (table 2) [14, 17, 52–58]. Comparable to BAL fluid, iron and zinc are those metals in highest concentration and are measurable in all EBC samples. However, other metals in the EBC samples are much lower in concentration and the majority of the samples are below the level of detection [17, 55]. Gender, age, and smoking can impact EBC metal concentrations [16, 17, 23, 25, 53, 59]. Differences between genders as well as aging can reflect the volume of collected condensate which is proportional to the ventilatory volume (pulmonary function in

healthy non-smokers is determined by gender, age, and height/weight). Tobacco smoke includes metals, and smoking can be predicted to introduce some quantity of metals into the respiratory tract [60]. Lung tissue from smokers demonstrated elevated concentrations of iron [46]. Current healthy smokers can have higher EBC concentrations of lead and cadmium relative to healthy non-smokers [51]. In smokers, iron concentrations in EBC were significantly increased when compared to healthy controls [61]. Cadmium, lead, and aluminum levels in EBC were higher among smokers and smokers with chronic obstructive pulmonary disease (COPD) [62]. When patients with COPD were subdivided into smokers versus ex-smokers and non-smokers, the smokers exhibited higher EBC lead, cadmium, and aluminum levels [53]. Ex-smokers with COPD who had quit smoking for more than two years continued to demonstrate elevations in EBC metals relative to non-smokers [51]. Interestingly, reduced concentrations of iron and nickel can also be observed in the EBC of smokers [16, 53]. Finally, an investigation has demonstrated that exposure to cigarette smoke does not always impact metal concentrations in EBC [16].

6. Metals in EBC in diseased populations

The physiologic consequences of lung disease itself can potentially impact metal concentrations in EBC. Following the evaporation of water from the epithelial surfaces of the respiratory tract, the dispersal of non-volatile molecules into the water vapor can be dependent on air flow. In diseased individuals, flow can be significantly diminished. Accordingly, the quantification of metal concentrations in EBC collected from diseased cohorts may have an additional limitation.

Metals in EBC samples have been measured in patients diagnosed with asthma, COPD, and interstitial lung disease. Among 50 healthy subjects and 30 asthmatics, the measurement of metal concentrations supported lower iron levels in the group of asthmatics [53]. There were no differences in the concentrations

of lead, aluminum, cadmium, copper, and manganese. Comparable results were observed in EBC collected from 22 healthy children and 17 asthmatics, with the latter having a statistically significantly lower iron concentration [56]. Another study with a small number of participants observed no significant differences in EBC iron concentration between asthmatics ($n = 10$) and non-asthmatics ($n = 16$) [63]. Using a 'bleomycin technique for measurement of pro-oxidant iron', there was an increase in EBC iron levels post-exposure to city environments among severe asthmatics [61]. These studies do not clearly define a participation of metals in the pathogenesis of asthma but do suggest that further investigation into the role of iron in the induction of asthma is warranted.

EBC obtained from patients with stable COPD ($n = 50$) revealed higher concentrations of lead, cadmium, and aluminum, and lower levels of iron and copper relative to samples collected from healthy subjects ($n = 50$) and healthy smokers ($n = 30$) [53]. No correlations were observed between indices of pulmonary function and EBC metal concentrations. When the COPD patients were classified on the basis of disease severity, associations between the differences in EBC metals were not demonstrated. Decreased EBC iron levels in COPD patients, relative to smokers with no obstruction, were attributed to a failure of the diseased lung to excrete iron [64]. EBC collected in 28 COPD patients presenting mild to moderate exacerbation revealed increased manganese concentrations relative to samples at recovery [65]. Based on this investigation, it is uncertain if metals participate in the etiology or contribute to the exacerbation of COPD. Further studies regarding an association between metals and COPD are needed, particularly with a focus on iron and copper.

Levels of metals were quantified in EBC from patients with interstitial lung diseases [55]. Among patients with sarcoid ($n = 22$), non-specific interstitial pneumonia ($n = 15$), and idiopathic pulmonary fibrosis ($n = 19$), concentrations of chromium and nickel could be increased relative to levels observed among healthy subjects ($n = 33$). Elevated EBC concentrations of chromium and nickel among the interstitial lung patients disagreed with observations in the COPD patients, implying that the two groups were different, despite smoking being a major risk factor for both. In contrast, both EBC iron and copper were decreased in the groups with interstitial lung disease. The observed decrements in the metal concentrations in EBC could reflect a reaction comparable to the hypoferremic response observed in serum with inflammation.

7. Metals in EBC in occupational and environmental studies

The analysis of EBC endpoints has been described as one of the most promising methods available for the study of pulmonary biomarkers of exposure, effect, and susceptibility in occupational settings [59]. Metals in EBC have been measured as biomarkers to evaluate occupational exposure, including those among welders and workers in the hard metal, chrome plating, lead processing, and aluminum production industries.

Samples of EBC from 45 welders showed elevated levels of aluminum, nickel, and chromium relative to 24 non-exposed control subjects [66]. EBC from welders revealed high iron and nickel concentrations [23]. Concentrations of manganese and nickel in EBC were significantly higher among 17 welders compared to 16 unexposed control subjects after five days' exposure [14]. Welders showed significantly higher concentrations of iron and nickel in EBC relative to non-exposed volunteers [67]. Dissimilar working conditions between different companies impacted elevations in iron and nickel concentrations in the EBC collected from 36 welders [16].

A second occupation with metal exposure which has been investigated employing EBC endpoints is the hard metal industry. Thirty-three workers in workshops producing either diamond tools or hard-metal mechanical parts showed detectable cobalt levels in EBC while tungsten was undetectable [22]. In contrast, EBC concentrations of cobalt and tungsten were reported to be measurable but not significantly elevated among 62 workers at a hard metal processing plant [68].

Metal concentrations in EBC have been utilized as endpoints in studies of several other industries, including chrome plating, lead processing, and aluminum production. EBC from groups of 10 and 24 chrome platers supported measurable chromium levels [52, 69]. Chromium levels measured in EBC correlated with those measured in the red blood cells among 14 non-smoking, male chrome-platers [70]. EBC samples collected from a cohort of 58 workers occupationally exposed to chromium compounds and 22 unexposed volunteers showed significantly higher levels of chromium in the former [58]. In a group of workers from two lead processing plants, lead concentrations in EBC reflected the levels of the metal in the work environment settings [71]. Among workers in an aluminum production plant, EBC concentrations of beryllium and aluminum were higher in pot room workers when compared with the controls [24].

Metal concentrations have also been quantified in EBC collected from workers (1) exposed to titanium dioxide (TiO_2) and (2) at an airport. EBC collected from 20 workers exposed to TiO_2 demonstrated higher concentrations of titanium relative to 20 controls in which levels were below detectable limits [57].

Cadmium concentrations in EBC provided by those working in the airport apron area were higher relative to office workers [25].

These studies of occupationally exposed workers confirm significant exposure to specific metals. The cohorts used are small, reflect uncommon conditions or exposures, and document data that may not be applicable to larger groups. To document an occupational exposure, it may be more efficient to measure filters collected for environmental monitoring rather than to quantify EBC metal concentrations. In addition, the lung is not a passive filter but actively metabolizes and transports metals. The EBC metal concentrations should not automatically be interpreted to support participation, or lack of participation, in an observed biological effect.

The measurement of metals in samples, including EBC, from occupationally and environmentally exposed individuals, supports some metal import by the lung with systemic translocation, including blood and urine [70]. This movement of metal from the lung to systemic sites is comparable to that described following metal exposure in humans and other animals [11, 72]. The amount of metal transported appears dependent on (1) the magnitude of the exposure and (2) time with some elevations in the blood persisting for days.

8. Conclusion and recommendations

The results of the investigation demonstrate that transition and post-transition metals can be detectable in EBC. Currently, the measurement of metal concentrations in EBC is a research tool in the investigation of the participation of metals in the pathogenesis of disease and injury following exposure. Measurements of EBC metals may be especially valuable in research to define associations between human disease and smoking. However, the considerable dilution of the EBC sample currently limits which metals can be accurately quantified; those which are normally measurable currently include iron, zinc, and copper. Individuals with a specific history of occupational and environmental exposure to metal-abundant particles are the exception, with additional metals being measurable. Furthermore, the failure to measure and address the variability in the dilution of airway aerosols in EBC poses limitations to analysis. The measurement of EBC metal concentrations is also a potentially effective research technique in delineating the translocation of metals from the lung.

The quantification of EBC metal concentrations is currently not reliable for clinical use in either supporting or determining any diagnosis. Issues that must be addressed prior to the increased use of EBC metal measurements include the establishment of both standardized collection and measurement techniques. After this is achieved, reference values for metals in

EBC collected from healthy normal populations can be compiled. Regardless of the current limitations of the methodological approaches, metals have been repeatedly demonstrated to potentially impact numerous different lung diseases and injuries. The procurement of EBC from both healthy subjects and diseased individuals with subsequent measurement of metals is strongly advocated in research efforts. Reports of patient cohorts with such a quantification will undoubtedly contribute to defining the basis for various lung diseases and injuries.

9. Disclaimer

This report has been reviewed by the National Health and Environmental Effects Research Laboratory, United States Environmental Protection Agency, and approved for publication. Approval does not signify that the contents necessarily reflect the views and policies of the Agency, nor does the mention of trade names or commercial products constitute endorsement or recommendations for use.

ORCID iDs

Andrew J Ghio  <https://orcid.org/0000-0003-0502-6448>

References

- [1] Hasleton P S 1972 The internal surface area of the adult human lung *J. Anat.* **112** 391–400
- [2] Rosias P 2012 Methodological aspects of exhaled breath condensate collection and analysis *J. Breath Res.* **6** 027102
- [3] Johnson G R and Morawska L 2009 The mechanism of breath aerosol formation *J. Aerosol. Med. Pulm. Drug. Deliv.* **22** 229–37
- [4] Wang X, Ghio A J, Yang F, Dolan K G, Garrick M D and Piantadosi C A 2002 Iron uptake and Nramp2/DMT1/DCT1 in human bronchial epithelial cells *Am. J. Physiol. Lung Cell Mol. Physiol.* **282** L987–95
- [5] Heilig E A, Thompson K J, Molina R M, Ivanov A R, Brain J D and Wessling-Resnick M 2006 Manganese and iron transport across pulmonary epithelium *Am. J. Physiol. Lung Cell Mol. Physiol.* **290** L1247–59
- [6] Yang F, Haile D J, Wang X, Dailey L A, Stonehuerner J G and Ghio A J 2005 Apical location of ferroportin 1 in airway epithelia and its role in iron detoxification in the lung *Am. J. Physiol. Lung Cell Mol. Physiol.* **289** L14–23
- [7] Nairz M, Theurl I, Swirski F K and Weiss G 2017 ‘Pumping iron’—how macrophages handle iron at the systemic, microenvironmental, and cellular levels *Pflugers Arch.* **469** 397–418
- [8] Kreyling W G, Godleski J J, Kariya S T, Rose R M and Brain J D 1990 *In vitro* dissolution of uniform cobalt oxide particles by human and canine alveolar macrophages *Am. J. Respir. Cell Mol. Biol.* **2** 413–22
- [9] Corhay J L, Weber G, Bury T, Mariz S, Roelandts I and Radermecker M F 1992 Iron content in human alveolar macrophages *Eur. Respir. J.* **5** 804–9
- [10] Puxeddu E *et al* 2014 Iron laden macrophages in idiopathic pulmonary fibrosis: the telltale of occult alveolar hemorrhage? *Pulm. Pharmacol. Ther.* **28** 35–40
- [11] Wallenborn J G, Kovalcik K D, McGee J K, Landis M S and Kodavanti U P Systemic translocation of (70)zinc: kinetics

- following intratracheal instillation in rats *Toxicol. Appl. Pharmacol.* 2009 **234** 25–32
- [12] Wallenborn J G, McGee J K, Schladweiler M C, Ledbetter A D and Kodavanti U P 2007 Systemic translocation of particulate matter-associated metals following a single intratracheal instillation in rats *Toxicol. Sci.* **98** 231–9
- [13] American Thoracic S European Respiratory S 2005 ATS/ERS recommendations for standardized procedures for the online and offline measurement of exhaled lower respiratory nitric oxide and nasal nitric oxide, 2005 *Am. J. Respir. Crit. Care Med.* **171** 912–30
- [14] Hulo S *et al* 2014 Manganese in exhaled breath condensate: a new marker of exposure to welding fumes *Toxicol. Lett.* **226** 63–9
- [15] Winters B R, Pleil J D, Angrish M M, Stiegel M A, Risby T H and Madden M C 2017 Standardization of the collection of exhaled breath condensate and exhaled breath aerosol using a feedback regulated sampling device *J. Breath Res.* **11** 047107
- [16] Hoffmeyer F *et al* 2011 Increased metal concentrations in exhaled breath condensate of industrial welders *J. Environ. Monit.* **13** 212–8
- [17] Fox J R *et al* 2013 Characterization of a portable method for the collection of exhaled breath condensate and subsequent analysis of metal content *Environ. Sci. Process Impacts* **15** 721–9
- [18] Schroeder W H, Dobson M, Kane D M and Johnson N D 1987 Toxic trace elements associated with airborne particulate matter: a review *JAPCA, J. Air Waste Manage. Assoc.* **37** 1267–85
- [19] Li R, Yang X, Fu H, Hu Q, Zhang L and Chen J 2017 Characterization of typical metal particles during haze episodes in Shanghai, China *Chemosphere* **181** 259–69
- [20] Bolann B J, Rahil-Khazen R, Henriksen H, Isrenn R and Ulvik R J 2007 Evaluation of methods for trace-element determination with emphasis on their usability in the clinical routine laboratory *Scand. J. Clin. Lab. Invest.* **67** 353–66
- [21] Felix P M *et al* 2015 The suitability of EBC-Pb as a new biomarker to assess occupational exposure to lead *Int. J. Environ. Health Res.* **25** 67–80
- [22] Goldoni M *et al* 2004 Exhaled breath condensate as a suitable matrix to assess lung dose and effects in workers exposed to cobalt and tungsten *Environ. Health Perspect.* **112** 1293–8
- [23] Hoffmeyer F *et al* 2012 Relation between biomarkers in exhaled breath condensate and internal exposure to metals from gas metal arc welding *J. Breath Res.* **6** 027105
- [24] Hulo S *et al* 2016 Beryllium in exhaled breath condensate as a biomarker of occupational exposure in a primary aluminum production plant *Int. J. Hyg. Environ. Health* **219** 40–7
- [25] Marie-Desvergne C *et al* 2016 Assessment of nanoparticles and metal exposure of airport workers using exhaled breath condensate *J. Breath Res.* **10** 036006
- [26] Rennard S I *et al* 1986 Estimation of volume of epithelial lining fluid recovered by lavage using urea as marker of dilution *J. Appl. Physiol.* **60** 532–8
- [27] Horvath I *et al* 2005 Exhaled breath condensate: methodological recommendations and unresolved questions *Eur. Respir. J.* **26** 523–48
- [28] Society A T 1989 Proteins in bronchoalveolar lavage fluid *Am. J. Respir. Crit. Care Med.* **141** S183–8
- [29] Bloemen K, Lissens G, Desager K and Schoeters G 2007 Determinants of variability of protein content, volume and pH of exhaled breath condensate *Respir. Med.* **101** 1331–7
- [30] Effros R M *et al* 2003 A simple method for estimating respiratory solute dilution in exhaled breath condensates *Am. J. Respir. Crit. Care Med.* **168** 1500–5
- [31] Esther C R Jr *et al* 2009 Mass spectrometric analysis of biomarkers and dilution markers in exhaled breath condensate reveals elevated purines in asthma and cystic fibrosis *Am. J. Physiol. Lung Cell Mol. Physiol.* **296** L987–93
- [32] Effros R M *et al* 2002 Dilution of respiratory solutes in exhaled condensates *Am. J. Respir. Crit. Care Med.* **165** 663–9
- [33] Syslova K *et al* 2010 LC-ESI-MS/MS method for oxidative stress multimarker screening in the exhaled breath condensate of asbestosis/silicosis patients *J. Breath Res.* **4** 017104
- [34] Effros R M, Dunning M B 3rd, Biller J and Shaker R 2004 The promise and perils of exhaled breath condensates *Am. J. Physiol. Lung Cell Mol. Physiol.* **287** L1073–80
- [35] Sabbioni E, Pietra R, Mousty F, Colombo F, Rizzato G and Scansetti G 1987 Trace-metals in human-lung as determined by neutron-activation analysis of bronchoalveolar lavage *J. Radioanal. Nucl. Chem. Artic.* **110** 595–601
- [36] Romeo L, Maranelli G, Malesani F, Tommasi I, Cazzadori A and Graziani M S 1992 Tentative reference values for some elements in bronchoalveolar lavage fluid *Sci. Total Environ.* **120** 103–10
- [37] Corhay J L *et al* 1995 Nonfibrous mineralogical analysis of bronchoalveolar lavage fluid from blast-furnace workers *Arch. Environ. Health* **50** 312–9
- [38] Nelson M E, O'Brien-Ladner A R and Wesselius L J 1996 Regional variation in iron and iron-binding proteins within the lungs of smokers *Am. J. Respir. Crit. Care Med.* **153** 1353–8
- [39] Harlyk C, McCourt J, Bordin G, Rodriguez A R and van der Eeckhout A 1997 Determination of copper, zinc and iron in broncho-alveolar lavages by atomic absorption spectroscopy *J. Trace Elem. Med. Biol.* **11** 137–42
- [40] Ghio A J, Carter J D, Richards J H, Brighton L E, Lay J C and Devlin R B 1998 Disruption of normal iron homeostasis after bronchial instillation of an iron-containing particle *Am. J. Physiol.* **274** L396–403
- [41] Stites S W, Plautz M W, Bailey K, O'Brien-Ladner A R and Wesselius L J 1999 Increased concentrations of iron and isoferitins in the lower respiratory tract of patients with stable cystic fibrosis *Am. J. Respir. Crit. Care Med.* **160** 796–801
- [42] Ghio A J, Carter J D, Richards J H, Richer L D, Grissom C K and Elstad M 2003 Iron and iron-related proteins in the lower respiratory tract of patients with acute respiratory distress syndrome *Crit. Care Med.* **31** 395–400
- [43] Bargagli E, Monaci F, Bianchi N, Bucci C and Rottoli P 2008 Analysis of trace elements in bronchoalveolar lavage of patients with diffuse lung diseases *Biol. Trace Elem. Res.* **124** 225–35
- [44] Ghio A J, Roggli V L, Soukup J M, Richards J H, Randell S H and Muhlebach M S 2013 Iron accumulates in the lavage and explanted lungs of cystic fibrosis patients *J. Cyst. Fibros.* **12** 390–8
- [45] Kollmeier H, Seemann J W, Rothe G, Muller K M and Wittig P 1990 Age, sex, and region adjusted concentrations of chromium and nickel in lung tissue *Br. J. Ind. Med.* **47** 682–7
- [46] Ghio A J, Pritchard R J, Dittrich K L and Samet J M 1997 Non-heme (Fe³⁺) in the lung increases with age in both humans and rats *J. Lab. Clin. Med.* **129** 53–61
- [47] Kim Y, Lobdell D T, Wright C W, Gocheva V V, Hudgens E and Bowler R M 2015 Blood metal concentrations of manganese, lead, and cadmium in relation to serum ferritin levels in Ohio residents *Biol. Trace Elem. Res.* **165** 1–9
- [48] Kim H J *et al* 2017 Determination of trace metal levels in the general population of Korea *Int. J. Environ. Res. Public Health* **29** E702
- [49] Jantzen C, Jorgensen H L, Duus B R, Sparring S L and Lauritzen J B 2013 Chromium and cobalt ion concentrations in blood and serum following various types of metal-on-metal hip arthroplasties: a literature overview *Acta Orthop.* **84** 229–36
- [50] Blazewicz A, Klatka M, Astel A, Partyka M and Kocjan R 2013 Differences in trace metal concentrations (Co, Cu, Fe, Mn, Zn, Cd, and Ni) in whole blood, plasma, and urine of obese and nonobese children *Biol. Trace Elem. Res.* **155** 190–200
- [51] Mutti A and Corradi M 2006 Recent developments in human biomonitoring: non-invasive assessment of target tissue dose and effects of pneumotoxic metals *Med. Lav.* **97** 199–206
- [52] Caglieri A *et al* 2006 The effect of inhaled chromium on different exhaled breath condensate biomarkers among chrome-plating workers *Environ. Health Perspect* **114** 542–6
- [53] Mutti A, Corradi M, Goldoni M, Vettori M V, Bernard A and Apostoli P 2006 Exhaled metallic elements and serum

- pneumoproteins in asymptomatic smokers and patients with COPD or asthma *Chest* **129** 1288–97
- [54] Goldoni M *et al* 2008 Chromium in exhaled breath condensate and pulmonary tissue of non-small cell lung cancer patients *Int. Arch. Occup. Environ. Health* **81** 487–93
- [55] Corradi M *et al* 2009 Metallic elements in exhaled breath condensate of patients with interstitial lung diseases *J. Breath Res.* **3** 046003
- [56] Vlasic Z *et al* 2009 Iron and ferritin concentrations in exhaled breath condensate of children with asthma *J. Asthma* **46** 81–5
- [57] Pelclova D *et al* 2015 Raman microspectroscopy of exhaled breath condensate and urine in workers exposed to fine and nano TiO₂ particles: a cross-sectional study *J. Breath Res.* **9** 036008
- [58] Leese E, Morton J, Gardiner P H E and Carolan V A 2017 The simultaneous detection of trivalent & hexavalent chromium in exhaled breath condensate: A feasibility study comparing workers and controls *Int. J. Hyg. Environ. Health* **220** 415–23
- [59] Corradi M, Gergelova P and Mutti A 2010 Use of exhaled breath condensate to investigate occupational lung diseases *Curr. Opin. Allergy Clin. Immunol.* **10** 93–8
- [60] Chiba M and Masironi R 1992 Toxic and trace elements in tobacco and tobacco smoke *Bull. World Health Organ.* **70** 269–75
- [61] Mumby S, Chung K F, McCreanor J E, Moloney E D, Griffiths M J and Quinlan G J 2011 Pro-oxidant iron in exhaled breath condensate: a potential excretory mechanism *Respir. Med.* **105** 1290–5
- [62] Rustemeier K, Stabbert R, Haussmann H J, Roemer E and Carmines E L 2002 Evaluation of the potential effects of ingredients added to cigarettes. Part 2: chemical composition of mainstream smoke *Food Chem. Toxicol.* **40** 93–104
- [63] Maloca Vuljanko I, Turkalj M, Nogalo B, Bulat Lokas S and Plavec D 2017 Diagnostic value of a pattern of exhaled breath condensate biomarkers in asthmatic children *Allergol. Immunopathol. (Madr.)* **45** 2–10
- [64] Mumby S, Saito J, Adcock I M, Chung K F and Quinlan G J 2016 Decreased breath excretion of redox active iron in COPD: a protective failure? *Eur. Respir. J.* **47** 1267–70
- [65] Corradi M *et al* 2009 Metallic elements in exhaled breath condensate and serum of patients with exacerbation of chronic obstructive pulmonary disease *Metallomics* **1** 339–45
- [66] Gube M *et al* 2010 Biological effect markers in exhaled breath condensate and biomonitoring in welders: impact of smoking and protection equipment *Int. Arch. Occup. Environ. Health* **83** 803–11
- [67] Hartmann L *et al* 2014 Assessment of the biological effects of welding fumes emitted from metal inert gas welding processes of aluminium and zinc-plated materials in humans *Int. J. Hyg. Environ. Health* **217** 160–8
- [68] Broding H C, Michalke B, Goen T and Drexler H Comparison between exhaled breath condensate analysis as a marker for cobalt and tungsten exposure and biomonitoring in workers of a hard metal alloy processing plant *Int. Arch. Occup. Environ. Health* 2009 **82** 565–73
- [69] Goldoni M *et al* 2006 Determination of hexavalent chromium in exhaled breath condensate and environmental air among chrome plating workers *Anal. Chim. Acta* **562** 229–35
- [70] Goldoni M *et al* 2010 Chromium in exhaled breath condensate (EBC), erythrocytes, plasma and urine in the biomonitoring of chrome-plating workers exposed to soluble Cr(VI) *J. Environ. Monit.* **12** 442–7
- [71] Felix P M *et al* 2013 Biomarkers of exposure to metal dust in exhaled breath condensate: methodology optimization *Arch. Environ. Occup. Health* **68** 72–9
- [72] Chamberlain A C, Clough W S, Heard M J, Newton D, Stott A N and Wells A C 1975 Uptake of inhaled lead from motor exhaust *Postgrad. Med. J.* **51** 790–4