

Modulation of GLP-1 Levels by a Genetic Variant That Regulates the Cardiovascular Effects of Intensive Glycemic Control in ACCORD

Diabetes Care 2018;41:348–355 | <https://doi.org/10.2337/dc17-1638>

Hetal S. Shah,^{1,2} Mario Luca Morieri,^{1,2} Santica M. Marcovina,³ Ronald J. Sigal,⁴ Hertz C. Gerstein,⁵ Michael J. Wagner,⁶ Alison A. Motsinger-Reif,⁷ John B. Buse,⁸ Peter Kraft,⁹ Josyf C. Mychaleckyj,¹⁰ and Alessandro Doria^{1,2}

OBJECTIVE

A genome-wide association study in the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial identified two markers (rs57922 and rs9299870) that were significantly associated with cardiovascular mortality during intensive glycemic control and could potentially be used, when combined into a genetic risk score (GRS), to identify patients with diabetes likely to derive benefit from intensive control rather than harm. The aim of this study was to gain insights into the pathways involved in the modulatory effect of these variants.

RESEARCH DESIGN AND METHODS

Fasting levels of 65 biomarkers were measured at baseline and at 12 months of follow-up in the ACCORD-Memory in Diabetes (ACCORD-MIND) MRI substudy ($n = 562$). Using linear regression models, we tested the association of the GRS with baseline and 12-month biomarker levels, and with their difference (Δ), among white subjects, with genotype data ($n = 351$) stratified by intervention arm.

RESULTS

A significant association was observed between GRS and Δ GLP-1 (glucagon-like peptide 1, active) in the intensive arm ($P = 3 \times 10^{-4}$). This effect was driven by rs57922 ($P = 5 \times 10^{-4}$). C/C homozygotes, who had been found to derive cardiovascular benefits from intensive treatment, showed a 22% increase in GLP-1 levels during follow-up. By contrast, T/T homozygotes, who had been found to experience increased cardiac mortality with intensive treatment, showed a 28% reduction in GLP-1 levels. No association between Δ GLP-1 and GRS or rs57922 was observed in the standard treatment arm.

CONCLUSIONS

Differences in GLP-1 axis activation may mediate the modulatory effect of variant rs57922 on the cardiovascular response to intensive glycemic control. These findings highlight the importance of GLP-1 as a cardioprotective factor.

With the global rise in type 2 diabetes, it is imperative to prevent its cardiovascular complications, since these are major contributors to the high mortality, morbidity, and socioeconomic burden associated with this disease (1). The Action to Control Cardiovascular Risk in Diabetes (ACCORD) randomized clinical trial aimed to study whether intensive as opposed to standard glycemic control could prevent cardiovascular disease (CVD) in type 2 diabetes (2). Despite a significant reduction in nonfatal myocardial

¹Research Division, Joslin Diabetes Center, Boston, MA

²Department of Medicine, Harvard Medical School, Boston, MA

³Department of Medicine, University of Washington, and Northwest Lipid Metabolism and Diabetes Research Laboratories, Seattle, WA

⁴Departments of Medicine, Cardiac Sciences, and Community Health Sciences, Faculties of Medicine and Kinesiology, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada

⁵Department of Medicine and the Population Health Research Institute, McMaster University, and Hamilton Health Sciences, Ontario, Hamilton, Canada

⁶Center for Pharmacogenomics and Individualized Therapy, University of North Carolina at Chapel Hill, Chapel Hill, NC

⁷Bioinformatics Research Center and Department of Statistics, North Carolina State University, Raleigh, NC

⁸Department of Medicine, University of North Carolina School of Medicine, Chapel Hill, NC

⁹Department of Epidemiology and Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA

¹⁰Center for Public Health Genomics, University of Virginia, Charlottesville, VA

Corresponding author: Alessandro Doria, alessandro.doria@joslin.harvard.edu.

Received 4 August 2017 and accepted 22 October 2017.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc17-1638/-/DC1>.

infarctions in subjects treated intensively, the trial was prematurely terminated owing to a paradoxical increase in cardiovascular mortality in the same arm (2).

Through a genome-wide association study (GWAS) on a subset of ACCORD participants, we have recently identified two genetic markers, rs9299870 on 10q26 and rs57922 on 5q13, that were associated with the excess cardiovascular mortality in the intensive glycemic arm (3). A genetic risk score (GRS) derived from these two variants significantly modulated the effect of glycemic treatment on cardiovascular outcomes. Intensive glycemic control prevented fatal and nonfatal cardiac events among subjects with a GRS = 0, whereas it increased cardiac mortality among those with a GRS ≥ 2 (3).

While the rs57922/rs9299870 GRS is a potential tool for precision medicine approaches to treat type 2 diabetes, regardless of the mechanism of its effect, understanding the pathways that are involved in this genetic modulation may provide new insights into the links between glycemic control and cardiovascular outcomes. To this end, we examined the association between the GRS and 65 biomarkers measured at baseline and at 12 months postrandomization as part of the ACCORD-Memory in Diabetes (ACCORD-MIND) MRI substudy (4). Our findings point to an unexpected link between one of the two GRS variants (rs57922) and the GLP-1 axis.

RESEARCH DESIGN AND METHODS

Study Population

In ACCORD, 10,251 individuals with type 2 diabetes and high cardiovascular risk from 77 clinical centers across the U.S. and Canada were randomized in a 1:1 ratio to intensive ($\text{HbA}_{1c} < 6.0\%$ [42 mmol/mol]) and standard ($\text{HbA}_{1c} 7\text{--}7.9\%$ [53–63 mmol/mol]) glycemic treatment, as well as to lipid and blood pressure subtrials, in a double 2×2 factorial design (2). GWAS data were generated for 8,084 participants, 5,360 of whom were self-reported whites (3). From the overall ACCORD cohort, 2,977 participants were enrolled in the ACCORD-MIND study, and 632 of these underwent MRI scans (4). ACCORD-MIND MRI subjects with available serum samples participated in an ancillary biomarker study ($n = 562$) (4). This subset did not differ significantly from the overall cohort, except for a

larger proportion of whites, females, and blood pressure subtrial participants and lower numbers with baseline CVD (Supplementary Table 1). The current study concerned 351 subjects corresponding to the overlap between ACCORD whites with GWAS data ($n = 5,360$) and the ACCORD-MIND MRI biomarker cohort ($n = 562$).

Genotyping

Genotyping and quality-control methods of the ACCORD-genetic data set have previously been described (3). Briefly, genotyping was carried out on two platforms at two sites based on the level of genetic consent. Those participants who had consented to genetic studies by any investigator were typed on an Illumina HumanOmniExpressExome-8, version 1.0, chip at the University of Virginia, and those who had consented only to studies by ACCORD investigators were genotyped on an Affymetrix Axiom-Biobank1 chip at the University of North Carolina (3). Two distinct sets emerged after various merge and quality-control procedures: one including 5,971 samples with any level of genetic consent (ANYSET), genotyped for 1,263,585 variants, and another one including 2,113 samples with only ACCORD investigator consent (ACCSET), genotyped for 572,192 variants. High-quality imputation performed by IMPUTE version 2.3.1 (Marchini Laboratory, Oxford) on both sets resulted in 24 million variants spanning the entire genome, of which 6.8 million common (minor allele frequency $\geq 5\%$) single nucleotide polymorphisms (SNPs) were tested in the GWAS (3). For the current study, posterior genotype probabilities of rs57922 and rs9299870 were extracted from the GWAS data.

Biomarker Measurement

Fasting blood samples were collected at baseline and at 12 months of follow-up (4). Serum sample aliquots (1 mL) were stored at -80°C at the ACCORD central laboratory (Northwest Lipid Metabolism and Diabetes Research Laboratories), and levels of 65 biomarkers were measured immediately on freshly thawed samples (4).

Apolipoproteins A-I and B, hs-CRP, and cystatin C were measured on a Siemens BNII nephelometer (Siemens Healthcare Diagnostics, Newark, DE) using Siemens reagents and in-house prepared calibrators and quality-control materials. The interassay coefficients of variation (CVs)

of the quality-control samples with low, medium, and high levels were consistently $< 3\%$ for all four biomarkers.

Nonesterified fatty acids were measured using Wako Diagnostics reagents (Wako Diagnostics, Richmond, VA) on a Roche Modular P autoanalyzer. The sensitivity of the method was 0.0014 mEq/L, and the linearity was 4.0 mEq/L. The interassay CVs, as determined on quality-control samples with low, medium, and high levels of nonesterified fatty acids, were 4.6%, 3.3%, and 3.3% respectively.

Total testosterone levels were determined by a Tosoh 2000 autoanalyzer (Tosoh Bioscience, Inc., San Francisco, CA). The assay sensitivity was 10 ng/dL. The interassay CVs on quality-control samples with high, medium, and low levels of testosterone were 2.48%, 2.69%, and 5.99%, respectively.

Estradiol (E-2) levels were measured by a Tosoh AIA-2000 analyzer in a competitive enzyme immunoassay. The assay was linear up to 3,000 pg/mL, and the sensitivity was 25 pg/mL. The interassay CVs on quality-control samples with high, medium, and low levels of estradiol were 4.2%, 4.9%, and 8.9%, respectively.

C-peptide levels were determined by a two-site immunoenzymometric assay using a Tosoh 2000 autoanalyzer calibrated against the World Health Organization's International Standard of 84/510. The assay had a sensitivity level of 0.05 ng/mL. The interassay CVs for low, medium, and high C-peptide control samples were 3.2%, 1.6%, and 1.8%, respectively.

Analyses for GAD-65 autoantibodies were based on a protocol provided by the National Institute of Diabetes and Digestive and Kidney Diseases Autoantibody Harmonization Committee. Based on the analysis of 550 samples provided by the Centers for Disease Control and Prevention Diabetes Autoantibody Standardization Program, positive/negative cutoff for the assay was determined to be 33 DK units. The GAD assay had 76% sensitivity and 92.2% specificity.

Analysis of glycated albumin was performed by means of the bromocresol purple method using a Lucica GA-L kit (Asahi Kasei Pharma, Tokyo, Japan) on a Roche Modular P autoanalyzer. The assay linear range was 3.2–68.1%, and the interassay CV was 3.0%.

The remaining biomarkers were measured by multiplexing kits from Millipore and Luminex using a single lot of reagent and

quality-control materials. These assays were grouped into the following panels.

1. Chemokine/cytokines: interleukins (IL-1 α , IL-1 β , IL-1ra, IL-5, IL-6, IL-8, IL-10, IL-12[p40], IL-12[p70], IL-15, and IL-17), epidermal growth factor (EGF), granulocyte-colony stimulating factor (G-CSF), interferon- γ (IFN γ), interferon-inducible protein 10 (IP-10), monocyte chemoattractant protein-1 (MCP-1), macrophage inflammatory proteins (MIP-1 α and MIP-1 β), soluble CD40 L (sCD40 L), transforming growth factor α (TGF α), tumor necrosis factor α (TNF α), vascular endothelial growth factor (VEGF), rotaxin, and fractalkine.
2. Endocrine: adiponectin, plasminogen activator inhibitor-1 (PAI-1) (active and total), resistin, hepatocyte growth factor (HGF), leptin, and nerve growth factor (NGF).
3. Metabolism: amylin (active and total), glucagon, gastric inhibitory polypeptide (GIP), GLP-1 (active), ghrelin (active), pancreatic polypeptide, and peptide YY (PYY).
4. CVD: matrix metalloproteinase-9 (MMP-9), myeloperoxidase (MPO), soluble E-selectin, soluble adhesion molecules sICAM and sVCAM, serum amyloid A, serum amyloid P, and N-terminal pro-b-type natriuretic peptide (NT-proBNP).
5. Sepsis/apoptosis: macrophage migration inhibitory factor (MIF), sFas, and sFasL.
6. Bone: osteoprotegerin (OPG) and osteocalcin.

Statistical Analysis

All statistical analyses were conducted using SAS, version 9.4 (SAS Institute, Cary, NC). Baseline characteristics of participants were described through means, medians, and proportions as appropriate. Minor allele dosages (ranging 0–2) for each of the polymorphisms, rs9299870 and rs57922, were calculated from the imputed posterior genotype probabilities for each individual, and these dosages were added up to form the GRS (scoring 0–4). Biomarkers with values above and below the limits of quantification were assigned the upper and lower limits of levels of quantification, respectively. The 65 biomarkers were log transformed for normalization of their distributions, and Δ values, were obtained by subtracting baseline from 12-month log-transformed levels.

Linear regression models tested the association of the GRS in each treatment arm with baseline and 12-month biomarker levels and their difference (Δ), adjusted by blood pressure and lipid subtrials, clinical center network, and genotyping set (ACCSET or ANYSET). In a principal component analysis including 65 biomarker measurements at baseline, 65 measurements at 12 months, and 65 Δ values, 100% of the variance of these 195 variables (65 \times 3) was explained by 70 principal components. On this basis, a Bonferroni-adjusted P value threshold of 3.6×10^{-4} (0.05/[70 \times 2 treatment arms]) was set as a significance threshold for our primary analysis. As the effect of GRS was significant for Δ GLP-1, the modulatory effects of polymorphisms and treatment on GLP-1 levels were evaluated by estimating least squares (LS) means of baseline, 12-month, and Δ GLP-1 levels within genotype and treatment groups. Further sensitivity analyses included adjustments for baseline CVD, age, sex, HbA_{1c}, fasting plasma glucose, heart rate, BMI, and glomerular filtration rate (GFR).

The associations between SNP rs57922 and the expression of selected genes suggested by the literature to be involved in GLP-1 synthesis or secretion (5–9) were tested using RNA-Seq data from up to 570 donors in the online Genotype-Tissue Expression (GTEx) database v6p (The Broad Institute of MIT and Harvard, Cambridge, MA). This publicly available database allows the identification of potential expression quantitative trait loci (eQTLs), that is, genetic variants affecting gene expression in different tissues. Random effects meta-analyses of rs57922 eQTLs across multiple tissues were conducted for each gene using Metasoft, version 2.0.1 (10).

RESULTS

Baseline characteristics of the ACCORD participants included in this study did not significantly differ between intensive and standard glycemic arms (Table 1). Supplementary Table 2 shows the results of the analysis for association between the rs57922/rs9299870 GRS, which had been found to be associated with cardiovascular mortality in the intensive arm, and 65 biomarkers measured at baseline and 12 months after randomization. No significant association was observed for any of the biomarkers at either time point in either treatment arm. However, when

biomarkers data were expressed as the difference (Δ) between 12-month and baseline levels, a highly significant association was observed in the intensive treatment arm for active GLP-1, with each GRS unit increment being associated with a 22% decrease in the change in active GLP-1 levels from baseline to 12 months ($P = 3 \times 10^{-4}$) (Table 2). No association was observed in the standard arm, resulting into a GRS \times treatment interaction P value of 0.016 (Table 2).

As shown in Figs. 1 and 2, the GRS effect on active GLP-1 levels was mostly driven by SNP rs57922. No significant differences were observed among rs57922 genotypes in active GLP-1 levels at baseline in either treatment arm (Fig. 1A). In the intensive arm, active GLP-1 levels increased from baseline to 12 months by 22% (95% CI 1–46) among C/C homozygotes, were unmodified in C/T heterozygotes, and decreased by 28% (95% CI 10–41) in T/T homozygotes ($P = 5 \times 10^{-4}$) (Fig. 2A). By contrast, no significant changes in active GLP-1 levels from baseline to 12 months were observed in the standard treatment arm for any of the genotype groups. Carriers of the rs9299870 minor (G) allele showed an overall tendency to have higher active GLP-1 at baseline (Fig. 1B), but no significant differences were observed in the change from baseline to 12 months (Fig. 2B) in any of the treatment/genotype groups. Adjustments for baseline CVD, diabetes duration, HbA_{1c}, fasting plasma glucose, heart rate, smoking, age, and sex did not alter these results (Supplementary Table 3). No significant correlations were observed between Δ GLP-1 levels and Δ HbA_{1c} ($r^2 = 0.01$) or between Δ GLP-1 and Δ fasting plasma glucose ($r^2 = 0.02$) within rs57922 genotypes in the intensive arm. In subgroup analyses, the association between active GLP-1 level changes and rs57922 genotype observed in the intensive arm was most pronounced among males ≥ 61 years and among individuals with baseline BMI < 35 kg/m² or BMI change in the upper tertile (Supplementary Tables 4 and 5).

There were only nine subjects on the GLP-1 analog exenatide during the 12 months of follow-up and none on dipeptidyl peptidase-4 (DPP-IV) inhibitors. Repeating the analysis by excluding the nine subjects on exenatide did not alter results. With regard to other antihyperglycemic treatment modalities, no statistical

Table 1—Baseline characteristics of white participants of the ACCORD-MIND MRI biomarker cohort (N = 351) within intensive and standard glycemc arms

Characteristic*	Intensive arm (N = 162)	Standard arm (N = 189)
Female sex, n (%)	64 (39.5)	84 (44.4)
Age, years, mean (SD)	62.3 (5.7)	63.3 (5.8)
Diabetes duration, years, median (IQR)	8.8 (5.0–11.0)	9.7 (5.0–13.0)
Previous cardiovascular event, n (%)	44 (27.2)	49 (25.9)
Current smoker, n (%)	16 (9.9)	22 (11.6)
HbA _{1c} , %		
Mean (SD)	8.2 (1.0)	8.0 (0.9)
Median (IQR)	8.0 (7.6–8.7)	7.9 (7.5–8.5)
Fasting serum glucose, mg/dL, mean (SD)	179.4 (52.6)	174.6 (46.5)
BMI, kg/m ² , mean (SD)	33.9 (4.7)	32.8 (4.7)
Waist circumference, cm, mean (SD)	111.2 (12.7)	107.9 (12.3)
Blood pressure, mmHg, mean (SD)		
Systolic	134.2 (16.3)	134.8 (17.4)
Diastolic	74.7 (9.5)	73.7 (9.7)
Serum creatinine, mg/dL, mean (SD)	0.9 (0.2)	0.9 (0.2)
eGFR (from MDRD), mL/min/1.73 m ² , mean (SD)	86.3 (20.0)	87.5 (19.6)
Lipids, mg/dL, mean (SD)		
Total cholesterol	182.7 (38.3)	184.9 (42.7)
LDL	100.0 (31.3)	101.4 (33.4)
HDL (women)	47.3 (12.1)	47.2 (11.4)
HDL (men)	38.1 (9.1)	39.9 (9.1)
Triglycerides	214.7 (129.7)	208.2 (124.2)
Blood pressure trial, n (%)		
Standard	102 (63.0)	116 (61.4)
Intensive	48 (29.6)	51 (26.9)
Lipid trial, n (%)		
Statin + placebo	54 (33.3)	65 (34.4)
Statin + fibrate	60 (37.0)	73 (38.6)
Statin + placebo	33 (20.4)	31 (16.4)
Statin + fibrate	27 (16.7)	42 (22.2)

eGFR, estimated glomerular filtration rate; IQR, interquartile range. *There were no significant ($P < 0.05$) differences between the two groups.

differences were observed in drug distributions across genotypes (Supplementary Table 6). Accordingly, adjustment for the different modalities yielded no or minimal attenuations of the effect of the SNP on GLP-1 change (Supplementary Table 7). Also, there was no evidence of interaction between the SNP and any of the treatment modalities (data not shown).

In a meta-analysis of multitissue eQTL data in GTEx, rs57922 was associated (random effects P values < 0.05) with the expression of four out of twenty genes selected for their involvement in the synthesis, secretion, and/or processing of GLP-1 (binomial $P = 0.01$) (Supplementary Table 8). Significant genes included *LEPR* (leptin receptor) ($P = 0.0007$), *GRPR* (gastrin-releasing peptide receptor) ($P = 0.0009$),

SLC5A1 (sodium/glucose cotransporter 1) ($P = 0.0039$), and *SLC2A5* (glucose transporter 5) ($P = 0.03$). In all four cases, the rs57922 T allele, which in ACCORD was associated with a decrease in GLP-1 levels during intensive treatment, was associated with lower expression levels of these genes within the terminal ileum and other tissues, with no evidence of significant heterogeneity across tissues. *cis*-eQTL analyses in intestinal tissue for genes located 2 MB upstream and downstream of rs57922 did not reveal any significant association with the expression of transcription factors that could explain the *trans* effects on *LEPR*, *GRPR*, or *GLUTs* (data not shown). However, specific expression of these genes within intestinal L cells could not be examined owing to lack of this data in the GTEx database.

CONCLUSIONS

Intensive glycemc control has been proposed as a strategy for preventing CVD in type 2 diabetes. However, this approach has remained controversial owing to the small overall benefit of this intervention and the suspicion, raised by ACCORD, that it may cause an increased risk of cardiovascular death. We recently identified two genetic variants that can potentially be used to identify a subset of subjects with type 2 diabetes who may derive greater benefit from intensive glycemc control without being exposed to its risks (3). Following up on those findings, we have now discovered a possible pathway that could mediate the modulatory effect of one of those variants (rs57922). Specifically, we have found that homozygotes for allele C of these variants, that is, those individuals who derived benefit from intensive glycemc control, responded to this intervention with an increase in fasting levels of active GLP-1 at 12 months from randomization. By contrast, T/T homozygotes, that is, those subjects who showed increased cardiovascular mortality in response to intensive glycemc control, were characterized by a significant reduction in 12-month active GLP-1 levels. While in the limited sample at our disposal we could not test the association of GLP-1 changes with cardiovascular mortality, one can hypothesize that the observed difference in GLP-1 levels may have contributed to the stark difference between the two genotype groups in the cardiovascular response to intensive glycemc control.

Table 2—Association of GRS with baseline and 12-month GLP-1 levels and with Δ GLP-1 levels in ACCORD intensive and standard arms

	Intensive glycemc arm		Standard glycemc arm		GRS \times treatment interaction P
	Fold change (95% CI)*	P	Fold change (95% CI)*	P	
Baseline GLP-1	1.15 (0.93–1.42)	0.193	1.03 (0.84–1.25)	0.810	0.457
12-month GLP-1	0.86 (0.70–1.06)	0.161	1.03 (0.85–1.24)	0.778	0.227
Δ GLP-1	0.78 (0.68–0.89)	0.0003‡	0.99 (0.86–1.15)	0.935	0.016

*Fold change of GLP-1 levels per unit of GRS obtained from linear regression models testing association of GRS with log-transformed active GLP-1 levels at baseline and 12 months, and the difference between the two, adjusted by clinical center networks and source of genetic data (ACCSET or ANYSET). ‡Significant at $P < 0.00036$ (Bonferroni adjusted).





Figure 1—A and B: Baseline and 12-month GLP-1 levels within glycaemic treatment arms and genotypes of rs57922 and rs9299870. LS means of GLP-1 obtained from model adjusted by trial covariates, clinical center network, and source of genetic data. P values are obtained from generalized linear regression for association with baseline or 12-month GLP-1 levels, using additive model of SNP, and adjusted for trial covariates, clinical center network, and source of genetic data. 12mths, 12 months.

How the interaction of intensive glycaemic control and the rs57922 variant may affect active GLP-1 plasma levels is unclear at this time. GLP-1 is secreted by

intestinal L cells after being cleaved from a precursor (preproglucagon) by the action of the enzyme prohormone convertase 1. Once secreted, active GLP-

1 is quickly inactivated by cleavage of its NH₂ terminus by the enzyme DPP-IV. SNP rs57922 is not placed in or within the vicinity of the genes coding for




Figure 2—A and B: Mean change in GLP-1 from baseline to 12 months within glycemic treatment arms and rs57922 genotypes. Twelve-month-to-baseline GLP-1 ratio derived from the difference (Δ) between the log-transformed baseline and 12-month GLP-1 levels; here, presented within intensive/standard glycemic treatment arms, are LS means of this ratio (from model adjusted for trial covariates, clinical center network, and source of genetic data) within genotypes of rs57922 on 5q13 (A) and rs9299870 on 10q23 (B). P values are obtained from generalized linear regression for association with Δ GLP-1 levels, using additive model of SNP, and adjusted for trial covariates, clinical center network, and source of genetic data. 12mth, 12-month.

preproglucagon (*GCG*), prohormone convertase 1 (*PCSK1*), and *DPP-IV* and was not found to be associated with the expression of those three genes in any of the tissues in the GTEx database. We found, however, an association of the variant with the expression of genes coding for proteins involved in the stimulation of GLP-1 secretion by L cells such as the receptor of gastrin-releasing peptide (a GLP-1 secretagogue) and two hexose transporters (sodium/glucose cotransporter 1 and glucose transporter 5) mediating the sensing of nutrients in the intestinal lumen by L cells. The variant was also associated with expression of the gene coding for leptin receptor, to which the cytokine leptin

binds, thereby inducing GLP-1 release via STAT (signal transducer and activator of transcription) phosphorylation (6,11). Since the allele associated with decreased GLP-1 levels during intensive glycemic control was associated with lower expression of these genes and vice versa, this appears to be a conceivable mechanism through which the variant may affect the GLP-1 response. Though *cis*-eQTL analyses in intestinal tissue could not explain the *trans* effects on LEPR, GRPR, or GLUTs, it should be considered that rs57922 is in close proximity to a number of long noncoding RNAs (LINC)s, which, in theory, could be responsible for *trans* effects. Unfortunately, this hypothesis cannot be readily

explored, as data on the expression of these long noncoding RNAs are not included in the GTEx database. Hence, whether the variant affects the expression of these genes directly through a long-range *trans* effect or indirectly by affecting other genes placed in its vicinity remains unclear at this time, as do the mechanisms of the interaction with intensive glycemic control. With regard to the latter, we did not find any difference among genotypes in the prevalence of glucose-lowering treatments that may influence GLP-1 levels such as metformin (12). None of the participants were on DPP-IV inhibitors, and only handfuls were on GLP-1 agonists during the 12 months of follow-up. We cannot exclude, however, that some drugs were used at a higher dosage in the intensive arm and that this was the feature having a permissive influence on the GLP-1 effects of the gene expression changes induced by the SNP. Unfortunately, we cannot explore this hypothesis, as we do not have access to individual level data about the dosage of each drug. Differences in HbA_{1c} or fasting plasma glucose levels do not seem to account for the interaction either. Given the evidence suggesting that intestinal microbiota can regulate GLP-1 secretion (13), one can also postulate a role of differences in microbiome induced by the intensive therapy, but again we do not have data to test this hypothesis.

In terms of mechanisms through which GLP-1 levels can be related to cardiovascular outcomes, it is well-known that GLP-1, in addition to its actions on glucose metabolism, has anti-inflammatory and antioxidative effects in cell types relevant to atherogenesis as well as direct cardioprotective effects (14–16). Consistent with these findings, treatment with the GLP-1 agonists liraglutide and semaglutide have been reported to reduce the incidence of major cardiovascular events (death from cardiovascular causes, nonfatal myocardial infarction, and nonfatal stroke) in individuals with type 2 diabetes independent of the effects of these drugs on glycemic control (17–19). Furthermore, by playing a regulatory role in the hypothalamic-pituitary-adrenal axis, GLP-1 modulates neuroendocrine and autonomic responses to acute and chronic stress, which may influence the ability of a subject to survive a harmful exposure (20–22). One must consider, however, that these anti-inflammatory,

antiatherogenic, and antistress effects have been observed with pharmacological doses of native GLP-1 or GLP-1 agonists (17,18,23–25) and/or in animal models with homozygous disruption of the GLP-1 receptor (20,21,26). Whether the relatively small changes in fasting GLP-1 levels that we observed in response to intensive glycemic control in carriers of specific rs57922 genotypes are sufficient to affect cardiovascular risk remains to be determined. A report from Sweden has described an association between fasting GLP-1 levels and left ventricular diastolic function in elderly men (27), but the same group failed to demonstrate an association between low GLP-1 levels and coronary artery disease (28). Further studies considering both fasting and postprandial GLP-1 concentrations in larger populations are necessary to clarify the role of nonpharmacological levels of GLP-1 activity in the modulation of cardiovascular risk.

Our study has several strengths, including 1) the meticulous design of ACCORD and its excellent randomization, follow-up, and adherence; 2) the excellent quality of DNA samples, genotyping, and imputation and the precise measurements of a large panel of biomarkers; and 3) the availability of longitudinal biomarker measurements. However, several limitations should be acknowledged. First, the sample size was relatively small, and we cannot exclude that smaller effects on other biomarkers were present but were missed owing to limited power. Also, as a result of the small sample size, we were unable to evaluate the association of GLP-1 with cardiovascular outcomes to determine how much of the modulatory effect of rs57922 was mediated by its association with GLP-1 level during the intervention. In fact, we cannot rule out the possibility of the variant being associated with a cardiovascular benefit via another mechanism not involving the GLP-1 axis, which in turn has a secondary effect on GLP-1 levels. Second, given the clinical characteristics of the ACCORD participants, our findings may not be generalizable to younger subjects, nonwhites, or those with low cardiovascular risk. Third, since biomarker measurements were limited to baseline and 12 months of follow-up, longer-term effects on GLP-1 or other biomarker levels could not be investigated. Finally, as only fasting samples were taken, we cannot

make any inference on a possible impact of rs57922 on postprandial GLP-1 levels.

In conclusion, our results suggest a possible role of diminished GLP-1 activity as a factor for increased cardiac mortality during intensive glycemic therapy in rs57922 T/T homozygotes, possibly via impaired cardio- and/or stress-protective mechanisms or enhanced inflammatory pathways. Further studies seeking replication of this observation in other data sets and exploring the underlying mechanisms are warranted.

Acknowledgments. The authors thank the investigators, staff, and participants of ACCORD for their support and contributions and for providing access to this rich data set. The authors also specially acknowledge Timothy M. Hughes, Jeff Williamson, and Laura Lovato from the Wake Forest University School of Medicine, Winston-Salem, NC, and the other members of the ACCORD Biomarker Study Group. Members of the ACCORD Data and Safety Monitoring Board are as follows: Antonio M. Gotto Jr. (chair), Kent Bailey, Dorothy Gohdes, Steven Haffner, Roland Hiss, Kenneth Jamerson, Kerry Lee, David Nathan, James Sowers, and LeRoy Walters.

The following companies provided study medications, equipment, or supplies: Abbott Laboratories (Abbott Park, IL); Amylin Pharmaceuticals (San Diego, CA); AstraZeneca (Wilmington, DE); Bayer HealthCare LLC (Tarrytown, NY); Closer Healthcare, Inc. (Tequesta, FL); GlaxoSmithKline (Philadelphia, PA); King Pharmaceuticals, Inc. (Bristol, TN); Merck & Co. (Whitehouse Station, NJ); Novartis Pharmaceuticals (East Hanover, NJ); Novo Nordisk (Princeton, NJ); Omron Healthcare, Inc. (Schaumburg, IL); Sanofi U.S. (Bridgewater, NJ); Schering-Plough Corporation (Kenilworth, NJ); and Takeda Pharmaceuticals (Deerfield, IL). None of these companies had an interest or bearing on the genome-wide analysis of the ACCORD data.

Funding. The ACCORD genome-wide association analysis was supported by National Institutes of Health (NIH) grants HL-110380 (to J.B.B.), HL-110400 (to A.D.), and DK-36836 (Advanced Genomics and Genetics Core of the Diabetes Research Center at the Joslin Diabetes Center). The project described was also supported by the National Center for Advancing Translational Sciences (NCATS), NIH, through grant UL1TR001111. J.B.B. was also supported by the NCATS, NIH, through grant UL1TR001111. M.L.M. was supported by a William Randolph Hearst Fellowship provided by the Hearst Foundation. ACCORD (ClinicalTrials.gov identifier: NCT00000620) was supported by National Heart, Lung, and Blood Institute contracts N01-HC-95178, N01-HC-95179, N01-HC-95180, N01-HC-95181, N01-HC-95182, N01-HC-95183, and N01-HC-95184 and IAA no. Y1-HC-9035 and IAA no. Y1-HC-1010. Other components of the NIH, including the National Institute of Diabetes and Digestive and Kidney Diseases, the National Institute on Aging, and the National Eye Institute, contributed funding. The Centers for Disease Control and Prevention funded substudies within ACCORD on cost-effectiveness and health-related quality of life. General Clinical Research Centers and

Clinical and Translational Science Awards provided support at many sites.

The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH or other funders.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. H.S.S. designed the study; acquired, analyzed, and interpreted data; and wrote the manuscript. M.L.M. analyzed and interpreted data and reviewed the manuscript. S.M.M. acquired data, wrote part of the RESEARCH DESIGN AND METHODS, and reviewed the manuscript. R.J.S. designed the study and reviewed the manuscript. H.C.G. designed the study and reviewed the manuscript. M.J.W. acquired data and reviewed the manuscript. A.A.M.-R. acquired data and reviewed the manuscript. J.B.B. acquired data and reviewed the manuscript. P.K. designed the study and reviewed the manuscript. J.C.M. designed the study, acquired data, and reviewed the manuscript. A.D. designed the study; acquired, analyzed, and interpreted data; and wrote the manuscript. A.D. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. International Diabetes Federation. *IDF Diabetes Atlas*. 7th ed. Brussels, Belgium, International Diabetes Federation, 2015
2. Gerstein HC, Miller ME, Byington RP, et al.; Action to Control Cardiovascular Risk in Diabetes Study Group. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545–2559
3. Shah HS, Gao H, Morieri ML, et al. Genetic predictors of cardiovascular mortality during intensive glycemic control in type 2 diabetes: findings from the ACCORD clinical trial. *Diabetes Care* 2016;39:1915–1924
4. Samaropoulos XF, Light L, Ambrosius WT, Marcovina SM, Probstfield J, Goff DC Jr. The effect of intensive risk factor management in type 2 diabetes on inflammatory biomarkers. *Diabetes Res Clin Pract* 2012;95:389–398
5. Sandoval DA, D'Alessio DA. Physiology of glucagon peptides: role of glucagon and GLP-1 in health and disease. *Physiol Rev* 2015;95:513–548
6. Lim GE, Brubaker PL. Glucagon-like peptide 1 secretion by the L-cell: the view from within. *Diabetes* 2006;55(Suppl. 2):S70–S7
7. Müssig K, Staiger H, Machicao F, Häring H-U, Fritsche A. Genetic variants affecting incretin sensitivity and incretin secretion. *Diabetologia* 2010; 53:2289–2297
8. Drucker DJ. The biology of incretin hormones. *Cell Metab* 2006;3:153–165
9. Ravassa S, Zudaire A, Díez J. GLP-1 and cardioprotection: from bench to bedside. *Cardiovasc Res* 2012;94:316–323
10. Han B, Eskin E. Random-effects model aimed at discovering associations in meta-analysis of genome-wide association studies. *Am J Hum Genet* 2011;88:586–598
11. Anini Y, Brubaker PL. Role of leptin in the regulation of glucagon-like peptide-1 secretion. *Diabetes* 2003;52:252–259
12. Preiss D, Dawed A, Welsh P, et al.; DIRECT Consortium Group. Sustained influence of metformin therapy on circulating glucagon-like peptide-1

levels in individuals with and without type 2 diabetes. *Diabetes Obes Metab* 2017;19:356–363

13. Zietek T, Rath E. Inflammation meets metabolic disease: gut feeling mediated by GLP-1. *Front Immunol* 2016;7:154

14. Shiraki A, Oyama J, Komoda H, et al. The glucagon-like peptide 1 analog liraglutide reduces TNF- α -induced oxidative stress and inflammation in endothelial cells. *Atherosclerosis* 2012;221:375–382

15. Krasner NM, Ido Y, Ruderman NB, Cacicedo JM. Glucagon-like peptide-1 (GLP-1) analog liraglutide inhibits endothelial cell inflammation through a calcium and AMPK dependent mechanism. *PLoS One* 2014;9:e97554

16. Lee YS, Park MS, Choung JS, et al. Glucagon-like peptide-1 inhibits adipose tissue macrophage infiltration and inflammation in an obese mouse model of diabetes. *Diabetologia* 2012;55:2456–2468

17. Marso SP, Daniels GH, Brown-Frandsen K, et al.; LEADER Steering Committee; LEADER Trial Investigators. Liraglutide and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2016;375:311–322

18. Marso SP, Poulter NR, Nissen SE, et al. Design of the Liraglutide Effect and Action in Diabetes: Evaluation of Cardiovascular Outcome Results (LEADER) trial. *Am Heart J* 2013;166:823–830.e5

19. Marso SP, Bain SC, Consoi A, et al.; SUSTAIN-6 Investigators. Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2016;375:1834–1844

20. MacLusky NJ, Cook S, Scrocchi L, et al. Neuroendocrine function and response to stress in mice with complete disruption of glucagon-like peptide-1 receptor signaling. *Endocrinology* 2000;141:752–762

21. Ghosal S, Packard AEB, Mahbod P, et al. Disruption of glucagon-like peptide 1 signaling in Sim1 neurons reduces physiological and behavioral reactivity to acute and chronic stress. *J Neurosci* 2017;37:184–193

22. Rinaman L. Interoceptive stress activates glucagon-like peptide-1 neurons that project to the hypothalamus. *Am J Physiol* 1999;277:R582–R590

23. Sokos GG, Nikolaidis LA, Mankad S, Elahi D, Shannon RP. Glucagon-like peptide-1 infusion improves left ventricular ejection fraction and functional status in patients with chronic heart failure. *J Card Fail* 2006;12:694–699

24. Liu Q, Anderson C, Broyde A, et al. Glucagon-like peptide-1 and the exenatide analogue AC3174 improve cardiac function, cardiac remodeling, and survival in rats with chronic heart failure. *Cardiovasc Diabetol* 2010;9:76

25. van Genugten RE, Möller-Goede DL, van Raalte DH, Diamant M. Extra-pancreatic effects of incretin-based therapies: potential benefit for cardiovascular-risk management in type 2 diabetes. *Diabetes Obes Metab* 2013;15:593–606

26. Gros R, You X, Baggio LL, et al. Cardiac function in mice lacking the glucagon-like peptide-1 receptor. *Endocrinology* 2003;144:2242–2252

27. Nathanson D, Zethelius B, Berne C, et al. Plasma levels of glucagon like peptide-1 associate with diastolic function in elderly men. *Diabet Med* 2011;28:301–305

28. Nathanson D, Zethelius B, Berne C, Holst JJ, Sjöholm A, Nyström T. Reduced plasma levels of glucagon-like peptide-1 in elderly men are associated with impaired glucose tolerance but not with coronary heart disease. *Diabetologia* 2010;53:277–280