

RESEARCH ISSUES: THE FOOD ENVIRONMENT AND OBESITY

Food Availability/Convenience and Obesity^{1–5}

Penny Gordon-Larsen*

Department of Nutrition, Gillings School of Global Public Health and School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC

ABSTRACT

Neighborhood environments have received considerable attention in recent local, state, and national obesity prevention initiatives, with a particular focus on food deserts, or areas with poor access to healthy foods. Yet, there are inconsistencies in the evidence base, suggesting a nuanced association between neighborhood environment, food availability, diet behaviors, and obesity. There is heterogeneity in associations between environmental exposures and health outcomes across race/ethnicity, gender, region, and urbanicity, which results in complexity in the interpretation of findings. There are several limitations in the literature, including a predominance of cross-sectional studies, reliance on commercial business listings, lack of attention to the process by which diet resources are established and expanded within neighborhoods and the potential for individuals to selectively migrate to locate near such facilities, a predominant focus on residential neighborhoods, and lack of information about the decision-making process underlying purchasing patterns. More research is needed to address the complexity of individual-level residential decision making as well as the purposeful placement of food environment resources across social and geographic space using longitudinal data and complex statistical approaches. In addition, improvements in data quality and depth related to food access and availability are needed, including behavioral data on purchase patterns and interactions with the food environment, and greater attention to heterogeneity across subpopulations. As policy changes to the food environment move forward, it is critical that there is rigorous and scientific evaluation of environmental changes and their impact on individual-level diet choices and behaviors, and their further influence on body weight. *Adv Nutr* 2014;5:809–817.

Define the Issue

Scope. The dimension of food availability/convenience that has received considerable recent attention in the literature is neighborhood access to healthy foods. Neighborhoods that

offer access to high-quality foods are theorized to improve individual-level diet and weight outcomes of individuals residing in those neighborhoods. There have been substantial changes in the food environment over the past several decades, which have been broadly linked to dietary behaviors and obesity (1,2). In addition, neighborhood environments have received considerable attention in recent local, state, and national obesity prevention initiatives, with a particular focus on food deserts, or areas with poor access to healthy foods (3–6). However, the largely cross-sectional literature consists of studies that vary in geographic coverage, in methods for assessing environmental exposures, and in statistical modeling of associations, making comparisons across studies complex. In addition, there is heterogeneity in associations between environmental exposures and health outcomes across race/ethnicity, gender, region, and urbanicity, which also results in complexity in the interpretation of findings.

Severity. The ubiquitous access to energy-dense foods and beverages affects the health of individuals around the globe

¹ This article was peer-reviewed and approved by the Reviews, Papers, and Guidelines Committee and Medical Nutrition Council of the American Society for Nutrition (ASN) as part of a series entitled "Research Issues: The Food Environment and Obesity." The article did not undergo a separate editorial peer review by the editors of *Advances in Nutrition* (AN). ASN's journals feature a section called "From the American Society for Nutrition," which is devoted to content that originates from the Society rather than the journals. This article is being published in this section because of its broad interest to the nutrition community and the need to widely disseminate the information contained in the article. Correspondence about articles published in the "From the American Society for Nutrition" section should be addressed directly to ASN (info@nutrition.org) rather than to the AN Editor-in-Chief.

² An introduction to this series of articles is being published concurrently in *The American Journal of Clinical Nutrition* and the journal *Obesity*.

³ Supported by the National Heart, Lung, and Blood Institute (R01-HL114091 and R01-HL104580). P. Gordon-Larsen receives general support from the Carolina Population Center, University of North Carolina at Chapel Hill (grant R24 HD050924 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development).

⁴ Author disclosures: P. Gordon-Larsen, no conflicts of interest.

⁵ The NIH had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; or preparation, review, or approval of the manuscript.

* Address correspondence to P. Gordon-Larsen, University of North Carolina at Chapel Hill, Carolina Population Center, University of North Carolina—Chapel Hill, Campus Box #8120, 137 East Franklin Street, Room 6124, Chapel Hill, NC 27516. E-mail: gordon_larsen@unc.edu.

(7). The relative costs of fruit and vegetables have increased greatly compared with prices of refined grains and sugar, making access to all sorts of processed foods progressively easier along a range of income levels (7–9), although individuals of low income are particularly affected by this price structure (10–13). Overall, food and beverages have become extremely easy to access, with an increasing number of sales from a variety of sources, including vending machines, drug stores, book stores, hardware stores, and big box stores. For example, point-of-sale Scan Track data from Nielsen indicate that 10 billion dollars were spent in 2008 on food items at U.S. drugstores, totaling an estimated 3 trillion calories (B.M. Popkin, unpublished data, 2011).

Characteristics of those affected. Although neighborhood food environments may have some degree of influence on all residents, vulnerable subpopulations may be particularly at risk (14–16), given the observed differential access to food resources (17–23). Yet, there are inconsistencies in the evidence base, suggesting a nuanced association between neighborhood environment and obesity (5,16,24). Differences in availability and costs of healthier food items as well as other individual-level factors may substantially limit the opportunity for healthy eating among less-wealthy individuals. For example, due to perceived cost or other barriers (25), low-income individuals may be more sensitive to cues related to the presence of fast-food restaurants (26). Yet, choice of stores, perceived quality of available produce, and consumer buying patterns also play important roles in the availability of foods in deprived neighborhoods (27). There are also strong differences in access to healthy foods across rural and urban settings (28), although this is a relatively understudied area. This relatively small literature suggests that rural and less-urban areas are just as much, if not more, underserved (29–34). The limited research on food environments of immigrant and Latino populations suggests that such populations are a concern as well (35–37).

Critical Review

The bulk of the literature on this topic comes from cross-sectional observational epidemiologic studies. Thus, this literature base predominantly deals with association as opposed to causation. Similarly, given the cross-sectional body of literature as well as the inherent difficulty of randomization and assignment of an intervention to a neighborhood, the assessment of efficacy and effectiveness is limited. In addition, this is a nascent area of research, with rapid development of data and methodology to deal with research questions related to food availability and convenience in relation to obesity. The literature is limited in several important ways, including the following: 1) lack of fine-grain, longitudinal data characterizing changes in the food environment and their association with individual-level diet behaviors and obesity; 2) measurement issues related to characterizing the food environment; 3) lack of complex statistical

models examining each piece of the time-dependent, complex system, accounting for the process by which diet resources are established and expanded within neighborhoods and the potential for individuals to selectively migrate to locate near such facilities; 4) lack of pathway-based analyses that examine environmental influences on individual-level dietary intake and through this pathway to obesity; and 5) lack of rigorous evaluation of policy implementation and other alternations to the food environment. These gaps in research limit a full understanding of the nature of the influence of food environments on obesity.

Although neighborhoods that offer access to high-quality foods are theorized to improve the diet, weight, and cardiometabolic profiles of individuals living in close proximity to resources that support a healthy diet, the largely cross-sectional literature has produced mixed results [see reviews and position papers (1,16,23,24,38–41)]. The most heavily studied element of the food environment is supermarkets. Several studies provided support for a positive association between neighborhood access to supermarkets and healthy diets (i.e., closer proximity and healthier diets) (42–46), whereas others did not (47–50). Similarly, some studies found an inverse association between neighborhood access to supermarkets and body weight (51–56), although others did not (55,57,58). Many of these same studies or others did not find positive associations between grocery stores (which are smaller than supermarkets) with diet or inverse associations with weight (46,47,51,52,59–61). There are fewer studies of the association between convenience stores and higher body weight, with some finding positive associations (51,54,56,62) and others finding null associations (52,59). Recent work has begun to address the presence and use of corner stores and the availability of relatively less healthy food options (63–66). It is important to recognize that this body of evidence on the topic of the food environment includes considerable heterogeneity in the observed associations by age, ethnicity, and degree of rural, suburban, or urban development, which makes comparisons across studies difficult.

Other research has focused on access to fast-food restaurants [see reviews (67,68)]. There is evidence of inequities in access to fast-food restaurants, with greater access in low-income (49,69) and minority urban areas (70,71). There has been relatively little study of access to fast food and restaurants with dietary consumption, with generally null results (42,47,72) but positive associations by subpopulation (49). Studies of access to fast food and body weight generally showed null results (52,53,55,59,73–75), although some positive associations between fast-food access and higher body weight were found (46,56,76–79). Again, there was heterogeneity in associations across populations and settings. This heterogeneity might be due to synergistic effects of urban development and neighborhood poverty levels, such that high-density areas have broad access to a wider range of resources across all income levels as a function

of general development and higher population density, whereas access in areas with less development may be quite different. Another possibility is that the access to specific types of foods is not well characterized when looking at access to stores and restaurants. Furthermore, the predominant focus on residential neighborhoods may mask the use of food resources in other settings, such as workplace neighborhoods or the travel corridor between home and work.

Few studies incorporated longitudinal measures of social, built, and economic factors and their relation to health, despite calls for life-course and longitudinal studies (80–82). A recent joint Institute of Medicine–National Academy of Sciences workshop on food deserts found that the cross-sectional research suggested sufficient rationale for the targeted placement of new supermarkets and farmers markets in urban and low socioeconomic status neighborhoods, but the workshop report also noted a lack of longitudinal research connecting targeted placement to improved diet and health (81). Yet, policy has moved forward as part of recent local, state, and national obesity prevention initiatives, with a particular focus on food deserts (3–5,83,84). Clearly, there is interest in ameliorating obesity disparities, but the conceptual and methodologic challenges underlying the evidence base make it difficult to fully understand the likely impact of these policy changes (85).

There are many barriers to achieving a healthy diet given the current food environment, yet individuals of racial/ethnic minority and low socioeconomic status face particularly strong barriers to achieving a healthy diet (19,86–90). Nonetheless, the concept of food deserts has been controversial (91,92). Although individuals of lower income pay more for food (10), several studies suggest that the evidence that impoverished areas have higher access to fast foods and reduced access to supermarkets is mixed (16,23,91,93–95). Indeed, the issue may be more complex than simply providing additional supermarkets and grocery stores, with no additional support for enhancing individual-level behaviors. It is possible that such efforts need to be coupled with promotion, education, and incentives for purchasing healthier foods (49).

Part of the methodologic limitations in this area of research relate to the measurement and characterization of the food environment. Most research defines neighborhoods on the basis of administrative boundaries, such as counties (96), census block groups (97–99), or residential buffers (49,100,101). Although these boundaries are readily and inexpensively available, they are somewhat arbitrary and may not correspond to what the population *in situ* may consider as a neighborhood. Neighborhood boundaries are complex to define and might be determined by transportation patterns, social networks, proximity to resources, and more. This heterogeneity in definitions of neighborhoods no doubt contributes to mixed findings in the literature. Similarly, the relatively small geographic range of many studies as

well as the lack of variation in environment measures (21,22,37,55,79,102–106) might underlie the mixed findings in the literature.

There is also considerable lack of consistency in statistical adjustment for factors correlated with neighborhood socio-demographic characteristics and independently related to food resource availability (16,102,104,107–109). For example, the number and distribution of food resources (e.g., restaurants, supermarkets) relate to general urban development. Scaling resources by population (i.e., resource counts per 10,000 population) (110) may address the placement of commercial establishments according to population density and sociodemographic characteristics. Alternatively, roadway-scaled measures (resource counts per roadway mile) (4,111,112) can represent the concentration of resources along access routes and may help adjust for overall commercial activity, whereas accounting for car ownership can clarify use of local resources (113), and the ratio of fast-food restaurants to other restaurants can clarify restaurant choices (70,76). Improvement of neighborhood environment measures to better isolate the influence of neighborhood food resources is needed.

Another methodologic limitation relates to the interdependence of neighborhoods, food resources, and individuals, which has been largely ignored. Cross-sectional neighborhood environment studies are particularly problematic because neighborhoods and individual behaviors evolve over time through complex, interrelated processes (114). The complexity relates to individual-level residential decision making as well as the purposeful placement of food environment resources across social and geographic space. Both processes have been understudied (115).

On the individual side, financial or social constraints and residential preferences shape residential choice and movement (116). Residential choices occur sequentially and purposefully, and it is possible that individual-level preferences for healthy lifestyles (including diet and physical activity) play into the residential decision-making process (117,118). Residential location choice is a function of consumption of amenities, such as the presence of parks, trails, and recreation facilities (119,120); hedonic property values and wages (121,122); and educational resources, crime, and proximity to stores within walking distance (123). Most research assumes that no selective migration occurs to take advantage of such amenities (115,124,125). In general, cross-sectional studies ignore these dynamic interactions and thus might result in biased estimates of associations between the food environment and obesity-related behaviors (126).

Although these individual-level residential preferences are important, the purposeful placement of food environment resources across social and geographic space is extremely relevant but understudied. City planners themselves selectively choose the locations where facilities will be placed on the basis of a complex set of factors, some of which are

demand-driven (127,128); restaurant site location favors aesthetically pleasing locations for full-service restaurants and high-traffic-volume areas for fast-food restaurants (129), as well as characteristics of nearby residents (130). Yet, within this process, individual-level dietary intake is generally not addressed (131). The issue is relevant in that restaurants and supermarkets may be placed in areas with high demand, thus creating the potential for reverse causality. Similarly, areas with infrastructure and resources may attract residents who are more prone to use these facilities, and if these mechanisms are not accounted for, findings may be contaminated in such a way to either overstate or understate their impact.

To address the complexity of individual-level residential decision making as well as the purposeful placement of food environment resources across social and geographic space, longitudinal data and complex statistical approaches are needed. If there are factors that determine both the outcome (e.g., dietary intake) and facility placement (e.g., grocery stores) and these factors are either unmeasured or not included in the empirical analysis (via incorporation of the selection process into estimation of effects or control for unobservable factors), estimates of the effectiveness related to the placement of diet facilities will be biased (132–134). Similarly, models must account for the possibility that people migrate to take advantage of facilities (132,134), which might similarly bias estimates. Longitudinal data are needed to address individual characteristics that may contribute to these dynamic interrelationships (115). One approach is to control for time-constant unmeasured characteristics (that drive location selection) by using fixed-effects longitudinal models, which condition on each individual, thereby analyzing variation observed within person, over time (115); in essence, each individual serves as his/her own control. Another approach is the use of complex sequential modeling that incorporates the decision-making process into the model (133,134). Ultimately, the field needs to move toward longitudinal data and complex analyses to fully understand the complex relation between the food environment, dietary behaviors, and obesity.

Proposed Future Research Agenda

As reviews on this topic have noted, many of the obstacles to progress relate to the nature and quality of neighborhood-level data (135–137). Given the lack of longitudinal food environment data linked with high-quality longitudinal individual-level health data, there is a clear need for better retrospective data on the introduction, renovation, and closure of food resources. In addition, high-quality data from food stores and restaurants related to purchasing patterns, location decisions, and overall sales would also be very useful in understanding these associations. Given the current focus on residential addresses, there is little understanding of multiple environments (e.g., neighborhoods around workplace or school, travel corridors between home and work)

that influence obesity and obesity-related behaviors. Broader studies of locations other than just the residential environment are sorely needed.

Given that most of the published data on the food environment are based on commercial databases, which focus on the presence or absence rather than the quality of services or foods provided, there is a wide gap in fully understanding how individuals use food resources. In addition, the commercial databases provide temporal snapshots of facilities; there is limited detail on the precise timing of opening and closure of facilities and the validity of these commercial sources ranges from good to questionable (138–140). Furthermore, the reliance on traditional facilities for food resources might miss opportunities for the purchase of fresh fruits and vegetables at other locales, such as dollar stores (141), corner stores (64,65,142), or *tiendas* (36), or from fast-food establishments outside of traditional chain locales (143), as well as a variety of understudied food resources, such as hardware stores or drug stores. Indeed, policy is surpassing research in this arena; for example, the White House effort with Walgreens aims to convert at least 1000 of its stores into food oasis stores, selling whole fruits and vegetables, precut fruit salads, green salads, breads, and ready-made meals (84).

Research on food environments and individual-level behaviors is limited by the lack of information about the decision-making process underlying purchasing patterns. The current literature identifies associations between stores and restaurants and obesity or obesity-related behaviors, but few identify the quality of foods offered (56,144–148), purchasing patterns (149), and the types of foods purchased and prepared once purchased. This lack of attention might also underlie the mixed findings for proximity to supermarkets and individual-level diet (81,150). Recent work with attention to both the community and consumer nutrition environment provides much greater detail on dimensions of access and availability of foods (151) and represents a strong direction for future work. There is also a great need for statistical methods that can account for the process by which food and activity facilities are established and expanded as well as selective migration, 2 sources of bias that are unaddressed in the current literature.

Given the complexity underlying the relation between the food environment, individual-level diet behavior, and obesity, a systems-oriented, multilevel framework is recommended (152). Ideally, randomized experimental studies would provide the detail necessary to understand the effects of neighborhood environments on health, yet such studies are unethical or unfeasible (153–155). Innovative study designs, such as natural experiments or randomized trials, are also critical for understanding causal effects of neighborhoods on health (126,156). Despite calls for rigorous experimental designs (1) and some randomized controlled trials in the United Kingdom (91,157–159), there is very little such literature. In 1 of few quasi-experimental studies,

Cummins et al. (160) studied changes in fruit and vegetable consumption after the opening of a supermarket-type store in the United Kingdom and found similar patterns in comparison to a control neighborhood. Cummins et al. (157) reported the greatest improvements in fruit and vegetable intake among those who adopted a new supermarket as their main food store, suggesting that the promotion of existing, new, or improved food resources is an important component of successful policies.

Observational approaches that can mimic randomized controlled trials are another potential direction (155,161). With better methods to precisely measure changes in food environments over time, it may be possible to assess whether environmental supports for healthy diet behaviors translate to improvements in behavioral, and weight, outcomes. Addressing key unmeasured predictors of location selection that vary over time, such as change in marital or employment status (162), requires instrumental variables (163) or simultaneous equation strategies (115) that use longitudinal data, which can provide deeper understanding of these complex relations. Similarly, innovative simulation studies can be useful in predicting changes in weight with policy implementation addressing access to food (164).

However, there are limitations of innovative methods, such as agent-based models, because these computational models are dependent on theoretical or simulated data and relations and typically provide a range of possible outcomes rather than specific predictions (155,165,166). Furthermore, these methods may not handle residential selectivity and measurement error and thus can produce misleading results (133). A hybrid approach that uses simulation methods to trace the effects of key explanatory variables on outcomes through time, and includes attention to parameterization, calibration of equations and algorithms, and transparency of the model (i.e., reporting standards and validation), may increase the utility of agent-based models.

With all of the methodologic advances it is still critical to address heterogeneity in environmental effects. There are clearly subpopulations at high risk of obesity, particularly low-income, ethnic minority, immigrant, and inner city residents. For example, for supermarkets to successfully improve diets, promoting existing resources within target groups may be necessary (158), as seen in recent efforts in New York City with the Healthy Bodegas program (167) and in Philadelphia with the Healthy Corner Store initiative (168). In addition, improving neighborhood resources should be weighed against (or in addition to) alternative approaches, such as taxation, subsidization, or incentives [e.g., reduced-cost exercise programs, sugar-sweetened beverage tax (169)]. Direct comparison of different policy strategies with the use of common health metrics may facilitate evidence-based policy making (170).

As policies, such as the Healthy Food Financing initiative (83), and the introduction of new food stores and changes to current food stores (84) move forward, it is critically

important for rigorous evaluation of these changes. Such evaluations have been lacking (81). A team of experts recently put forth a number of recommendations for improving methodologies for environment and policy research, with evaluation of “natural experiments” receiving particularly high recommendation (171). In another article, a set of researchers suggested a set of questions designed to evaluate complex public health and wider social interventions in terms of process and impacts of policy changes (172). Ultimately, these approaches are necessary to fully understand the impact of environmental policies and changes and to maximize the use of scarce resources, particularly in low-income neighborhoods.

In sum, there is a great need for high-quality longitudinal data and rigorous analytical methodologies to fully understand the relation between food availability/convenience and obesity. Improvements in data quality and depth related to food access and availability are clearly needed, as are behavioral data on purchasing patterns and interactions with the food environment, along with greater attention to heterogeneity across subpopulations. There is also a major need for innovative analytical tools for modeling these relations in a multilevel and systems-based approach that can account for residential selectivity and purposeful placement of facilities. As policy changes to the food environment move forward, rigorous and scientific evaluation of environmental changes and their impact on individual-level diet choices and behaviors, and further influence on body weight, is a necessary component.

Acknowledgments

The sole author had responsibility for all parts of the manuscript.

References

1. Sallis JF, Glanz K. Physical activity and food environments: solutions to the obesity epidemic. *Milbank Q* 2009;87:123–54.
2. Hill JO, Peters JC. Environmental contributions to the obesity epidemic. *Science* 1998;280:1371–4.
3. Let's Move. Accessing healthy and affordable food. 2010 [cited 2010 Jul 7]. Available from: <http://www.letsmove.gov/make-healthy-food-affordable-and-accessible>.
4. New York City Industrial Development Agency. New York City industrial development agency approves incentives for first two supermarkets under the fresh program. 2010 [cited 2010 Jul 7]. Available from: <http://www.nycdc.com/PressRoom/PressReleases/Pages/IDAApprovesIncentivesforSupermarkets.aspx>.
5. Sturm R, Cohen DA. Zoning For health? The year-old ban on new fast-food restaurants in South LA. *Health Aff (Millwood)* 2009;28:1088–97.
6. Walker RE, Keane CR, Burke JG. Disparities and access to healthy food in the United States: a review of food deserts literature. *Health Place* 2010;16:876–84.
7. Popkin BM. Agricultural policies, food and public health. *EMBO Rep* 2011;12:11–8.
8. Alston JM, Sumner DA, Vosti SA. Are agricultural policies making us fat? Likely links between agricultural policies and human nutrition and obesity, and their policy implications. *Rev Agric Econ* 2006;28:313–22.

9. Christian T, Rashad I. Trends in U.S. food prices, 1950–2007. *Econ Hum Biol* 2009;7:113–20.
10. Caplovitz D. *The poor pay more: consumer practices of low-income families*. New York: Free Press; 1967.
11. Chung C, Myers S. Do the poor pay more for food? An analysis of grocery store availability and food price disparities. *J Consum Aff* 1999;33:276–96.
12. Drewnowski A, Darmon N. Food choices and diet costs: an economic analysis. *J Nutr* 2005;135:900–4.
13. Maillot M, Darmon N, Vieux F, Drewnowski A. Low energy density and high nutritional quality are each associated with higher diet costs in French adults. *Am J Clin Nutr* 2007;86:690–6.
14. Pickett KE, Pearl M. Multilevel analyses of neighbourhood socioeconomic context and health outcomes: a critical review. *J Epidemiol Community Health* 2001;55:111–22.
15. Robert SA, Reither EN. A multilevel analysis of race, community disadvantage, and body mass index among adults in the US. *Soc Sci Med* 2004;59:2421–34.
16. Larson NI, Story MT, Nelson MC. Neighborhood environments disparities in access to healthy foods in the US. *Am J Prev Med* 2009;36:74–81.
17. Cubbin C, Hadden W, Winkleby M. Neighborhood context and cardiovascular disease risk factors: the contribution of material deprivation. *Ethn Dis* 2001;11:687–700.
18. Cummins S, Stafford M, Macintyre S, Marmot M, Ellaway A. Neighbourhood environment and its association with self rated health: evidence from Scotland and England. *J Epidemiol Community Health* 2005;59:207–13.
19. Moore LV, Diez Roux AV. Associations of neighborhood characteristics with the location and type of food stores. *Am J Public Health* 2006;96:325–31.
20. Reidpath DD, Burns C, Garrard J, Mahoney M, Townsend M. An ecological study of the relationship between social and environmental determinants of obesity. *Health Place* 2002;8:141–5.
21. Block JP, Scribner RA, DeSalvo KB. Fast food, race/ethnicity, and income: a geographic analysis. *Am J Prev Med* 2004;27:211–7.
22. Morland K, Wing S, Diez Roux A, Poole C. Neighborhood characteristics associated with the location of food stores and food service places. *Am J Prev Med* 2002;22:23–9.
23. Lovasi GS, Hutson MA, Guerra M, Neckerman KM. Built environments and obesity in disadvantaged populations. *Epidemiol Rev* 2009;31:7–20.
24. Macintyre S. Deprivation amplification revisited; or, is it always true that poorer places have poorer access to resources for healthy diets and physical activity? *Int J Behav Nutr Phys Act* 2007;4:32–8.
25. Dammann KW, Smith C. Factors affecting low-income women's food choices and the perceived impact of dietary intake and socioeconomic status on their health and weight. *J Nutr Educ Behav* 2009;41:242–53.
26. Paquet C, Daniel M, Knauper B, Gauvin L, Kestens Y, Dube L. Interactive effects of reward sensitivity and residential fast-food restaurant exposure on fast-food consumption. *Am J Clin Nutr* 2010;91:771–6.
27. Cummins S, Smith DM, Aitken Z, Dawson J, Marshall D, Sparks L, Anderson AS. Neighbourhood deprivation and the price and availability of fruit and vegetables in Scotland. *J Hum Nutr Diet* 2010;23:494–501.
28. Smith DM, Cummins S, Taylor M, Dawson J, Marshall D, Sparks L, Anderson AS. Neighbourhood food environment and area deprivation: spatial accessibility to grocery stores selling fresh fruit and vegetables in urban and rural settings. *Int J Epidemiol* 2010;39:277–84.
29. Dean WR, Sharkey JR. Rural and urban differences in the associations between characteristics of the community food environment and fruit and vegetable intake. *J Nutr Educ Behav* 2011;43:426–33.
30. Sharkey JR, Horel S. Neighborhood socioeconomic deprivation and minority composition are associated with better potential spatial access to the ground-truthed food environment in a large rural area. *J Nutr* 2008;138:620–7.
31. Sharkey JR. Measuring potential access to food stores and food-service places in rural areas in the U.S. *Am J Prev Med* 2009;36:S151–5.
32. Liese AD, Weis KE, Pluto D, Smith E, Lawson A. Food store types, availability, and cost of foods in a rural environment. *J Am Diet Assoc* 2007;107:1916–23.
33. Frost SS, Goins RT, Hunter RH, Hooker SP, Bryant LL, Kruger J, Pluto D. Effects of the built environment on physical activity of adults living in rural settings. *Am J Health Promot* 2010;24:267–83.
34. Hosler AS. Retail food availability, obesity, and cigarette smoking in rural communities. *J Rural Health* 2009;25:203–10.
35. Sharkey JR, Dean WR, Johnson CM. Association of household and community characteristics with adult and child food insecurity among Mexican-origin households in Colonias along the Texas-Mexico border. *Int J Equity Health* 2011;10:19–32.
36. Emond JA, Madanat HN, Ayala GX. Do Latino and non-Latino grocery stores differ in the availability and affordability of healthy food items in a low-income, metropolitan region? *Public Health Nutr* 2011;15:1–10.
37. Galvez MP, Morland K, Raines C, Kobil J, Siskind J, Godbold J, Brenner B. Race and food store availability in an inner-city neighbourhood. *Public Health Nutr* 2008;11:624–31.
38. Feng J, Glass TA, Curriero FC, Stewart WF, Schwartz BS. The built environment and obesity: a systematic review of the epidemiologic evidence. *Health Place* 2010;16:175–90.
39. Giskes K, van Lenthe F, Avendano-Pabon M, Brug J. A systematic review of environmental factors and obesogenic dietary intakes among adults: are we getting closer to understanding obesogenic environments? *Obes Rev* 2011;12:e95–106.
40. Holsten JE. Obesity and the community food environment: a systematic review. *Public Health Nutr* 2009;12:397–405.
41. Papas MA, Alberg AJ, Ewing R, Helzlsouer KJ, Gary TL, Klassen AC. The built environment and obesity. *Epidemiol Rev* 2007;29:129–43.
42. Morland K, Wing S, Diez Roux A. The contextual effect of the local food environment on residents' diets: the Atherosclerosis Risk in Communities Study. *Am J Public Health* 2002;92:1761–7.
43. Laraia BA, Siega-Riz AM, Kaufman JS, Jones SJ. Proximity of supermarkets is positively associated with diet quality index for pregnancy. *Prev Med* 2004;39:869–75.
44. Rose D, Richards R. Food store access and household fruit and vegetable use among participants in the US Food Stamp Program. *Public Health Nutr* 2004;7:1081–8.
45. Moore LV, Diez Roux AV, Nettleton JA, Jacobs DR Jr. Associations of the local food environment with diet quality—a comparison of assessments based on surveys and geographic information systems: the Multi-Ethnic Study of Atherosclerosis. *Am J Epidemiol* 2008;167:917–24.
46. Morland KB, Evenson KR. Obesity prevalence and the local food environment. *Health Place* 2009;15:491–5.
47. Jago R, Baranowski T, Baranowski JC, Cullen KW, Thompson D. Distance to food stores & adolescent male fruit and vegetable consumption: mediation effects. *Int J Behav Nutr Phys Act* 2007;4:35–44.
48. Ford PB, Dziewaltowski DA. Limited supermarket availability is not associated with obesity risk among participants in the Kansas WIC Program. *Obesity (Silver Spring)* 2010;18:1944–51.
49. Boone-Heinonen J, Gordon-Larsen P, Kiefe CI, Shikany JM, Lewis CE, Popkin BM. Fast food restaurants and food stores: longitudinal associations with diet in young to middle-aged adults: the CARDIA study. *Arch Intern Med* 2011;171:1162–70.
50. Jilcott SB, Keyserling T, Crawford T, McQuirt JT, Ammerman AS. Examining associations among obesity and per capita farmers' markets, grocery stores/supermarkets, and supercenters in US counties. *J Am Diet Assoc* 2011;111:567–72.
51. Morland K, Diez Roux AV, Wing S. Supermarkets, other food stores, and obesity: the Atherosclerosis Risk in Communities Study. *Am J Prev Med* 2006;30:333–9.
52. Liu GC, Wilson JS, Qi R, Ying J. Green neighborhoods, food retail and childhood overweight: differences by population density. *Am J Health Promot* 2007;21:317–25.

53. Lopez RP. Neighborhood risk factors for obesity. *Obesity* (Silver Spring) 2007;15:2111–9.
54. Powell LM, Auld MC, Chaloupka FJ, O'Malley PM, Johnston LD. Associations between access to food stores and adolescent body mass index. *Am J Prev Med* 2007;33:S301–7.
55. Wang MC, Kim S, Gonzalez AA, MacLeod KE, Winkleby MA. Socio-economic and food-related physical characteristics of the neighbourhood environment are associated with body mass index. *J Epidemiol Community Health* 2007;61:491–8.
56. Bodor JN, Rice JC, Farley TA, Swalm CM, Rose D. The association between obesity and urban food environments. *J Urban Health* 2010;87:771–81.
57. Gary-Webb TL, Baptiste-Roberts K, Pham L, Wesche-Thobaben J, Patricio J, Pi-Sunyer FX, Brown AF, Jones L, Brancati FL; Look AHEAD Research Group. Neighborhood and weight-related health behaviors in the Look AHEAD (Action for Health in Diabetes) study. *BMC Public Health* 2010;10:312–22.
58. Casagrande SS, Franco M, Gittelsohn J, Zonderman AB, Evans MK, Fanelli Kuczmarowski M, Gary-Webb TL. Healthy food availability and the association with BMI in Baltimore, Maryland. *Public Health Nutr* 2011;14:1001–7.
59. Sturm R, Datar A. Body mass index in elementary school children, metropolitan area food prices and food outlet density. *Public Health* 2005;119:1059–68.
60. Powell LM, Auld MC, Chaloupka FJ, O'Malley PM, Johnston LD. Access to fast food and food prices: relationship with fruit and vegetable consumption and overweight among adolescents. *Adv Health Econ Health Serv Res* 2007;17:23–48.
61. Inagami S, Cohen DA, Finch BK, Asch SM. You are where you shop: grocery store locations, weight, and neighborhoods. *Am J Prev Med* 2006;31:10–7.
62. Galvez MP, Hong L, Choi E, Liao L, Godbold J, Brenner B. Childhood obesity and neighborhood food-store availability in an inner-city community. *Acad Pediatr* 2009;9:339–43.
63. D'Angelo H, Suratkar S, Song H-J, Stauffer E, Gittelsohn J. Access to food source and food source use are associated with healthy and unhealthy food-purchasing behaviours among low-income African-American adults in Baltimore City. *Public Health Nutr* 2011;14:1632–9.
64. Lucan SC, Karpyn A, Sherman S. Storing empty calories and chronic disease risk: snack-food products, nutritive content, and manufacturers in Philadelphia corner stores. *J Urban Health* 2010;87:394–409.
65. Borradaile KE, Sherman S, Vander Veur SS, McCoy T, Sandoval B, Nachmani J, Karpyn A, Foster GD. Snacking in children: the role of urban corner stores. *Pediatrics* 2009;124:1293–8.
66. Gittelsohn J, Song HJ, Suratkar S, Kumar MB, Henry EG, Sharma S, Mattingly M, Anliker JA. An urban food store intervention positively affects food-related psychosocial variables and food behaviors. *Health Educ Behav* 2010;37:390–402.
67. Fleischhacker SE, Evenson KR, Rodriguez DA, Ammerman AS. A systematic review of fast food access studies. *Obes Rev* 2011;12:e460–71.
68. Fraser LK, Edwards KL, Cade J, Clarke GP. The geography of fast food outlets: a review. *Int J Environ Res Public Health* 2010;7:2290–308.
69. Moore LV, Diez Roux AV, Nettleton JA, Jacobs DR, Franco M. Fast-food consumption, diet quality, and neighborhood exposure to fast food: the Multi-Ethnic Study of Atherosclerosis. *Am J Epidemiol* 2009;170:29–36.
70. Powell LM, Chaloupka FJ, Bao Y. The availability of fast-food and full-service restaurants in the United States: associations with neighborhood characteristics. *Am J Prev Med* 2007;33:S240–5.
71. Macdonald L, Cummins S, Macintyre S. Neighbourhood fast food environment and area deprivation—substitution or concentration? *Appetite* 2007;49:251–4.
72. Richardson AS, Boone-Heinonen J, Popkin BM, Gordon-Larsen P. Neighborhood fast food restaurants and fast food consumption: a national study. *BMC Public Health* 2011;11:543–50.
73. Burdette HL, Whitaker RC. Neighborhood playgrounds, fast food restaurants, and crime: relationships to overweight in low-income preschool children. *Prev Med* 2004;38:57–63.
74. Simmons D, McKenzie A, Eaton S, Cox N, Khan MA, Shaw J, Zimmet P. Choice and availability of takeaway and restaurant food is not related to the prevalence of adult obesity in rural communities in Australia. *Int J Obes (Lond)* 2005;29:703–10.
75. Jeffery RW, Baxter J, McGuire M, Linde J. Are fast food restaurants an environmental risk factor for obesity? *Int J Behav Nutr Phys Act* 2006;3:2–7.
76. Mehta NK, Chang VW. Weight status and restaurant availability: a multilevel analysis. *Am J Prev Med* 2008;34:127–33.
77. Spence JC, Cutumisu N, Edwards J, Raine KD, Smoyer-Tomic K. Relation between local food environments and obesity among adults. *BMC Public Health* 2009;9:192–7.
78. Li F, Harmer P, Cardinal BJ, Bosworth M, Johnson-Shelton D. Obesity and the built environment: does the density of neighborhood fast-food outlets matter? *Am J Health Promot* 2009;23:203–9.
79. Currie J, Della Vigna S, Moretti E, Pathania V. The effect of fast food restaurants on obesity and weight gain. *Am Econ J Econ Policy* 2010;2:32–63.
80. Diez-Roux AV. Multilevel analysis in public health research. *Annu Rev Public Health* 2000;21:171–92.
81. Whitacre PT, Tsai P, Mulligan J; National Research Council. The public health effects of food deserts: workshop summary. Washington: National Academies Press; 2009.
82. Story M, Giles-Corti B, Yaroch AL, Cummins S, Frank LD, Huang TT, Lewis B. Work group IV: future directions for measures of the food and physical activity environments. *Am J Prev Med* 2009;36:S182–8.
83. U.S. Department of Health and Human Services. Obama Administration details healthy food financing initiative. 2010 [cited 2011 Jul 25]. Available from: <http://www.usda.gov/wps/portal/usda/usdahome?contentid=2010/02/0077.xml>.
84. The White House. First Lady Michelle Obama announces nationwide commitments to provide millions of people access to healthy, affordable food in underserved communities. 2011 [cited 2011 Jul 25, 2011]. Available from: <http://www.whitehouse.gov/the-press-office/2011/07/20/first-lady-michelle-obama-announces-nationwide-commitments-provide-milli>.
85. Diez Roux AV, Mair C. Neighborhoods and health. *Ann N Y Acad Sci* 2010;1186:125–45.
86. Zenk SN, Lachance LL, Schulz AJ, Mentz G, Kannan S, Ridella W. Neighborhood retail food environment and fruit and vegetable intake in a multiethnic urban population. *Am J Health Promot* 2009;23:255–64.
87. Horowitz CR, Colson KA, Hebert PL, Lancaster K. Barriers to buying healthy foods for people with diabetes: evidence of environmental disparities. *Am J Public Health* 2004;94:1549–54.
88. Morland K, Filomena S. Disparities in the availability of fruits and vegetables between racially segregated urban neighbourhoods. *Public Health Nutr* 2007;10:1481–9.
89. Satia JA, Galanko JA, Siega-Riz AM. Eating at fast-food restaurants is associated with dietary intake, demographic, psychosocial and behavioural factors among African Americans in North Carolina. *Public Health Nutr* 2004;7:1089–96.
90. Zenk SN, Schulz AJ, Israel BA, James SA, Bao S, Wilson ML. Neighborhood racial composition, neighborhood poverty, and the spatial accessibility of supermarkets in metropolitan Detroit. *Am J Public Health* 2005;95:660–7.
91. Cummins S, Macintyre S. "Food deserts"—evidence and assumption in health policy making. *BMJ* 2002;325:436–8.
92. Beaulac J, Kristjansson E, Cummins S. A systematic review of food deserts, 1966–2007. *Prev Chronic Dis* 2009;6:A105–14.
93. Pearson T, Russell J, Campbell MJ, Do Barker ME. 'Food deserts' influence fruit and vegetable consumption?—a cross-sectional study. *Appetite* 2005;45:195–7.

94. Cummins S, Macintyre S. A systematic study of an urban foodscape: the price and availability of food in greater Glasgow. *Urban Stud* 2002; 39:2115–30.
95. Powell LM, Slater S, Mirtcheva D, Bao Y, Chaloupka FJ. Food store availability and neighborhood characteristics in the United States. *Prev Med* 2007;44:189–95.
96. Ewing R, Brownson RC, Berrigan D. Relationship between urban sprawl and weight of United States youth. *Am J Prev Med* 2006;31: 464–74.
97. Gordon-Larsen P, Nelson MC, Page P, Popkin BM. Inequality in the built environment underlies key health disparities in physical activity and obesity. *Pediatrics* 2006;117:417–24.
98. Diez Roux AV, Merkin SS, Arnett D, Chambless L, Massing M, Nieto FJ, Sorlie P, Szklo M, Tyroler HA, Watson RL. Neighborhood of residence and incidence of coronary heart disease. *N Engl J Med* 2001; 345:99–106.
99. Rundle A, Roux AV, Free LM, Miller D, Neckerman KM, Weiss CC. The urban built environment and obesity in New York City: a multi-level analysis. *Am J Health Promot* 2007;21:326–34.
100. Diez Roux AV, Evenson KR, McGinn AP, Brown DG, Moore L, Brines S, Jacobs DR Jr. Availability of recreational resources and physical activity in adults. *Am J Public Health* 2007;97:493–9.
101. Cohen DA, Ashwood JS, Scott MM, Overton A, Evenson KR, Staten LK, Porter D, McKenzie TL, Catellier D. Public parks and physical activity among adolescent girls. *Pediatrics* 2006;118:e1381–9.
102. Zenk SN, Powell LM. US secondary schools and food outlets. *Health Place* 2008;14:336–46.
103. Sturm R. Disparities in the food environment surrounding US middle and high schools. *Public Health* 2008;122:681–90.
104. Hurvitz P, Moudon A, Rehm C, Streichert L, Drewnowski A. Arterial roads and area socioeconomic status are predictors of fast food restaurant density in King County, WA. *Int J Behav Nutr Phys Act* 2008;6: 1–21.
105. Timperio A, Ball K, Roberts R, Campbell K, Andrianopoulos N, Crawford D. Children's fruit and vegetable intake: associations with the neighbourhood food environment. *Prev Med* 2008;46:331–5.
106. Simon PA, Kwan D, Angelescu A, Shih M, Fielding JE. Proximity of fast food restaurants to schools: do neighborhood income and type of school matter? *Prev Med* 2008;47:284–8.
107. Powell LM, Bao Y. Food prices, access to food outlets and child weight. *Econ Hum Biol* 2009;7:64–72.
108. Rundle A, Neckerman KM, Freeman L, Lovasi GS, Purciel M, Quinn J, Richards C, Sircar N, Weiss C. Neighborhood food environment and walkability predict obesity in New York City. *Environ Health Perspect* 2009;117:442–7.
109. Li F, Harmer PA, Cardinal BJ, Bosworth M, Acock A, Johnson-Shelton D, Moore JM. Built environment, adiposity, and physical activity in adults aged 50–75. *Am J Prev Med* 2008;35:38–46.
110. Boone-Heinonen J, Gordon-Larsen P. Life stage and sex specificity in relationships between the built and socioeconomic environments and physical activity. *J Epidemiol Community Health* 2011;65:847–52.
111. Richardson AS, Boone-Heinonen J, Popkin BM, Gordon-Larsen P. Neighborhood fast food restaurants and fast food consumption: a national study. *BMC Public Health* 2011;11:543. <http://www.biomedcentral.com/1471-2458/11/543>.
112. Romley JA, Cohen D, Ringel J, Sturm R. Alcohol and environmental justice: the density of liquor stores and bars in urban neighborhoods in the United States. *J Stud Alcohol Drugs* 2007;68:48–55.
113. Inagami S, Cohen DA, Brown AF, Asch SM. Body mass index, neighborhood fast food and restaurant concentration, and car ownership. *J Urban Health* 2009;86:683–95.
114. Plantinga AJ, Bernell S. A spatial economic analysis of urban land use and obesity. *J Reg Sci* 2005;45:473–92.
115. Boone-Heinonen J, Gordon-Larsen P, Guilkey D, Jacobs DR, Song YM, Popkin BM. Environment and physical activity dynamics: the role of residential self selection. *Psychol Sport Exerc* 2011;12:54–60.
116. Sampson RJ, Sharkey P. Neighborhood selection and the social reproduction of concentrated racial inequality. *Demography* 2008;45:1–29.
117. Pagliara F, Wilson A. The state-of-the-art in building residential location models. In: Pagliara F, Preston J, Simmonds D. (Eds.) *Residential Location Choice: Models and Applications*. New York: Springer, 2010, pp. 1–20.
118. Dowding K, John P, Biggs S. Tiebout: a survey of the empirical literature. *Urban Stud* 1994;31:767–98.
119. Levinson DM. Accessibility and the journey to work. *J Transp Geogr* 1998;6:11–21.
120. Cho EJ, Rodriguez DA, Song Y. The role of employment subcenters in residential location decisions. *J Transport Land Use* 2008;1:121–51.
121. Bartik TJ, Smith VK. Urban amenities and public policy. In: Mills ES, editor. *Urban economics*. Amsterdam: North-Holland; 1987:1207–49.
122. Song Y, Sohn J. Valuing spatial accessibility to retailing: a case study of the single family housing market in Hillsboro, Oregon. *J Retailing Con Serv* 2007;14:279–88.
123. Song Y, Knaap G. New urbanism and housing values: a disaggregate assessment. *J Urban Econ* 2003;54:218–38.
124. Boone-Heinonen J, Guilkey D, Evenson KR, Gordon-Larsen P. Residential self-selection bias in the estimation of built environment effects on physical activity between adolescence and young adulthood. *Int J Behav Nutr Phys Act* 2010;7:70–81.
125. Mokhtarian PL, Cao X. Examining the impacts of residential self-selection on travel behavior: a focus on methodologies. *Transp Res Part B* 2008;42:204–28.
126. Oakes JM. The (mis)estimation of neighborhood effects: causal inference for a practicable social epidemiology. *Soc Sci Med* 2004;58:1929–52.
127. Berke P, Godschalk D, Kaiser E, Rodriguez DA. *Urban land use planning*. 5th ed. Urbana-Champaign (IL): University of Illinois Press; 2006.
128. Brown S. Retail location theory: evolution and evaluation. *Int Rev Retail Distrib Consum Res* 1993;3:185–229.
129. Pillsbury R. From Hamburger Alley to Hedgerose Heights: toward a model of restaurant location dynamics. *Prof Geogr* 1987;39:326–44.
130. Zwolak R. (2010). *IBISWorld Industry Report 72221. Fast Food Restaurants in the US*. Available from <http://www.ibisworld.com>. [Cited 2011 Nov 15].
131. Smith H. Supermarket choice and supermarket competition in market equilibrium. *Rev Econ Stud* 2004;71:235–63.
132. Strauss J, Thomas D. Human resources: empirical modeling of household and family decisions. In: Srinivasan TN, Behrman J, eds. *Handbook of development economics*. Amsterdam: North Holland Press; 1995:1883–2023.
133. Angeles G, Guilkey DK, Mroz TA. Purposive program placement and the estimation of family program effects in Tanzania. *J Am Stat Assoc* 1998;93:884–99.
134. Rindfuss RR, Guilkey D, Morgan SP, Kravdal O, Guzzo KB. Child care availability and first-birth timing in Norway. *Demography* 2007;44: 345–72.
135. Lytle LA. Measuring the food environment: state of the science. *Am J Prev Med* 2009;36:S134–44.
136. Porsyth A, Lytle LA, Van Riper D. Finding food: issues and challenges in using Geographic Information Systems (GIS) to measure food access. *J Transp Land Use* 2010;3:43–65.
137. McKinnon RA, Reedy J, Morrisette MA, Lytle LA, Yaroch AL. Measures of the food environment: a compilation of the literature, 1990–2007. *Am J Prev Med* 2009;36:S124–33.
138. Liese AD, Colabianchi N, Lamichhane AP, Barnes TL, Hibbert JD, Porter DE, Nichols MD, Lawson AB. Validation of 3 food outlet databases: completeness and geospatial accuracy in rural and urban food environments. *Am J Epidemiol* 2010;172:1324–33.
139. Longacre MR, Primack BA, Owens PM, Gibson L, Beauregard S, Mackenzie TA, Dalton MA. Public directory data sources do not accurately characterize the food environment in two predominantly rural states. *J Am Diet Assoc* 2011;111:577–82.

140. Cummins S, Macintyre S. Are secondary data sources on the neighbourhood food environment accurate? Case-study in Glasgow, UK. *Prev Med* 2009;49:527–8.
141. Sharkey JR, Horel S, Dean WR. Neighborhood deprivation, vehicle ownership, and potential spatial access to a variety of fruits and vegetables in a large rural area in Texas. *Int J Health Geogr* 2010; 9:26–52.
142. Ghirardelli A, Quinn V, Foerster SB. Using geographic information systems and local food store data in California's low-income neighborhoods to inform community initiatives and resources. *Am J Public Health* 2010;100:2156–62.
143. Creel JS, Sharkey JR, McIntosh A, Anding J, Huber JC, Jr. Availability of healthier options in traditional and nontraditional rural fast-food outlets. *BMC Public Health* 2008;8:395–404.
144. Andreyeva T, Blumenthal DM, Schwartz MB, Long MW, Brownell KD. Availability and prices of foods across stores and neighborhoods: the case of New Haven, Connecticut. *Health Aff (Millwood)* 2008;27:1381–8.
145. Rose D, Bodor JN, Hutchinson PL, Swalm CM. The importance of a multi-dimensional approach for studying the links between food access and consumption. *J Nutr* 2010;140:1170–4.
146. Azuma AM, Gilliland S, Vallianatos M, Gottlieb R. Food access, availability, and affordability in 3 Los Angeles communities, Project CAFE, 2004–2006. *Prev Chronic Dis* 2010;7:A27–35.
147. Glanz K, Sallis JF, Saelens BE, Frank LD. Nutrition Environment Measures Survey in stores (NEMS-S): development and evaluation. *Am J Prev Med* 2007;32:282–9.
148. Saelens BE, Glanz K, Sallis JF, Frank LD. Nutrition Environment Measures Study in restaurants (NEMS-R): development and evaluation. *Am J Prev Med* 2007;32:273–81.
149. Blanck HM, Thompson OM, Nebeling L, Yaroch AL. Improving fruit and vegetable consumption: use of farm-to-consumer venues among US adults. *Prev Chronic Dis* 2011;8:A49–53.
150. Ver Ploeg M. Access to affordable, nutritious food is limited in "food deserts". *Amber Waves* 2010;20–7. [Cited 2011 Nov 15]. Available from: <http://www.ers.usda.gov/AmberWaves/March10/Features/FoodDeserts.htm>.
151. Cerin E, Frank LD, Sallis JF, Saelens BE, Conway TL, Chapman JE, Glanz K. From neighborhood design and food options to residents' weight status. *Appetite* 2011;56:693–703.
152. Huang TT, Drewnowski A, Kumanyika S, Glass T. A systems-oriented multilevel framework for addressing obesity in the 21st century. *Prev Chronic Dis* 2009;6:A82–91.
153. Oakes JM. Causal inference and the relevance of social epidemiology. *Soc Sci Med* 2004;58:1969–71.
154. Subramanian SV. The relevance of multilevel statistical methods for identifying causal neighborhood effects. *Soc Sci Med* 2004;58:1961–7.
155. Auchincloss AH, Diez Roux AV. A new tool for epidemiology: the usefulness of dynamic-agent models in understanding place effects on health. *Am J Epidemiol* 2008;168:1–8.
156. Glymour MM. Natural experiments and instrumental variable analysis in social epidemiology. In: Oakes JM, Kaufman JS, editors. *Methods in social epidemiology*. (pp. 429–460). San Francisco: Jossey-Bass; 2006.
157. Cummins S, Findlay A, Petticrew M, Sparks L. Healthy cities: the impact of food retail-led regeneration on food access, choice and retail structure. *Built Environ* 2005;31:288–301.
158. Cummins S, Petticrew M, Higgins C, Findlay A, Sparks L. Large scale food retailing as an intervention for diet and health: quasi-experimental evaluation of a natural experiment. *J Epidemiol Community Health* 2005;59:1035–40.
159. Petticrew M, Cummins S, Ferrell C, Findlay A, Higgins C, Hoy C, Kearns A, Sparks L. Natural experiments: an underused tool for public health? *Public Health* 2005;119:751–7.
160. Cummins S, Petticrew M, Sparks L, Findlay A. Large scale food retail interventions and diet. *BMJ* 2005;330:683–4.
161. Auchincloss AH, Riolo RL, Brown DG, Cook J, Diez Roux AV. An agent-based model of income inequalities in diet in the context of residential segregation. *Am J Prev Med* 2011;40:303–11.
162. Geist C, McManus PA. Geographical mobility over the life course: motivations and implications. *Popul Space Place* 2008;14:283–303.
163. Hernán MA, Robins JM. Instruments for causal inference: an epidemiologist's dream? *Epidemiology* 2006;17:360–72.
164. Chen SE, Florax RJ. Zoning for health: the obesity epidemic and opportunities for local policy intervention. *J Nutr* 2010;140:1181–4.
165. Bankes SC. Tools and techniques for developing policies for complex and uncertain systems. *Proc Natl Acad Sci USA* 2002;99 Suppl 3:7263–6.
166. Hammond RA. Complex systems modeling for obesity research. *Prev Chronic Dis* 2009;6:A97–106.
167. New York City Department of Health and Mental Hygiene. Healthy Bodegas initiative: 2010 report. New York: Department of Health and Mental Hygiene Center for Economic Opportunity; 2010.
168. Healthy Corner Store Initiative. The food trust, 2010 [cited 2011 Jul 25]. Available from: http://www.thefoodtrust.org/php/programs/corner_store.campaign.php.
169. Duffey KJ, Gordon-Larsen P, Shikany JM, Guilkey D, Jacobs DR Jr., Popkin BM. Food price and diet and health outcomes: 20 years of the CARDIA study. *Arch Intern Med* 2010;170:420–6.
170. Carter R, Moodie M, Markwick A, Magnus A, Vos T, Swinburn B, Haby MM. Assessing cost-effectiveness in obesity (ACE-obesity): an overview of the ACE approach, economic methods and cost results. *BMC Public Health* 2009;9:419–29.
171. Sallis JF, Story M, Lou D. Study designs and analytic strategies for environmental and policy research on obesity, physical activity, and diet: recommendations from a meeting of experts. *Am J Prev Med* 2009;36: S72–7.
172. Ogilvie D, Cummins S, Petticrew M, White M, Jones A, Wheeler K. Assessing the evaluability of complex public health interventions: five questions for researchers, funders, and policymakers. *Milbank Q* 2011;89:206–25.