

**HHS PUBLIC ACCESS**

Author manuscript

Gastroenterology. Author manuscript; available in PMC 2017 August 01.

Published in final edited form as:

Gastroenterology. 2012 August ; 143(2): 336–346. doi:10.1053/j.gastro.2012.04.032.**Consensus Statements for Management of Barrett's Dysplasia and Early-Stage Esophageal Adenocarcinoma, Based on a Delphi Process***A full list of authors and affiliations appears at the end of the article.***Abstract**

BACKGROUND & AIMS—Esophageal adenocarcinoma (EA) is increasingly common among patients with Barrett's esophagus (BE). We aimed to provide consensus recommendations based on the medical literature that clinicians could use to manage patients with BE and low-grade dysplasia, high-grade dysplasia (HGD), or early-stage EA.

Address requests for reprints to: Janusz A. Z. Jankowski, MSc, MBChB, MD, PhD, FRCP, FACP, AGAF, Centre for Digestive Diseases, Blizard Institute, Queen Mary, University of London, 4 Newark Street, Whitechapel, London EC1 2AT, UK. j.a.jankowski@qmul.ac.uk; fax: + 44 (0) 116 252 3294 or Paul Moayyedi, BSc, MbChB, PhD, MPH, FRCP (London), FRCPC, AGAF, FACP, Department of Medicine, McMaster University Medical Centre, 1200 Main Street West, HSC 4W8E, Hamilton, ON, L8N 3Z5, Canada. moayyep@mcmaster.ca; fax: (905) 518-8544.

Supplementary Material

Note: To access the supplementary material accompanying this article, visit the online version of *Gastroenterology* at www.gastrojournal.org, and at <http://dx.doi.org/10.1053/j.gastro.2012.04.032>.

Conflicts of interest

These authors disclose the following: Janusz Jankowski is a paid consultant to AstraZeneca UK and Amiral and a grant holder from FALK. He is Chief Investigator for the ASPeCT and CHoPIN trials, which are supported by AstraZeneca. Cathy Bennett is the proprietor of Systematic Research Ltd and received a consultancy fee for her work on this consensus document. Paul Moayyedi is a consultant to AstraZeneca. Nimish Vakil is a consultant to AstraZeneca, Takeda, Ironwood, Restech, and Orexo. Robert Ganz is the primary inventor and the cofounder of BARRX Medical, holds equity in the company, and serves as a paid consultant. Peter Kahrilas performs ad hoc consulting for AstraZeneca, Eisai, EndoGastric Solutions, and Ironwood, and serves on advisory boards for Torax and Reckitt Benckiser. Michio Hongo is a consultant to Abbott Japan, AstraZeneca Japan, AstellasPharma, Daiichi-Sankyo, Dainippon Sumitomo Pharma, Eisai, Kissei Pharmaceutical, Takeda Pharmaceutical, Scampo Pharma, and Zelia Pharmaceutical. Yvonne Romero is a consultant to AstraZeneca, Santarus, Takeda, Kala, Pfizer, and Aptalis. David Armstrong has received one or more of the following: educational and research grants, honoraria, consulting fees, and related travel expenses from Abbott Laboratories, AltanaPharma, AstraZeneca, Axcan, Eisai Limited, Gilead, Janssen Ortho Inc, Merck, NPS Pharmaceuticals, Nycomed, Olympus Canada Inc, Pentax Medical Inc, Pfizer, Proctor & Gamble, Schering-Plough, Shire Canada, Takeda Canada, Warner-Chilcott, and XenoPort Inc. Richard Sampliner received a BARRX research grant. Oliver Pech is a consultant to Hitachi Medical, Fujinon, Norgine, and AstraZeneca. Jaroslaw Regula is a consultant to Abbott, Astellas, AstraZeneca, Krka, MSD, Polpharma, Sandoz, and Warner-Chilcott. M. Brian Fennerty is a consultant to Aptalis, Oncoscope, and Meridian Bioscience. Nicholas Talley has had grant support from Falk, Forest, Janssen, and Takeda, has been a consultant for ARYx, Astellas, AstraZeneca, Boehringer Ingelheim, Care Capital, ConCERT, Edusa, Falk, Focus Medical Communications, Forest, Ironwood, Janssen, Johnson & Johnson, Meritage, NicOx, Novartis, Prometheus, Salix, Sanofi-Adventis, Shire, Tranzyme, Theravance, XenoPort, and Zeria, and is a key opinion leader for Doyen Medical Inc. John de Caestecker is Chair of ASPeCT Trial Management Group, which is AstraZeneca supported. Jacques Bergman is a consultant for Boston Scientific and has research support from BARRX Medical, Olympus, and Cook. Stephen Attwood is on the aspect trial management committee, which is AstraZeneca supported. JeanPaul Galmiche is a consultant and speaker for Given Imaging, Mauna Kea Technologies, Shire, Norgine, and Xenoport. His institution has received research grants from AstraZeneca, Janssen Cilag France, ADDEX, and Pentax. Laurence Lovat is on the Advisory Board of Ninepoint Medical and performed ad hoc consulting for Given Imaging and research support for Axcan Pharma, DUSA Pharmaceuticals, and BARRX. Peter Watson is a member of ASPeCT Trial Management Group, which is AstraZeneca sponsored. Kenneth Wang is a consultant to BARRX, Ironwood Pharma, CDX Diagnostics, Pinnacle Pharma, and CSA. David Johnston has received speaker's fees and support to attend educational meetings from AstraZeneca. Krish Raguath received research support, educational grants and speaker honoraria from Olympus Keymed, Cook Medical and BARRX Medical. Stuart Gittens is managing director of ECD solutions web data handling company. The remaining authors disclose no conflicts.

METHODS—We performed an international, multidisciplinary, systematic, evidence-based review of different management strategies for patients with BE and dysplasia or early-stage EA. We used a Delphi process to develop consensus statements. The results of literature searches were screened using a unique, interactive, Web-based data-sifting platform; we used 11,904 papers to inform the choice of statements selected. An a priori threshold of 80% agreement was used to establish consensus for each statement.

RESULTS—Eighty-one of the 91 statements achieved consensus despite generally low quality of evidence, including 8 clinical statements: (1) specimens from endoscopic resection are better than biopsies for staging lesions, (2) it is important to carefully map the size of the dysplastic areas, (3) patients that receive ablative or surgical therapy require endoscopic follow-up, (4) high-resolution endoscopy is necessary for accurate diagnosis, (5) endoscopic therapy for HGD is preferred to surveillance, (6) endoscopic therapy for HGD is preferred to surgery, (7) the combination of endoscopic resection and radiofrequency ablation is the most effective therapy, and (8) after endoscopic removal of lesions from patients with HGD, all areas of BE should be ablated.

CONCLUSIONS—We developed a data-sifting platform and used the Delphi process to create evidence-based consensus statements for the management of patients with BE and early-stage EA. This approach identified important clinical features of the diseases and areas for future studies.

Keywords

BADCAT; Esophageal Cancer; Treatment Strategy; Systematic Analysis

Barrett's esophagus (BE) is defined as the replacement of distal esophageal squamous mucosa with metaplastic columnar epithelium.¹ It occurs in 4% of patients undergoing an upper gastrointestinal endoscopy, and in 9% of men over 50 years of age.² BE is more common in developed countries, affecting 2% of the population, because it is strongly associated with gastroesophageal re-flux disease^{3,4} and this disease incidence is increasing in developing countries.⁵ The main concern with BE is the associated increased risk for esophageal adenocarcinoma (EA). EA is the fastest growing cause of cancer mortality,⁶ and it is estimated that patients with BE have at least a 20-fold increased risk of developing EA.⁷⁻⁹ Most guidelines^{10,11} recommend surveillance endoscopy every 2 to 5 years in patients with BE to detect early, treatable neoplasia and early signs of high-grade dysplasia (HGD). If progression to low-grade dysplasia (LGD), HGD, or EA can be detected early in its course, cancer can either be prevented or treated at a curable stage.^{12,13}

There is a lack of agreement concerning optimal management of dysplasia and early EA and, therefore, management practice patterns vary considerably among BE experts. The classification and recognition of dysplasia, both by endoscopy and histology, are variable among and within countries, and between some medical centers. There remains heterogeneity in the management of HGD/early EA throughout the world; the primary alternatives include managing HGD with surveillance alone, endoscopic therapy to remove HGD or early EA, or surgical resection of the BE (esophagectomy).¹⁴ Innovations have taken place in the endoscopic management of EA.¹⁵ In this rapidly changing field, a rational consensus approach to BE patients with LGD, HGD, and early EA is necessary to help inform the practicing clinician. Previously, several consensus papers have had some impact

on clinical management^{1,10} but have focused on BE in general; the focus of this guideline is LGD, HGD, and early EA.

Methods

The specific population under consideration consisted of adults aged 18 years or older with a diagnosis of BE plus LGD, HGD, or early EA, the latter being defined as intramucosal EA (T1m) or superficial submucosal EA (T1sm1). We used a Delphi process to develop consensus statements for LGD/HGD/early EA. This approach combines the principles of evidence-based medicine supported by systematic literature reviews with the use of an iterative anonymous voting process. This software program permitted anonymous individual feedback and changes of views during the process, together with controlled feedback of evidence regulated by the coordinator (CB) and the consensus chair (JJ). The Delphi process¹⁶ is now increasingly used in health care as a rigorous means of determining consensus in a defined clinical area^{1,17} and is reliable.¹⁸

The principal steps in the process were: (1) selection of the consensus group; (2) development of draft statements by panels; (3) systematic literature reviews to identify evidence to support each statement (search key words, Appendix 1); (4) 4 rounds of repeated anonymous voting on iterations of the statements (with feedback at each round) until consensus was reached (Appendix 2) (Figure 1); and (5) grading of the strength and quality of the evidence and strength of the recommendations using accepted criteria^{19,20} (Appendix 2). Details are listed in Appendix 3.²¹

Results

The initial stage was development of statements followed by a comprehensive literature review. Eventually, 4 in-person meetings followed by 4 rounds of consensus voting resulted in consensus (80% of respondents strongly agree or agree with reservation) being achieved in 81 of 91 statements. The respondents were asked to choose 1 of the following for each statement; agree strongly (A+), agree with reservation (A), undecided (U), disagree (D) or disagree strongly (D+). Although evidence-based explanations with key references were provided when relevant, it was the statement on which people voted. Consistent with principles of the Delphi process,¹⁹ the level of agreement increased with each round of voting (Figure 1). This high level of consensus was also exemplified by a post hoc analysis, where if >50% of respondents strongly agreed with the statement, it was accepted as a measure of agreement (Figure 1). Overall, the proportion of participants voting for each statement increased with each round of voting.

We selected 20 statements that represent the following key clinically relevant areas: diagnosis, epidemiology, methods of surveillance, approaches to treatment, and prevention of HGD and early adenocarcinoma in patients with BE. A description of any concerns about the statement is provided from the online comments of the respondents. We focused on HGD and early EA, as this area has the most evidence. All the remaining statements are outlined in Appendix 2.

Diagnosis of BE and HGD

Histologically, there is poor inter-observer agreement among pathologists in distinguishing HGD from intra-mucosal adenocarcinoma

Agreement: A+ 62%, A 33%, U 4%, D 1%, D+ 0%. Evidence: Moderate: The extent and severity of the dysplastic changes distinguishes HGD from LGD.²² Expert pathologists have found distinguishing HGD from intramucosal EA remains problematic.²³ The widely accepted definition of intramucosal EA is a lesion in which neoplastic cells have penetrated the basement membrane and invaded the lamina propria, but have not yet penetrated through the muscularis mucosae. However, reliable histologic recognition of lamina propria invasion is difficult due to the absence of objective and validated criteria. Kappa statistics for distinguishing between HGD and intramucosal EA vary between 0.21 and 0.47, suggesting poor, or at best, fair agreement.²³⁻²⁵

At least 2 experienced gastrointestinal pathologists should evaluate all Barrett's biopsies when a diagnosis of dysplasia is considered

Agreement: A+ 79%, A 15%, U 4%, D 1%, D+ 1%. Evidence: Moderate: It has long been recognized that there is inter-observer variability between pathologists in differentiating HGD from intramucosal EA as described here. Five studies²⁶⁻³⁰ have shown that the prediction of progression of esophageal dysplasia is improved if at least 2 expert pathologists agree on a diagnosis of dysplasia, and increases when more pathologists concur with the diagnosis.^{27,29,30}

Risk of Progression to Esophageal Adenocarcinoma

Non-goblet columnar metaplasia of the esophagus can progress to cancer, but the magnitude of risk is unknown

Agreement: A+ 59%, A 33%, U 6%, D 2%, D+ 0%. Evidence: Low: The US definition of BE requires that intestinal metaplasia is present in the salmon-colored esophageal columnar-lined mucosa of the tubular esophagus. There is, however, evidence that non-goblet columnar metaplasia of the distal esophagus shows biological features of intestinal differentiation, and possesses molecular abnormalities consistent with a risk of malignancy of neoplasia precursor lesions.³¹⁻³⁴ Two retrospective studies^{35,36} evaluated the risk of neoplasia in patients with columnar metaplasia of the esophagus either with or without goblet cells. There were 991 patients with intestinal metaplasia and 631 without intestinal metaplasia. The incidence of cancer progression from BE was similar in the 2 patient groups (4.5% vs 3.6% in one study³⁵ and 3.1% vs 3.2% in the other³⁶). Non-goblet cell columnar metaplasia has malignant potential, although the relative risk is unclear.

Extent of dysplasia can correlate with progression to cancer in BE

Agreement: A+ 52%, A 44%, U 4%, D+ 0%, D 0%; Evidence: Very low: The majority of participants agreed with this statement and 3 articles have evaluated whether the extent of dysplasia is a risk factor for EA in BE.³⁷⁻³⁹ Two studies^{37,38} (total of 177 patients) concluded that the extent of dysplasia was correlated with the risk of progression.^{37,38} However, one retrospective study³⁹ of 42 patients from a pathology database with BE and HGD who underwent esophagectomy failed to show a significant association between extent

of dysplasia and the risk of malignancy. Each of the 3 studies used different criteria and definitions for dysplasia.

Ulcers in BE that fail to heal with proton pump inhibitor therapy are a very suspicious finding and should be monitored closely for development of carcinoma

Agreement: A+ 66%, A 24%, U 10%, D 0%, D+ 0%. Evidence: Very

low: Unfortunately, there are no good case series on ulcerating lesions in BE that do not heal with proton pump inhibitor therapy,⁴⁰ although experts would suggest that BE-related ulcers are associated with malignancy.

Visible lumps or nodules consisting of HGD suggest a more advanced lesion with invasion might be present

Agreement: A+ 73%, A 26%, U 1%, D 0%, D+ 0%. Evidence: Low: Endoscopic mucosal resection (EMR/ER) of visible lumps with HGD on endoscopic biopsy results in upgrading the final diagnosis to cancer in 40% of cases.^{41,42} In a series of esophagectomies performed for presumed HGD identified by endoscopic biopsies, coexisting EA was found in 7 of 9 patients (78%) with a visible lesion and 7 of 22 patients (32%) without a visible lesion ($P = .02$).⁴³

Risk of progression from HGD to EA is approximately 10% per year (range 6%–19%)

Agreement: A+ 45%, A 40%, U 5%, D 6%, D+ 4%. Evidence: Low: This statement achieved consensus based on a systematic review,⁴⁴ which identified 4 studies^{45–48} involving 236 BE patients with HGD that suggested a conversion rate of 6% per year, contrasting with a large randomized controlled trial demonstrating conversion from HGD to EA of 19% in 1 year.⁴⁹ This risk estimate assumes that no endoscopic or surgical intervention takes place and that there are no macroscopically visible lesions. The issue of concomitant EA in patients who are diagnosed with BE and HGD is another consideration. In the absence of visible lesions in BE, the prevalence of EA in patients who underwent esophagectomy was 3%.^{50,51}

Methods of Surveillance for Patients With BE and With HGD

For evaluation of patients with BE, the use of high-resolution endoscopes and targeted biopsies of every suspicious lesion followed by 4-quadrant biopsies every 1–2 cm are recommended

Agreement: A+ 60%, A 38%, U 1%, D 0%, D+ 1%. Evidence: Very low: A high-resolution endoscope (>850,000 pixels) should be used to evaluate patients with BE. Standard-resolution endoscopes are not recommended, although there is scant scientific evidence for this recommendation. Evidence that greater resolution improves diagnosis is only available and supports narrow band imaging,⁵² but for chromoendoscopy there was no superiority to chromoendoscopy over standard endoscopy, although acetic acid spraying can improve visualization of lesions.^{53,54} Even with high-resolution endoscopes, 4-quadrant biopsies are still necessary after careful evaluation of the BE segment to exclude

synchronous neoplastic lesions. They should be performed with 4 biopsies at 1–2-cm intervals throughout the entire BE segment. There are no data demonstrating superiority of 1-cm intervals compared with 2-cm intervals.^{55,56}

Treatment of HGD and Early EA

Endoscopic treatment should be preferred over endoscopic surveillance for management of most BE patients with HGD/T1m Barrett's esophagus

Agreement: A+ 78%, A 19%, U 4%, D 0%, D+ 0%. Evidence: Moderate: There was strong agreement with this statement among the group. It is difficult to exclude EA complicating HGD based on biopsies only. Endoscopic surveillance can lead to under-diagnosis of cancer at baseline, especially when HGD is located in the area of BE that is endoscopically unremarkable.^{23,42} Endoscopic therapy (initially EMR for visible lesions) aimed at removing all BE mucosa should treat all areas of HGD and early EA that might have been missed by surveillance alone. Two randomized sham-controlled studies^{46,49} of ablation therapy (after initial EMR where appropriate) vs endoscopic surveillance have shown a significantly higher progression rate to cancer in the surveillance arm. Endoscopic treatment can cause complete remission of neoplasia in 80%–100% of cases and complete removal of BE with intestinal metaplasia in >75% of cases.^{40,49,57–60} Severe complications (such as bleeding, perforation, or stricture) are uncommon.^{40,49,57–60}

For patients with HGD in an endoscopically visible abnormality, endoscopic resection is essential for proper diagnosis and staging

Agreement: A+ 79%, A 16%, U 3%, D 1%, D+ 1%. Evidence: Moderate: EMR can lead to a significant change in diagnosis compared with a previous biopsy diagnosis.^{42,61–63} EMR provides a larger tissue specimen that is generally better orientated, allowing easier interpretation by pathologists.⁶² In addition, when an area of HGD is endoscopically visible, it is more likely to harbor EA.^{42,63,64} If EA is found in the EMR specimen, the risk of local lymph node metastasis has been shown to correlate with the depth of invasion,^{65,66} allowing better selection of therapy.^{67,68}

Endoscopic treatment should be preferred over surgical treatment for management of most patients with HGD in BE

Agreement: A+ 64%, A 29%, U 3%, D 2%, D+ 2%. Evidence: Low: There was strong consensus for this approach. HGD in BE is rarely associated with lymph node involvement, provided that deeper invasion has been ruled out by EMR (as described in statement 10).^{57,58,69,70} Two case series^{40,57,58} reported that survival after EMR was high, similar to that expected in a surgical cohort. One cohort study⁷¹ reported that the disease-specific survival rate after endoscopic treatment was not different from surgical therapy. The case series reported a lower morbidity than might be expected in surgical patients.^{40,57,58} Endoscopic treatment is associated with a higher rate of HGD recurrence,^{40,57,58,71} although this can usually be treated endoscopically.^{40,57,58,72} Finally, on the rare occasion that endoscopic treatment fails, surgical resection is still possible and generally curative.^{40,57,58}

Widespread EMR can cause strictures (especially when more than two thirds of the circumference is removed)

Agreement: A+ 74%, A 21%, U 4%, D 1%, D+ 0%. Evidence: Low: The intention of EMR/ER) should be to remove all visible dysplasia. It should ideally be restricted to less than two thirds of the esophageal circumference in order to reduce the risk of strictures, but all visible lesions should be resected. Strictures resulting from EMR respond well to dilation.^{73–75}

Endoscopic treatment of HGD/T1m should only be performed in tertiary referral care centers after proper training of the endoscopists and pathologists involved

Agreement: A+ 57.5%, A 34%, U 2.5%, D 6%, D+ 0%. Evidence: Very low: There are no studies that have shown that centers with expertise, or those that have high case volumes, provide better quality care for BE patients with HGD/early EA. The consensus group voted positively for this statement because in other areas of gastroenterology, expertise and case volumes are associated with better outcomes.^{76,77} Adequate management of these patients encompasses a wide range of experience, equipment, and a certain case volume (which we arbitrarily defined as >10 cases per year).^{78–82}

After EMR has removed visible lesions with HGD/T1m, the remaining BE segment should be eradicated regardless of whether or not it includes the presence or absence of dysplasia

Agreement: A+ 54%, A 30%, U 13%, D 3%, D+ 0%. Evidence: Very low: Statement 10 recommended EMR for visible abnormalities with HGD. If EMR is the only modality that is used and the remaining BE mucosa is left untreated, case series have reported recurrence of neoplasia. Rates vary from 11% to 30% (mean follow-up of 3 years).^{57,83} Ablation of the remaining BE is associated with a lower recurrence rate.^{40,49,59,60,84,85}

Radiofrequency ablation is currently the best available ablation technique for treatment of flat HGD and for eradication of residual BE mucosa after focal EMR

Agreement: A+ 59%, A 25%, U 11%, D 1%, D+ 4%. Evidence: Low: Statement 14 recommended endoscopic ablation of BE after EMR for visible lesions. The question remains, what is the most appropriate endoscopic technique? The alternatives that have been most frequently studied include photodynamic therapy, radiofrequency ablation (RFA), and/or stepwise EMR of all BE. A systematic review⁸⁶ of photodynamic therapy for HGD of BE mucosa esophagus suggests that this approach reduces the risk of progression to cancer compared with surveillance alone.^{46,87} However, complications remain a problem with this technique,⁴⁶ and HGD dysplasia persists in 33%–50% of patients.^{87,88} Other therapeutic modalities include cryotherapy and argon plasma coagulation. Cryotherapy has not been evaluated in randomized controlled trials and argon plasma coagulations has only been reported in small randomized controlled trials, although there are anecdotal high-success rates.⁸⁹ One systematic review⁹⁰ suggests that success rates with RFA are superior, with

approximately 90% of patients having no HGD after therapy and this seems to be maintained.^{49,59,91–93}

In patients with superficial submucosal cancer in BE and low-risk characteristics (invasion <500 μm; G1–G2 cancers, no lymph–vascular invasion), endoscopic treatment is a valid alternative to esophagectomy

Agreement: A+ 41%, A 35%, U 8%, D 11%, D+ 5%. Evidence: Very low: This statement failed to achieve consensus. The paradigm comes from studies from Japan on early gastric cancer, which have shown that well to moderately differentiated cancers that invade into the submucosa <500 μm and have no lymphovascular invasion, have virtually no risk of lymph node metastases.⁹⁴ Furthermore, in a prospective series of 21 BE patients meeting these low-risk criteria,⁹⁵ no lymph node metastases were found in any of the patients after a median follow-up period of 62 months. The implications of lymph node spread are so important that more data are needed before this statement can be supported.

Successful surgery/intervention for early cancer can be determined by long-term (5 years or longer) survival

Agreement: A+ 73%, A 23%, U 3%, D 1%, D+ 0%. Evidence: Very low: The group reached consensus that successful surgery is determined by 5-year survival. However, most patients with HGD should receive EMR and/or RFA because it is safer and carries a similar efficacy rate,⁹⁶ although more studies are needed (see statements 11, 14, and 15). Surgery is still considered the treatment of choice for early EA that has extended into the submucosa.^{97,98} Case series suggest that the 5-year survival rates range from 80% to 90%.^{99,100}

Reported operative mortality rate for esophagectomy for HGD and T1m generally ranges from 0% to 4%, with a mean overall operative mortality of 2%

Agreement: A+ 65%, A 30%, U 3%, D 1%, D+ 1%. Evidence: Very low: Operative mortality for patients undergoing esophagectomy for HGD or early EA is difficult to generalize because data are primarily from self-selected high-volume centers and analysis is retrospective. We identified 10 case series^{71,98,101–108} evaluating a total of 567 HGD or early EA patients (Table 1). Operative mortality rate for esophagectomy for HGD and early EA ranges from 0% to 4%, with an overall operative 30-day mortality rate of approximately 2%.¹⁰¹

Operative mortality is improved if surgery is undertaken in specialist surgical centers

Agreement: A+ 90%, A 8%, U 2%, D 0%, D+ 0%. Evidence: Moderate: In contrast to the evidence for endoscopic therapy (statement 13), there are good observational data to support the performance of esophageal surgery in specialist centers for treatment of EA. Results for individual surgeons improve with experience¹⁰⁹ and patient outcomes have consistently been shown to be better in high-volume centers.^{110–113}

After eradication of HGD by endoscopic therapy or surgery, endoscopic follow-up is required

Agreement: A+ 72.5%, A 20%, U 5%, D 0%, D+ 2.5%. Evidence: Very low: There are 2^{114,115} surgical follow-up series involving 57 BE patients that support this statement. Both studies report that new BE occur occurs after curative subtotal esophagectomy with gastric conduit reconstruction for either EA, squamous cell carcinoma, or HGD.^{114,115} Development of BE occurs in half of patients studied^{114,115} and can recur from 6 months or less after surgery¹¹⁴ to 10 years after surgery.^{115,115} The risk of developing dysplasia or malignancy in the “neosquamous” epithelium is unknown, but goblet cells are detected with increasing frequency as follow-up continues.¹¹⁵ Based on available evidence, a suggested strategy for post-esophagectomy surveillance is to perform screening endoscopy at 2, 5, and 10 years after surgery. If the risk of dysplasia is assumed to be similar to patients with de novo BE, it is reasonable to recommend every 2-year surveillance endoscopies once BE has been detected.¹¹⁶ The surveillance interval for patients that have BE ablated with RFA is unclear, but a 5-year follow-up study evaluated patients every 2.5 years without any recurrence of dysplasia and a low recurrence of BE.⁹¹

Discussion

We focused on statements concerning HGD and EA as evidence relating to LGD is particularly weak. The management of HGD and EA of the esophagus is heterogeneous and the clinician’s perception of the available evidence is one major determinant of this variation in practice. The relatively poor quality of data relating to dysplasia in BE is emphasized by 46 statements having a very low or low level and 38 having moderate or high levels of evidence. However, in many cases, it is unlikely that large, well-designed randomized trials will ever be done and in this information vacuum there is a need for an authoritative consensus on areas where there is good agreement. Our multidisciplinary international group has developed consensus to help the practicing clinician with the diagnosis and management of HGD and early EA in BE. We focused on patient populations with high-risk disease rather than including those statements about LGD, a condition for which there are even less objective data in the literature.

The literature search technique used for this consensus process was unique in a number of ways. It was much more inclusive than more focused searches, and permitted inclusion of additional articles during the consensus process that might have been missed during initial searches. Before including articles for citation, the articles were reviewed by panel members and a panel chair and were ultimately reviewed and graded by a single senior author, resulting in consistency in assessment of the evidence. This mechanism resulted in the largest number of articles ever captured in a literature review for gastrointestinal diseases. We found that the overall quality of evidence related to the statements was low.

The consensus process resulted in a high level of consensus for most statements, which suggests that many results are appropriate for clinical application at this time. The relationship of highly relevant clinically applicable consensus findings regarding EMR is appreciable. First, EMR provides better staging for visible lesions than do biopsies alone. Second, careful mapping of the size of the dysplastic areas by EMR is important to assess

the prognosis and risk of progression. Third, EMR combined with RFA is the most proven ablative therapy for visible HGD and for ablation of BE in patients with HGD (Figure 2). HGD should be managed by RFA with or without EMR, and surgery can be considered for early EA.

In defining early cancer, we chose T1sm1 as being the extent of early cancer, as beyond this point metastases increases from ~1% to >10% for T1sm2. Including T1sm1 could be controversial, but if low-risk sm1 (differentiation grades 1 and 2, without lymphovascular invasion and with a negative deep resection margin) tumors are selected, they might be more amenable to successful endoscopic therapy. We recognize that evidence from larger series is still required to conclude that sm1 are to be considered amenable to endoscopic therapy. Using a multidisciplinary approach, surgical treatment should still be considered for early cancer (as opposed to HGD) for all patients fit for surgery.

The consensus process also identified several areas where urgent research is needed (Appendix 2), including evaluation of genetic markers to determine cancer risk.^{117,118} Determining the true risk of progression from dysplasia to EA has implications in term of the cost effectiveness of surveillance strategies and impacts on the provision of effective treatments. There are no randomized controlled trials, but large epidemiology studies^{119–121} and one meta-analysis¹²² have reinforced the low conversion rate to cancer for nondysplastic Barrett's^{123,124} and short Barrett's.^{120,121}

There are a number of potential shortcomings of this study. First, some geographical areas were under-represented. We did not use meta-analysis techniques in a more rigorous approach to evaluating the literature, as we believed that the relevant literature was relatively scant in quality (even though 11,000 articles were assessed) and diverse in approaches and reporting styles, both of which would have severely limited the applicability of these techniques to our process. Finally, a template was not used to standardize comments for statements, which might have resulted in some unevenness in the presentation of clinical view points.

This work represents the most far-reaching, inclusive, and informative consensus process on evaluation and management of BE with HGD/early cancer published to date. Most of the findings are clinically relevant and the high degree of consensus achieved for most of the questions indicates that many of the statements are appropriate for immediate use in guiding clinical activity. In addition, areas in which consensus was not achieved are identified, helping to guide areas in which future clinical research is likely to be productive (Table 2).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Authors

CATHY BENNETT¹, NIMISH VAKIL², JACQUES BERGMAN³, REBECCA HARRISON⁴, ROBERT ODZE⁵, MICHAEL VIETH⁶, SCOTT SANDERS⁷, LAURA GAY⁸, OLIVER PECH⁶, GAIUS LONGCROFT–WHEATON⁹, YVONNE ROMERO¹⁰,

JOHN INADOMI¹¹, JAN TACK¹², DOUGLAS A. CORLEY¹³, HENDRIK MANNER¹⁴, SUSI GREEN⁹, DAVID AL DULAIMI¹⁵, HAYTHEM ALI¹⁶, BILL ALLUM¹⁷, MARK ANDERSON¹⁸, HOWARD CURTIS¹⁹, GARY FALK²⁰, M. BRIAN FENNERTY²¹, GRANT FULLARTON²², KAUSILIA KRISHNADATH³, STEPHEN J. MELTZER²³, DAVID ARMSTRONG²⁴, ROBERT GANZ²⁵, GIANPAOLO CENGIA²⁶, JAMES J. GOING²², JOHN GOLDBLUM²⁷, CHARLES GORDON²⁸, HEIKE GRABSCH³⁰, CHRIS HAIGH³¹, MICHIO HONGO³², DAVID JOHNSTON³³, RICKY FORBES–YOUNG³⁴, ELAINE KAY³⁵, PHILIP KAYE³⁶, TONI LERUT¹², LAURENCE B. LOVAT³⁷, LARS LUNDELL³⁸, PHILIP MAIRS³⁹, TADAKUZA SHIMODA⁴⁰, STUART SPECHLER⁴¹, STEPHEN SONTAG⁴², PETER MALFERTHEINER⁴³, IAIN MURRAY⁴⁴, MANOJ NANJI⁸, DAVID POLLER⁹, KRISH RAGUNATH³⁶, JAROSLAW REGULA⁴⁵, RENZO CESTARI²⁶, NEIL SHEPHERD⁴⁶, RAJVINDER SINGH⁴⁷, HUBERT J. STEIN⁴⁸, NICHOLAS J. TALLEY⁴⁹, JEAN–PAUL GALMICHE⁵⁰, TONY C. K. THAM⁵¹, PETER WATSON¹, LISA YERIAN²⁷, MASSIMO RUGGE²⁹, THOMAS W. RICE²⁷, JOHN HART⁵², STUART GITTENS⁵³, DAVID HEWIN⁴⁶, JUERGEN HOCHBERGER⁵⁴, PETER KAHRILAS⁵⁵, SEAN PRESTON⁵⁶, RICHARD SAMPLINER⁵⁷, PRATEEK SHARMA⁵⁸, ROBERT STUART⁵⁹, KENNETH WANG¹⁰, IRVING WAXMAN⁵², CHRIS ABLEY⁴, DUNCAN LOFT⁶⁰, IAN PENMAN³⁴, NICHOLAS J. SHAHEEN⁶¹, AMITABH CHAK⁶², GARETH DAVIES⁶³, LORNA DUNN⁶⁴, YNGVE FALCK–YTTER⁶⁵, JOHN DECAESTECKER⁴, PRADEEP BHANDARI⁹, CHRISTIAN ELL⁶, S. MICHAEL GRIFFIN⁶⁴, STEPHEN ATTWOOD⁶⁶, HUGH BARR⁴⁶, JOHN ALLEN⁶⁷, MARK K. FERGUSON⁵², PAUL MOAYYEDI²⁴, and JANUSZ A. Z. JANKOWSKI^{4,8,68}

Affiliations

¹Queens University, Belfast, UK ²University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin ³Amsterdam Medical Center, Amsterdam, The Netherlands ⁴University Hospitals of Leicester, Leicester, UK ⁵Harvard Medical School, Boston, Massachusetts ⁶Klinikum Bayreuth, Bayreuth, Germany ⁷Warwick Medical School, Coventry, UK ⁸Queen Mary University London, London, UK ⁹Queen Alexandra Hospital, Portsmouth, UK ¹⁰Mayo Clinic, Rochester, Minnesota ¹¹Washington University, Seattle, Washington ¹²Leuven University, Leuven, Belgium ¹³Kaiser Permanente, San Francisco, California ¹⁴HSK Hospital, Wiesbaden, Germany ¹⁵Redditch General Hospital, Redditch, UK ¹⁶Maidstone and Tunbridge Wells NHS trust, Maidstone, UK ¹⁷Royal Marsden Hospital, London, UK ¹⁸City Hospital, Birmingham, UK and Sandwell Hospital, West Midlands, UK ¹⁹Queen Marys Hospital, Sidcup, UK ²⁰University of Pennsylvania, Philadelphia, Pennsylvania ²¹Oregon Health & Science University, Portland, Oregon ²²Royal Infirmary, Glasgow, UK ²³The Johns Hopkins University School of Medicine, Baltimore, Maryland ²⁴McMaster University, Hamilton, Ontario, Canada ²⁵Bloomington Medical Centre, Bloomington, Minnesota ²⁶University of Brescia, Brescia, Italy ²⁷Anatomic Pathology, The Cleveland Clinic, Cleveland, Ohio ²⁸The Royal Bournemouth Hospital, Bournemouth, UK ²⁹University of Padova, Padua, Italy ³⁰University of Leeds, Leeds, UK ³¹Wansbeck General Hospital, Northumbria, UK ³²Tohoku University Hospital, Tohoku, Japan ³³Ninewells Hospital, Dundee, UK

³⁴Royal Infirmary, Edinburgh, UK ³⁵Trinity College, Dublin, Ireland ³⁶Digestive Diseases Centre, Nottingham University Hospital, Nottingham, UK ³⁷University College London, London, UK ³⁸Karolinska Institutet, CLINTEC, Stockholm, Sweden ³⁹Darrent Valley Hospital, Kent, UK ⁴⁰National Cancer Center, Tokyo, Japan ⁴¹University of Dallas, Dallas, Texas ⁴²Hines Veterans Affairs Hospital, Chicago, Illinois ⁴³Magdeburg University, Magdeburg, Germany ⁴⁴Royal Cornwall Hospitals, Truro, UK ⁴⁵Institute of Oncology, Warsaw, Poland ⁴⁶Gloucestershire Royal Hospitals, Gloucestershire, UK ⁴⁷Lyell McEwin Hospital, University of Adelaide, Adelaide, Australia ⁴⁸Polyclinic Klinikum München, München, Germany ⁴⁹University of Newcastle, Newcastle, Australia ⁵⁰Department of Gastroenterology, CHU and University of Nantes, Nantes, France ⁵¹Ulster Hospital, Belfast, UK ⁵²University of Chicago, Chicago, Illinois ⁵³ECD Solutions, PO Box 862, Bridgetown, St. Michael, Barbados ⁵⁴St. Bernward Hospital, Hildesheim, Germany ⁵⁵Northwestern University, Chicago, Illinois ⁵⁶Barts Health NHS Trust, London, UK ⁵⁷University of Arizona, Tucson, Arizona ⁵⁸Veterans Affairs Medical Center and University of Kansas ⁵⁹Oesophageal Cancer Fund, Dublin, Ireland ⁶⁰Walsgrave Hospital, Coventry, UK ⁶¹University of North Carolina School of Medicine, Chapel Hill, North Carolina ⁶²Case Western Reserve University School of Medicine, Cleveland, Ohio ⁶³Harrogate District Hospital, Harrogate, UK ⁶⁴Northern Oesophagogastric Cancer Unit Royal Victoria Infirmary, Newcastle upon Tyne, UK ⁶⁵Louis Stokes Medical Center, Cleveland, Ohio ⁶⁶Durham University, Durham, UK ⁶⁷University of Minnesota School of Medicine, Minneapolis, Minnesota ⁶⁸University of Oxford, Oxford, UK

Acknowledgments

We thank Marion Lawlor of University Hospitals of Leicester, National Health Service Trust for administrative support and accounts. In addition, others provided essential core support, including Jan Lilleyman (administration). We would also like to thank the various funders for their contributions, which enabled this process to occur completely independently of pharmaceutical support. Literature searches were designed by LUCID Health, University of Leeds, UK (Pat Spoor and Ros Dunlevey). Peer review comments were offered by Hanna Lewin (National Institute for Health and Clinical Excellence), Sara Clarke (National Health Service evidence–gastroenterology and liver diseases), Karin Dearness (Cochrane UGPD group), and Iris Gordon (Cochrane Collaboration Eyes and Vision Group).

A total of 104 people were approached, of whom 95 took active part in the process, 92 completed their conflict of interest forms. Three contributors of the 95 active participants did not complete a declaration of conflicts of interest at any time and are not authors, 9 respondents took no part in the process or withdrew at an early stage (4 from Europe, 3 from the United States, 1 from Australia and 1 from Asia).

Funding

Funding has been received from the International Society of Diseases of the Esophagus (\$3500), British Society of Gastroenterology (£2500), American College of Gastroenterology (\$2000), American Gastroenterological Association (\$2000), American Society for Gastrointestinal Endoscopy (\$2000), Association of Upper Gastrointestinal Surgeons (£1000), Fight Oesophageal Reflux Together (£1000), German Society of Endoscopy (€2000), Netherlands Association of Hepatogastroenterologists (€100), and Oesophageal Cancer Fund of Ireland (€3000).

Abbreviations used in this paperd

BAD CAT	Barrett's dysplasia and cancer task force
BE	Barrett's esophagus
EA	esophageal adenocarcinoma
EMR	endoscopic mucosal resection
HGD	high-grade dysplasia
LGD	low-grade dysplasia
RFA	radiofrequency ablation

References

1. Vakil NJ, van Zanten SV, Kahrilas P, et al. The Montreal definition and classification of gastroesophageal reflux disease: a global evidence-based consensus. *Am J Gastroenterol.* 2006; 101:1900–1920. [PubMed: 16928254]
2. Ford AC, Forman D, Reynolds PD, et al. Ethnicity, gender, and socioeconomic status as risk factors for esophagitis and Barrett's esophagus. *Am J Epidemiol.* 2005; 162:454–460. [PubMed: 16076833]
3. Taylor JB, Rubenstein JH. Meta-analyses of the effect of symptoms of gastroesophageal reflux on the risk of Barrett's esophagus. *Am J Gastroenterol.* 2010; 105:1730–1737.
4. Ronkainen J, Talley NJ, Storskrubb T, et al. Erosive esophagitis is a risk factor for Barrett's esophagus: a community-based endoscopic follow-up study. *Am J Gastroenterol.* 2011; 106:1946–1952. [PubMed: 21946284]
5. Moayyedi P, Axon AT. Review article: gastro-esophageal reflux disease—the extent of the problem. *Aliment Pharmacol Ther.* 2005; 22(Suppl 1):11–19.
6. National Cancer Institute. [Accessed May 15, 2011] Fast stats: esophagus cancer. 2008. Available from <http://seer.cancer.gov/faststats/selections.php>
7. Jung KW, Talley NJ, Romero Y, et al. Epidemiology and natural history of intestinal metaplasia of the gastroesophageal junction and Barrett's esophagus: a population-based study. *Am J Gastroenterol.* 2011; 106:1447–1455. [PubMed: 21483461]
8. Hvid-Jensen F, Pedersen L, Drewes AM, et al. Incidence of adenocarcinoma among patients with Barrett's esophagus. *N Engl J Med.* 2011; 365:1375–1383. [PubMed: 21995385]
9. Solaymani-Dodaran M, Logan RF, West J, et al. Risk of oesophageal cancer in Barrett's oesophagus and gastro-oesophageal reflux. *Gut.* 2004; 53:1070–1074. [PubMed: 15247170]
10. Spechler SJ, Sharma P, Souza RF, et al. American Gastroenterological Association medical position statement on the management of Barrett's esophagus. *Gastroenterology.* 2011; 140:1084–1091. [PubMed: 21376940]
11. Wang K, Sampliner R. Updated guidelines 2008 for the diagnosis, surveillance and therapy of Barrett's esophagus. *Am J Gastroenterol.* 2008; 103:788–797. [PubMed: 18341497]
12. Spechler SJ, Sharma P, Souza RF, et al. American Gastroenterological Association technical review on the management of Barrett's esophagus. *Gastroenterology.* 2011; 140:e18–e52. [PubMed: 21376939]
13. Prasad GA, Bansal A, Sharma P, et al. Predictors of progression in Barrett's esophagus: current knowledge and future directions. *Am J Gastroenterol.* 2010; 105:1490–1502. [PubMed: 20104216]
14. Spechler SJ. Barrett's esophagus: clinical issues. *Gastrointest Endosc Clin N Am.* 2011; 21:1–7. [PubMed: 21112493]
15. Dent J. Barrett's esophagus: a historical perspective, an update on core practicalities and predictions on future evolutions of management. *J Gastroenterol Hepatol.* 2011; 26(Suppl 1):11–30. [PubMed: 21199510]

16. Evans C. The use of consensus methods and expert panels in pharmacoeconomic studies. Practical applications and methodological shortcomings. *Pharmacoeconomics*. 1997; 12:121–129. [PubMed: 10169665]
17. Powell C. The Delphi technique: myths and realities. *J Adv Nurs*. 2003; 41:376–382. [PubMed: 12581103]
18. Murphy MK, Black NA, Lamping DL, et al. Consensus development methods, and their use in clinical guideline development. *Health Technol Assess*. 1998; 2(3)
19. Schunemann HJ, Best D, Vist G, et al. Letters, numbers, symbols and words: how to communicate grades of evidence and recommendations. *CMAJ*. 2003; 169:677–680. [PubMed: 14517128]
20. Guyatt G, Gutterman D, Baumann MH, et al. Grading strength of recommendations and quality of evidence in clinical guidelines: report from an American College of Chest Physicians Task Force. *Chest*. 2006; 129:174–181. [PubMed: 16424429]
21. Talley NJ, Abreu MT, Achkar JP, et al. An evidence-based systematic review on medical therapies in inflammatory bowel disease. *Am J Gastroenterol*. 2011; 106(Suppl 1):S2–S25. [PubMed: 21472012]
22. Schlemper RJ, Riddell RH, Kato Y, et al. Vienna classification of gastrointestinal epithelial neoplasia. *Gut*. 2000; 47:251–255. [PubMed: 10896917]
23. Ormsby AH, Petras RE, Henricks WH, et al. Observer variation in the diagnosis of superficial oesophageal adenocarcinoma. *Gut*. 2002; 51:671–676. [PubMed: 12377805]
24. Montgomery E, Bronner MP, Goldblum JR, et al. Reproducibility of the diagnosis of dysplasia in Barrett esophagus: a reaffirmation. *Hum Pathol*. 2001; 32:368–378. [PubMed: 11331953]
25. Downs-Kelly E, Mendelin JE, Bennett AE, et al. Poor interobserver agreement in the distinction of high-grade dysplasia and adenocarcinoma in pretreatment Barrett's esophagus biopsies. *Am J Gastroenterol*. 2008; 103:2333–2340. [PubMed: 18671819]
26. Reid BJ, Haggitt RC, Rubin CE, et al. Observer variation in the diagnosis of dysplasia in Barrett's esophagus. *Hum Pathol*. 1988; 19:166–178. [PubMed: 3343032]
27. Skacel M, Petras RE, Gramlich TL, et al. The diagnosis of low grade dysplasia in Barrett's esophagus and its implication for disease progression. *Am J Gastroenterol*. 2000; 95:3383–3387. [PubMed: 11151865]
28. Montgomery E, Goldblum JR, Greenson JK, et al. Dysplasia as a predictive marker for invasive carcinoma in Barrett esophagus: a follow-up study based on 138 cases from a diagnostic variability study. *Hum Pathol*. 2001; 32:379–388. [PubMed: 11331954]
29. Kaye PV, Haider SA, Ilyas M, et al. Barrett's dysplasia and the Vienna classification: reproducibility, prediction of progression and impact of consensus reporting and p53 immunohistochemistry. *Histopathology*. 2009; 54:699–712. [PubMed: 19438745]
30. Curvers W, ten Kate F, Krishnadath K, et al. Low-grade dysplasia in Barrett's esophagus: overdiagnosed and underestimated. *Am J Gastroenterol*. 2010; 105:1523–1530. [PubMed: 20461069]
31. Riddell RH, Odze RD. Definition of Barrett's esophagus: time for a rethink-is intestinal metaplasia dead? *Am J Gastroenterol*. 2009; 104:2588–2594. [PubMed: 19623166]
32. Chaves P, Crespo M, Ribeiro C, et al. Chromosomal analysis of Barrett's cells: demonstration of instability and detection of the metaplastic lineage involved. *Mod Pathol*. 2007; 20:788–796. [PubMed: 17529926]
33. Romagnoli S, Roncalli M, Graziani D, et al. Molecular alterations of Barrett's esophagus on microdissected endoscopic biopsies. *Lab Invest*. 2001; 81:241–247. [PubMed: 11232646]
34. Liu W, Hahn H, Odze RD, et al. Metaplastic esophageal columnar epithelium without goblet cells shows DNA content abnormalities similar to goblet cell containing epithelium. *Am J Gastroenterol*. 2009; 104:816–824. [PubMed: 19293780]
35. Kelty CJ, Gough MD, Van Wyk Q, et al. Barrett's oesophagus: intestinal metaplasia is not essential for cancer risk. *Scand J Gastroenterol*. 2007; 42:1271–1274. [PubMed: 17852872]
36. Gatenby PA, Ramus JR, Caygill CP, et al. Relevance of the detection of intestinal metaplasia in non-dysplastic columnar-lined oesophagus. *Scand J Gastroenterol*. 2008; 43:524–530. [PubMed: 18415743]

37. Srivastava A, Hornick JL, Li X, et al. Extent of low-grade dysplasia is a risk factor for the development of esophageal adenocarcinoma in Barrett's esophagus. *Am J Gastroenterol.* 2007; 102:483–493. [PubMed: 17338734]
38. Buttar NS, Wang KK, Sebo TJ, et al. Extent of high-grade dysplasia in Barrett's esophagus correlates with risk of adenocarcinoma. *Gastroenterology.* 2001; 120:1630–1639. [PubMed: 11375945]
39. Dar MS, Goldblum JR, Rice TW, et al. Can extent of high grade dysplasia in Barrett's oesophagus predict the presence of adenocarcinoma at oesophagectomy? *Gut.* 2003; 52:486–489. [PubMed: 12631655]
40. Pech O, Behrens A, May A, et al. Long-term results and risk factor analysis for recurrence after curative endoscopic therapy in 349 patients with high-grade intraepithelial neoplasia and mucosal adenocarcinoma in Barrett's oesophagus. *Gut.* 2008; 57:1200–1206. [PubMed: 18460553]
41. Pech O, Gossner L, Manner H, et al. Prospective evaluation of the macroscopic types and location of early Barrett's neoplasia in 380 lesions. *Endoscopy.* 2007; 39:588–593. [PubMed: 17611912]
42. Peters FP, Brakenhoff KP, Curvers WL, et al. Histologic evaluation of resection specimens obtained at 293 endoscopic resections in Barrett's esophagus. *Gastrointest Endosc.* 2008; 67:604–609. [PubMed: 18155214]
43. Tharavej C, Hagen JA, Peters JH, et al. Predictive factors of coexisting cancer in Barrett's high-grade dysplasia. *Surg Endosc.* 2006; 20:439–443. [PubMed: 16437272]
44. Rastogi A, Puli S, El-Serag HB, et al. Incidence of esophageal adenocarcinoma in patients with Barrett's esophagus and high-grade dysplasia: a meta-analysis. *Gastrointest Endosc.* 2008; 67:394–398. [PubMed: 18045592]
45. Schnell TG, Sontag SJ, Chejfec G, et al. Long-term nonsurgical management of Barrett's esophagus with high-grade dysplasia. *Gastroenterology.* 2001; 120:1607–1619. [PubMed: 11375943]
46. Overholt BF, Lightdale CJ, Wang KK, et al. Photodynamic therapy with porfimer sodium for ablation of high-grade dysplasia in Barrett's esophagus: international, partially blinded, randomized phase III trial. *Gastrointest Endosc.* 2005; 62:488–498. [PubMed: 16185958]
47. Reid BJ, Levine DS, Longton G, et al. Predictors of progression to cancer in Barrett's esophagus: baseline histology and flow cy-tometry identify low- and high-risk patient subsets. *Am J Gastroenterol.* 2000; 95:1669–1676. [PubMed: 10925966]
48. Weston AP, Sharma P, Topalovski M, et al. Long-term follow-up of Barrett's high-grade dysplasia. *Am J Gastroenterol.* 2000; 95:1888–1893. [PubMed: 10950031]
49. Shaheen NJ, Sharma P, Overholt BF, et al. Radiofrequency ablation in Barrett's esophagus with dysplasia. *N Engl J Med.* 2009; 360:2277–2288. [PubMed: 19474425]
50. Wani S, Puli SR, Shaheen NJ, et al. Esophageal adenocarcinoma in Barrett's esophagus after endoscopic ablative therapy: a meta-analysis and systematic review. *Am J Gastroenterol.* 2009; 104:502–513. [PubMed: 19174812]
51. Wang VS, Hornick JL, Sepulveda JA, et al. Low prevalence of submucosal invasive carcinoma at esophagectomy for high-grade dysplasia or intramucosal adenocarcinoma in Barrett's esophagus: a 20-year experience. *Gastrointest Endosc.* 2009; 69:777–783. [PubMed: 19136106]
52. Wolfsen HC, Crook JE, Krishna M, et al. Prospective, controlled tandem endoscopy study of narrow band imaging for dysplasia detection in Barrett's Esophagus. *Gastroenterology.* 2008; 135:24–31. [PubMed: 18442484]
53. Pohl J, May A, Rabenstein T, et al. Comparison of computed virtual chromoendoscopy and conventional chromoendoscopy with acetic acid for detection of neoplasia in Barrett's esophagus. *Endoscopy.* 2007; 39:594–598. [PubMed: 17611913]
54. Longcroft-Wheaton G, Duku M, Mead R, et al. Acetic acid spray is an effective tool for the endoscopic detection of neoplasia in patients with Barrett's esophagus. *Clin Gastroenterol Hepatol.* 2010; 8:843–847. [PubMed: 20601133]
55. Peters FP, Curvers WL, Rosmolen WD, et al. Surveillance history of endoscopically treated patients with early Barrett's neoplasia: nonadherence to the Seattle biopsy protocol leads to sampling error. *Dis Esophagus.* 2008; 21:475–479. [PubMed: 18430186]

56. Kariv R, Plesec TP, Goldblum JR, et al. The Seattle protocol does not more reliably predict the detection of cancer at the time of esophagectomy than a less intensive surveillance protocol. *Clin Gastroenterol Hepatol*. 2009; 7:653–658. [PubMed: 19264576]
57. Peters FP, Kara MA, Rosmolen WD, et al. Endoscopic treatment of high-grade dysplasia and early stage cancer in Barrett's esophagus. *Gastrointest Endosc*. 2005; 61:506–514. [PubMed: 15812401]
58. Ell C, May A, Pech O, et al. Curative endoscopic resection of early esophageal adenocarcinomas (Barrett's cancer). *Gastrointest Endosc*. 2007; 65:3–10. [PubMed: 17185072]
59. Pouw RE, Wirths K, Eisendrath P, et al. Efficacy of radiofrequency ablation combined with endoscopic resection for Barrett's esophagus with early neoplasia. *Clin Gastroenterol Hepatol*. 2010; 8:23–29. [PubMed: 19602454]
60. Pouw RE, Seewald S, Gondrie JJ, et al. Stepwise radical endoscopic resection for eradication of Barrett's oesophagus with early neoplasia in a cohort of 169 patients. *Gut*. 2010; 59:1169–1177. [PubMed: 20525701]
61. Hull MJ, Mino-Kenudson M, Nishioka NS, et al. Endoscopic mucosal resection: an improved diagnostic procedure for early gastroesophageal epithelial neoplasms. *Am J Surg Pathol*. 2006; 30:114–118. [PubMed: 16330950]
62. Mino-Kenudson M, Hull MJ, Brown I, et al. EMR for Barrett's esophagus-related superficial neoplasms offers better diagnostic reproducibility than mucosal biopsy. *Gastrointest Endosc*. 2007; 66:660–666. [PubMed: 17905005]
63. Moss A, Bourke MJ, Hourigan LF, et al. Endoscopic resection for Barrett's high-grade dysplasia and early esophageal adenocarcinoma: an essential staging procedure with long-term therapeutic benefit. *Am J Gastroenterol*. 2010; 105:1276–1283. [PubMed: 20179694]
64. Curvers WL, Singh R, Song LM, et al. Endoscopic tri-modal imaging for detection of early neoplasia in Barrett's oesophagus: a multi-centre feasibility study using high-resolution endoscopy, autofluorescence imaging and narrow band imaging incorporated in one endoscopy system. *Gut*. 2008; 157:67–172.
65. Stein HJ, Feith M, Bruecher BLD, et al. Early esophageal squamous cell and adenocarcinoma: pattern of lymphatic spread and prognostic factors for long term survival after surgical resection. *Ann Surg*. 2005; 242:566–573. [PubMed: 16192817]
66. Bollschweiler E, Baldus SE, Schröder W, et al. High rate of lymph-node metastasis in submucosal esophageal squamous-cell carcinomas and adenocarcinomas. *Endoscopy*. 2006; 38:149–156. [PubMed: 16479422]
67. Pech O, May A, Gunter E, et al. The impact of endoscopic ultrasound and computed tomography on the TNM staging of early cancer in Barrett's esophagus. *Am J Gastroenterol*. 2006; 101:2223–2229. [PubMed: 17032186]
68. Pech O, Günter E, Dusemund F, et al. Value of high-frequency miniprobe and conventional radial endoscopic ultrasound in the staging of early Barrett's carcinoma. *Endoscopy*. 2010; 42:98–103. [PubMed: 20140826]
69. Vieth M, Ell C, Gossner L, et al. Histological analysis of endoscopic resection specimens from 326 patients with Barrett's esophagus and early neoplasia. *Endoscopy*. 2004; 36:776–781. [PubMed: 15326572]
70. Westerterp M, Koppert LB, Buskens CJ, et al. Outcome of surgical treatment for early adenocarcinoma of the esophagus or gastro-esophageal junction. *Virchows Arch*. 2005; 446:497–504. [PubMed: 15838647]
71. Prasad GA, Wu TT, Wigle DA, et al. Endoscopic and surgical treatment of mucosal (T1a) esophageal adenocarcinoma in Barrett's esophagus. *Gastroenterology*. 2009; 137:815–823. [PubMed: 19524578]
72. Badreddine RJ, Prasad GA, Wang KK, et al. Prevalence and predictors of recurrent neoplasia after ablation of Barrett's esophagus. *Gastrointest Endosc*. 2010; 71:697–703. [PubMed: 19959164]
73. Lopes CV, Hela M, Pesenti C, et al. Circumferential endoscopic resection of Barrett's esophagus with high-grade dysplasia or early adenocarcinoma. *Surg Endosc*. 2007; 21:820–824. [PubMed: 17294308]

74. Giovannini M, Bories E, Pesenti C, et al. Circumferential endoscopic mucosal resection in Barrett's esophagus with high-grade intraepithelial neoplasia or mucosal cancer. Preliminary results in 21 patients. *Endoscopy*. 2004; 36:782–787. [PubMed: 15326573]
75. Seewald S, Akaraviputh T, Seitz U, et al. Circumferential EMR and complete removal of Barrett's epithelium: a new approach to management of Barrett's esophagus containing high-grade intraepithelial neoplasia and intramucosal carcinoma. *Gastrointest Endosc*. 2003; 57:854–859. [PubMed: 12776032]
76. Metzger R, Bollschweiler E, Vallbohmer D, et al. High volume centers for esophagectomy: what is the number needed to achieve low postoperative mortality? *Dis Esophagus*. 2004; 17:310–314. [PubMed: 15569369]
77. Testoni PA, Mariani A, Giussani A, et al. SEIFRED Group. Risk factors for post-ERCP pancreatitis in high- and low-volume centers and among expert and non-expert operators: a prospective multicenter study. *Am J Gastroenterol*. 2010; 105:1753–1761. [PubMed: 20372116]
78. Bergman JJ. Radiofrequency ablation—great for some or justified for many? *N Engl J Med*. 2009; 360:2353–2355. [PubMed: 19474433]
79. Pech O, Ell C. Endoscopic therapy of Barrett's esophagus. *Curr Opin Gastroenterol*. 2009; 25:405–411. [PubMed: 19474724]
80. Pech O, May A, Rabenstein T, et al. Endoscopic resection of early oesophageal cancer. *Gut*. 2007; 56:1625–1634. [PubMed: 17938435]
81. Pouw RE, Bergman JJ. Endoscopic resection of early oesophageal and gastric neoplasia. *Best Pract Res Clin Gastroenterol*. 2008; 22:929–943. [PubMed: 18790439]
82. Pouw RE, Sharma VK, Bergman JJ, et al. Radiofrequency ablation for total Barrett's eradication: a description of the endoscopic technique, its clinical results and future prospects. *Endoscopy*. 2008; 40:1033–1040. [PubMed: 19065488]
83. May A, Gossner L, Pech O, et al. Local endoscopic therapy for intraepithelial high-grade neoplasia and early adenocarcinoma in Barrett's oesophagus: acute-phase and intermediate results of a new treatment approach. *Eur J Gastroenterol Hepatol*. 2002; 14:1085–1091. [PubMed: 12362099]
84. Gondrie JJ, Pouw RE, Sondermeijer CM, et al. Stepwise circumferential and focal ablation of Barrett's esophagus with high-grade dysplasia: results of the first prospective series of 11 patients. *Endoscopy*. 2008; 40:359–369. [PubMed: 18494131]
85. Gondrie JJ, Pouw RE, Sondermeijer CM, et al. Effective treatment of early Barrett's neoplasia with stepwise circumferential and focal ablation using the HALO system. *Endoscopy*. 2008; 40:370–379. [PubMed: 18494132]
86. Fayter D, Corbett M, Heirs M, et al. A systematic review of photodynamic therapy in the treatment of pre-cancerous skin conditions, Barrett's oesophagus and cancers of the biliary tract, brain, head and neck, lung, oesophagus and skin. *Health Technol Assess*. 2010; 14:1–288.
87. Overholt BF, Wang KK, Burdick JS, et al. International Photodynamic Group for High-Grade Dysplasia in Barrett's Esophagus. Five-year efficacy and safety of photodynamic therapy with Photofrin in Barrett's high-grade dysplasia. *Gastrointest Endosc*. 2007; 66:460–468. [PubMed: 17643436]
88. Hage M, Siersema PD, van Dekken H, et al. 5-aminolevulinic acid photodynamic therapy versus argon plasma coagulation for ablation of Barrett's oesophagus: a randomised trial. *Gut*. 2004; 53:785–790. [PubMed: 15138203]
89. Li YM, Li L, Yu CH, et al. A systematic review and meta-analysis of the treatment for Barrett's esophagus. *Digest Dis Sci*. 2008; 53:2837–2846. [PubMed: 18427992]
90. Semlitsch T, Jeitler K, Schoefl R, et al. A systematic review of the evidence for radiofrequency ablation for Barrett's esophagus. *Surg Endosc*. 2010; 24:2935–2943. [PubMed: 20464420]
91. Fleischer DE, Overholt BF, Sharma VK, et al. Endoscopic radiofrequency ablation for Barrett's esophagus: 5-year outcomes from a prospective multicenter trial. *Endoscopy*. 2010; 42:781–789. [PubMed: 20857372]
92. Beaumont H, Gondrie JJ, McMahon BP, et al. Stepwise radiofrequency ablation of Barrett's esophagus preserves esophageal inner diameter, compliance, and motility. *Endoscopy*. 2009; 41:2–8. [PubMed: 19160152]

93. Pouw RE, Gondrie JJ, Rygiel AM, et al. Properties of the neosquamous epithelium after radiofrequency ablation of Barrett's esophagus containing neoplasia. *Am J Gastroenterol.* 2009; 104:1366–1373. [PubMed: 19491850]
94. Gotoda T, Yanagisawa A, Sasako M, et al. Incidence of lymph node metastasis from early gastric cancer: estimation with a large number of cases at two large centers. *Gastric Cancer.* 2000; 3:219–225. [PubMed: 11984739]
95. Manner H, May A, Pech O, et al. Early Barrett's carcinoma with "low-risk" submucosal invasion: long-term results of endoscopic resection with a curative intent. *Am J Gastroenterol.* 2008; 103:2589–2597. [PubMed: 18785950]
96. Menon D, Stafinski T, Wu H, et al. Endoscopic treatments for Barrett's esophagus: a systematic review of safety and effectiveness compared to esophagectomy. *BMC Gastroenterol.* 2010; 10:111. [PubMed: 20875123]
97. DeMeester SR. Evaluation and treatment of superficial esophageal cancer. *J Gastrointest Surg.* 2010; 14(Suppl 1):S94–S100. [PubMed: 19760303]
98. Prasad GA, Wang KK, Buttar NS, et al. Long-term survival following endoscopic and surgical treatment of high-grade dysplasia in Barrett's esophagus. *Gastroenterology.* 2007; 132:1226–1233. [PubMed: 17408660]
99. McAllaster JD, Buckles D, Al-Kasspoles M. Treatment of Barrett's esophagus with high-grade dysplasia. *Expert Rev Anticancer Ther.* 2009; 9:303–316. [PubMed: 19275509]
100. Altorki NK, Lee PC, Liss Y, et al. Multifocal neoplasia and nodal metastases in T1 esophageal carcinoma: implications for endoscopic treatment. *Ann Surg.* 2008; 247:434–439. [PubMed: 18376186]
101. Tseng EE, Wu TT, Yeo CJ, et al. Barrett's esophagus with high grade dysplasia: surgical results and long-term outcome-an update. *J Gastrointest Surg.* 2003; 7:164–170. [PubMed: 12600440]
102. Reed MF, Tolis G Jr, Edil BH, et al. Surgical treatment of esophageal high-grade dysplasia. *Ann Thorac Surg.* 2005; 79:1110–1115. [PubMed: 15797034]
103. Chang LC, Oelschlager BK, Quiroga E, et al. Long-term outcome of esophagectomy for high-grade dysplasia or cancer found during surveillance for Barrett's esophagus. *J Gastrointest Surg.* 2006; 10:341–346. [PubMed: 16504878]
104. Rice TW. Pro: esophagectomy is the treatment of choice for high-grade dysplasia in Barrett's esophagus. *Am J Gastroenterol.* 2006; 101:2177–2179. [PubMed: 17032178]
105. Moraca RJ, Low DE. Outcomes and health-related quality of life after esophagectomy for high-grade dysplasia and intramucosal cancer. *Arch Surg.* 2006; 141:545–549. [PubMed: 16785354]
106. Peyre CG, DeMeester SR, Rizzetto C, et al. Vagal-sparing esophagectomy: the ideal operation for intramucosal adenocarcinoma and Barrett with high-grade dysplasia. *Ann Surg.* 2007; 246:665–671. [PubMed: 17893503]
107. Williams VA, Watson TJ, Herbella FA, et al. Esophagectomy for high grade dysplasia is safe, curative, and results in good alimentary outcome. *J Gastrointest Surg.* 2007; 11:1589–1597. [PubMed: 17909921]
108. Mirnezami R, Rohatgi A, Sutcliffe RP, et al. Transhiataloesoph-agectomy: treatment of choice for high-grade dysplasia. *Eur J Cardiothorac Surg.* 2009; 36:364–367. [PubMed: 19464917]
109. Sutton DN, Wayman J, Griffin SM. Learning curve for oesophageal cancer surgery. *Br J Surg.* 1998; 85:1399–1402. [PubMed: 9782024]
110. Begg CB, Cramer LD, Hoskins WJ, et al. Impact of hospital volume on operative mortality for major cancer surgery. *JAMA.* 1998; 280:1747–1751. [PubMed: 9842949]
111. Wenner J, Zilling T, Bladstrom A, et al. The influence of surgical volume on hospital mortality and 5-year survival for carcinoma of the oesophagus and gastric cardia. *Anticancer Res.* 2005; 25:419–424. [PubMed: 15816605]
112. Wouters MW, Karim-Kos HE, Le Cessie S, et al. Centralization of esophageal cancer surgery: does it improve clinical outcome? *Ann Surg Oncol.* 2009; 16:1789–1798. [PubMed: 19370377]
113. Skipworth RJ, Parks RW, Stephens NA, et al. The relationship between hospital volume and post-operative mortality rates for upper gastrointestinal cancer resections: Scotland 1982–2003. *Eur J Surg Oncol.* 2010; 36:141–147. [PubMed: 19879717]

114. Hamilton SR, Yardley JH. Regenerative of cardiac type mucosa and acquisition of Barrett mucosa after esophagogastronomy. *Gastroenterology*. 1977; 72:669–675. [PubMed: 838221]
115. Dresner SM, Griffin SM, Wayman J, et al. Human model of duodenogastro-oesophageal reflux in the development of Barrett's metaplasia. *Br J Surg*. 2003; 90:1120–1128. [PubMed: 12945080]
116. Watson, A., Heading, RC., Shepherd, NA. [Accessed May 4, 2010] Guidelines for the diagnosis and management of Barrett's columnar-lined oesophagus. A Report of the Gastroenterology Working Party of the British Society of Gastroenterology. Aug. 2005 Available at: <http://www.bsg.org.uk/clinical-guidelines/oesophageal/guidelines-forthe-diagnosis-and-management-of-barrett-s-columnar-lined-oesophagus.html>
117. Nicholson A, Jankowski J. Editorial: one small step for metaplasia, but one giant leap for biomarkers is needed. *Am J Gastroenterol*. 2009; 104:2681–2683. [PubMed: 19888234]
118. Cronin J, McAdam E, Danikas A, et al. Epidermal growth factor receptor (EGFR) is overexpressed in high-grade dysplasia and adenocarcinoma of the esophagus and may represent a bio-marker of histological progression in Barrett's esophagus (BE). *Am J Gastroenterol*. 2011; 106:46–56. [PubMed: 21157443]
119. Wani S, Falk GW, Post J, et al. Risk factors for progression of low-grade dysplasia in patients with Barrett's esophagus. *Gastroenterology*. 2011; 141:1179–1186. [PubMed: 21723218]
120. Sikkema M, Looman CWN, Steyerberg EW, et al. Predictors for neoplastic progression in patients with Barrett 's esophagus: a prospective cohort study. *Am J Gastroenterol*. 2011; 106:1231–1238. [PubMed: 21577245]
121. Desai TK, Krishnan K, Samala N, et al. The incidence of oesophageal adenocarcinoma in non-dysplastic Barrett's oesophagus: a meta-analysis. *Gut*. 2011 Oct 13. Epub ahead of print.
122. De Jonge PJF, van Blankenstein M, Looman CWN, et al. Risk of malignant progression in patients with Barrett's oesophagus: a Dutch nationwide cohort study. *Gut*. 2010; 59:1030–1036. [PubMed: 20639249]
123. Hvid-Jensen F, Pedersen L, Drewes AM, et al. Incidence of adenocarcinoma among patients with Barrett's esophagus. *N Engl J Med*. 2011; 365:1375–1383. [PubMed: 21995385]
124. Bhat S, Coleman H, Yousef F, et al. Risk of malignant progression in Barrett's esophagus patients: results from a large population-based study. *J Nat Cancer Inst*. 2011; 103:1049–1057. [PubMed: 21680910]




Figure 1. Proportion of statements achieving consensus with each round of voting. With each round voting improved with iterative changes to the question and supporting evidence.




Figure 2. Management of HGD and/or mucosal cancer (stage T1m) in BE. This consensus has allowed the development of a care pathway for HGD and early adenocarcinoma.

Table 1

Operative Mortality for Surgical Series in Patients With HGD or Early EA With BE

First author	Year	HGD	T1m	Operative mortality
Tseng ¹⁰¹	2003	60	0	1
Reed ¹⁰²	2005	49	0	1
Chang ¹⁰³	2006	9	16	0
Rice ¹⁰⁴	2006	111	0	0
Moraca ¹⁰⁵	2006	23	1	0
Peyre ¹⁰⁶	2007	24	85	7
Williams ¹⁰⁷	2007	38	0	0
Prasad ⁹⁸	2007	70	0	1
Mirnezami ¹⁰⁸	2009	23	0	0
Prasad ⁷¹	2009	0	46	1
		407	160	11/567 (1.9%)

Mean operative mortality is 2%.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Areas Ready to Be Applied to Clinical Management

Pathology	
1	At least 2 experienced gastrointestinal pathologists should evaluate all Barrett's biopsies when a diagnosis of dysplasia is considered.
Endoscopy	
1	The Prague C&M ^a Criteria is the best available tool for grading the endoscopic extent of BE.
2	Visible lumps in nodules consisting of HGD suggest a more advanced lesion with invasion might be present.
Populations at risk	
1	Men have approximately twice the rate of developing HGD or esophageal cancer compared with women, and the rate at which EA is increasing in Western populations is twice as high in men as it is in women.
2	Non-Hispanic white patients with BE are at higher risk for development of HGD/cancer compared with other racial/ethnic groups with BE.
3	Obesity is an independent risk factor for development of EA.
Therapy	
1	Endoscopic treatment should be preferred over endoscopic surveillance for management of most patients with HGD/T1m BE.
2	RFA is currently the best available ablation technique for treatment of flat HGD and for eradication of residual BE after focal EMR.
3	The operative mortality is improved if surgery is undertaken in specialist surgical centers.

NOTE. Several areas that can be applied to clinical practice now include use of Prague Criteria, recognition of subtle masses and use of ER to stage lesions.

^aC, circumferential length, M, maximal length.