

RESEARCH PAPER

Structure-activity relationships of pentamidine-affected ion channel trafficking and dofetilide mediated rescue

R Varkevisser¹, M J C Houtman¹, T Linder², K C G de Git¹,
H D M Beekman¹, R R Tidwell³, A P IJzerman⁴, A Stary-Weinzinger²,
M A Vos¹ and M A G van der Heyden¹

¹Department of Medical Physiology, University Medical Center Utrecht, Utrecht, The Netherlands,

²Department of Pharmacology and Toxicology, University of Vienna, Vienna, Austria,

³Department of Pathology and Laboratory Medicine, School of Medicine, The University of North Carolina, Chapel Hill, NC, USA, and ⁴Division of Medicinal Chemistry, Leiden/Amsterdam Center for Drug Research, Leiden University, Leiden, The Netherlands

Correspondence

M A G van der Heyden,
Department of Medical
Physiology, Yalelaan 50, 3584 CM
Utrecht, the Netherlands. E-mail:
m.a.g.vanderheyden@umcutrecht.nl

Keywords

ion channel trafficking;
pentamidine; dofetilide;
drug-induced trafficking defects;
hERG; K_{IR}2.1; chemical
substructures; Long QT
syndrome; safety pharmacology

Received

29 August 2012

Revised

13 February 2013

Accepted

4 April 2013

BACKGROUND AND PURPOSE

Drug interference with normal hERG protein trafficking substantially reduces the channel density in the plasma membrane and thereby poses an arrhythmic threat. The chemical substructures important for hERG trafficking inhibition were investigated using pentamidine as a model drug. Furthermore, the relationship between acute ion channel block and correction of trafficking by dofetilide was studied.

EXPERIMENTAL APPROACH

hERG and K_{IR}2.1 trafficking in HEK293 cells was evaluated by Western blot and immunofluorescence microscopy after treatment with pentamidine and six pentamidine analogues, and correction with dofetilide and four dofetilide analogues that displayed different abilities to inhibit *I_{Kr}*. Molecular dynamics simulations were used to address mode, number and type of interactions between hERG and dofetilide analogues.

KEY RESULTS

Structural modifications of pentamidine differentially affected plasma membrane levels of hERG and K_{IR}2.1. Modification of the phenyl ring or substituents directly attached to it had the largest effect, affirming the importance of these chemical residues in ion channel binding. PA-4 had the mildest effects on both ion channels. Dofetilide corrected pentamidine-induced hERG, but not K_{IR}2.1 trafficking defects. Dofetilide analogues that displayed high channel affinity, mediated by pi-pi stacks and hydrophobic interactions, also restored hERG protein levels, whereas analogues with low affinity were ineffective.

CONCLUSIONS AND IMPLICATIONS

Drug-induced trafficking defects can be minimized if certain chemical features are avoided or 'synthesized out'; this could influence the design and development of future drugs. Further analysis of such features in hERG trafficking correctors may facilitate the design of a non-blocking corrector for trafficking defective hERG proteins in both congenital and acquired LQTS.

Abbreviations

ER, endoplasmic reticulum; hERG, human ether-a-go-go-related gene; *I_{K1}*, cardiac inward rectifying K⁺ current; *I_{Kr}*, rapid component of the delayed rectifier K⁺ current; LQTS, long QT syndrome; MD, molecular dynamics; TdP, Torsade de Pointes

Introduction

The rapid component of the delayed rectifier K^+ current (I_{Kr}) plays a major role in cardiomyocyte repolarization. The current is carried by pore-forming α -subunits encoded by the human ether-a-go-go-related gene (hERG or KCNH2) (Warmke and Ganetzky, 1994; Sanguinetti and Tristani-Firouzi, 2006). Gene mutations and accompanying hERG ($K_v11.1$) channel current reduction (Curran *et al.*, 1995), or a gain of function mechanism leading to loss of K^+ selectivity and inward rectification (Lees-Miller *et al.*, 2000b) have been linked to human hereditary long QT syndrome (LQTS), which is characterized by a prolonged QT interval on the ECG and an increased risk of sudden cardiac death due to potentially life-threatening Torsade de Pointes (TdP) arrhythmias (Dessertenne, 1966). The acquired form of LQTS can arise as a result of antiarrhythmic drug therapy or as a rare unintended side effect of structurally diverse medications, such as pentamidine, cisapride, terfenadine or sertindole (Thomsen *et al.*, 2003; Kannankeril *et al.*, 2010). Consequently, hERG blockade, QT prolongation and proarrhythmia have become an important component of cardiac safety screening of new pharmacological entities (Haverkamp *et al.*, 2000; Fenichel *et al.*, 2004). The list of (potentially) pro-arrhythmic compounds is still growing and consists mainly of I_{Kr} blockers (Haverkamp *et al.*, 2000; Roden and Viswanathan, 2005) (up-to-date list: <http://www.torsades.org>).

For years, malfunctioning of the hERG channel was presumed to be mainly caused by mutations or drugs that interfered with normal ion conduction, leading to a loss of ion channel function. However, more recently, an entirely different mechanism has been postulated; disruption of normal ion channel trafficking can lead to severely reduced functional protein levels at the plasma membrane, thereby posing an arrhythmic threat. The process of trafficking comprises both anterograde (forward transport towards the plasma membrane) and retrograde (internalization from the plasma membrane) transport of ion channel proteins. Loss of hERG channel function in type 2 LQTS (LQTS2) has been shown to be caused by different mechanisms; mutations result in channels that either gate abnormally, are retained in the endoplasmic reticulum (ER) due to abnormal intracellular processing, or are non-functional (Zhou *et al.*, 1998). Remarkably, it has been shown more recently that the majority of the mutations found in LQTS2 lead to a trafficking defective hERG channel that is trapped in the ER (Anderson *et al.*, 2006). Similarly, a growing number of drugs have been shown to interfere with normal hERG protein trafficking, thereby reducing the number of channels in the cell membrane (van der Heyden *et al.*, 2008). Of note, this chronic drug effect is not detected in conventional cardiac safety screening, and can thus pose a potential arrhythmic threat.

The antiprotozoal agent pentamidine has been known to cause QT prolongation and TdP when used clinically (Jha, 1983; Wharton *et al.*, 1987; Bibler *et al.*, 1988; Girgis *et al.*, 1997). It severely reduces hERG cell surface expression without interfering with ion conduction (Cordes *et al.*, 2005; Kuryshv *et al.*, 2005), acutely blocks the I_{K1} current, and interferes with normal $K_{IR2.1}$ protein expression (de Boer *et al.*, 2010; Nalos *et al.*, 2011). Recently, it was shown that

pentamidine inhibits hERG ER exit and consequently maturation; an effect that can be corrected by application of the class III agent astemizole (Dennis *et al.*, 2012). Interestingly, several trafficking deficient LQTS2 mutants can also be rescued by application of class III agents (Balijepalli *et al.*, 2010).

Currently, it is not known which chemical substructures are responsible for drug-induced hERG trafficking inhibition or the restoration of both acquired and inherited LQTS. Therefore, we aimed to (i) determine the chemical features within pentamidine that are important for acquired trafficking defects, (ii) determine whether dofetilide can restore the acquired trafficking defects of hERG and $K_{IR2.1}$, and (iii) determine the relationship between acute ion channel block and correction of trafficking. We found that structural modifications of pentamidine differentially affect hERG and/or $K_{IR2.1}$ trafficking. Dofetilide and its analogues corrected pentamidine-induced hERG trafficking defects and this effect was found to be dependent on their affinity for the channel.

Methods

Animal experiments

Animal care and handling was performed in accordance with the 'European Directive for the Protection of Vertebrate animals used for Experimental and Scientific Purpose, European Community Directive 86/609/CEE'. All experiments were approved by the Committee for Experiments on Animals of the Utrecht University, The Netherlands. Three adult purpose-bred mongrel dogs (Marshall, North Rose, NY, USA) were used. General anaesthesia was induced by pentobarbital (25 mg·kg⁻¹ i.v.) and maintained by isoflurane (1.5% in O₂ : NO₂ 1:2). Atrioventricular block was induced by radiofrequency ablation in all dogs, and ventricular activation was controlled using a DDDR pacemaker (Vitatron, Arnhem, The Netherlands). Detailed descriptions of procedures, pacing protocols and data analysis were published previously (Winckels *et al.*, 2007). After a minimum of 4 weeks (4–6 weeks), pentamidine (10 mg·kg⁻¹ for 60 min, i.v.) was administered under full anaesthesia. Standard six-lead ECG recordings were made regularly over a period of 6 weeks to determine the long-term electrophysiological effects of pentamidine. All studies involving animals are reported in accordance with the ARRIVE guidelines for reporting experiments involving animals (Kilkenny *et al.*, 2010; McGrath *et al.*, 2010).

Cell culture

HEK-293 cells stably expressing GFP-tagged murine wildtype $K_{IR2.1}$ (HEK-KWGF) were generated as described previously (de Boer *et al.*, 2006). HEK-hERG cells were obtained from C. January (Zhou *et al.*, 1998). Cells were maintained in DMEM supplemented with 10% fetal calf serum, 2 mM l-glutamine, 50 U·mL⁻¹ penicillin, and 50 µg·mL⁻¹ streptomycin (Verviers, Belgium). For experiments cells were seeded and incubated in complete DMEM containing different drugs at various concentrations as indicated. Variable incubation times were achieved by replacing control medium with drug-containing medium at appropriate time points. During the experiments,

drug-containing medium was refreshed every 24 h. The amount of DMSO added to the cells was kept below 0.01%.

Chemical compounds

Pentamidine-isethionate (Pentacrit® 300, Sanofi Aventis, Gouda, The Netherlands) was dissolved in water at a concentration of 0.1 M, sterilized by filtration (22 µM), aliquoted and stored at -20°C until further use. Pentamidine analogues (PA; see Table 1, PA-1 to PA-7) were synthesized as described previously (Jones *et al.*, 1990; Tidwell *et al.*, 1990; Bakunova *et al.*, 2009a,b), and were dissolved in water (PA-1 to PA-5) or DMSO (PA-7) at the highest concentration possible (≤25 mM). Dofetilide and its analogues were dissolved in DMSO at a concentration of 10 mM. Dofetilide analogues (DA) were synthesized as described previously (Shagufra *et al.*, 2009).

Western blot

Cell lysates were prepared in buffer D (20 mM HEPES, 125 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1 mM dithiothrei-


tol, 10% glycerol, and 1% Triton X-100) supplemented with 1 mM PMSF and 6.8 µg·mL⁻¹ aprotinin (Sigma-Aldrich, Zwijndrecht, The Netherlands). Fifteen micrograms of protein lysate was mixed with Laemmli sample buffer, separated by 7% (hERG) or 10% (K_{IR2.1}) SDS-PAGE and subsequently electroblotted onto a nitrocellulose membrane (Biorad, Veenendaal, The Netherlands). Reversible Ponceau staining was used to reveal equal protein loading and subsequently quantification. K_{IR2.1}-GFP was detected by monoclonal anti-GFP (cat. no. Sc-9996; Santa Cruz Biotechnology, Santa Cruz, CA, USA), hERG protein was detected by polyclonal anti-hK_v11.1 primary antibody (cat. no. APC-062; Alomone Labs, Jerusalem, Israel). Peroxidase-conjugated secondary antibodies were used to facilitate final detection with a standard ECL procedure (Santa Cruz Biotechnology).

Immunofluorescence microscopy

HEK-hERG cells were cultured on poly-L-lysine (Sigma)-coated 15 Ø glass cover slips (Smethwick, Warley, UK),

Table 1

Chemical structures of pentamidine (P) and dofetilide (D) and their derivatives used in this study



fixed with 3% paraformaldehyde dissolved in PBS. Cells were quenched with 50 mM glycine-PBS after permeabilization with 0.5% Triton X-100, and subsequently blocked with NET-gel (150 mM NaCl, 5 mM EDTA, 50 mM Tris-HCl, pH 7.4, 0.05% igepal, 0.25% gelatin, 0.02% NaN₃). Cells were then incubated overnight with polyclonal anti-hK_v11.1 primary antibody (cat. no. APC-062; Alomone Labs), followed by incubation with anti-rabbit FITC-conjugated secondary antibody (Jackson ImmunoResearch) for 2 h. Coverslips were mounted with Vectashield (Vector Laboratories Inc., Burlingame, CA, USA), and confocal images were obtained using a Zeiss Axiovert 200 M confocal microscope (Carl Zeiss Microscopy GmbH, Jena, Germany) equipped with a $\times 40$ water immersion objective (NA 1.2) plus $\times 2$ digital zoom. Excitation was performed with an air-cooled argon ion laser (LASOS, RMC 7812Z, 488 nm).

Drug docking and molecular dynamics (MD) simulations

Docking was performed with the program Gold 4.0.1 (GOLD, 2008) using the Gold scoring functions. The recently published hERG homology model termed 'model 6' (Stary *et al.*, 2010) after 50 ns MD simulations was used as a template for docking. Coordinates of the geometric centre calculated among the Y652 and F656 residues were taken as binding site. The binding site radius was set to 10 Å. The 10 best-ranked poses of each docking run were used for visual analysis. The most frequent binding mode was used as a starting conformation for MD simulations.

MD simulations were performed with Gromacs v. 4.5.4. (Hess *et al.*, 2008) as described previously (Knape *et al.*, 2011). Briefly, the hERG model was embedded in an equilibrated membrane consisting of 280 dioleoylphosphatidylcholine (DOPC) lipids by making use of the *g_membed* tool (Wolf *et al.*, 2010). The amber99sb force field (Hornak *et al.*, 2006) with lipid parameters taken from Siu *et al.* (2008), and the TIP3P water model (Jorgensen *et al.*, 1983) were utilized. Geometry optimization and topology generation of the drugs was carried out with HF/3-21G, implemented in Gaussian09 (Frisch *et al.*, 2009) and antechamber (Case *et al.*, 2010) respectively. Prior to simulations, 1000 conjugate gradient energy-minimization steps were performed. Subsequently, the systems were equilibrated for 1 ns for each docked compound. During equilibration, the protein and the ligand were restrained with a force constant of 1000 kJ·mol⁻¹·nm⁻², while the lipids, the ions, and the water were allowed to move freely. Each drug was simulated for 50 ns.

Statistics

Group averages are presented as mean \pm SEM, unless indicated otherwise. Differences between groups were tested using one-way ANOVA with a Bonferroni *post hoc* test. A statistically significant difference was considered if $P < 0.05$. All analyses were carried out using SPSS (SPSS 19.0 IBM, Armonk, NY, USA).

Drugs and molecular target nomenclature conform to the British Journal of Pharmacology's *Guide to Receptors and Channels* (Alexander *et al.*, 2011).

Results

A single dose of pentamidine lengthens the QT interval

The time course of QT interval changes after a single dose of pentamidine is summarized in Figure 1. The QT interval did not change during and directly after pentamidine administration in any of the three dogs. After 7 days, the QT interval prolongs progressively, which persists until day 21, demonstrating the typical chronic effect of pentamidine on repolarization.

Pentamidine and its analogues differentially inhibit hERG and/or K_{IR2.1} trafficking

Pentamidine inhibited anterograde hERG trafficking, as can be seen in Figure 2A. The mature protein level (upper band) was severely reduced compared with control levels after 48 h treatment with 10 μ M pentamidine (0.2 ± 0.02 vs. 1.0 ± 0.0 , $P < 0.05$). In addition, K_{IR2.1} protein levels were also analysed. As expected, 10 μ M pentamidine for 48 h significantly decreased K_{IR2.1} protein levels compared with control (Figure 2B). At present, the chemical characteristics and substructures within trafficking inhibiting drugs that specifically interfere with the normal trafficking process of these potassium ion channels have not been identified. To identify potentially trafficking inhibiting substructures, pentamidine was used as a model drug, and several analogues were tested for ion channel trafficking inhibiting capacity. The molecular structure of pentamidine is depicted in Table 1. Compared with the lead compound, PA-1, 2 and 3 contain modifications in the linker between the benzamidine groups, PA-4 has pyridine instead of phenyl rings, and PA-5 and 7 contain substituted benzamidine moieties (Table 1).

Both HEK-hERG and HEK-KWGF cells were treated with 10 μ M pentamidine or its analogues for 48 h (Figure 2). Com-




Figure 1

Electrophysiological effects of 10 mg·kg⁻¹ pentamidine over 60 min in AV-blocked dogs paced from the high septum. A delayed effect of pentamidine on the QT interval was observed. QT prolongation became apparent 4 days after i.v. pentamidine administration that persisted for 17 days, after which the prolongation dissipated gradually. Data are presented as mean \pm SD ($n = 3$).




Figure 2

Pentamidine and its analogues differentially affect hERG and $K_{IR}2.1$ protein levels. (A) Western blot showing the effects of pentamidine or its structural analogues (10 μ M, 48 h) on hERG forward trafficking ($n = 12$ for control and pentamidine, $n = 3$ for PA-4, $n = 4$ for all other analogues). (B) Western blot showing the effects of pentamidine or its structural analogues (10 μ M, 48 h) on $K_{IR}2.1$ protein expression ($n = 3$ for PA-4 and 7, $n = 4$ for control, pentamidine, and all other analogues). Total protein staining (Ponceau) was used as a loading control. Control protein levels (untreated cells) were designated as 100% after correction; *indicates $P < 0.05$ versus control, †indicates $P < 0.05$ versus pentamidine treatment.

pared with pentamidine, either shortening or lengthening the carbon linker between the aromatic rings had slight effects on hERG protein maturation. PA-1 was a slightly less effective trafficking inhibitor (0.30 ± 0.03 vs. P 0.20 ± 0.02 ,

ns), while PA-2 was slightly more potent than pentamidine (0.15 ± 0.02 vs. P 0.20 ± 0.02 , ns). Interestingly, substitution of the oxygen directly attached to the aromatic residues by sulphur considerably increased hERG trafficking inhibition

(PA-3 0.04 ± 0.02 vs. P, PA-1, PA-4, PA-5, or PA-7, $P < 0.05$). In contrast, PA-5 ($P < 0.05$ vs. P, and PA-2, 3, 4) and PA-7 ($P < 0.05$ vs. P, and PA-2, 3, 4) were significantly less potent than pentamidine, although protein levels comparable with control were not reached. PA-4 – in which the phenyl rings are replaced by pyridine – was the least potent hERG trafficking inhibiting compound ($P < 0.05$ for PA-4 vs. P and PA-1 to 7). Although control protein levels were not reached, mature hERG protein levels were substantially increased when PA-4 and pentamidine treatment were compared (Figure 2A). All these data indicate that an unaltered amidine substituent and the nature of the aromatic rings are important determinants of the interaction of pentamidine and the hERG channel during the trafficking process. It should be kept in mind though, that the compounds were tested at one concentration only, and that full concentration–response curves would allow a more definitive and quantitative SAR analysis.

In accordance with their effect on hERG trafficking, the effect of PA-1 and 2 on $K_{IR2.1}$ protein levels differed slightly when compared with pentamidine (Figure 2B). In contrast, PA-4, 5 and 7 hardly affected $K_{IR2.1}$ protein levels. Interestingly, PA-4 also slightly affected hERG trafficking. When comparing the effects of the pentamidine analogues on hERG and $K_{IR2.1}$, the differential effect of PA-3 on these proteins represents an obvious discrepancy. While hERG trafficking was severely inhibited, $K_{IR2.1}$ protein levels were hardly affected (0.90 ± 0.2 vs. control 1.0 ± 0.0 , ns), which could imply that the mechanism by which pentamidine inhibits trafficking differs between these ion channels.

Dofetilide rescues pentamidine-induced hERG, but not $K_{IR2.1}$, trafficking defects

In order to reduce the pentamidine-induced hERG channel trafficking block, we applied the high affinity class III agent dofetilide in different concentrations to HEK-hERG cells in the continuous presence of $10 \mu\text{M}$ pentamidine. Application of even $0.03 \mu\text{M}$ dofetilide significantly increased mature hERG levels compared with pentamidine (0.7 ± 0.1 vs. 0.3 ± 0.03 , $P < 0.05$, Figure 3A), and at a concentration of $1 \mu\text{M}$, dofetilide completely restored anterograde hERG trafficking. Interestingly, higher concentrations of dofetilide seemed to increase hERG trafficking efficiency, since significantly higher mature protein levels were reached compared with control levels.

The timeframe in which the pharmacological correction of drug-induced trafficking was taking place is unknown. Therefore, HEK-hERG cells were first exposed to pentamidine ($10 \mu\text{M}$) for 48 h; subsequently, dofetilide ($1 \mu\text{M}$) was added in the continued presence of pentamidine and mature protein levels were determined after different incubation times. Mature protein levels increased time- dependently, and were completely restored after 4–6 h of dofetilide treatment (Figure 3B). This timeframe strongly suggests that the correction of pentamidine-induced trafficking defects occurred post-transcriptionally, most likely via the relief of ER retention.

Since dofetilide is a high affinity class III agent, we hypothesized that its correction of ion channel trafficking defects could be ion channel-specific, and the pharmacological chaperone used has to bind directly to the particular ion channel involved. To test this hypothesis, we evaluated

whether pentamidine-affected $K_{IR2.1}$ protein levels could be corrected by application of dofetilide. As expected, treatment with dofetilide up to $30 \mu\text{M}$ did not restore $K_{IR2.1}$ protein levels (0.4 ± 0.06 vs. P 0.25 ± 0.05 , ns, $n = 3$, Figure 3C).

Correction of pentamidine-induced hERG trafficking defects depends on an affinity for the channel

Since dofetilide has a high affinity for the hERG channel and specifically restores drug-induced hERG trafficking defects, we wondered which substructures that affect channel affinity are important in the molecule. The structure of dofetilide is depicted in Table 1; its protonated tertiary nitrogen, phenyl rings and methanesulphonamide substituents are known to be important for high affinity binding (Shagufta *et al.*, 2009). Furthermore, the spacing between the central basic nitrogen and the aromatic residues might also be of significance for efficient binding (Pearlstein *et al.*, 2003). In all the dofetilide analogues (DA) used in this study, the methanesulphonamide substituent was replaced; in DA-2 by $-\text{NH}_2$, in DA-1, 3 and 4 by NO_2 groups. In addition, DA-3 was differently substituted at the central nitrogen, and in DA-4, the length of the chain connecting the two phenyl groups was modified. DA-1 was used as a reference compound for synthesis of DA-3 and 4. Compared with dofetilide (K_i 4.9 ± 0.9 nM), DA-1 had a comparable affinity (K_i 5.7 ± 3.0 nM) for the hERG channel, DA-3 and 4 had moderate affinity (K_i 278 ± 73 nM and K_i 202 ± 4.9 nM), and DA-2 had a very low affinity ($K_i > 10 \mu\text{M}$) for the channel (Shagufta *et al.*, 2009).

Dofetilide and the four analogues were tested for their ability to correct drug-induced trafficking defects (Figure 4A). DA-1, in which a nitro-substituent replaces the methanesulphonamide group, was almost as effective as dofetilide. In contrast, substitution with a polar NH_2 group (DA-2) completely abolished the trafficking correction capacity ($30 \mu\text{M}$ DA-2 0.40 ± 0.04 vs. control 1.0 ± 0.0 , $P < 0.05$; $30 \mu\text{M}$ DA-2 0.40 ± 0.04 vs. $30 \mu\text{M}$ dofetilide 1.9 ± 0.4 , $P < 0.05$) in association with a substantial decrease in hERG binding affinity (Shagufta *et al.*, 2009). In DA-3, the basicity of the central nitrogen was abolished by acylation; as a consequence, not only was hERG affinity decreased considerably, but also hERG trafficking defects were only partially corrected ($30 \mu\text{M}$ DA-3 0.60 ± 0.1 vs. control 1.0 ± 0.0 , $P < 0.05$; $30 \mu\text{M}$ DA-3 0.60 ± 0.1 vs. $30 \mu\text{M}$ dofetilide 1.9 ± 0.4 , $P < 0.05$). DA-4, which had a moderate hERG affinity due to a variation in chain length, also partially corrected the pentamidine-induced trafficking defect. It remains puzzling, however, why higher concentrations of the lower-affinity compounds DA-3 and DA-4 do not fully restore mature protein levels.

The subcellular localization of the hERG protein after trafficking inhibition with pentamidine and correction thereof with dofetilide and its analogues ($1 \mu\text{M}$) was studied by immunofluorescence staining of hERG (Figure 4B). In untreated control cells, hERG was distributed throughout the cell and clear membrane staining was detected, as was the case after only dofetilide treatment. As expected, membrane staining was lost after pentamidine treatment, and intracellular staining had a more restricted perinuclear pattern. The localization of the hERG protein after application of the correctors correlated with their efficacy and was in agreement with the results of the Western blots. Correction of




Figure 3

Dofetilide corrects pentamidine-induced hERG trafficking defects. (A) Western blot showing the concentration-dependent correction of hERG protein levels in the presence of pentamidine. Dofetilide 1 μ M completely restored mature hERG protein levels, higher concentrations increased mature hERG protein levels even further ($n = 24$ for control and pentamidine, $n = 6$ for all dofetilide concentrations). (B) Dofetilide time-dependently corrects mature hERG protein levels. Treatment of pentamidine-exposed (10 μ M, 48 h) HEK-hERG cells with 1 μ M dofetilide restored mature hERG levels within 4–6 h ($n = 24$ for control and pentamidine, $n = 3$ dofetilide). (C) In contrast to the beneficial effect on hERG trafficking, pentamidine-mediated downregulation of K_{IR}2.1 could not be corrected by application of dofetilide ($n = 3$ for all conditions). Total protein staining (Ponceau) was used as a loading control. Control protein levels (untreated cells) were designated as 100% after correction; *indicates $P < 0.05$ versus control, ‡indicates $P < 0.05$ versus pentamidine treatment.




Figure 4

Correction of pentamidine-induced hERG trafficking defects is influenced by structural modifications of dofetilide (A). Compared with dofetilide, the high affinity analogue DA-1 was an equally effective corrector, while the other analogues were significantly less effective when tested at 1 μ M ($n = 24$ for control and pentamidine, $n = 6$ for dofetilide, $n = 3$ for DA-1 and 2, $n = 5$ for DA-3 and 4). Total protein staining (Ponceau) was used as a loading control. Control protein levels (untreated cells) were designated as 100% after correction; *indicates $P < 0.05$ versus control, ‡indicates $P < 0.05$ versus pentamidine treatment. (B) In control conditions, the hERG protein is present in the cytoplasm and at the plasma membrane. A similar staining pattern was seen after only dofetilide treatment. After pentamidine treatment, only cytoplasmic staining was apparent. Staining correction patterns after correction of these effects with dofetilide or its structural analogues were in agreement with the Western blot results; effective correction showed hERG staining both at the plasma membrane and intracellularly, while ineffective correctors showed only intracellular staining. The hERG protein is shown in green.

pentamidine-induced trafficking defects with dofetilide or DA-1 revealed a pattern comparable with control cells, as both compounds completely restored mature hERG protein levels. In contrast, DA-2- treated cells displayed a staining similar to cells treated with pentamidine alone. After treatment with DA-3 or DA-4, which are both partial correctors, the hERG protein was scattered throughout the cell but

membrane staining was considerably less when compared to control cells.

Finally, to gain further insight in the relationship between drug binding affinity and correction of hERG trafficking defects, drug docking and molecular dynamic simulations were performed. All dofetilide derivatives in our hERG model displayed considerable stability during the course of 50 ns MD

Figure 5

MD poses and 2D interaction profiles of dofetilide (A,B), DA-1 (C,D), DA-2 (E,F), DA-3 (G,H) and DA-4 (I,J). Snapshots after 50 ns are shown. Binding determinants Y652 and F656 are shown as raspberry sticks. Drug molecules are coloured orange, with nitrogen atoms coloured blue, oxygen atoms coloured red and sulfur groups coloured yellow. 2D interaction profiles were generated with PoseView (poseview.zbh.uni-hamburg.de). Green dotted lines indicate pi-pi interactions between aromatic rings, green solid lines represent hydrophobic interactions. Hydrogen bonds are shown as black dotted lines.

Table 2

Number of interactions for dofetilide derivatives


Compound	Σ of hydrophobic IAs (pi-pi stacking)	HB IAs
D	3 (1)	–
DA-1	4 (2)	–
DA-2	–	1
DA-3	2 (1)	–
DA-4	1 (1)	–

HB, hydrogen bonds; IAs, interactions.

simulations. Key interactions, involving Y652 and F656, are shown in Figure 5. The numbers of hydrophobic and aromatic interactions a compound has (Table 2) are in good agreement with its ability to correct pentamidine-induced trafficking defects. While the compounds dofetilide and DA-1 (Figure 5A–D) had a large number of favourable interactions, DA-2, DA-3 and DA-4 had very few interactions with the hERG model. Our modelling studies suggest that dofetilide and DA-1 stabilize the hERG structure via pi-pi stacking, while the low affinity compound DA-2 cannot provide sufficient stabilizing interactions to correct pentamidine trafficking defects. DA-3 and DA-4 take an intermediate position between D and DA-1 on the one, and DA-2 on the other hand.

Discussion

Detrimental drug interactions with the hERG potassium channel constitute a major pharmacological safety concern during drug development. In recent years, it has become recognized that, in addition to hERG current inhibition, chronic drug effects should also be considered as a potential safety hazard (van der Heyden *et al.*, 2008), as the dangerous effects of trafficking inhibiting agents like pentamidine and arsenic trioxide are not always detected in conventional hERG safety screening assays (Katchman *et al.*, 2006). Although ion channel trafficking assays are available (HERG-Lite® and CHAN-Lite®, ChanTest™, Cleveland, Ohio), they have not been adopted by the current guidelines. In order to avoid toxicity, it is imperative to understand the structural requirements needed not only for hERG binding, but also for hERG trafficking inhibition, and this would be of great benefit for the drug design and development process.



At present, it is not known which chemical substructures in pentamidine are important for selective disruption of hERG ion channel trafficking. Our results indicate that the aromatic residues, in particular, are essential, as modifications of these groups (PA-3 and 4) have a large effect on hERG ion channel trafficking. PA-4, which had the most favourable trafficking safety profile, was evaluated in a mouse model of trypanosomiasis. Unfortunately, the compound was not effective, although it was well tolerated (Bakunova *et al.*, 2009b). Interestingly, both PA-3 and 4 mildly affected $K_{IR}2.1$ density, while PA-3 severely affected hERG maturation. PA-2 had a large effect on both $K_{IR}2.1$ and hERG density. The differential effect of these compounds suggests that the mechanism by which pentamidine exerts protein downregulation differs between these potassium channels. In line with this, we found that the specific I_{Kr} inhibitor dofetilide corrected hERG channel but not $K_{IR}2.1$ channel density after pentamidine treatment. Previously, we showed that inhibition of lysosomal degradation restores $K_{IR}2.1$ protein levels in pentamidine-treated HEK-KWGF cells, which strongly suggests that pentamidine increases $K_{IR}2.1$ retrograde trafficking and degradation (Nalos *et al.*, 2011). In addition, pharmacological interference at different levels in the clathrin-dependent internalization route and thus retrograde trafficking, increase pentamidine-affected $K_{IR}2.1$ protein levels (Varkevisser *et al.*, 2013).

In the present study, we showed that dofetilide also corrects pentamidine-induced hERG anterograde trafficking defects and the correction efficacy depends on the compound's affinity for the channel. A fair agreement between affinity and correction was observed; a high affinity analogue (DA-1) completely restored mature hERG protein levels in the presence of pentamidine while analogues with moderate (DA-3 and 4) or low affinity (DA-2) restored mature protein levels partially or not at all. Our results favour the hypothesis that pentamidine and its correctors compete for the same binding site within the hERG channel. While pentamidine supposedly binds to an emerging hERG drug binding site in a conformation folding intermediate leading to an arrest in channel maturation (Dennis *et al.*, 2012), displacement by a high affinity corrector stabilizes the channel, thereby alleviating the block in maturation. However, based on their chemical structure, it seems unlikely that dofetilide and pentamidine share a common binding site. Alternatively, pentamidine could bind to a different binding site in the immature hERG protein, which is not available in the mature protein. When performing MD simulations with pentamidine, no consistent interactions between the drug and the hERG model were found, in contrast to dofetilide and its analogues (Figure 5). This finding would be consistent with pentamidine binding to a channel intermediate instead of the mature channel. Presumably, pentamidine binding destabilizes the immature hERG protein, leading to its retention in the ER by quality control mechanisms; the binding of a high affinity class III agent at an allosteric site will then stabilize the immature protein and promote maturation. If this hypothesis is correct, dofetilide would operate similarly in the presence of other destabilizing agents. Mutation of amino acid residues F656 and Y652 that affect pentamidine sensitivity (Dennis *et al.*, 2012) could, therefore, also have an indirect effect, as a mutation of these residues can affect the characteristics of the

current (Lees-Miller *et al.*, 2000a; Fernandez *et al.*, 2004; Kamiya *et al.*, 2006) possibly by small conformational changes. This could either directly obscure the binding site or reduce pentamidine binding because of steric hindrance. The direct hERG block and trafficking inhibition induced by several other drugs were found to be mediated by the drugs binding to different sites. Ketoconazole, fluconazole and fluoxetine, which are all direct I_{Kr} blockers, as well as hERG trafficking inhibitors, do not seem to bind to the canonical drug binding site in the immature hERG protein, as a mutation of residues Phe⁶⁵⁶ and Tyr⁶⁵² abolished the direct I_{Kr} block but had no effect on hERG trafficking inhibition (Rajamani *et al.*, 2006; Takemasa *et al.*, 2008; Han *et al.*, 2011). Furthermore, application of the high affinity I_{Kr} blocker E-4031 did not restore the hERG trafficking inhibition induced by either fluoxetine or norfluoxetine (Rajamani *et al.*, 2006).

The mechanism by which pentamidine arrests hERG channel maturation is not known. It has been suggested that disruption of the channel-chaperone interactions could play a role. Wild-type hERG proteins have been shown to interact with the cytosolic chaperones Hsp70 and Hsp90 and the ER chaperone calnexin (Ficker *et al.*, 2003; Gong *et al.*, 2006). Not surprisingly, drugs like geldanamycin and As_2O_3 that inhibit Hsp70 and 90, arrested hERG maturation (Ficker *et al.*, 2003; 2004). However, it should be noted that these drugs have complex actions and could potentially inhibit the correct folding of other proteins, including other (cardiac) ion channels. On the other hand, it has been shown that pentamidine does not interfere with the actions of Hsp70 or Hsp90 (Dennis *et al.*, 2012), but it could disrupt the association of hERG with other chaperones, like calnexin. Calnexin associates transiently with wild-type immature hERG channels for up to 8 h, whereas its association with the trafficking deficient LQTS2 mutant N470D was prolonged (Gong *et al.*, 2006). Association times between calnexin and this mutant channel approached the normal time frame in the presence of E-4031 and accordingly the protein levels of this mutant hERG were completely restored within 8 h (Zhou *et al.*, 1999). In agreement with these findings, we showed that dofetilide completely restored hERG protein levels within 8 h in the presence of pentamidine. Further research is needed to elucidate the precise mechanism by which pentamidine arrests hERG maturation.

In several human diseases mutant proteins fold or assemble improperly, resulting in defective protein trafficking (Cohen and Kelly, 2003). Correction of defective trafficking by protein-specific modulators or drug substrates was first shown for misprocessed human P-glycoprotein mutants (Loo and Clarke, 1997). Using this approach as potential clinical therapy, great progress has been achieved in the treatment of cystic fibrosis. A promising small molecule (VX-809) that corrects anterograde trafficking of $\Delta F508$ -CFTR mutant proteins has entered clinical trials (Clancy *et al.*, 2012). Pharmacological correctors of either mutant or drug-induced trafficking deficient hERG channels are typically also direct hERG channel blockers, and their efficacy appears to be related to the domain in which the mutation is located and the affinity of the class III agent used (Ficker *et al.*, 2002; Balijepalli *et al.*, 2010; Dennis *et al.*, 2012). In agreement with earlier results in LQTS2 mutants (Ficker *et al.*, 2002), in the present study, we showed that correction of drug-induced

hERG trafficking defects is dependent on channel affinity and is channel specific. In particular, modifications near a compound's aromatic residues greatly affect its ability to restore the drug-induced hERG trafficking. Further research into the difference between direct hERG channel blockade and correction of defective trafficking is necessary and may be of benefit for the treatment of defective hERG trafficking.

In conclusion, structural modifications of pentamidine differentially affected plasma membrane levels of hERG and Kir2.1. Modification of the phenyl ring or substituents directly attached to it had the largest effect, highlighting the importance of these features in ion channel binding. Thus, if certain residues are avoided or 'synthesized out', the ability to inhibit trafficking can be abolished, as was shown with PA-4, which had mild effects on both ion channels. Correction of defective protein trafficking is ion channel-specific and requires high affinity binding. Further analysis of important features in other hERG channel trafficking correctors may facilitate the design of a non-blocking corrector for trafficking defective hERG proteins in both congenital and acquired LQTS.

Acknowledgements

We would like to thank Dr C. January (University of Wisconsin) for providing HEK-hERG cells. Part of the research leading to the results has received funding from the European Community's Seventh Framework Programme FP7/2007–2013 under grant agreement No. HEALTH-F2-2009-241526, EUTrigTreat. Part of this work was supported by The Austrian Science Fund (FWF; Grants P22395, W1232). T. L. was supported by a research fellowship 2013 from the University of Vienna.

Conflict of interest

None.

References

- Alexander SPH, Mathie A, Peters JA (2011). Guide to Receptors and Channels (GRAC), 5th edn. *Br J Pharmacol* 164 (Suppl. 1): S1–S324.
- Anderson CL, Delisle BP, Anson BD, Kilby JA, Will ML, Tester DJ *et al.* (2006). Most LQT2 mutations reduce Kv11.1 (hERG) current by a class 2 (trafficking-deficient) mechanism. *Circulation* 113: 365–373.
- Bakunova SM, Bakunov SA, Patrick DA, Kumar EV, Ohemeng KA, Bridges AS *et al.* (2009a). Structure-activity study of pentamidine analogues as antiprotozoal agents. *J Med Chem* 52: 2016–2035.
- Bakunova SM, Bakunov SA, Wenzler T, Barszcz T, Werbovetz KA, Brun R *et al.* (2009b). Synthesis and antiprotozoal activity of pyridyl analogues of pentamidine. *J Med Chem* 52: 4657–4667.
- Balijepalli SY, Anderson CL, Lin EC, January CT (2010). Rescue of mutated cardiac ion channels in inherited arrhythmia syndromes. *J Cardiovasc Pharmacol* 56: 113–122.

Bibler MR, Chou TC, Toltzis RJ, Wade PA (1988). Recurrent ventricular tachycardia due to pentamidine-induced cardiotoxicity. *Chest* 94: 1303–1306.

de Boer TP, van Veen TA, Houtman MJ, Jansen JA, van Amersfoorth SC, Doevendans PA *et al.* (2006). Inhibition of cardiomyocyte automaticity by electrotonic application of inward rectifier current from Kir2.1 expressing cells. *Med Biol Eng Comput* 44: 537–542.

de Boer TP, Nalos L, Stary A, Kok B, Houtman MJ, Antoons G *et al.* (2010). The anti-protozoal drug pentamidine blocks Kir2.x-mediated inward rectifier current by entering the cytoplasmic pore region of the channel. *Br J Pharmacol* 159: 1532–1541.

Case DA, Darden TA, Cheatham TE III, Simmerling CL, Wang J, Duke RE *et al.* (2010). AMBER 11. University of California: San Francisco, CA.

Clancy JP, Rowe SM, Accurso FJ, Aitken ML, Amin RS, Ashlock MA *et al.* (2012). Results of a phase IIa study of VX-809, an investigational CFTR corrector compound, in subjects with cystic fibrosis homozygous for the F508del-CFTR mutation. *Thorax* 67: 12–18.

Cohen FE, Kelly JW (2003). Therapeutic approaches to protein-misfolding diseases. *Nature* 426: 905–909.

Cordes JS, Sun Z, Lloyd DB, Bradley JA, Opsahl AC, Tengowski MW *et al.* (2005). Pentamidine reduces hERG expression to prolong the QT interval. *Br J Pharmacol* 145: 15–23.

Curran ME, Splawski I, Timothy KW, Vincent GM, Green ED, Keating MT (1995). A molecular basis for cardiac arrhythmia: HERG mutations cause long QT syndrome. *Cell* 80: 795–803.

Dennis AT, Wang L, Wan H, Nassal D, Deschenes I, Ficker E (2012). Molecular determinants of pentamidine-induced hERG trafficking inhibition. *Mol Pharmacol* 81: 198–209.

Dessertenne F (1966). [Ventricular tachycardia with 2 variable opposing foci]. *Arch Mal Coeur Vaiss* 59: 263–272.

Fenichel RR, Malik M, Antzelevitch C, Sanguinetti M, Roden DM, Priori SG *et al.* (2004). Drug-induced torsades de pointes and implications for drug development. *J Cardiovasc Electrophysiol* 15: 475–495.

Fernandez D, Ghanta A, Kauffman GW, Sanguinetti MC (2004). Physicochemical features of the hERG channel drug binding site. *J Biol Chem* 279: 10120–10127.

Ficker E, Obejero-Paz CA, Zhao S, Brown AM (2002). The binding site for channel blockers that rescue misprocessed human long QT syndrome type 2 ether-a-gogo-related gene (HERG) mutations. *J Biol Chem* 277: 4989–4998.

Ficker E, Dennis AT, Wang L, Brown AM (2003). Role of the cytosolic chaperones Hsp70 and Hsp90 in maturation of the cardiac potassium channel hERG. *Circ Res* 92: e87–100.

Ficker E, Kuryshv YA, Dennis AT, Obejero-Paz C, Wang L, Hawryluk P *et al.* (2004). Mechanisms of arsenic-induced prolongation of cardiac repolarization. *Mol Pharmacol* 66: 33–44.

Frisch MJ, Trucks CW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR *et al.* (2009). Gaussian 09. Gaussian, Inc.: Wallingford, CT.

Girgis I, Gualberti J, Langan L, Malek S, Mustaciulo V, Costantino T *et al.* (1997). A prospective study of the effect of I.V. pentamidine therapy on ventricular arrhythmias and QTc prolongation in HIV-infected patients. *Chest* 112: 646–653.

- GOLD (2008). GOLD, version 4.0: Cambridge Crystallographic Data Centre; Cambridge, UK.
- Gong Q, Jones MA, Zhou Z (2006). Mechanisms of pharmacological rescue of trafficking-defective hERG mutant channels in human long QT syndrome. *J Biol Chem* 281: 4069–4074.
- Han S, Zhang Y, Chen Q, Duan Y, Zheng T, Hu X *et al.* (2011). Fluconazole inhibits hERG K(+) channel by direct block and disruption of protein trafficking. *Eur J Pharmacol* 650: 138–144.
- Haverkamp W, Breithardt G, Camm AJ, Janse MJ, Rosen MR, Antzelevitch C *et al.* (2000). The potential for QT prolongation and pro-arrhythmia by non-anti-arrhythmic drugs: clinical and regulatory implications. Report on a Policy Conference of the European Society of Cardiology. *Cardiovasc Res* 47: 219–233.
- Hess B, Kutzner C, van der Spoel D, Lindahl E (2008). GROMACS 4: algorithms for highly efficient, load-balanced, and scalable molecular simulation. *J Chem Theory Comput* 4: 435–447.
- van der Heyden MA, Smits ME, Vos MA (2008). Drugs and trafficking of ion channels: a new pro-arrhythmic threat on the horizon? *Br J Pharmacol* 153: 406–409.
- Hornak V, Abel R, Okur A, Strockbine B, Roitberg A, Simmerling C (2006). Comparison of multiple amber force fields and development of improved protein backbone parameters. *Proteins* 65: 712–725.
- Jha TK (1983). Evaluation of diamidine compound (pentamidine isethionate) in the treatment resistant cases of kala-azar occurring in North Bihar, India. *Trans R Soc Trop Med Hyg* 77: 167–170.
- Jones SK, Hall JE, Allen MA, Morrison SD, Ohemeng KA, Reddy VV *et al.* (1990). Novel pentamidine analogs in the treatment of experimental *Pneumocystis carinii* pneumonia. *Antimicrob Agents Chemother* 34: 1026–1030.
- Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML (1983). Comparison of simple potential functions for simulating liquid water. *J Chem Phys* 79: 926–935.
- Kamiya K, Niwa R, Mitcheson JS, Sanguinetti MC (2006). Molecular determinants of HERG channel block. *Mol Pharmacol* 69: 1709–1716.
- Kannankeril P, Roden DM, Darbar D (2010). Drug-induced long QT syndrome. *Pharmacol Rev* 62: 760–781.
- Katchman AN, Koerner J, Tosaka T, Woosley RL, Ebert SN (2006). Comparative evaluation of HERG currents and QT intervals following challenge with suspected torsadogenic and nontorsadogenic drugs. *J Pharmacol Exp Ther* 316: 1098–1106.
- Kilkenny C, Browne W, Cuthill IC, Emerson M, Altman DG (2010). NC3Rs Reporting Guidelines Working Group. *Br J Pharmacol* 160: 1577–1579.
- Knappe K, Linder T, Wolschann P, Beyer A, Strydom A (2011). In silico analysis of conformational changes induced by mutation of aromatic binding residues: consequences for drug binding in the hERG K+ channel. *Plos ONE* 6: e28778.
- Kuryshv YA, Ficker E, Wang L, Hawryluk P, Dennis AT, Wible BA *et al.* (2005). Pentamidine-induced long QT syndrome and block of hERG trafficking. *J Pharmacol Exp Ther* 312: 316–323.
- Lees-Miller JP, Duan Y, Teng GQ, Duff HJ (2000a). Molecular determinant of high-affinity dofetilide binding to HERG1 expressed in *Xenopus* oocytes: involvement of S6 sites. *Mol Pharmacol* 57: 367–374.
- Lees-Miller JP, Duan Y, Teng GQ, Thorstad K, Duff HJ (2000b). Novel gain-of-function mechanism in K(+) channel-related long-QT syndrome: altered gating and selectivity in the HERG1 N629D mutant. *Circ Res* 86: 507–513.
- Loo TW, Clarke DM (1997). Correction of defective protein kinesin of human P-glycoprotein mutants by substrates and modulators. *J Biol Chem* 272: 709–712.
- McGrath J, Drummond G, McLachlan E, Kilkenny C, Wainwright C (2010). Guidelines for reporting experiments involving animals: the ARRIVE guidelines. *Br J Pharmacol* 160: 1573–1576.
- Nalos L, de Boer TP, Houtman MJ, Rook MB, Vos MA, van der Heyden MA (2011). Inhibition of lysosomal degradation rescues pentamidine-mediated decreases of K(IR)2.1 ion channel expression but not that of K(v)11.1. *Eur J Pharmacol* 652: 96–103.
- Pearlstein R, Vaz R, Rampe D (2003). Understanding the structure-activity relationship of the human ether-a-go-go-related gene cardiac K+ channel. A model for bad behavior. *J Med Chem* 46: 2017–2022.
- Rajamani S, Eckhardt LL, Valdivia CR, Klemens CA, Gillman BM, Anderson CL *et al.* (2006). Drug-induced long QT syndrome: hERG K+ channel block and disruption of protein trafficking by fluoxetine and norfluoxetine. *Br J Pharmacol* 149: 481–489.
- Roden DM, Viswanathan PC (2005). Genetics of acquired long QT syndrome. *J Clin Invest* 115: 2025–2032.
- Sanguinetti MC, Tristani-Firouzi M (2006). hERG potassium channels and cardiac arrhythmia. *Nature* 440: 463–469.
- Shagufra GD, Guo D, Klaasse E, de Vries H, Brussee J, Nalos L *et al.* (2009). Exploring chemical substructures essential for HERG k(+) channel blockade by synthesis and biological evaluation of dofetilide analogues. *ChemMedChem* 4: 1722–1732.
- Siu SW, Vacha R, Jungwirth P, Bockmann RA (2008). Biomolecular simulations of membranes: physical properties from different force fields. *J Chem Phys* 128: 125103.
- Strydom A, Wacker SJ, Boukharta L, Zachariae U, Karimi-Nejad Y, Aqvist J *et al.* (2010). Toward a consensus model of the hERG potassium channel. *ChemMedChem* 5: 455–467.
- Takemasa H, Nagatomo T, Abe H, Kawakami K, Igarashi T, Tsurugi T *et al.* (2008). Coexistence of hERG current block and disruption of protein trafficking in ketoconazole-induced long QT syndrome. *Br J Pharmacol* 153: 439–447.
- Thomsen MB, Volders PG, Stengl M, Spatjens RL, Beekman JD, Bischoff U *et al.* (2003). Electrophysiological safety of sertindole in dogs with normal and remodeled hearts. *J Pharmacol Exp Ther* 307: 776–784.
- Tidwell RR, Jones SK, Geratz JD, Ohemeng KA, Cory M, Hall JE (1990). Analogues of 1,5-bis(4-aminophenoxy)pentane (pentamidine) in the treatment of experimental *Pneumocystis carinii* pneumonia. *J Med Chem* 33: 1252–1257.
- Varkevisser R, Houtman MJ, Waasdorp M, Man JC, Heukers R, Takanari H *et al.* (2013). Inhibiting the clathrin-mediated endocytosis pathway rescues K(IR)2.1 downregulation by pentamidine. *Pflugers Arch* 465: 247–259.
- Warmke JW, Ganetzky B (1994). A family of potassium channel genes related to eag in *Drosophila* and mammals. *Proc Natl Acad Sci U S A* 91: 3438–3442.
- Wharton JM, Demopoulos PA, Goldschlager N (1987). Torsade de pointes during administration of pentamidine isethionate. *Am J Med* 83: 571–576.

Winckels SK, Thomsen MB, Oosterhoff P, Oros A, Beekman JD, Attevelt NJ *et al.* (2007). High-septal pacing reduces ventricular electrical remodeling and proarrhythmia in chronic atrioventricular block dogs. *J Am Coll Cardiol* 50: 906–913.

Wolf MG, Hoefling M, Aponte-Santamaria C, Grubmuller H, Groenhof G (2010). g_membed: efficient insertion of a membrane protein into an equilibrated lipid bilayer with minimal perturbation. *J Comput Chem* 31: 2169–2174.

Zhou Z, Gong Q, Epstein ML, January CT (1998). HERG channel dysfunction in human long QT syndrome. Intracellular transport and functional defects. *J Biol Chem* 273: 21061–21066.

Zhou Z, Gong Q, January CT (1999). Correction of defective protein trafficking of a mutant HERG potassium channel in human long QT syndrome. Pharmacological and temperature effects. *J Biol Chem* 274: 31123–31126.