



NIH PUBLIC ACCESS

Author Manuscript

Int J Eat Disord. Author manuscript; available in PMC 2011 January 1.

Published in final edited form as:

Int J Eat Disord. 2010 January ; 43(1): 14–21. doi:10.1002/eat.20670.**Substance Use Disorders in Women with Anorexia Nervosa**

Tammy Root, Ph.D.¹, Andréa Poyastro Pinheiro, M.D., Ph.D.¹, Laura Thornton, Ph.D.¹, Michael Strober, Ph.D.², Fernando Fernandez-Aranda, Ph.D.^{3,4}, Harry Brandt, M.D.⁵, Steve Crawford, M.D.⁵, Manfred M. Fichter, M.D.⁶, Katherine A. Halmi, M.D.⁷, Craig Johnson, Ph.D.⁸, Allan S. Kaplan, M.D., F.R.A.C.P.⁹, Kelly L. Klump, Ph.D.¹⁰, Maria La Via, M.D.¹, James Mitchell, M.D.¹¹, D. Blake Woodside, M.D.⁹, Alessandro Rotondo, M.D.¹², Wade H. Berrettini, M.D.¹³, Walter H. Kaye, M.D.¹⁴, and Cynthia M. Bulik, Ph.D.^{1,15}

¹ Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC² Semel Institute for Neuroscience and Human Behavior, and Resnick Neuropsychiatric Hospital, David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, CA³ Department of Psychiatry, University Hospital of Bellvitge, Feixa Llarga s/n, PC: 08907 Barcelona, Spain⁴ Ciber Fisiopatología Obesidad y Nutrición (CIBEROBN), Instituto Salud Carlos III., Hospital Clínico Santiago de Compostela, Choupana s/n, PC: 15706 Santiago de Compostela, Spain⁵ Department of Psychiatry, University of Maryland School of Medicine, Baltimore, MD⁶ Department of Psychiatry, University of Munich (LMU), Munich, Germany and Roseneck Hospital for Behavioral Medicine, Prien, Germany⁷ New York Presbyterian Hospital-Westchester Division, Weill Medical College of Cornell University, White Plains, NY⁸ Laureate Psychiatric Clinic and Hospital, Tulsa, OK⁹ Department of Psychiatry, The Toronto Hospital, Toronto, Canada¹⁰ Department of Psychology, Michigan State University, East Lansing, MI¹¹ Department of Psychiatry, Neuropsychiatric Research Institute, Fargo, ND¹² Neuropsychiatric Research Biotechnologies, University of Pisa, Italy¹³ Department of Psychiatry, University of Pennsylvania, School of Medicine, Philadelphia, PA¹⁴ Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA¹⁵ Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, NC**Abstract**

Objective—We examined prevalence of substance use disorders (SUD) in women with: (1) anorexia nervosa (AN) restricting type (RAN); (2) AN with purging only (PAN); (3) AN with binge eating only (BAN); and (4) lifetime AN and bulimia nervosa (ANBN). Secondary analyses examined SUD related to lifetime purging behavior and lifetime binge eating.

Method—Participants ($N = 731$) were drawn from the International Price Foundation Genetic Studies.

Results—The prevalence of SUD differed across AN subtypes, with more in the ANBN group reporting SUD than those in the RAN and PAN groups. Individuals who purged were more likely to report substance use than those who did not purge. Prevalence of SUD differed across lifetime binge eating status.

Conclusion—SUD are common in AN and are associated with bulimic symptomatology. Results underscore the heterogeneity in AN, highlighting the importance of screening for SUD across AN subtypes.

Keywords

eating disorders; anorexia nervosa; bulimia nervosa; drug use; alcohol related disorders; cannabis

Strong evidence demonstrates that eating disorders and substance use disorders (SUD) co-occur among individuals.^{1–5} The prevalence of drug and alcohol abuse is approximately 50% in individuals with an eating disorder, compare with a prevalence of approximately 9% in the general population.¹ Similarly, among individuals with SUD, over 35% report having an eating disorder compared with a 1–3% prevalence of eating disorders in the general population.^{1, 2} However, of theoretical importance is that among persons with some form of an eating disorder, nicotine, alcohol, and illicit drug use are more common among those who binge eat.¹ Thus, inconsistencies across studies in reported rates of co-occurrence of eating disorders and SUD³ along with inter-study differences in sample characteristics and general methodology may partly influence this variation.

The suggestion of greater liability to drug use and abuse among persons who binge eat compared to those with restricting type anorexia nervosa^{2, 3, 4} is intriguing, if not counterintuitive, in that among laboratory animals, food restriction enhances, while food-satiation reduces, self-administration of nearly every licit and illicit substance with abuse potential.⁵ Interestingly, risk of substance abuse may nevertheless be moderated by differences in consummatory patterns across AN subtypes. Strober et al.⁶ demonstrated that the 10-year prospective risk of incident cases of SUD in persons hospitalized for AN was six-times greater among individuals who reported binge eating while underweight compared with those with no history of binge eating up to the time of index hospitalization. This risk was also associated with elevated alcohol use disorder among first-degree relatives. Moreover, alcohol abuse among those who developed binge eating subsequent to weight restoration did not differ significantly from those who maintained a restricting profile throughout the study period. Bulik et al.⁷ similarly reported that alcohol use disorders were significantly more prevalent among women with AN with binge eating compared with those with AN without binge eating, although this pattern of results has not been universally replicated.^{8–11} Finally, little is known regarding the extent to which purging behavior, in the absence of binge eating,^{12, 13} is associated with risk for SUD in AN.

The motivation for the current study stems from: 1) discrepancies in the extant literature regarding the frequency with which lifetime SUD co-occurs with AN; 2) the dearth of studies examining SUD across well-defined AN subtypes; 3) reported findings suggesting the need to examine the extent to which SUD risk is exclusively associated with binge eating in the low weight state;⁶ and 4) the importance of examining a purging-only subtype of AN¹³ with reference to SUD. To address these questions, the aims of the current study are: 1) to examine the prevalence of SUD in a large sample of diagnostically well-categorized women with AN; 2) to compare the prevalence of SUD across AN subtypes; 3) to determine whether SUD are more common in those who report binge eating in the underweight state compared to those who develop binge eating at normal weight (i.e., not during episodes of AN); and, 4) to examine the specific associations of SUD with binge eating and purging. A large international collaboration involving sites in North America and Europe whose goal is to identify

susceptibility genes associated with eating disorders¹⁴ served as the sample for the current study, and subsequently represents the largest study to date examining the phenotypic comorbidity of SUD in AN.

METHODS

Participants

Participants were drawn from the International Price Foundation Genetic Study of AN Trios sample, which recruited male and female probands affected with AN from nine sites in North America and Europe. Self-report, clinical, and blood sample data were collected on affected probands. The study was approved by the appropriate ethical review boards and all participants completed written informed consent before participation. A detailed description of the study design is described elsewhere.¹⁵

For the current study, only female participants were included as the number of males was too small for independent analysis or for inclusion as a control for sex effects, resulting in a final sample size of $N = 731$. Data on eating disorders and substance use were collected from self-report and clinical assessments. Blood sample data were not used.

Measures

Demographic and Clinical Variables—Demographic information included age at time of interview, duration of eating disorders, and highest and lowest self-reported lifetime body mass index (BMI kg/m²).

Eating Disorder Diagnosis—Modified lifetime history of eating disorders (i.e., amenorrhea criterion not required) was assessed using the Structured Inventory of Anorexia Nervosa and Bulimic Syndromes (SIAB-EX), a semi-structured clinical interview designed to establish DSM-IV and ICD-10 eating disorder diagnoses¹⁶ and with an expanded version of Module H of the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID).¹⁷ Information regarding recovery status was obtained from the SCID and information regarding presence or absence of component features of eating disorder psychopathology (e.g., restrained eating, binge eating, purging) was obtained from the SIAB-EX.

Participants were classified into one of four AN diagnostic subgroups based on lifetime history: (1) AN – restricting (RAN) - restricted food intake without purging behavior; (2) AN –purging (PAN) - met diagnostic criteria for AN and reported purging behavior without binge eating; (3) AN - binge eating (BAN) - met criteria for AN and reported binge eating with or without purging; and (4) AN - bulimia nervosa (ANBN) - lifetime history of AN and normal weight BN.

In addition to diagnostic subtypes, we assessed the relation between SUD and purging behavior by dividing the sample by presence or absence of lifetime purging as determined by responses to the SCID and SIAB-EX. Individuals defined as reporting lifetime purging reported engaging in vomiting, or use of enemas, ipecac, laxatives, and/or diuretics at the subthreshold (less than twice a week) or threshold level (at least twice a week for at least three months).

Substance Use—Lifetime alcohol abuse and dependence were assessed using the SCID.¹⁸ Two groups were created based on DSM-IV criteria for alcohol abuse or dependence:¹⁹ (1) no alcohol abuse/dependence; and (2) alcohol abuse/dependence.

Lifetime drug abuse and dependence were assessed using the following categories of items from the SCID: cannabis, sedatives, stimulants, opiates, hallucinogens, cocaine, and other substances. Abuse and dependence were assessed if participants reported using a substance

more than 10 times per month (no duration criterion required). Three groups were created: (1) those who reported never having tried drugs or only having tried a substance once were classified in the “no use or experimenters” group; (2) those engaging in drug use more than 5 times but not meeting DSM-IV criteria for abuse/dependence were categorized in the “drug use” group; and, (3) individuals meeting criteria for drug abuse or dependence were categorized in the “drug abuse/dependence” group.

Statistical Analyses

Analyses were conducted using SAS version 9.1^{20, 21} with the GENMOD procedure. Age at time of interview was entered as a covariate. Prevalence of substance use was calculated across AN subtypes. Logistic regression analysis was used to test for significant differences in prevalence of substance use across AN subtypes. Substance use was examined with reference to purging status and binge eating status. All reported *p*-values were adjusted for multiple testing using the method of false discovery rate.²²

RESULTS

Demographic Variables

Table 1 describes the demographic variables across AN subtypes. Participants ranged in age from 13–58 years with a mean age of 26.7 years. The mean for highest BMI ranged from 20.9 to 22.9 kg/m² and the mean for lowest BMI ranged from 13.5 to 14.4 kg/m². Participants in the RAN group were the youngest at the time of interview (i.e., 24.7 years) and those in the ANBN group were the oldest at time of interview (i.e., 28.5 years). Average eating disorder duration was longest for those in the ANBN group (i.e., 11.2 years) and shortest for those in the RAN group (i.e., 7.7 years).

Eating Disorder Subtypes

The RAN group (*n* = 328) comprised 44.9% of the sample, the PAN group (*n* = 184) comprised 25.2% of the sample, 14.9% was comprised of the BAN (*n* = 109) group, and 15.0% of the sample was comprised of the ANBN group (*n* = 110).

Substance Use across AN Subtypes

Table 2 presents prevalence of alcohol and drug use and abuse/dependence across AN subtypes and, due to space limitations, Table 3 presents only the statistically significant odds ratios for AN subtypes pairwise comparisons.

Alcohol Use—Across the total sample, 19.8% met criteria for lifetime history of alcohol abuse/dependence. Prevalence was highest in ANBN (35.5%) and lowest in RAN (13.7%).

Risk for alcohol abuse/dependence was 3.20 times greater in the ANBN group and 1.85 times greater in the BAN group than the RAN group. Risk for alcohol abuse/dependence was 2.24 times greater in the ANBN group compared with the PAN group (χ^2 (3, *N* = 725) = 20.29, *p* < 0.001).

Drug Use—Across the total sample, 25.9% reported lifetime drug use and an additional 13.8% met criteria for lifetime history of drug abuse/dependence. The RAN group had the lowest percentage of drug use (i.e., 23.2%) and drug abuse/dependence (i.e., 6.4%), the PAN group had the highest percentage of drug use (i.e., 29.9%), and the ANBN group had the highest percentage of drug abuse/dependence (31.8%; Table 2).

Across AN subtypes, statistically significant differences emerged in the proportion of individuals meeting criteria for drug use, χ^2 (3, *N* = 730) = 37.62, *p* < 0.001. Individuals in the

ANBN group were 1.93 times more likely to report any drug use in comparison to individuals in the RAN group. The risk for drug abuse/dependence was 6.25 times greater in individuals with ANBN, 2.84 times greater in individuals with BAN, and 2.27 times greater in individuals with PAN, compared with those in the RAN group. Also, risk for drug abuse/dependence was 2.78 and 2.17 times greater in those with ANBN compared to those with PAN and BAN, respectively.

Drug Use Category—Table 2 presents the prevalence for each drug category across AN subtypes. Because abuse/dependence prevalence was low for the drug categories across subtypes, we combined the drug use and abuse/dependence into one group. Across AN subtypes, cannabis was the most frequently reported drug used. Analyses revealed statistically significant differences in all drug use categories across AN subtypes (Table 2).

AN subtypes pairwise comparison odds ratio by drug use category are presented in Table 3. Risk for all drug use categories was greatest in the ANBN group and lowest in the RAN group. Those in the ANBN group were 6.55 times more likely to report stimulant use, 5.89 times more likely to report cocaine use, 4.72 times more likely to report sedative use, and 5.44 times more likely to report ‘other’ drug use compared to those in the RAN group. See Table 3 for a complete list of all pairwise comparisons.

Substance Use by Lifetime Purging Status

Table 4 presents prevalence and odds ratios of substance use by purging status. Lifetime purging behavior was endorsed by 53.5% of the sample. Lifetime purging was associated with greater alcohol abuse/dependence, $\chi^2(1, N = 720) = 9.70, p = 0.002$. Those in the purging group were 1.83 times as likely to meet criteria for alcohol abuse/dependence compared to the no purging group. The relation between drug abuse/dependence and purging behavior was also statistically significant, $\chi^2(1, N = 725) = 32.25, p < 0.001$, with those in the purging group 1.67 times more likely to be at risk for drug use and 3.79 times more likely to meet criteria for drug abuse/dependence compared to the no purging group.

Substance Use by Lifetime Binge Eating Status

Table 5 presents prevalence of substance use by lifetime binge eating status. The relation between substance use categories and binge eating was examined by creating a binge-eating grouping variable: (1) no history of binge eating ($n = 522$; 71.4% of the sample); (2) history of binge eating at low weight ($n = 151$; 20.6% of the sample); and (3) history of binge eating only at normal weight (i.e., not during an episode of AN; $n = 22$; 3.0% of the sample).

Statistically significant differences ($\chi^2(2, N = 689) = 10.85, p = 0.005$) emerged across the binge eating groups for alcohol abuse/dependence. Compared to the no binge eating group, risk for alcohol abuse/dependence was 1.99 times more likely in the binge eating at low weight group. No significant differences in alcohol abuse/dependence were found for the binge eating at low weight group and the binge eating at normal weight group.

Those in the binge eating at low weight group were approximately 2.79 times more likely to report drug abuse/dependence and those in the binge eating at normal weight group were 4.03 times more likely to report drug use compared with the no history of binge eating group ($\chi^2(2, N = 668) = 14.54, p = 0.001$). There were no statistically significant differences for drug use between the two binge eating groups.

DISCUSSION

This is the first study to examine the comorbidity between alcohol and drug use disorders and AN by investigating the prevalence of SUD across AN subtypes, by comparing individuals who report binge eating in the underweight state versus those who develop binge eating at normal weight (i.e., not during episodes of AN), and across the entire sample stratified by the presence of binge eating and purging. Our findings indicate that: (1) SUD are most common among individuals with the ANBN subtype; (2) those who endorse purging behavior have higher rates of SUD compared to those who do not report purging; and (3) prevalence of drug use differs across binge eating status.

While our observed prevalence of SUD was higher than other AN samples,^{3, 9, 10, 23–25} which could be explained by definitional issues (i.e., we used a subthreshold, broader definition for SUD) or cross-cultural differences in SUD — our findings are consistent with several previous studies reporting that individuals with BN or with a history of bulimic symptoms during the course of AN were more likely to report SUD compared to those with RAN.^{8–11, 26–30} Specifically, in the current study, ANBN participants reported significantly higher levels of drug abuse/dependence than all other AN subtypes, while the RAN group had the lowest prevalence of drug abuse/dependence. This finding supports previous research suggesting that ratio of alcohol abuse/dependence across a sample of inpatient females with BN, ANBN, and RAN was 9:5:1,¹⁰ as well as findings from a community sample with Canadian adolescents reporting that binge eaters, particularly those who compensated, were more likely to report substance use.²⁶ Likewise, dietary restraint and bulimic symptoms in Latina adolescents were positively correlated with alcohol, tobacco and illicit drug use.²⁹ Our finding that greater substance use occurred in the purging group suggests that there may be meaningful characteristics associated with classification based on the presence or absence of purging.^{12, 13} We also examined the relation of drug category across AN subtypes and found differences in the prevalence of the drug use category across AN subtypes. In the current study, the most frequently used drug was cannabis, followed by hallucinogens. These findings support previous research suggesting that greater pathological eating behavior is associated with not just alcohol and tobacco but also marijuana and other hard substances.²⁶ The prevalence of cannabis use being the highest is consistent with population norms³¹ and previous research suggesting that cannabis is the most frequently reported illicit drug among those with restricting and binge eating/purging symptomatology.^{8–11} Our findings are not surprising given that epidemiological data indicate that cannabis is the most frequently reported illicit drug.³¹ What is surprising is that results from this study challenge the commonly held belief that individuals with RAN report lower levels of drug use.³ Nonetheless it is important to note that the prevalence reported in this paper may partly reflect normative experimentation with alcohol, tobacco, and illicit substances, particularly during adolescence and young adulthood. Therefore, results should be interpreted with caution. Unexpectedly, hallucinogen use was the second most commonly reported substance used among those with RAN. While little research exists on the relation between hallucinogen use and eating disorder symptoms, it is possible that the appetite-suppressing effect of hallucinogens,³² along with the physiological response of changes in perceptions and thoughts which might allow one to “escape” the anxiety associated with the eating disorder, are motivating factors among those who restrict food intake. These motivations may be further encouraged given the availability of some hallucinogens on the internet.³³

Elaborating on the association between SUD and AN within the current study, findings also suggest that a higher prevalence of individuals in the BAN and ANBN groups reported greater sedative, stimulant and cocaine use compared to those in the RAN group, with those in the ANBN reporting the greatest use. This finding supports previous research suggesting that cocaine and amphetamine use is greater primarily in individuals with the binge eating/purging

subtype of AN.³⁰ The use of stimulants and cocaine among the BAN and ANBN groups may be due to the appetite suppressant effects these drugs may have; thus, their use may be an effort to avoid the consequences of overeating. It has also been hypothesized that an association between SUD and eating disorders reflects an underlying influence of personality traits such as heightened impulsivity.^{34, 35} Our study was not, however, designed to identify mechanisms underlying the relation between SUD and binge eating/purging AN.

Of final note, our findings suggest that binge eating was related to substance use under certain conditions. Risk for alcohol abuse/dependence and drug abuse/dependence was higher in those who reported binge eating at low weight relative to those who did not binge eat, but differences did not emerge between those who binge ate at low weight versus binge ate only after weight restoration. These findings are inconsistent with previous research suggesting that risk for excessive alcohol consumption is higher in individuals who exhibit binge eating at low weight compared to those who develop binge eating after restoration of normal weight.⁶ This inconsistency could be due to differences across study design (i.e., longitudinal vs. cross-sectional), and to our smaller sample size. It has also been suggested that reward hypersensitivity may be a common vulnerability factor for both hazardous drinking and disordered eating.^{36–38}

Limitations

Limitations to our study must be considered. First, participants are primarily of European ancestry and therefore cannot be generalized to other ancestry groups or to males. Second, a healthy comparison group was not available for comparison and our primary interest was examining differences in SUD across the AN subtypes. Third, we did not make distinctions between current and lifetime diagnoses for AN. It is possible that differences in substance use patterns would have emerged had such information been available. Fourth, substance use data also focused on lifetime use only, not frequency and duration of use which could differentiate AN subgroups in unique ways. Additionally, the categorization of drug use is a potential limitation in that the “drug use” group included those who used a substance at least twice but not enough to meet criteria for abuse/dependence. Thus, results related to the “drug use” group need to be interpreted given this knowledge. Finally, causal conclusions pertaining to the development of either eating disorders or SUD cannot be discussed. The nature of any casual relation between AN and SUD is unknown, thus we are unable to ascertain if the eating disorder lead to SUD or if the SUD lead to eating disorder symptomatology. Further, we are not aware if certain substances were specifically used for weight-loss. Additional unexamined factors may have influenced findings. Of particular relevance are data suggesting that depression, negative affect, and anxiety are associated with eating disorders and SUD.^{39, 40} Future studies should consider depression, negative affect, and anxiety as each relates to the comorbidity of eating disorders and SUD.

Conclusions and Implications

Results from this and other studies highlight the need to assess alcohol and drug use behavior when screening and treating individuals with AN, particularly those with bulimic symptomatology. Our findings support previous research indicating that individuals with lifetime diagnoses of both AN and BN (ANBN group in the current analysis) and those who engage in bulimic behaviors report more alcohol abuse/dependence and drug abuse/dependence than those who only engage in restricting behaviors. These findings are of clinical importance because prior research indicates that individuals presenting with both an eating disorder and SUD may be at heightened risk for physical health complications, including increased lethality,⁴¹ and additional psychological comorbidities.⁷ Given the comorbidity between AN and SUD, particularly among those with ANBN and BAN, it would appear

prudent and necessary to combine prevention and treatment efforts in order to better avert the emergence or advancement of these disorders.

Acknowledgments

Dr. Root was supported by National Institute of Health grant T32MH076694. The authors wish to thank the Price Foundation for the support of the clinical collection of participants and support of data analysis. The authors acknowledge the staff of the Price Foundation Collaborative Group for their efforts in participant screening and clinical assessments. The authors also wish to thank Xiaofei Mo, M.D. for her assistance in the preparation of this manuscript. The authors are indebted to the participating families for their contribution of time and effort in support of this study. This study was also supported by grant MH-66117 from the National Institutes of Health, Bethesda, MD.

References

1. CASA. Food for Thought: Substance Abuse and Eating Disorders. Report. New York: The National Center on Addiction and Substance Abuse (CASA) at Columbia University; 2003.
2. Krug I, Treasure J, Anderluh M, Bellodi L, Cellini E, di Bernardo M, et al. Present and lifetime comorbidity of tobacco, alcohol and drug use in eating disorders: a European multicenter study. *Drug and alcohol dependence* 2008 Sep 1;97(1–2):169–79. [PubMed: 18571341]
3. Holderness CC, Brooks-Gunn J, Warren MP. Co-morbidity of eating disorders and substance abuse review of the literature. *Int J Eat Disord* 1994 Jul;16(1):1–34. [PubMed: 7920577]
4. Salbach-Andrae H, Lenz K, Simmendinger N, Klinkowski N, Lehmkuhl U, Pfeiffer E. Psychiatric comorbidities among female adolescents with anorexia nervosa. *Child psychiatry and human development* 2008 Sep;39(3):261–72. [PubMed: 17987378]
5. Carr KD. Chronic food restriction: enhancing effects on drug reward and striatal cell signaling. *Physiol Behav* 2007 Aug 15;91(5):459–72. [PubMed: 17081571]
6. Strober M, Freeman R, Bower S, Rigali J. Binge eating in anorexia nervosa predicts later onset of substance use disorder: A ten-year prospective, longitudinal follow-up of 95 adolescents. *J Youth Adol* 1995;25:519–32.
7. Bulik CM, Klump KK, Thornton L, Kaplan AS, Devlin B, Fichter MM, et al. Alcohol use disorder comorbidity in eating disorders: a multicenter study. *Journal of Clinical Psychiatry* 2004;65(7):1000–6. [PubMed: 15291691]
8. Wiederman MW, Pryor T. Substance use and impulsive behaviors among adolescents with eating disorders. *Addict Behav* 1996 Mar-Apr;21(2):269–72. [PubMed: 8730530]
9. Corcos M, Nezelof S, Speranza M, Topa S, Girardon N, Guilbaud O, et al. Psychoactive substance consumption in eating disorders. *Eat Behav* 2001 Spring;2(1):27–38. [PubMed: 15001048]
10. Blinder BJ, Cumella EJ, Sanathara VA. Psychiatric comorbidities of female inpatients with eating disorders. *Psychosom Med* 2006 May-Jun;68(3):454–62. [PubMed: 16738079]
11. Stock SL, Goldberg E, Corbett S, Katzman DK. Substance use in female adolescents with eating disorders. *J Adolesc Health* 2002 Aug;31(2):176–82. [PubMed: 12127388]
12. Keel PK, Fichter M, Quadflieg N, Bulik CM, Baxter MG, Thornton L, et al. Application of a latent class analysis to empirically define eating disorder phenotypes. *Archives of General Psychiatry* 2004;61:192–200. [PubMed: 14757596]
13. Wade TD. A retrospective comparison of purging type disorders: eating disorder not otherwise specified and bulimia nervosa. *Int J Eat Disord* 2007 Jan;40(1):1–6. [PubMed: 16958118]
14. Kaye WH, Lilenfeld LR, Berrettini WH, Strober M, Devlin B, Klump KL, et al. A search for susceptibility loci for anorexia nervosa: methods and sample description. *Biol Psychiatry* 2000;47(9):794–803. [PubMed: 10812038]
15. Reba L, Thornton L, Tozzi F, Klump KL, Brandt H, Crawford S, et al. Relationships between features associated with vomiting in purging-type eating disorders. *Int J Eat Disord* 2005 Dec;38(4):287–94. [PubMed: 16261604]
16. Fichter MM, Herpertz S, Quadflieg N, Herpertz-Dahlmann B. Structured Interview for Anorexic and Bulimic disorders for DSM-IV and ICD-10: updated (third) revision. *Int J Eat Disord* 1998 Nov;24(3):227–49. [PubMed: 9741034]

17. First, M.; Spitzer, R.; Gibbon, M.; Williams, J. Structured Clinical Interview for DSM-IV Axis I Disorders, Research Version, Patient Edition. New York: Biometrics Research, New York State Psychiatric Institute; 1997.
18. First, M.; Gibbon, M.; Spitzer, R.; Williams, J. Users guide for the structured clinical interview for DSM IV Axis I disorders - research version (SCID-I, version 2.0). New York: New York State Psychiatric Institute; 1996.
19. APA APA. Diagnostic and Statistical Manual of Mental Disorders. Vol. 4. Washington, DC: American Psychiatric Press; 1994.
20. SAS Institute Inc. SAS/STAT=AE Software: Changes and Enhancements. Cary, NC: SAS Institute, Inc; 1996.
21. SAS Institute Inc. SAS/STAT® Software: Version 9. Cary, NC: SAS Institute, Inc.; 2004.
22. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society* 1995;57:289–300.
23. Herzog DB, Keller MB, Sacks NR, Yeh CJ, Lavori PW. Psychiatric comorbidity in treatment-seeking anorexics and bulimics. *J Am Acad Child Adolesc Psychiatry* 1992 Sep;31(5):810–8. [PubMed: 1400111]
24. Braun DL, Sunday SR, Halmi KA. Psychiatric comorbidity in patients with eating disorders. *Psychol Med* 1994 Nov;24(4):859–67. [PubMed: 7892354]
25. Jordan J, Joyce PR, Carter FA, Horn J, McIntosh VV, Luty SE, et al. Anxiety and psychoactive substance use disorder comorbidity in anorexia nervosa or depression. *Int J Eat Disord* 2003 Sep;34(2):211–9. [PubMed: 12898557]
26. Ross HE, Ivis F. Binge eating and substance use among male and female adolescents. *Int J Eat Disord* 1999 Nov;26(3):245–60. [PubMed: 10441240]
27. Nagata T, Kawarada Y, Ohshima J, Iketani T, Kiriike N. Drug use disorders in Japanese eating disorder patients. *Psychiatry Res* 2002 Mar 15;109(2):181–91. [PubMed: 11927143]
28. von Ranson KM, Iacono WG, McGue M. Disordered eating and substance use in an epidemiological sample: I. Associations within individuals. *Int J Eat Disord* 2002 May;31(4):389–403. [PubMed: 11948644]
29. Granillo T, Jones-Rodriguez G, Carvajal SC. Prevalence of eating disorders in Latina adolescents: associations with substance use and other correlates. *J Adolesc Health* 2005 Mar;36(3):214–20. [PubMed: 15737777]
30. Herzog DB, Franko DL, Dorer DJ, Keel PK, Jackson S, Manzo MP. Drug abuse in women with eating disorders. *Int J Eat Disord* 2006 Jul;39(5):364–8. [PubMed: 16565976]
31. Compton WM, Thomas YF, Stinson FS, Grant BF. Prevalence, correlates, disability, and comorbidity of DSM-IV drug abuse and dependence in the United States: results from the national epidemiologic survey on alcohol and related conditions. *Arch Gen Psychiatry* 2007 May;64(5):566–76. [PubMed: 17485608]
32. Curran HV, Robjant K. Eating attitudes, weight concerns and beliefs about drug effects in women who use ecstasy. *J Psychopharmacol* 2006 May;20(3):425–31. [PubMed: 16574717]
33. Halpern JH, Pope HG Jr. Hallucinogens on the Internet: a vast new source of underground drug information. *Am J Psychiatry* 2001 Mar;158(3):481–3. [PubMed: 11229993]
34. Bardone-Cone AM, Abramson LY, Vohs KD, Heatherton TF, Joiner TE Jr. Predicting bulimic symptoms: an interactive model of self-efficacy, perfectionism, and perceived weight status. *Behav Res Ther* 2006 Jan;44(1):27–42. [PubMed: 16301012]
35. Bardone-Cone AM, Wonderlich SA, Frost RO, Bulik CM, Mitchell JE, Uppala S, et al. Perfectionism and eating disorders: current status and future directions. *Clin Psychol Rev* 2007 Apr;27(3):384–405. [PubMed: 17267086]
36. Loxton NJ, Dawe S. Alcohol abuse and dysfunctional eating in adolescent girls: the influence of individual differences in sensitivity to reward and punishment. *Int J Eat Disord* 2001 May;29(4):455–62. [PubMed: 11285583]
37. Loxton NJ, Dawe S. Reward and punishment sensitivity in dysfunctional eating and hazardous drinking women: associations with family risk. *Appetite* 2006 Nov;47(3):361–71. [PubMed: 16846665]

38. Davis C, Claridge G. The eating disorders as addiction: a psychobiological perspective. *Addict Behav* 1998 Jul-Aug;23(4):463–75. [PubMed: 9698975]
39. Measelle JR, Stice E, Springer DW. A prospective test of the negative affect model of substance abuse: moderating effects of social support. *Psychol Addict Behav* 2006 Sep;20(3):225–33. [PubMed: 16938060]
40. Measelle JR, Stice E, Hogansen JM. Developmental trajectories of co-occurring depressive, eating, antisocial, and substance abuse problems in female adolescents. *J Abnorm Psychol* 2006 Aug;115(3):524–38. [PubMed: 16866592]
41. Keel PK, Dorer DJ, Eddy KT, Franko D, Charatan DL, Herzog DB. Predictors of mortality in eating disorders. *Arch Gen Psychiatry* 2003 Feb;60(2):179–83. [PubMed: 12578435]

Table 1

Demographic Characteristics across AN Subtypes.

| | RAN (n = 328) Mean (SD) | PAN (n=184) Mean (SD) | BAN (n=109) Mean (SD) | ANBN (n=110) Mean (SD) |
|---|--|--------------------------------------|--------------------------------------|---------------------------------------|
| Age at interview (years) | 24.7 (7.4) | 26.4 (7.0) | 27.5 (8.5) | 28.5 (8.8) |
| Highest lifetime BMI (kg/m ²) | 20.9 (2.3) | 21.3 (2.4) | 21.0 (2.2) | 22.9 (2.1) |
| Lowest lifetime BMI (kg/m ²) | 13.5 (1.8) | 13.8 (1.9) | 13.5 (2.0) | 14.4(1.8) |
| Eating disorder duration (years) | 7.7 (6.7) | 9.5 (6.7) | 10.2 (8.6) | 11.2 (8.9) |

Table 2
Prevalence of Alcohol Abuse/dependence, Drug Use and Abuse/Dependence, and Drug Use by Drug Category across AN Subtypes.

| | RAN (n = 328) n (%) | PAN (n = 184) n (%) | BAN (n = 109) n (%) | ANBN (n = 110) n (%) | χ^2 (p-value) df = 3 |
|---|---------------------------|---------------------------|---------------------------|----------------------------|------------------------------|
| <u>Alcohol Use</u> | | | | | |
| No Abuse/dependence | 281 (85.7) | 147 (79.9) | 83 (76.1) | 70 (63.6) | 20.29 (<.001) |
| Abuse/dependence | 45 (13.7) | 35 (19.0) | 26 (23.9) | 39 (35.5) | |
| <u>Drug Use and Drug Abuse/dependence</u> | | | | | |
| No use/experimenters | 231 (70.4) | 103 (56.0) | 62 (56.9) | 45 (40.9) | |
| Drug Use | 76 (23.2) | 55 (29.9) | 28 (25.7) | 30 (27.3) | 37.62 (<.001) |
| Abuse/dependence | 21 (6.4) | 26 (14.1) | 19 (17.4) | 35 (31.8) | |
| <u>Drug Use by Drug Category*</u> | | | | | |
| Sedatives | 10 (3.0) | 13 (7.1) | 13 (11.9) | 17 (15.5) | 16.55 (.001) |
| Cannabis | 93 (28.4) | 64 (34.8) | 40 (36.7) | 60 (54.5) | 20.44 (<.001) |
| Stimulants | 9 (2.7) | 11 (6.0) | 12 (11.0) | 19 (17.3) | 23.19 (<.001) |
| Opioids | 9 (2.7) | 10 (5.4) | 11 (10.1) | 11 (10.0) | 11.26 (.010) |
| Cocaine | 11 (3.4) | 17 (9.2) | 13 (11.9) | 18 (16.4) | 18.50 (<.001) |
| Hallucinogens | 26 (7.9) | 23 (12.5) | 13 (11.9) | 25 (22.7) | 15.82 (.002) |
| Other | 10 (3.0) | 16 (8.7) | 7 (6.4) | 17 (15.5) | 17.96 (<.001) |

* Drug use by drug category is defined as having used the substance more than once regardless of abuse or dependence. Only endorsement values are listed.

Table 3

Statistically Significant Odds Ratios for Alcohol Abuse/Dependence, Drug Use, Drug Abuse/Dependence, and Drug Use by Drug Category by AN subtypes.

| Substance | AN subtypes comparison | Odds Ratio (95% CI) |
|---|------------------------|---------------------|
| <u>Alcohol</u> | | |
| Alcohol Abuse/Dependence | BAN > RAN | 1.85 (1.80, 2.00) |
| Alcohol Abuse/Dependence | ANBN > RAN | 3.20 (1.92, 5.33) |
| Alcohol Abuse/Dependence | ANBN > PAN | 2.24 (1.30, 3.84) |
| <u>Drug Use and Drug Abuse/dependence</u> | | |
| Drug Use (no abuse/dependence) | ANBN > RAN | 1.93 (1.13, 3.29) |
| Drug Abuse/Dependence | PAN > RAN | 2.27 (1.22, 4.17) |
| Drug Abuse/Dependence | BAN > RAN | 2.84 (1.47, 5.56) |
| Drug Abuse/Dependence | ANBN > RAN | 6.25 (3.45, 11.11) |
| Drug Abuse/Dependence | ANBN > PAN | 2.78 (1.53, 5.00) |
| Drug Abuse/Dependence | ANBN > BAN | 2.17 (1.14, 4.17) |
| <u>Drug Use by Drug Category</u> | | |
| Sedatives* | BAN > RAN | 3.69 (1.54, 8.79) |
| Sedatives* | ANBN > RAN | 4.72 (2.06, 10.81) |
| Cannabis* | ANBN > RAN | 2.82 (1.79, 4.43) |
| Cannabis* | ANBN > PAN | 2.17 (1.33, 3.52) |
| Cannabis* | ANBN > BAN | 2.04 (1.18, 3.51) |
| Stimulants* | BAN > RAN | 4.00 (1.62, 9.34) |
| Stimulants* | ANBN > RAN | 6.55 (2.84, 15.12) |
| Stimulants* | ANBN > PAN | 3.05 (1.38, 6.74) |
| Opiates* | BAN > RAN | 3.76 (1.50, 9.39) |
| Opiates* | ANBN > RAN | 3.64 (1.44, 9.15) |
| Cocaine* | PAN > RAN | 2.78 (1.27, 6.09) |
| Cocaine* | BAN > RAN | 3.51 (1.51, 8.15) |
| Cocaine* | ANBN > RAN | 5.89 (2.21, 10.85) |
| Hallucinogens* | ANBN > RAN | 3.51 (1.91, 6.46) |
| Hallucinogens* | ANBN > PAN | 2.09 (1.12, 3.92) |
| Hallucinogens* | ANBN > BAN | 2.19 (1.05, 4.56) |
| Other* | PAN > RAN | 2.94 (1.30, 6.64) |
| Other* | ANBN > RAN | 5.44 (2.38, 12.42) |
| Other* | ANBN > BAN | 2.62 (1.04, 6.61) |

* Drug use by drug category is defined as having used the substance more than once regardless of abuse or dependence. Only endorsement values are listed.

Table 4

Prevalence of Alcohol and Drug Use by Purging Status and Odds Ratios with 95% Confidence Intervals.

| | No Purging Behavior <i>n</i> = 335 <i>n</i> (%) | Purging Behavior <i>n</i> = 386 <i>n</i> (%) | χ^2 (p-value) df = 1 | Odds Ratio (95% CI) |
|----------------------|---|--|------------------------------|------------------------|
| <u>Alcohol Use</u> | | | | |
| No Abuse/dependence | 287 (85.7) | 290 (75.1) | 9.70 (.002) | ----- |
| Abuse/dependence | 48 (14.3) | 96 (24.9) | | 1.83 (1.24, 2.70) |
| <u>Drug Use</u> | | | | |
| No use/experimenters | 238 (70.6) | 199 (51.2) | 32.25 (<.001) | ----- |
| Use | 77 (22.9) | 112 (28.8) | | 1.67* (1.18, 2.37) |
| Abuse/dependence | 22 (6.5) | 78 (20.0) | | 3.79* (2.26, 6.35) |

* Comparisons are made with respect to the No use/experimenters group.

Note. Dashes indicate reference group.

Table 5
Prevalence of Alcohol Abuse/Dependence, Drug Use Group and Abuse/Dependence of Specific Drug Categories across Binge Eating Status.

| | No Binge Eating (n (%)) | Binge Eating at Low Weight n (%) | Binge Eating at Normal Weight n (%) | χ^2 (p-value) df = 12 | Odds Ratio (95% CI) |
|---|----------------------------|-------------------------------------|--|-------------------------------|------------------------|
| <u>Alcohol Use</u> | | | | | |
| No abuse/dependence | 435 (84.0) | 16 (72.7) | 107 (71.3) | | ----- |
| Alcohol abuse/dependence | 83 (16.0) | 43 (28.3) | 6 (27.3) | 10.85 (.005) | 1.99 (1.29, 3.06) |
| <u>Drug Use and Drug Abuse/dependence</u> | | | | | |
| No use/experimenters | 341 (65.3) | 11 (50.0) | 177 (51.0) | | ----- |
| Use | 133 (25.5) | 5 (22.7) | 80 (26.5) | 14.54 (.001) | 4.03 (1.41, 11.53) |
| Abuse/dependence | 48 (9.2) | 6 (27.3) | 34 (22.5) | | 2.79 (1.67, 4.07) |

Note. Dashes indicate reference group.