



Published in final edited form as:

J Trauma Dissociation. 2014 ; 15(5): 527–547. doi:10.1080/15299732.2014.908805.

Incidence and Predictors of Acute Psychological Distress and Dissociation after Motor Vehicle Collision: a Cross-Sectional Study

Gemma C. Lewis^{1,2}, Timothy F. Platts-Mills^{1,2}, Israel Liberzon³, Eric Bair⁴, Robert Swor⁵, David Peak⁶, Jeffrey Jones⁷, Niels Rathlev⁸, David Lee⁹, Robert Domeier¹⁰, Phyllis Hendry¹¹, and Samuel A. McLean^{1,2}

¹TRYUMPH Research Program, University of North Carolina, Chapel Hill, North Carolina

²School of Medicine, Departments of Anesthesiology and Emergency Medicine, University of North Carolina, Chapel Hill, North Carolina

³Department of Psychiatry, University of Michigan, Ann Arbor, Michigan

⁴School of Dentistry, University of North Carolina, Chapel Hill, North Carolina

⁵Department of Emergency Medicine, William Beaumont Hospital, Royal Oak, Michigan

⁶Department of Emergency Medicine, Massachusetts General Hospital, Boston, Massachusetts

⁷Department of Emergency Medicine, Spectrum Health – Butterworth Campus, Grand Rapids, Michigan

⁸Department of Emergency Medicine, Baystate Medical Center, Springfield, Massachusetts

⁹Department of Emergency Medicine, North Shore University Hospital, Manhasset, New York

¹⁰Department of Emergency Medicine, St. Joseph Mercy Hospital, Ann Arbor, Michigan

¹¹Department of Emergency Medicine and Pediatrics, University of Florida-Jacksonville, Jacksonville, Florida

Abstract

Objective—The authors examined the incidence and predictors of peritraumatic distress and dissociation after one of the most common forms of civilian trauma exposure: motor vehicle collision (MVC).

Methods—In this study, patients presenting to the emergency department after MVC who were without serious injury and discharged to home after evaluation (n = 935) completed an emergency department interview evaluating sociodemographic, collision-related, and psychological characteristics.

Results—The incidence and predictors of distress (Peritraumatic Distress Inventory score ≥ 23) and dissociation (Michigan Critical Events Perception Scale score >3) were assessed. Distress was

present in 355 of 935 patients (38%) and dissociation was present in 260 of 942 patients (28%). These outcomes showed only moderate correlation ($r = 0.45$), and had both shared and distinct predictors. Female gender, anxiety symptoms prior to MVC, and vehicle damage severity predicted both distress and dissociation. Higher socioeconomic status (higher education, higher income, full time employment) had a protective effect against distress but not dissociative symptoms. Better physical health and worse overall mental health were associated with increased risk of dissociation, but not distress. Distress but not dissociation was associated with lower patient confidence in recovery and a longer expected duration of recovery.

Conclusion—There are unique predictors of peritraumatic distress and dissociation. Further work is needed to better understand the neurobiology of peritraumatic distress and dissociation, and the influence of these peritraumatic outcomes on persistent psychological sequelae.

Keywords

Distress; Dissociation; Post-Traumatic Stress Disorder; Trauma; Motor Vehicle Collision

Motor vehicle collisions (MVCs) are among the most common life-threatening experiences, resulting in 50 million injuries worldwide and almost four million US emergency department (ED) visits each year (Peden, 2004; Niska, Bhuiya, & Xu, 2010). More than 90% of patients presenting to US EDs following MVC do not have injuries requiring hospitalization and are discharged to home following evaluation (Platts-Mills, Hunold, Esserman, Sloane, & McLean, 2012). However, persistent psychological sequelae such as posttraumatic stress disorder (PTSD) are common in this population (Buitenhuis, 2006; Ehlers, Mayou, & Bryant, 1998; Nishi et al., 2010) and result in substantial morbidity and health status decline (Ehlers et al., 1998; Kuch, 1996; Mayou, 1993).

Understanding which patients are at risk for PTSD is an important priority. Three distinct categories of risk factors have been described: extant patient characteristics, stressor characteristics, and response characteristics (Green, 1994; Shalev, Peri, Canetti, & Schreiber, 1996). The extant patient characteristics include socio-demographics, mental health, and past experiences (McFarlane, 2008). Stressor characteristics include severity of harm or physical threat and other dimensions of the event (Green, 1990). Individual response characteristics include psychological sequelae immediately following the event, in particular distress and dissociation. Importantly, acute psychological response predicts PTSD independent of the severity of the traumatic stressor (Feinstein & Dolan, 1991; Lawyer et al., 2006; Perry, Difede, Musngi, Frances, & Jacobsberg, 1992). Although there is no currently known mechanism by which PTSD can be prevented in all individuals (Shalev, 2009), developing evidence suggests that a window exists shortly after the exposure during which therapeutic intervention may decrease the risk of PTSD development in susceptible patients (Bryant, 2003; Litz, 2008). Indeed, early interventions in rape victims and soldiers exposed to traumatic events have been shown to reduce prevalence of PTSD (Foa, Rothbaum, Riggs, & Murdock, 1991; Solomon & Benbenishty, 1986). A better understanding of immediate response characteristics may facilitate early identification and referral for treatment of those at high risk of PTSD.

Distress symptoms include perception of life threat; fear; feelings of helplessness, horror, guilt and shame; anger; loss of bowel and bladder control; shaking or trembling; and increased heart rate (Brunet et al., 2001; Jehel, Brunet, Paterniti, & Guelfi, 2005; Nishi et al., 2009). Higher levels of peritraumatic distress have been shown to correlate with increased likelihood of acute stress disorders in older patients (Bui et al., 2010), assault victims (Jehel, Paterniti, Brunet, Louville, & Guelfi, 2006), earthquake survivors and rescue workers (Nishi et al., 2012; Nishi & Matsuoaka, 2013), sudden death survivors (Hargrave, Leathem, & Long, 2012), and in patients who have been involved in a severe MVC (Nishi et al., 2010); while the absence of peritraumatic distress strongly predicts absence of subsequent PTSD (Nishi et al., 2010). The association between peritraumatic distress and PTSD weakens as time between the traumatic event and the PTSD assessment increases (Thomas, Saumier, & Brunet, 2012), suggesting the involvement of other interim mediators.

Dissociative symptoms are characterized by alterations in the experience of time, place, and/or person during the course of the trauma exposure, characterized as derealization, numbing, detachment, reduced awareness of surroundings, and depersonalization (Marmar et al., 1994). It has been suggested that this alteration in perception may yield failures in memory processing and storage, thereby creating a subsequent sense of unreality surrounding recall of the traumatic event (Michaels, Michaels, Moon, et al., 1999). Dissociation in response to trauma is more common in patients reporting a prior history of exposure to combat (Branscomb, 1991), fires and other natural disasters (Cardena & Spiegel, 1993; Koopman, Classen, & Spiegel, 1994), child abuse (Chu, Frey, Ganzel, & Matthews, 1999), and interpersonal violence (Feeny, Zoellner, Fitzgibbons, & Foa, 2000), although whether this association results from increased reporting of prior trauma among those experiencing dissociation is unclear (Merckelbach & Muris, 2001). It has been proposed that dissociation is an adaptive mechanism to reduce the immediate psychological impact of a traumatic event (van der Kolk & van der Hart, 1989). Although dissociation may diminish the immediate psychological impact of trauma, observational data indicate that dissociative symptoms are associated with increased risk of PTSD development in victims of MVCs (Ehlers et al., 1998), emergency service workers (Marmar, Weiss, Metzler, & Delucchi, 1996), police officers (Martin, Marchand, Boyer, & Martin, 2009), Vietnam veterans (Bremner et al., 1992), and patients injured by other mechanisms (Michaels, Michaels, Moon, et al., 1999; Michaels, Michaels, Zimmerman, et al., 1999). A meta-analysis indicates that dissociation is the psychological response which most strongly predicts PTSD (Ozer, Best, Lipsey, & Weiss, 2003). Extant PTSD in turn predicts dissociative symptoms during subsequent traumatic events (Hetzl-Riggin & Wilber, 2010; Kaufman et al., 2002).

The relationship between peritraumatic dissociation and PTSD is, however, complex, and causality cannot be established based upon studies to date (Lensvelt-Mulders et al., 2008). Peritraumatic dissociation may not be predictive of PTSD when controlled for symptoms of persistent dissociative symptoms after the event (Briere, Scott, & Weathers, 2005; van der Velden & Wittmann, 2008). Additionally the relationship between peritraumatic dissociation and eventual development of PTSD may be partially mediated by avoidant coping (Pacella et al., 2011) and/or depression (Hodgson & Webster, 2011) in the subacute period following trauma, or may simply be reflective of underlying experiential avoidance as a predictor of

both dissociative symptoms and post-traumatic stress symptoms (Kumpula, Orcutt, Bardeen, & Varkovitzky, 2011).

There are several proposed relationships between peritraumatic distress and dissociation. Some authors have hypothesized that dissociation is a protective mechanism in which depersonalization and derealization occur to prevent the real-time experience of severe distress in response to trauma (Spiegel, Hunt, & Dondershine, 1988). Conversely, others have hypothesized that dissociation actually occurs as an epiphenomenon of high levels of distress (Bernat, Ronfeldt, Calhoun, & Arias, 1998; Friedman, 2000). It has been observed that dissociative symptoms often occur in the context of high levels of distress (Fikretoglu et al., 2006) and that dissociation may act as a mediator between the relationship between distress and PTSD development rather than being a distinct predictor (Otis, Marchand, & Courtois, 2012). Another possibility is that dissociation plays an important role in PTSD development in patients who are predisposed to dissociative reactions but not in others (Kihlstrom, Glisky, & Angiulo, 1994); dissociation is not necessary for PTSD to occur (Otis et al., 2012). Distress but not dissociation is associated with the development of PTSD in the elderly, further suggesting differences in underlying pathogenesis and relationship to PTSD between these two peritraumatic phenomena (Brunet et al., 2013). Little is known about the pathogenesis of these phenomena or what risk factors may predict distress and/or dissociation in response to trauma, and what the relationship between the two may entail. Prior investigation into risk factors for peritraumatic dissociation and acute distress symptoms has been minimally elucidative (Jaycox, Marshall, & Orlando, 2003). Understanding patient characteristics predictive of these acute responses may clarify their etiology and influence on PTSD development.

In this study we sought to characterize the incidence of distress and dissociation among a large cohort of patients presenting to the ED for care during the hours after MVC. We also assessed the strength of association between distress and dissociation in the early aftermath of MVC, compared sociodemographic, collision-related, and psychological characteristics that predict these outcomes, and evaluated the association between these outcomes and patients' expectations of recovery. Such information may improve understanding of the experience of individuals experiencing MVC, and provide further information regarding both the etiology of distress and dissociation as well as the association of these outcomes with patient expectations that may influence the recovery process (Castro, 2001; Ferrari, 2003).

Method

Patients were recruited between February 2009 and December 2010 from eight EDs located in four states (Michigan, Massachusetts, New York, and Florida). The study was approved by the institutional review boards of all participating hospitals. Each participant provided written informed consent.

A detailed description of study methods has been published. (Platts-Mills et al., 2011) In brief, patients age 18 to 65 who presented to the ED within 24 hours of MVC were screened for eligibility. Patients who required hospitalization, had fractures other than phalangeal

fractures, had more than 4 lacerations requiring sutures or a single laceration more than 20 cm in length, or had intracranial or vertebral injuries were excluded. Patients who were not alert and oriented were also excluded, as were pregnant patients, prisoners, patients unable to read and understand English, patients taking a β -blockers, or patients taking opioids above a total daily dose of 20 mg of oxycodone or equivalent. Enrollment was also limited to non-Hispanic whites (the most common ethnicity at study sites), because the study included the collection of genetic samples and ethnic heterogeneity increases the risk of population stratification bias. (Diatchenko et al., 2007)

A research assistant stationed in each of the study site EDs monitored the chief complaint and injury history of presenting patients. Potentially eligible patients were approached to determine eligibility and eligible patients were offered enrollment into the study. In order to assess patient competency to give informed consent, eligible patients were required to describe the essential elements of the study back to the research assistant. Following signing a local IRB-approved informed consent, study participants completed an ED interview using a web-based survey. Before enrolling patients in the ED, each research assistant completed a study training module followed by an interview with a standardized mock ED patient. Comparison of mock ED patient data across research assistants demonstrated high concordance (error rate 1.3%). Study participants were compensated for their time at the amount of \$80.00 for completing the ED interview. After study participant discharge from the ED, a study site research assistant completed a web-based data extraction form which collected information from ED and hospital medical records.

Demographic information was assessed using standardized questionnaire items. Depressive symptoms during the week prior to the MVC were assessed using the Center for Epidemiologic Studies Depression Scale (CES-D). (Radloff, 1977) Based on previous evidence, CES-D scores between 16 and 25 were defined as mild depressive symptoms and scores greater than or equal to 26 defined severe depressive symptoms. (Weissman, 1977; Zich, 1990) Trait anger and anxiety were assessed using the State-Trait Personality Inventory form Y (STPI-Y) anger and anxiety subscales. (Spielberger, 2000) Physical and mental health status during the 4 weeks prior to the MVC was assessed using the Mental Component Summary and Physical Component Summary scores of the Short-Form Health Survey version 2 (SF-12). (Ware, 1996) At-risk alcohol use was evaluated using the TWEAK alcohol screening instrument. TWEAK score has been shown to have high sensitivity (87%) and specificity (76%) for identifying individuals who meet ICD-10 criteria for alcohol dependence or harmful drinking. (Cherpitel, 1995) Marijuana use during the 3 months prior to the MVC was also assessed. Overall pain at the time of the ED interview and average overall pain during the month prior to the MVC were assessed using a 0 to 10 numeric rating scale. Severity of 21 individual somatic symptoms (e.g. fatigue, dizziness, blurry vision) in the month prior to MVC were assessed using a 0–10 numerical scale. These scores were compiled into an aggregate somatic symptom score were between 0 and 210. Research participant interview has been shown to be an accurate means of obtaining MVC history (Lee, 2011) and was used to assess collision characteristics.

Distress was assessed using the Peritraumatic Distress Inventory (PDI). This measure has high internal consistency (0.75–0.76) and test-retest reliability (0.74). (Brunet et al., 2001) A

PDI cut-off score of 23 was used to define marked distress symptoms (“distress”). (Nishi et al., 2010) Dissociation was defined by the Michigan Critical Events Perception Scale (MCEPS). (Michaels, Michaels, Moon, et al., 1999) This measure has high internal consistency (0.81) and strong correlation with the Peritraumatic Dissociative Experiences Questionnaire ($r=0.5$), a previously validated measure of dissociation. A MCEPS cut-off score of greater than 3.0 is indicative that the patient experienced dissociation. (Michaels, Michaels, Moon, et al., 1999) To assess confidence in recovery, participants were asked “How certain, or sure, are you that you will fully recover from this accident” on a 0–10, where 0 = “certain you will not recover” and 10 = “certain you will recover fully”. Participants were also asked to estimate how many days they expected it would take for them to recover physically and how many days they expected it would take for them to recover emotionally from the accident.

To evaluate the strength of association between individual predictors and distress and dissociation outcomes, log binomial regression models (adjusted for study site) were used to generate relative risks with 95% confidence intervals. To select a set of variables that together best predicted distress and dissociation outcomes, multivariable lasso regression was performed using the “glmnet” package in R. (Tibshirani, 1996) Lasso is a form of penalized regression that performs variable selection while guarding against overfitting. All candidate predictors with $p < 0.1$ in bivariate analyses were included as candidate predictors for developing lasso models. The tuning parameter for each model was chosen using cross-validation. The overall predictive utility of each model was evaluated using the area under the receiver-operator characteristic curve (AUROC).

Since coefficient estimates from multivariable models are biased, they are not a reliable estimate of the effect size of an individual model variables. (Tibshirani, 1996) Therefore to evaluate the strength of association between individual model variables and study outcomes while adjusting for potential confounders (study site, gender, age, and education), log binomial regression modeling was used to generate relative risks with 95% confidence intervals for each variable with a nonzero lasso coefficient.

Results

A total of 949 individuals were evaluated in the ED in the early aftermath of MVC (mean time from MVC to ED presentation 2.2 +/- 2.8 hours). Most enrolled MVC survivors were women, had some education past high school, and worked full time (Table 1). Less than one third of participants were smokers and only about one in five reported mild or moderate depressive symptoms during the week prior to MVC. The great majority of patients were driving the vehicle at the time of the collision and over half of collisions resulted in severe vehicle damage.

Peritraumatic distress was experienced by 355/935 (38%) MVC survivors. Peritraumatic dissociation was less common, occurring in 260/942 (28%) MVC survivors. Raw scores for distress and dissociation were only moderately correlated ($n = 930$, Pearson correlation coefficient 0.45, $p < 0.001$). Among patients within the highest quartile of dissociative symptoms (MCEPS score 3.4), only 36% were distressed. Similarly, among individuals

within the highest quartile of distress symptoms (PDI symptoms score ≥ 27), only 51% dissociated.

Bivariate associations between sociodemographic and collision characteristics and distress and dissociation outcomes are shown in Figure 1A. Female sex and severity of vehicle damage were associated with both distress and dissociation. Those 50–65 years of age were less likely to experience either distress or dissociation than those aged 18–34. A history of marijuana use and higher vehicle speed at the time of the collision were associated with increased risk of distress (but not dissociation); levels of family income above twenty thousand dollars and rear end collision were protective against distress (but not dissociation). BMI within categories of 25–30 or >30 was protective against dissociation (versus BMI < 25), but was not protective against distress.

Bivariate associations between health status, psychological, and cognitive characteristics and distress and dissociation outcomes in the hours after MVC are shown in Figure 1B. Trait anxiety (highest versus lowest tertile) predicted both distress and dissociation and better mental health prior to MVC decreased risk of both outcomes. Increased depressive symptoms prior to MVC (≥ 6 versus < 6), increased trait anger, past somatic symptom score (high versus lowest and middle versus lowest tertile), and participant feeling that collision was their fault were associated with increased risk of distress but not dissociation. Patients with better physical health were more likely to have dissociation, but not distress.

From the bivariate analyses shown in Figures 1A and 1B, a parsimonious set of predictors of distress and dissociation were identified using Lasso regression. Female sex, trait anxiety, severity of vehicle damage, and depressive symptoms prior to MVC together most efficiently predicted distress symptoms (Table 2, mean AUROC=0.69). Female sex, trait anxiety, and severity of vehicle damage, and worse overall mental health prior to MVC were most predictive of dissociation (Table 2, mean AUROC=0.57).

Table 3 displays associations between distress and dissociation after MVC and patient confidence in recovery and expected time to physical and emotional recovery. Distressed patients were 30% less likely to feel confident in achieving a full recovery than individuals who were not distressed, and were 40% less likely to estimate a time to emotional recovery of less than 30 days (Table 3). In contrast, the association between dissociation and patient expectations was weaker and did not reach statistical significance.

Discussion

We find that adverse psychological responses are common among patients treated in the ED after MVC and discharged to home. More than 1 in 3 individuals in our study experienced substantial distress, and more than 1 in 4 experienced dissociation. These outcomes were only moderately correlated ($r = 0.45$) and were less correlated than seen in a prior study (Fikretoglu et al., 2006). Among patients within the highest quartile of dissociative symptoms, only 36% were distressed. Similarly, among individuals within the highest quartile of distress symptoms, only 51% experienced dissociation. These results imply that distress and dissociation are separate entities and that, rather than dissociation being an

epiphenomenon of high levels of distress or protective against distress, distress and dissociation are distinct psychological processes. Although we do not directly examine the relationships between dissociation and long-term health or psychological outcomes, the absence of a relationship between dissociation and expectations for recovery in our sample does not support the hypothesis that dissociation indicates a more severe response or confers a negative prognosis (Holm, Carroll, Cassidy, Skillgate, & Ahlbom, 2008).

A previous study which sent mail questionnaires to police officers and civilian peers and asked about symptoms in response to past stressful life events (critical incidents) found a much stronger association between distress and dissociation symptoms (Fikretoglu et al., 2006). In that study 57% of police and 51% of civilians who reported dissociative symptoms in the highest quartile of respondents after a past critical incident also reported distress symptoms in the highest quartile. Differences in study results between that study and the present one could be due in part to differences in the populations, time elapsed since the traumatic event, and type of trauma exposure. An advantage of the present study is that distress and dissociation symptoms were assessed only hours after the MVC, reducing the likelihood of recall bias.

Consistent with weak correlation between distress and dissociation in our sample, predictors of these reactions also differed. In particular, higher socioeconomic status (higher education, higher income, full time employment) consistently had a significant protective effect against distress but not dissociative symptoms. Cigarette use was also associated with increased distress but not dissociation, but whether this relationship is a result of the direct effects of chronic nicotine exposure or confounding by socioeconomic status is unknown. Depressive and anxiety symptoms, and the feeling that the MVC was the participant's fault, were also more predictive of distress than dissociation. Other factors, such as reduced mental health prior to MVC, were more strongly predictive of dissociation. Interestingly, better physical health status had different effects on risk of distress and dissociation: better physical health was associated with greater risk of dissociation, but not associated with risk of distress. The distinct risk factors for distress and dissociation suggest that these phenomena are distinct clinical entities with distinct etiologies.

The relationship of several other study findings to previous literature are also worthy of comment. A prior investigation found that, as in our study, female gender was predictive of both dissociation and distress as characterized by "emotional reactions" (Lawyer et al., 2006). This is consistent with other reported findings (Brunet et al., 2001; Bryant & Harvey, 2003). Our results differ, however, in that our study identified greater age as being protective against both dissociation and distress, whereas in the study by Brunet et al. age greater than 55 predicted dissociative response. Our findings are consistent with two prior reports (Fullerton et al., 2000; Ursano et al., 1999) which found that education, income, and marital status were associated with distress/emotional response. Several exposure-related factors were found to be predictive of trauma response patterns in the studies by Fullerton et al. and Ursano et al.; fear of death/injury, friend or family member killed, location relative to the attack, and media exposure in the prior study as compared to vehicle damage, vehicle speed, location in vehicle, and collision type in ours. Other risk factors identified in these

analyses such as involvement in rescue efforts, race or ethnicity, loss of possessions, and fear of death were not investigated in our study.

Alcohol misuse has been reported as a possible risk factor for the development of PTSD (Blanchard et al., 1996). However at-risk drinking during the past 3 months was not associated with either distress or dissociation in the present study. Nicotine dependence is associated with increased risk of both PTSD (Koenen et al., 2005) and panic disorders (Isensee, Wittchen, Stein, Hofler, & Lieb, 2003), and cigarette smoking was associated with risk of peritraumatic distress in the present study. This association between tobacco use and distress has been described previously (Feldner, Babson, & Zvolensky, 2007), and could be due to the association between tobacco use and anxiety disorders (Morissette, Tull, Gulliver, Kamholz, & Zimering, 2007), self-medication among those with worse mental health, (Khantzian, 1997) or the association between smoking status and other sociodemographic markers (Laaksonen, Rahkonen, Karvonen, & Lahelma, 2005) that increase distress vulnerability. These same factors could also account for the association between frequency of marijuana use and peritraumatic distress (Bonn-Miller, Vujanovic, Feldner, Bernstein, & Zvolensky, 2007; Bremner, Southwick, Darnell, & Charney, 1996). Marijuana use has also been shown to cause dissociative symptoms in vulnerable individuals (Johnson, 1990).

Prior investigations have focused on traumatic exposures or stressors as risk factors for acute peritraumatic responses (Branscomb, 1991; Feeny et al., 2000; Lawyer et al., 2006). This is the first study to date to identify self-reported baseline psychological traits including depression, anxiety, anger, and poor overall mental health as predictors of peritraumatic distress and dissociation, as well as the first to relate peritraumatic responses to expectations of recovery. Recovery expectations are strong predictors of recovery and disability after MVC (Carroll, 2011; Holm et al., 2008), and may help shape cognitions and behaviors during the weeks after trauma (Carroll, 2011). Negative cognitions and altered behaviors during the weeks after MVC may be important mediators of the influence of distress on post-MVC outcomes. In our sample, distressed individuals were much less likely to be confident of full recovery and to believe that they would recover emotionally within one month of MVC. Point estimates suggest that dissociation has less influence on recovery expectations, and the associations between dissociation and recovery expectations did not reach statistical significance.

A number of study limitations should be considered when interpreting our study findings. First, our study was cross-sectional, therefore health characteristics and psychological traits prior to MVC were assessed retrospectively. The patient's emotional state at the time of the ED visit may have influenced responses to questionnaires, particularly for psychological assessments of pre-trauma anxiety, anger, and depression. Assessment of these characteristics prior to induction of traumatic response is possible in experimental studies (e.g. CO₂ inhalation to incite distress and cold exposure or pharmacologic production of dissociation) but much more difficult in non-experimental studies. To try to reduce this influence, questions about current psychological symptoms were assessed completely prior to turning to questions regarding past psychological symptoms. The fact that our evaluations were performed only hours after MVC is also a strength of our study, as peritraumatic distress and dissociation symptoms were assessed in the peritraumatic period rather than

days or weeks after trauma exposure. This contemporaneous assessment may have provided us with a more accurate assessment of the peritraumatic psychological response. In addition, we did not adjust for multiple comparisons in our bivariate analyses, which increases the risk of false positive associations. However, many of the candidate predictor variables assessed were correlated, therefore adjusting for multiple tests would most likely result in a significant decrease in power with very little corresponding decrease in the risk of type I error. In addition, the need for such adjustment is a matter of considerable debate, particularly in the setting of studies such as this one in which candidate predictors were selected because of substantial pre-test association with study outcomes (Rothman, 1990; Savitz & Olshan, 1995). Another study limitation is that our study only assessed individuals experiencing one very common traumatic exposure, MVC. The incidence and association of distress and dissociation symptoms after other types of trauma is unknown. In addition, only patients discharged home from the ED after MVC were included. However, only about 10% of patients who seek ED evaluation after MVC are admitted to the hospital (Nishi et al., 2010) and evaluating a homogenous outcome (in which the degree of tissue injury is relatively constant) likely increased our ability to identify predictors of distress and dissociation separate from those related to tissue injury. Finally, because of genetic analyses performed as part of this project, enrollment was restricted to European Americans only. Non-European Americans may have different rates and predictors of distress and dissociation after MVC.

In summary, we found that distress and dissociation are common among individuals discharged home from the ED after MVC, with each outcome occurring in approximately one third of patients. These outcomes are only moderately correlated, and have both shared and distinct predictors including baseline psychological traits. Perhaps most notably in this regard, higher socioeconomic status had a consistent protective effect against distress but not dissociation. Similarly, better physical health was a risk factor for dissociation but not distress. Together these data suggest that distress and dissociation during the peritraumatic period are distinct phenomena with distinct biologic underpinnings. Further work is needed to understand the neurobiology of distress and dissociation and the relationship between these acute peritraumatic responses and persistent psychological sequelae. Such improved understanding may inform the development of early interventions to prevent long term psychological consequences of trauma exposure.

Acknowledgments

FUNDING

This research was supported by NIAMS R01AR056328 from the NIH awarded to Samuel A. McLean. The authors would like to thank the participants of the study.

References

- Bernat JA, Ronfeldt HM, Calhoun KS, Arias I. Prevalence of traumatic events and peritraumatic predictors of posttraumatic stress symptoms in a nonclinical sample of college students. *Journal of Traumatic Stress*. 1998; 11(4):645–664. [PubMed: 9870219]

- Blanchard EB, Hickling EJ, Barton KA, Taylor AE, Loos WR, Jones-Alexander J. One-year prospective follow-up of motor vehicle accident victims. *Behavioral Research and Therapy*. 1996; 34(10):775–786.
- Bonn-Miller MO, Vujanovic AA, Feldner MT, Bernstein A, Zvolensky MJ. Posttraumatic stress symptom severity predicts marijuana use coping motives among traumatic event-exposed marijuana users. *Journal of Traumatic Stress*. 2007; 20(4):577–586. [PubMed: 17721963]
- Branscomb LB. Dissociation in combat-related post-traumatic stress disorder. *Dissociation*. 1991; 4:13–20.
- Bremner JD, Southwick S, Brett E, Fontana A, Rosenheck R, Charney DS. Dissociation and posttraumatic stress disorder in Vietnam combat veterans. *American Journal of Psychiatry*. 1992; 149(3):328–332. [PubMed: 1536269]
- Bremner JD, Southwick SM, Darnell A, Charney DS. Chronic PTSD in Vietnam combat veterans: course of illness and substance abuse. *American Journal of Psychiatry*. 1996; 153(3):369–375. [PubMed: 8610824]
- Briere J, Scott C, Weathers F. Peritraumatic and persistent dissociation in the presumed etiology of PTSD. *American Journal of Psychiatry*. 2005; 162(12):2295–2301. [PubMed: 16330593]
- Brunet A, Sanche S, Manetti A, Aouizerate B, Ribereau-Gayon R, Charpentier S, Arbus C. Peritraumatic distress but not dissociation predicts posttraumatic stress disorder in the elderly. *International Psychogeriatrics*. 2013; 25(6):1007–1012. [PubMed: 23433477]
- Brunet A, Weiss DS, Metzler TJ, Best SR, Neylan TC, Rogers C, Marmar CR. The Peritraumatic Distress Inventory: a proposed measure of PTSD criterion A2. *American Journal of Psychiatry*. 2001; 158(9):1480–1485. [PubMed: 11532735]
- Bryant RA. Early predictors of posttraumatic stress disorder. *Biological Psychiatry*. 2003; 53(9):789–795. [PubMed: 12725971]
- Bryant RA, Harvey AG. Gender differences in the relationship between acute stress disorder and posttraumatic stress disorder following motor vehicle accidents. *Australian and New Zealand Journal of Psychiatry*. 2003; 37(2):226–229. [PubMed: 12656964]
- Bui E, Joubert S, Manetti A, Camassel C, Charpentier S, Ribereau-Gayon R, Arbus C. Peritraumatic distress predicts posttraumatic stress symptoms in older people. *International Journal of Geriatric Psychiatry*. 2010; 25(12):1306–1307. [PubMed: 21086541]
- Buitenhuis J, de Jong PJ, Jaspers JPC, Groothoff JW. Relationship between posttraumatic stress disorder symptoms and the course of whiplash complaints. *Journal of Psychosomatic Research*. 2006; 61(5):681–689. [PubMed: 17084147]
- Cardena E, Spiegel D. Dissociative reactions to the San Francisco Bay Area earthquake of 1989. *American Journal of Psychiatry*. 1993; 150(3):474–478. [PubMed: 8434665]
- Carroll LJ. Beliefs and expectations for recovery, coping, and depression in whiplash-associated disorders: lessening the transition to chronicity. *Spine (Phila Pa 1976)*. 2011; 36(25 Suppl):S250–256. [PubMed: 22020620]
- Castro WH, Meyer SJ, Becke ME, Nentwig CG, Hein MF, Ercan BI, Thomann S, Wessels U, Du Chesne AE. No stress - no whiplash? Prevalence of “whiplash” symptoms following exposure to a placebo rear-end collision. *International Journal of Legal Medicine*. 2001; 114(6):316–322. [PubMed: 11508796]
- Cherpitel CJ. Screening for alcohol problems in the emergency room: a rapid alcohol problems screen. *Drug and Alcohol Dependence*. 1995; 40(2):133–137. [PubMed: 8745135]
- Chu JA, Frey LM, Ganzel BL, Matthews JA. Memories of childhood abuse: dissociation, amnesia, and corroboration. *American Journal of Psychiatry*. 1999; 156(5):749–755. [PubMed: 10327909]
- Diatchenko L, Nackley AG, Slade GD, Belfer I, Max MB, Goldman D, Maixner W. Responses to Drs. Kim and Dionne regarding comments on Diatchenko, et al. Catechol-O-methyltransferase gene polymorphisms are associated with multiple pain-evoking stimuli. *Pain*. 2007; 129(3):366–370. [PubMed: 17851590]
- Ehlers A, Mayou RA, Bryant B. Psychological predictors of chronic posttraumatic stress disorder after motor vehicle accidents. *Journal of Abnormal Psychology*. 1998; 107(3):508–519. [PubMed: 9715585]

- Feeny NC, Zoellner LA, Fitzgibbons LA, Foa EB. Exploring the roles of emotional numbing, depression, and dissociation in PTSD. *Journal of Traumatic Stress*. 2000; 13(3):489–498. [PubMed: 10948488]
- Feinstein A, Dolan R. Predictors of post-traumatic stress disorder following physical trauma: an examination of the stressor criterion. *Psychological Medicine*. 1991; 21(1):85–91. [PubMed: 2047509]
- Feldner MT, Babson KA, Zvolensky MJ. Smoking, traumatic event exposure, and post-traumatic stress: a critical review of the empirical literature. *Clinical Psychology Review*. 2007; 27(1):14–45. [PubMed: 17034916]
- Ferrari R, Constantovannis C, Papadakis N. Laypersons' expectation of the sequelae of whiplash injury: a cross-cultural comparative study between Canada and Greece. *Medical Science Monitor*. 2003; 9(3):CR120–124. [PubMed: 12640340]
- Fikretoglu D, Brunet A, Best S, Metzler T, Delucchi K, Weiss DS, Marmar C. The relationship between peritraumatic distress and peritraumatic dissociation: an examination of two competing models. *Journal of Nervous and Mental Disease*. 2006; 194(11):853–858. [PubMed: 17102710]
- Foa EB, Rothbaum BO, Riggs DS, Murdock TB. Treatment of posttraumatic stress disorder in rape victims: a comparison between cognitive-behavioral procedures and counseling. *Journal of Consulting and Clinical Psychology*. 1991; 59(5):715–723. [PubMed: 1955605]
- Friedman MJ. What might the psychobiology of posttraumatic stress disorder teach us about future approaches to pharmacotherapy? *Journal of Clinical Psychiatry*. 2000; 61(Suppl 7):44–51. [PubMed: 10795609]
- Fullerton CS, Ursano RJ, Epstein RS, Crowley B, Vance KL, Kao TC, Baum A. Peritraumatic dissociation following motor vehicle accidents: relationship to prior trauma and prior major depression. *Journal of Nervous and Mental Disease*. 2000; 188(5):267–272. [PubMed: 10830563]
- Green BL. Defining trauma: terminology and generic stressor dimensions. *Journal of Applied Social Psychology*. 1990; 20:1632–1642.
- Green BL. Psychosocial research in traumatic stress: an update. *Journal of Traumatic Stress*. 1994; 7(3):341–362. [PubMed: 8087399]
- Hargrave PA, Leathem JM, Long NR. Peritraumatic distress: its relationship to posttraumatic stress and complicated grief symptoms in sudden death survivors. *Journal of Traumatic Stress*. 2012; 25(3):344–347. [PubMed: 22685096]
- Hetzl-Riggin MD, Wilber EL. To dissociate or suppress? Predicting automatic vs. conscious cognitive avoidance. *Journal of Trauma and Dissociation*. 2010; 11(4):444–457. [PubMed: 20938868]
- Hodgson RC, Webster RA. Mediating role of peritraumatic dissociation and depression on post-MVA distress: path analysis. *Depression and Anxiety*. 2011; 28(3):218–226. [PubMed: 21328635]
- Holm LW, Carroll LJ, Cassidy JD, Skillgate E, Ahlbom A. Expectations for recovery important in the prognosis of whiplash injuries. *PLoS Medicine*. 2008; 5(5):e105. [PubMed: 18479182]
- Isensee B, Wittchen HU, Stein MB, Hofler M, Lieb R. Smoking increases the risk of panic: findings from a prospective community study. *Archives of General Psychiatry*. 2003; 60(7):692–700. [PubMed: 12860773]
- Jaycox LH, Marshall GN, Orlando M. Predictors of acute distress among young adults injured by community violence. *Journal of Traumatic Stress*. 2003; 16(3):237–245. [PubMed: 12816335]
- Jehel L, Brunet A, Paterniti S, Guelfi JD. Validation of the Peritraumatic Distress Inventory's French translation. *Canadian Journal of Psychiatry*. 2005; 50(1):67–71.
- Jehel L, Paterniti S, Brunet A, Louville P, Guelfi JD. Peritraumatic distress prospectively predicts PTSD symptoms in assault victims. *L'Encephale*. 2006; 32(6 Pt 1):953–956.
- Johnson BA. Psychopharmacological effects of cannabis. *British Journal of Hospital Medicine*. 1990; 43(2):114–116. 118–120, 122. [PubMed: 2178712]
- Kaufman ML, Kimble MO, Kaloupek DG, McTeague LM, Bachrach P, Forti AM, Keane TM. Peritraumatic dissociation and physiological response to trauma-relevant stimuli in Vietnam combat veterans with posttraumatic stress disorder. *Journal of Nervous and Mental Disease*. 2002; 190(3):167–174. [PubMed: 11923651]

- Khantzian EJ. The self-medication hypothesis of substance use disorders: a reconsideration and recent applications. *Harvard Review of Psychiatry*. 1997; 4(5):231–244. [PubMed: 9385000]
- Kihlstrom JF, Glisky ML, Angiulo MJ. Dissociative tendencies and dissociative disorders. *Journal of Abnormal Psychology*. 1994; 103(1):117–124. [PubMed: 8040473]
- Koenen KC, Hitsman B, Lyons MJ, Niaura R, McCaffery J, Goldberg J, Tsuang M. A twin registry study of the relationship between posttraumatic stress disorder and nicotine dependence in men. *Archives of General Psychiatry*. 2005; 62(11):1258–1265. [PubMed: 16275813]
- Koopman C, Classen C, Spiegel D. Predictors of posttraumatic stress symptoms among survivors of the Oakland/Berkeley, Calif., firestorm. *American Journal of Psychiatry*. 1994; 151(6):888–894. [PubMed: 8184999]
- Kuch K, Cox BJ, Evans RJ. Posttraumatic stress disorder and motor vehicle accidents: a multidisciplinary overview. *Canadian Journal of Psychiatry*. 1996; 41:429–434.
- Kumpula MJ, Orcutt HK, Bardeen JR, Varkovitzky RL. Peritraumatic dissociation and experiential avoidance as prospective predictors of posttraumatic stress symptoms. *Journal of Abnormal Psychology*. 2011; 120(3):617–627. [PubMed: 21604826]
- Laaksonen M, Rahkonen O, Karvonen S, Lahelma E. Socioeconomic status and smoking: analysing inequalities with multiple indicators. *European Journal of Public Health*. 2005; 15(3):262–269. [PubMed: 15755781]
- Lawyer SR, Resnick HS, Galea S, Ahern J, Kilpatrick DG, Vlahov D. Predictors of peritraumatic reactions and PTSD following the September 11th terrorist attacks. *Psychiatry*. 2006; 69(2):130–141. [PubMed: 16822192]
- Lee YM, Platts-Mills TF, MacWilliams J, Jones JS, Domeier RM, Swor RA, Sochor MA, Schneider J, McLean SA. Description of Motor Vehicle Collisions by Participants in Emergency Department-Based Studies: Are they Accurate? *Western Journal of Emergency Medicine*. 2011; 13(4):329–334. [PubMed: 22942933]
- Lensvelt-Mulders G, van der Hart O, van Ochten JM, van Son MJ, Steele K, Breeman L. Relations among peritraumatic dissociation and posttraumatic stress: a meta-analysis. *Clinical Psychology Review*. 2008; 28(7):1138–1151. [PubMed: 18502549]
- Litz BT. Early intervention for trauma: where are we and where do we need to go? A commentary. *Journal of Traumatic Stress*. 2008; 21(6):503–506.10.1002/jts.20373 [PubMed: 19107726]
- Peden, M.; Scurfield, R.; Sleet, D.; Mohan, D.; Hyder, AA.; Jarawan, E.; Mathers, C. *The fundamentals World Report on Road Traffic Injury Prevention*. Geneva, Switzerland: World Health Organization; 2004.
- Marmar CR, Weiss DS, Metzler TJ, Delucchi K. Characteristics of emergency services personnel related to peritraumatic dissociation during critical incident exposure. *American Journal of Psychiatry*. 1996; 153(7 Suppl):94–102. [PubMed: 8659646]
- Marmar CR, Weiss DS, Schlenger WE, Fairbank JA, Jordan BK, Kulka RA, Hough RL. Peritraumatic dissociation and posttraumatic stress in male Vietnam theater veterans. *American Journal of Psychiatry*. 1994; 151(6):902–907. [PubMed: 8185001]
- Martin M, Marchand A, Boyer R, Martin N. Predictors of the development of posttraumatic stress disorder among police officers. *Journal of Trauma and Dissociation*. 2009; 10(4):451–468. [PubMed: 19821179]
- Mayou R, Bryant B, Duthrie R. Psychiatric consequences of road traffic accidents. *British Medical Journal*. 1993; 307:647–651. [PubMed: 8401049]
- McFarlane AC. Predictors of post-traumatic stress disorder after major injury. *ANZ Journal of Surgery*. 2008; 78(7):533–534. [PubMed: 18593405]
- Merckelbach H, Muris P. The causal link between self-reported trauma and dissociation: a critical review. *Behavioral Research and Therapy*. 2001; 39(3):245–254.
- Michaels AJ, Michaels CE, Moon CH, Smith JS, Zimmerman MA, Taheri PA, Peterson C. Posttraumatic stress disorder after injury: impact on general health outcome and early risk assessment. *Journal of Trauma*. 1999; 47(3):460–466. discussion 466–467. [PubMed: 10498298]
- Michaels AJ, Michaels CE, Zimmerman MA, Smith JS, Moon CH, Peterson C. Posttraumatic stress disorder in injured adults: etiology by path analysis. *Journal of Trauma*. 1999; 47(5):867–873. [PubMed: 10568714]

- Morissette SB, Tull MT, Gulliver SB, Kamholz BW, Zimering RT. Anxiety, anxiety disorders, tobacco use, and nicotine: a critical review of interrelationships. *Psychological Bulletin*. 2007; 133(2):245–272. [PubMed: 17338599]
- Nishi D, Koido Y, Nakaya N, Sone T, Noguchi H, Hamazaki K, Matsuoka Y. Peritraumatic distress, watching television, and posttraumatic stress symptoms among rescue workers after the Great East Japan earthquake. *PLoS One*. 2012; 7(4):e35248. [PubMed: 22558130]
- Nishi D, Matsuoka Y. Peritraumatic distress after an earthquake: a bridge between neuroimaging and epidemiology. *Molecular Psychiatry*. 2013; 18(7):743–744. [PubMed: 22751489]
- Nishi D, Matsuoka Y, Noguchi H, Sakuma K, Yonemoto N, Yanagita T, Kim Y. Reliability and validity of the Japanese version of the Peritraumatic Distress Inventory. *General Hospital Psychiatry*. 2009; 31(1):75–79. [PubMed: 19134513]
- Nishi D, Matsuoka Y, Yonemoto N, Noguchi H, Kim Y, Kanba S. Peritraumatic Distress Inventory as a predictor of post-traumatic stress disorder after a severe motor vehicle accident. *Psychiatry and Clinical Neurosciences*. 2010; 64(2):149–156. [PubMed: 20447011]
- Niska R, Bhuiya F, Xu J. National Hospital Ambulatory Medical Care Survey: 2007 emergency department summary. *National Health Statistics Report*. 2010; 26:1–31.
- Otis C, Marchand A, Courtois F. Peritraumatic dissociation as a mediator of peritraumatic distress and PTSD: a retrospective, cross-sectional study. *Journal of Trauma and Dissociation*. 2012; 13(4):469–477. [PubMed: 22651680]
- Ozer EJ, Best SR, Lipsey TL, Weiss DS. Predictors of posttraumatic stress disorder and symptoms in adults: a meta-analysis. *Psychology Bulletin*. 2003; 129(1):52–73.
- Pacella ML, Irish L, Ostrowski SA, Sledjeski E, Ciesla JA, Fallon W, Delahanty DL. Avoidant coping as a mediator between peritraumatic dissociation and posttraumatic stress disorder symptoms. *Journal of Traumatic Stress*. 2011; 24(3):317–325. [PubMed: 21618289]
- Perry S, Difede J, Musngi G, Frances AJ, Jacobsberg L. Predictors of posttraumatic stress disorder after burn injury. *American Journal of Psychiatry*. 1992; 149(7):931–935. [PubMed: 1609874]
- Platts-Mills TF, Ballina L, Bortsov AV, Soward A, Swor RA, Jones JS, McLean SA. Using emergency department-based inception cohorts to determine genetic characteristics associated with long term patient outcomes after motor vehicle collision: methodology of the CRASH study. *BMC Emergency Medicine*. 2011; 11:14. [PubMed: 21943293]
- Platts-Mills TF, Hunold KM, Esserman DA, Sloane PD, McLean SA. Motor vehicle collision-related emergency department visits by older adults in the United States. *Academic Emergency Medicine*. 2012; 19(7):821–827.
- Radloff LS. The CES-D Scale. A self-report depression scale for research in the general population. *Applied Psychological Measurement*. 1977; 1:385–401.
- Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology*. 1990; 1(1):43–46. [PubMed: 2081237]
- Savitz DA, Olshan AF. Multiple comparisons and related issues in the interpretation of epidemiologic data. *American Journal of Epidemiol*. 1995; 142(9):904–908.
- Shalev AY. Posttraumatic stress disorder and stress-related disorders. *Psychiatry Clinics of North America*. 2009; 32(3):687–704.
- Shalev AY, Peri T, Canetti L, Schreiber S. Predictors of PTSD in injured trauma survivors: a prospective study. *American Journal of Psychiatry*. 1996; 153(2):219–225. [PubMed: 8561202]
- Solomon Z, Benbenishty R. The role of proximity, immediacy, and expectancy in frontline treatment of combat stress reaction among Israelis in the Lebanon War. *American Journal of Psychiatry*. 1986; 143(5):613–617. [PubMed: 3963249]
- Spiegel D, Hunt T, Dondershine HE. Dissociation and hypnotizability in posttraumatic stress disorder. *American Journal of Psychiatry*. 1988; 145(3):301–305. [PubMed: 3344845]
- Spielberger, CD. Preliminary manual for the State - Trait Personality Inventory (STPI). 2000.
- Thomas E, Saumier D, Brunet A. Peritraumatic distress and the course of posttraumatic stress disorder symptoms: a meta-analysis. *Canadian Journal of Psychiatry*. 2012; 57(2):122–129.
- Tibshirani R. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B*. 1996; 58(1):267–288.

- Ursano RJ, Fullerton CS, Epstein RS, Crowley B, Vance K, Kao TC, Baum A. Peritraumatic dissociation and posttraumatic stress disorder following motor vehicle accidents. *American Journal of Psychiatry*. 1999; 156(11):1808–1810. [PubMed: 10553747]
- van der Kolk BA, van der Hart O. Pierre Janet and the breakdown of adaptation in psychological trauma. *American Journal of Psychiatry*. 1989; 146(12):1530–1540. [PubMed: 2686473]
- van der Velden PG, Wittmann L. The independent predictive value of peritraumatic dissociation for PTSD symptomatology after type I trauma: a systematic review of prospective studies. *Clinical Psychology Review*. 2008; 28(6):1009–1020. [PubMed: 18406027]
- Ware J, Kosinski M, Keller SD. SF-12: A 12-item Short-Form Health Survey: Construction of scales and preliminary tests of reliability and validity. *Medical Care*. 1996; 34:220–233. [PubMed: 8628042]
- Weissman MM, Sholomskas D, Pottenger M, Prusoff BA, Locke BZ. Assessing depressive symptoms in five psychiatric populations: a validation study. *American Journal of Epidemiology*. 1977; 106:203–214. [PubMed: 900119]
- Zich JM, Attkisson CC, Greenfield TK. Screening for depression in primary care clinics: the CES-D and the BDI. *International Journal of Psychiatry Medicine*. 1990; 20(3):259–277.

Figure 1A.




Figure 1B.




Figure 1.

Figure 1A. Association of Sociodemographic and Collision Characteristics with Distress and Dissociation

Figure 1B. Association of Health Status and Psychologic and Cognitive Characteristics with Distress and Dissociation

Table 1

Study participant characteristics (n = 949).

Predictor	Categories	N	%
Age	18–34	506	54.1
	35–49	251	26.8
	50–65	178	19.0
Gender	Male	377	39.7
	Female	572	60.3
Education	K-12 or less	227	24.0
	Some college	369	39.0
	College/more	351	37.1
Full Time Employment	No	396	41.7
	Yes	553	58.3
Income	<\$20k	117	13.9
	\$20k to \$40k	176	20.9
	\$40k to \$80k	277	32.9
	>\$80k	273	32.4
Relationship Status	Not serious	298	31.8
	Not living together	146	15.6
	Living together	121	12.9
	Married	371	39.6
BMI	<25	356	38.7
	25–30	287	31.2
	>30	277	30.1
Cigarette Use	No	691	73.0
	Yes	256	27.0
Depressive Symptoms Prior to MVC*	<16	755	79.7
	16–25	123	13.0
	>=26	69	7.3
Collision Type	Other	608	64.1
	Rear-end	341	35.9
Extent of Damage [†]	None-Minor	128	14.0
	Moderate	282	30.8
	Severe	507	55.3
Vehicle Speed	Stopped	257	27.3
	1–40mph	492	52.2
	41–80mph	194	20.6
Location in Vehicle	Passenger	137	14.4
	Driver	812	85.6
Fault	Nobody's fault	164	17.3
	Subject's fault	119	12.6
	Another's fault	664	70.1

* As measured by CES-D Score

† Severe damage defined as car not being drivable

Table 2

Final Lasso regression models for distress and dissociation.

	Predictor	RR	95% CI	p
Distress (PDI >3)	Female Gender	1.87	1.49–2.35	<0.001
	Trait Anxiety [*]	1.63	1.35–1.96	<0.001
	Vehicle Damage [†]	1.67	1.35–2.07	<0.001
	Depressive Symptoms in Month Prior to MVC [‡]	1.80	1.50–2.15	<0.001
Dissociation (MCEPS>3)	Female Gender	1.52	1.16–1.99	0.003
	Trait Anxiety [*]	1.44	1.13–1.83	0.004
	Vehicle Damage [†]	1.39	1.08–1.79	0.011
	Mental Health Score [±]	0.62	0.45–0.85	0.003

Each variable adjusted for study site, age, gender, and education status. Relative risks for individual variables identified by model.

^{*} STPI-Y Anxiety subscale score in highest tertile (raw score 17–37) versus lowest tertile (raw score <13)

[†] Severe damage versus None-Minor damage

[‡] CES-D score in highest tertile (raw score 26–53) versus lowest tertile (raw score <16)

[±] SF-12 Mental Health Subscale Score in highest tertile (raw score 57.1–71.4) versus lowest tertile (raw score <47.8)

Associations between distress and dissociation and patient confidence in recovery and expected time to physical and emotional recovery after MVC*

Table 3

Expectation	Distress*			Dissociation†		
	RR	95% CI	P	RR	95% CI	P
Confident in Full Recovery	0.70	0.57–0.88	0.002	0.84	0.64–1.09	0.198
Expected Time to Physical Recovery <30 days	0.76	0.58–1.00	0.053	0.81	0.58–1.13	0.221
Expected Time to Emotional Recovery <30 days	0.62	0.49–0.79	<0.001	0.80	0.60–1.06	0.133

Each variable adjusted for age, education, pain severity, and study site

* As measured by PDI 23

† As measured by MCEPS >3