

**HHS PUBLIC ACCESS**

Author manuscript

Mol Genet Metab. Author manuscript; available in PMC 2015 April 13.

Published in final edited form as:

Mol Genet Metab. 2014 January ; 111(1): 46–51. doi:10.1016/j.ymgme.2013.11.004.**Maternal-Fetal Metabolic Gene-Gene Interactions and Risk of Neural Tube Defects****Philip J. Lupo^a, Laura E. Mitchell^b, Mark A. Canfield^c, Gary M. Shaw^d, Andrew F. Olshan^e, Richard H. Finnell^f, Huiping Zhu^f, and The National Birth Defects Prevention Study**^aDepartment of Pediatrics, Section of Hematology-Oncology, Baylor College of Medicine, Houston, Texas^bHuman Genetics Center, Division of Epidemiology, Human Genetics and Environmental Sciences, University of Texas School of Public Health, Houston, Texas^cTexas Department of State Health Services, Austin, Texas^dDepartment of Pediatrics, Stanford University School of Medicine, Stanford, California^eDepartment of Epidemiology, University of North Carolina, Chapel Hill, North Carolina^fDell Pediatric Research Institute, Department of Nutritional Sciences, University of Texas at Austin, Austin, Texas**Abstract**

Single-gene analyses indicate that maternal genes associated with metabolic conditions (e.g., obesity) may influence the risk of neural tube defects (NTDs). However, to our knowledge, there have been no assessments of maternal-fetal metabolic gene-gene interactions and NTDs. We investigated 23 single nucleotide polymorphisms among 7 maternal metabolic genes (*ADRB3*, *ENPP1*, *FTO*, *LEP*, *PPARG*, *PPARGC1A*, and *TCF7L2*) and 2 fetal metabolic genes (*SLC2A2* and *UCP2*). Samples were obtained from 737 NTD case-parent triads included in the National Birth Defects Prevention Study for birth years 1999–2007. We used a 2-step approach to evaluate maternal-fetal gene-gene interactions. First, a case-only approach was applied to screen all potential maternal and fetal interactions (n=76), as this design provides greater power in the assessment of gene-gene interactions compared to other approaches. Specifically, ordinal logistic regression was used to calculate the odds ratio (OR) and 95% confidence interval (CI) for each maternal-fetal gene-gene interaction, assuming a log-additive model of inheritance. Due to the number of comparisons, we calculated a corrected p-value (q-value) using the false discovery rate. Second, we confirmed all statistically significant interactions (q<0.05) using a log-linear approach among case-parent triads. In step 1, there were 5 maternal-fetal gene-gene interactions with q<0.05. The “top hit” was an interaction between maternal *ENPP1* rs1044498 and fetal *SLC2A2*

Corresponding Author: Huiping Zhu, MD, PhD, Dell Pediatric Research Institute, Department of Nutritional Sciences, University of Texas at Austin, Campus Mail Code: R1800, 1400 Barbara Jordan Blvd, Austin, Texas 78723, Tel: 512-495-3003, FAX: 512-495-4805, hzhu@austin.utexas.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

rs6785233 (interaction OR=3.65, 95% CI: 2.32–5.74, $p=2.09\times 10^{-8}$, $q=0.001$), which was confirmed in step 2 ($p=0.00004$). Our findings suggest that maternal metabolic genes associated with hyperglycemia and insulin resistance and fetal metabolic genes involved in glucose homeostasis may interact to increase the risk of NTDs.

Keywords

Birth defects; gene-gene interactions; maternal genetics; metabolic genes; neural tube defects; obesity; pre-gestational diabetes

1. INTRODUCTION

Neural tube defects (NTDs) are among the most common, costly, and deadly of all human congenital anomalies whose etiologies remain largely unknown [1, 2]. Maternal pre-gestational diabetes and pre-pregnancy obesity are two well-established risk factors for NTDs [3–19]. While the exact mechanisms behind these associations are unknown, it is believed that glucose homeostasis plays an important role. At the time of neural tube closure (approximately the fourth week of gestation), mothers with poorly regulated glucose levels are likely to have an altered intrauterine environment leading to abnormal organogenesis. Several genes related to glucose homeostasis have been previously identified in human and animal studies. Furthermore, genes related to glucose homeostasis have been associated with type 2 diabetes and obesity risk in genome-wide association studies (GWAS) [20–23]. Work from our group indicated an association between inherited (i.e., fetal) variation in the *UCP2* gene and NTDs [24]. *SLC2A2* is an important glucose transporter during embryonic neural tube development [25]. Additionally, we found associations between maternal genotypes in *FTO*, *TCF7L2*, and *LEP* and NTDs suggesting maternal genetic effects may cause changes in intrauterine environment and play a role in disease risk [24]. The Hyperglycemia and Adverse Pregnancy Outcome (HAPO) study has demonstrated that common genetic variants in genes such as *TCF7L2* are associated with fasting and post-challenge glucose levels during pregnancy [26]. Because of these findings, we sought to evaluate the interactions between maternal and fetal genes related to glucose homeostasis and the risk of NTDs.

2. MATERIALS AND METHODS

2.1 Subjects

The study population included NTD case-parent triads ($n=737$) from the National Birth Defects Prevention Study (NBDPS), with estimated dates of delivery between January 1, 1999 and December 31, 2007. Details of the NBDPS have been published elsewhere [27]. In brief, the NBDPS is a population-based case-control study of major structural birth defects. For the period 1999–2007, case infants with one or more congenital anomalies were ascertained through ten birth defects surveillance systems throughout the United States (Arkansas, California, Georgia, Iowa, Massachusetts, New Jersey, New York, North Carolina, Texas, and Utah) and included live births, stillbirths, and induced pregnancy terminations. NTDs included in the NBDPS had British Pediatric Association (BPA) codes for the diagnoses anencephaly (740.0), craniorachischisis (740.1), spina bifida (741.0), and encephalocele (742.0). Abstracted data for all NTD case infants were reviewed by clinical

geneticists using specific criteria, including standardized case definitions and confirmatory diagnostic procedures [28]. Infants/fetuses with known single gene disorders or chromosomal abnormalities were excluded from the NBDPS. Mothers completed a one-hour computer assisted telephone interview (CATI) in English or Spanish between 6 weeks and 2 years after the estimated date of delivery. The interview included sections on maternal conditions and illnesses, lifestyle and behavioral factors, and multivitamin use.

2.2 Maternal and Fetal Candidate Genes and Single Nucleotide Polymorphisms (SNPs)

The selection criterion for candidate genes and SNPs were reported previously [24]. Briefly, genes and SNPs selected were those identified as being associated with type 2 diabetes or obesity in multiple GWAS studies, or those with supporting evidence from both candidate gene studies and animal models. Maternal candidate genes included in the current study were *ADRB3*, *ENPP1*, *FTO*, *LEP*, *PPARG*, *PPARGC1A*, and *TCF7L2*. Fetal candidate genes analyzed were *UCP2* and *SLC2A2* [20, 25, 29–34]. Information on the SNPs evaluated and the selection criteria used is listed in Table 1.

2.3 DNA Samples and Genotyping Analysis

Buccal brushes from mothers, fathers, and infants were collected as part of the NBDPS [35]. DNA was extracted from buccal cells and a standard quality control procedure was applied to each sample before they were submitted to the NBDPS sample repository [35]. To assure genotyping proficiency, high quality, and high concordance among all NBDPS laboratories, annual evaluations are conducted to confirm the performance of each laboratory (See Supplemental Material). Our laboratory at the University of Texas at Austin, Dell Pediatric Research Institute has passed all of these evaluations with a score of 100%. SNPs were assayed using TaqMan method (Life Technologies Corporation, Carlsbad, CA) and genotypes were read and discriminated on the ABI PRISM[®] 7900HT Sequence Detection System (Life Technologies Corporation, Carlsbad, CA).

2.4 Statistical Analysis

The characteristics of cases and case mothers were summarized using counts and proportions for the following variables: phenotype (spina bifida, anencephaly, encephalocele); infant sex (male, female); maternal age (<20, 20–34, 35 years); maternal race/ethnicity (non-Hispanic White, non-Hispanic Black, Hispanic, other); maternal education (<12, 12, 13–15, >15 years); maternal folic acid supplementation during three months before conception through the first month of pregnancy (no, yes); maternal pre-pregnancy body mass index or BMI (underweight [$<18.5 \text{ kg/m}^2$], average weight [$18.5\text{--}24.9 \text{ kg/m}^2$], overweight [$25.0\text{--}29.9 \text{ kg/m}^2$], and obese [$\geq 30.0 \text{ kg/m}^2$]); and maternal pre-pregnancy diabetes (no, yes). For each analyzed polymorphism, samples for which a genotype could not be assigned and triads that had genotype combinations that were inconsistent with Mendelian inheritance were determined. For each subject, the number of genotyping failures (i.e., genotypes that could not be assigned) was determined. These analyses were performed using Intercooled Stata, version 12.1 (StataCorp LP, College Station, TX).

We utilized a 2-step approach to evaluate maternal-fetal gene-gene interactions [36]. For step 1, a case-only approach was used to screen all potential interactions ($n=76$), as this design provides greater power in the assessment of gene-gene interactions compared to a case-control design or case-parent triads [37]. The case-only design has been described elsewhere [38] and has been used extensively for the assessment of gene-environment and gene-gene interactions [36, 38–43]. Specifically, ordinal logistic regression was used to calculate the odds ratio (OR) and 95% confidence interval (CI) for each maternal-fetal gene-gene interaction, assuming a log-additive model of inheritance. The genotypes for each SNP were classified according to the number of minor alleles present (i.e., 0, 1, 2). In the ordinal logistic regression model, the maternal genotype was treated as the dependent variable and the fetal genotype was treated as the independent variable [36, 37, 43]. Due to the number of comparisons, we calculated a corrected p-value (q-value) to control for the false discovery rate (FDR) at 0.05 [44, 45]. These analyses were conducted using Intercooled Stata version 12.1 (StataCorp LP, College Station, TX). All interactions where $q < 0.05$ were included in step 2.

For step 2 (i.e., case-parent triad approach), maternal-fetal gene-gene interactions that were associated with NTDs in the case-only analyses (i.e., $q < 0.05$) were investigated using log-linear models for joint effects [46]. To test the no-interaction null hypothesis, we calculated a 2-degrees-of-freedom likelihood ratio test (LRT) statistic as twice the difference of the log likelihoods for the log-linear model that included two parameters indexing the inherited genotype (SNP1), two parameters indexing the maternal genotype (SNP1), and two interaction terms representing the product of maternal-fetal SNP1-SNP2 pairwise genotypes (SNP2 being the fetal “interacting” SNP) and a reduced model that excluded the interaction terms [36, 46]. These analyses were run using LEM [47], a program for log-linear analysis with missing data that allows information from case-parent triads that have not been completely genotyped (e.g., father not available) to be included in the analysis for any given variant [48]. To reduce concerns regarding possible mating stratification bias [49, 50], we also examined interactions among case-parent triads in which both parents were reported to be non-Hispanic White. Additionally, analyses were conducted in three subgroups: 1) those case-parent triads with spina bifida only (to reduce the potential for phenotypic heterogeneity); 2) those case-parent triads where mothers did not have pre-gestational diabetes (in order to determine if these effects were independent; and 3) those case-parent triads where mothers were not obese (in order to determine if these effects were independent of obesity).

3. RESULTS

Participation in the NBDPS for the period 1999–2007 was 74% among NTD case mothers, yielding 1,553 families available for analysis. Among those, 759 (49%) provided buccal brushes (1,787 individuals). Genotyping was performed on DNA samples derived from these 759 families. Based on quality control checks, 18 families (2% of families) were excluded for being inconsistent with Mendelian inheritance at more than two genotypes. Additionally, 47 subjects were excluded for failure at more than 11 genotypes (>50%), leaving a total of 737 case-parent triads (97% of the original sample). Of those, 317 were complete triads, 313 were dyads, and 107 were monads with only one person in the family.

After these quality control measures were applied, at least 95% of the samples for each variant were available; therefore the genotypes were considered of sufficiently high quality for analysis.

The distributions of key characteristics among NTD case-parent triads are presented in Table 2. Spina bifida was the most common phenotype among case subjects (n=449, 60.9%). Furthermore, a majority of case mothers were non-Hispanic White (n=439, 59.8%). Among case mothers, 176 were obese (25.4%), 13 had pre-pregnancy diabetes (1.8%), and 28 had gestational diabetes (4.2%). The only characteristics presented in Table 2 that were significantly different between interviewed case mothers who provided buccal brushes and those who did not were race/ethnicity (those who provided buccal brushes were more likely to be non-Hispanic White compared to those who did not) and education (those who provided buccal brushes were more likely to have >12 years education compared to those who did not), data not shown.

Of the 76 interactions evaluated, five had $q < 0.05$ (Table 3). Among these five, four were confirmed using log-linear models among case-parent triads (step 2). Our results were similar when restricted to non-Hispanic white mating combinations (data not shown). Specifically, the following interactions were confirmed: maternal *ENPP1* rs1044498-fetal *SLC2A2* rs6785233 (interaction OR=3.65, 95% CI: 2.32–5.74, $q=0.001$, LRT $p=0.00004$); maternal *LEP* rs12706831-fetal *SLC2A2* rs6785233 (interaction OR=0.45, 95% CI: 0.29–0.71, $q=0.016$, LRT $p=0.00001$); maternal *ENPP1* rs1044498-fetal *SLC2A2* rs5400 (interaction OR=1.98, 95% CI: 1.34–2.92, $q=0.016$, LRT $p=0.001$); and maternal *LEP* rs2071045-fetal *SLC2A2* rs5400 (interaction OR=0.50, 95% CI: 0.32–0.77, $q=0.03$, LRT $p=0.008$). As in our previous assessment [24], our results were similar when our analyses were restricted to 1) those case-parent triads with spina bifida only; 2) those case-parent triads where mothers did not have pre-gestational diabetes; and 3) those case-parent triads where mothers were not obese (Table 4 for results among non-obese mothers) [24].

4. DISCUSSION

To our knowledge, this is the first study reporting maternal-fetal gene-gene interactions in metabolic genes and their associations with NTD risk. Significant interactions were identified between the fetal *SLC2A2* gene and maternal variants in *LEP* and *ENPP1* genes. Specifically, four of the 76 interactions were $q < 0.05$ and were confirmed in step 2 of our analysis. The minor alleles of maternal *ENPP1* and fetal *SLC2A2* were associated with increased risk of NTDs, whereas the minor alleles of *LEP* and fetal *SLC2A2* were inversely associated with NTD risk. The direction of these associations is consistent with our previous single locus analysis [24]. Interestingly the maternal (*ENPP1* rs1044498, *LEP* rs12706831, and *LEP* rs2071045) and fetal (*SLC2A2* rs6785233 and *SLC2A2* rs5400) SNPs identified as being significant in these analyses were not significant in single locus analyses [24], suggesting the importance of evaluating factors that may not have significant “main” effects. It is noteworthy and *SLC2A2* gene is known to contribute to impaired glucose tolerance and type 2 diabetes [51]; however, we have previously evaluated potential maternal effect of *SLC2A2* SNPs and no significant association was observed [24].

Leptin is a hormone produced and secreted by white adipose tissue and has profound effects on eating behavior, metabolic rate, endocrine function, and glucose homeostasis. Leptin deficiency in both mice and humans causes morbid obesity and diabetes, and replacement treatment leads to decreased food intake, normalized glucose homeostasis, and increased energy expenditure [32, 52–55]. Two genetic markers adjacent to human *LEP* gene have been found to be modestly associated with NTDs possibly via an inherited effect, irrespective of maternal BMI [56]. In our previous study, we observed a modest increase of NTD risk (though not statistically significant) among women who carried the minor allele of SNP rs2071045 [24]; however, the functionality of this SNP is unknown.

Ectonucleotide pyrophosphate phosphodiesterase (*ENPP1*) is a membrane-bound glycoprotein that inhibits insulin receptor signaling. *ENPP1* is the same protein as liver nucleotide phosphodiesterase and liver alkaline phosphodiesterase 1 and a member of a family of five enzymes (*ENPP1–5*) that regulate nucleotide metabolism [57]. The K121Q polymorphism (rs1044498) in exon 4 of *ENPP1* gene has been associated with insulin resistance in some populations [31] but not others [58–60]. There is evidence suggesting that this variant interacts with adiposity in modulating glucose homeostasis [61, 62]; however, the possible effect of this variant on obesity remains unclear, with variable results [62–64]. We did not find a significant association between the *ENPP1* gene and NTD risk when evaluating main genetic effects in our previous assessment [24].

At the time of neural tube closure (approximately the 4th week of gestation), an embryo receiving excessive amounts of glucose may not be able to regulate these levels, which subsequently leads to abnormal organogenesis and birth defects [25, 65, 66]. In mice, *Glut2* is expressed from the 8-cell stage onward [67]. Under the condition of maternal hyperglycemia, inactivation of the *Glut2* gene in mouse can protect embryos from maternal diabetes-induced NTDs [25]. Our previous study shows that fetal variants in *SLC2A2* (the human homolog of mouse *Glut2*) alone does not significantly influence NTD risk [24]. However, in this analysis, it appears as though *SLC2A2* may interact with maternal *LEP* and *ENPP1* genes to modify the risk of NTDs, suggesting *SLC2A2* may confer sensitivity of the developing embryos under compromised intrauterine environment.

An important strength of our study is the use of data from the NBDPS, the largest population-based study of birth defects, which provided a unique opportunity to examine the interactions between maternal and fetal genes on NTD risk. The case-parent triad design is immune to population stratification bias in the assessment of fetal genotypes [50]. The log-linear modeling approach to analyses also allowed us to include data from incomplete triads (i.e., genotype data is missing for one or two individuals) [48, 68]. An additional strength of the NBDPS is the extensive and standardized case review employed by clinical geneticists, which maximizes homogeneity among case groups. The main weakness of this study was the limited proportion of families with biologic samples (49%), which may limit the generalizability of our findings. In addition, clinical data such as insulin resistance or fasting blood glucose levels are not available as part of the NBDPS; therefore it is not possible to exclude mechanisms other than their associations with maternal obesity or impaired glucose homeostasis that alter the intrauterine environment. In conclusion, our findings suggest that maternal metabolic genes associated with hyperglycemia and insulin resistance and fetal

metabolic genes involved in glucose homeostasis may interact to increase the risk of NTDs. Replication of these findings in other populations and investigation of additional genes is warranted. Furthermore, since maternal obesity and diabetes are also risk factors for other malformations [5, 8, 69], assessing the maternal-fetal gene-gene interactions in other birth defects will broaden our understanding of diabetes and obesity-related teratogenicity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This study was supported by the National Institute of Child Health and Development (NICHD) (H. Zhu: R21 HD 058912). It was also supported by the Centers for Disease Control and Prevention Centers for Excellence Award No. U50/CCU925286 (California), U01/DD000494 (Texas), and NIH R01 NS 050249. This research was also supported in part by a grant from the National Institute of Environmental Health Sciences (P30ES010126). We thank the California Department of Public Health, Maternal Child and Adolescent Health Division for providing surveillance data from California for this study. We also thank the families who participated in this study.

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention or the California Department of Public Health.

Abbreviations

BMI	body mass index
CATI	computer assisted telephone interview
CI	confidence interval
FDR	false discovery rate
GWAS	genome-wide association study
LRT	likelihood ratio test
NBDPS	National Birth Defects Prevention Study
NTDs	neural tube defects
SNP	single nucleotide polymorphism
RR	risk ratio

References

1. Campbell LR, Dayton DH, Sohal GS. Neural tube defects: a review of human and animal studies on the etiology of neural tube defects. *Teratology*. 1986; 34:171–187. [PubMed: 3535149]
2. Ouyang L, Grosse SD, Armour BS, Waitzman NJ. Health care expenditures of children and adults with spina bifida in a privately insured U.S. population. *Birth Defects Res A Clin Mol Teratol*. 2007; 79:552–558. [PubMed: 17335056]
3. Shaw GM, Todoroff K, Finnell RH, Lammer EJ. Spina bifida phenotypes in infants or fetuses of obese mothers. *Teratology*. 2000; 61:376–381. [PubMed: 10777833]
4. Shaw GM, Velie EM, Schaffer D. Risk of neural tube defect-affected pregnancies among obese women. *Jama*. 1996; 275:1093–1096. [PubMed: 8601928]
5. Watkins ML, Rasmussen SA, Honein MA, Botto LD, Moore CA. Maternal obesity and risk for birth defects. *Pediatrics*. 2003; 111:1152–1158. [PubMed: 12728129]

6. Watkins ML, Scanlon KS, Mulinare J, Khoury MJ. Is maternal obesity a risk factor for anencephaly and spina bifida? *Epidemiology*. 1996; 7:507–512. [PubMed: 8862982]
7. Soler NG, Walsh CH, Malins JM. Congenital malformations in infants of diabetic mothers. *Q J Med*. 1976; 45:303–313. [PubMed: 781716]
8. Waller DK, Shaw GM, Rasmussen SA, Hobbs CA, Canfield MA, Siega-Riz AM, Gallaway MS, Correa A. Prepregnancy obesity as a risk factor for structural birth defects. *Arch Pediatr Adolesc Med*. 2007; 161:745–750. [PubMed: 17679655]
9. Waller DK, Mills JL, Simpson JL, Cunningham GC, Conley MR, Lassman MR, Rhoads GG. Are obese women at higher risk for producing malformed offspring? *Am J Obstet Gynecol*. 1994; 170:541–548. [PubMed: 8116710]
10. Hendricks KA, Nuno OM, Suarez L, Larsen R. Effects of hyperinsulinemia and obesity on risk of neural tube defects among Mexican Americans. *Epidemiology*. 2001; 12:630–635. [PubMed: 11679789]
11. Werler MM, Louik C, Shapiro S, Mitchell AA. Prepregnant weight in relation to risk of neural tube defects. *Jama*. 1996; 275:1089–1092. [PubMed: 8601927]
12. Kallen K. Maternal smoking, body mass index, and neural tube defects. *Am J Epidemiol*. 1998; 147:1103–1111. [PubMed: 9645788]
13. Cabrera RM, Hill DS, Etheredge AJ, Finnell RH. Investigations into the etiology of neural tube defects. *Birth Defects Res C Embryo Today*. 2004; 72:330–344. [PubMed: 15662706]
14. Andreasen KR, Andersen ML, Schantz AL. Obesity and pregnancy. *Acta Obstet Gynecol Scand*. 2004; 83:1022–1029. [PubMed: 15488115]
15. Carmichael SL, Rasmussen SA, Shaw GM. Prepregnancy obesity: a complex risk factor for selected birth defects. *Birth Defects Res A Clin Mol Teratol*. 2010; 88:804–810. [PubMed: 20973050]
16. King JC. Maternal obesity, metabolism, and pregnancy outcomes. *Annu Rev Nutr*. 2006; 26:271–291. [PubMed: 16704347]
17. Ray JG, Thompson MD, Vermeulen MJ, Meier C, Wyatt PR, Wong PY, Summers AM, Farrell SA, Cole DE. Metabolic syndrome features and risk of neural tube defects. *BMC Pregnancy Childbirth*. 2007; 7:21. [PubMed: 17880716]
18. Scialli AR. Teratology Public Affairs Committee position paper: maternal obesity and pregnancy. *Birth Defects Res A Clin Mol Teratol*. 2006; 76:73–77. [PubMed: 16463272]
19. Reece EA. Diabetes-induced birth defects: what do we know? What can we do? *Current diabetes reports*. 2012; 12:24–32. [PubMed: 22167469]
20. Zeggini E, McCarthy MI. TCF7L2: the biggest story in diabetes genetics since HLA? *Diabetologia*. 2007; 50:1–4. [PubMed: 17096114]
21. Tung YC, Yeo GS. From GWAS to biology: lessons from FTO. *Ann N Y Acad Sci*. 2011; 1220:162–171. [PubMed: 21388413]
22. Barker A, Langenberg C, Wareham NJ. Genetic determinants of glucose homeostasis Best practice & research. *Clinical endocrinology & metabolism*. 2012; 26:159–170. [PubMed: 22498246]
23. Fall T, Ingelsson E. Genome-wide association studies of obesity and metabolic syndrome. *Molecular and cellular endocrinology*. 2012
24. Lupo PJ, Canfield MA, Chapa C, Lu W, Agopian AJ, Mitchell LE, Shaw GM, Waller DK, Olshan AF, Finnell RH, Zhu H. Diabetes and obesity-related genes and the risk of neural tube defects in the national birth defects prevention study. *Am J Epidemiol*. 2012; 176:1101–1109. [PubMed: 23132673]
25. Li R, Thorens B, Loeken MR. Expression of the gene encoding the high-Km glucose transporter 2 by the early postimplantation mouse embryo is essential for neural tube defects associated with diabetic embryopathy. *Diabetologia*. 2007; 50:682–689. [PubMed: 17235524]
26. Freathy RM, Hayes MG, Urbanek M, Lowe LP, Lee H, Ackerman C, Frayling TM, Cox NJ, Dunger DB, Dyer AR, Hattersley AT, Metzger BE, Lowe WL Jr, Group HSCR. Hyperglycemia and Adverse Pregnancy Outcome (HAPO) study: common genetic variants in GCK and TCF7L2 are associated with fasting and postchallenge glucose levels in pregnancy and with the new consensus definition of gestational diabetes mellitus from the International Association of Diabetes and Pregnancy Study Groups. *Diabetes*. 2010; 59:2682–2689. [PubMed: 20682688]

27. Yoon PW, Rasmussen SA, Lynberg MC, Moore CA, Anderka M, Carmichael SL, Costa P, Druschel C, Hobbs CA, Romitti PA, Langlois PH, Edmonds LD. The National Birth Defects Prevention Study. *Public Health Rep.* 2001; 116(Suppl 1):32–40. [PubMed: 11889273]
28. Rasmussen SA, Olney RS, Holmes LB, Lin AE, Keppler-Noreuil KM, Moore CA. Guidelines for case classification for the National Birth Defects Prevention Study. *Birth Defects Res A Clin Mol Teratol.* 2003; 67:193–201. [PubMed: 12797461]
29. Scott LJ, Mohlke KL, Bonnycastle LL, Willer CJ, Li Y, Duren WL, Erdos MR, Stringham HM, Chines PS, Jackson AU, Prokunina-Olsson L, Ding CJ, Swift AJ, Narisu N, Hu T, Pruim R, Xiao R, Li XY, Conneely KN, Riebow NL, Sprau AG, Tong M, White PP, Hetrick KN, Barnhart MW, Bark CW, Goldstein JL, Watkins L, Xiang F, Saramies J, Buchanan TA, Watanabe RM, Valle TT, Kinnunen L, Abecasis GR, Pugh EW, Doheny KF, Bergman RN, Tuomilehto J, Collins FS, Boehnke M. A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. *Science.* 2007; 316:1341–1345. [PubMed: 17463248]
30. Volcik KA, Shaw GM, Zhu H, Lammer EJ, Finnell RH. Risk factors for neural tube defects: associations between uncoupling protein 2 polymorphisms and spina bifida. *Birth Defects Res A Clin Mol Teratol.* 2003; 67:158–161. [PubMed: 12797456]
31. Pizzuti A, Frittitta L, Argiolas A, Baratta R, Goldfine ID, Bozzali M, Ercolino T, Scarlato G, Iacoviello L, Vigneri R, Tassi V, Trischitta V. A polymorphism (K121Q) of the human glycoprotein PC-1 gene coding region is strongly associated with insulin resistance. *Diabetes.* 1999; 48:1881–1884. [PubMed: 10480624]
32. Ohnuma H, Yamatani K, Daimon M, Igarashi M, Manaka H, Sasaki H, Kato T. Impaired neural regulation of insulin secretion related to the leptin receptor gene mutation in Wistar fatty rats. *Physiol Behav.* 2000; 70:527–532. [PubMed: 11111007]
33. Zhang Y, Wat N, Stratton IM, Warren-Perry MG, Orho M, Groop L, Turner RC. UKPDS 19: heterogeneity in NIDDM: separate contributions of IRS-1 and beta 3-adrenergic-receptor mutations to insulin resistance and obesity respectively with no evidence for glycogen synthase gene mutations. *UK Prospective Diabetes Study. Diabetologia.* 1996; 39:1505–1511. [PubMed: 8960833]
34. Nitz I, Ewert A, Klapper M, Doring F. Analysis of PGC-1alpha variants Gly482Ser and Thr612Met concerning their PPARgamma2-coactivation function. *Biochem Biophys Res Commun.* 2007; 353:481–486. [PubMed: 17187763]
35. Rasmussen SA, Lammer EJ, Shaw GM, Finnell RH, McGehee RE Jr, Gallagher M, Romitti PA, Murray JC. Integration of DNA sample collection into a multi-site birth defects case-control study. *Teratology.* 2002; 66:177–184. [PubMed: 12353214]
36. Lupo PJ, Goldmuntz E, Mitchell LE. Gene-gene interactions in the folate metabolic pathway and the risk of conotruncal heart defects. *Journal of Biomedicine and Biotechnology.* 2010; 2010:630940. [PubMed: 20111745]
37. Gauderman WJ. Sample size requirements for association studies of gene-gene interaction. *American Journal of Epidemiology.* 2002; 155:478–484. [PubMed: 11867360]
38. Khoury MJ, Flanders WD. Nontraditional epidemiologic approaches in the analysis of gene-environment interaction: case-control studies with no controls! *American Journal of Epidemiology.* 1996; 144:207–213. [PubMed: 8686689]
39. Gatto NM, Campbell UB, Rundle AG, Ahsan H. Further development of the case-only design for assessing gene-environment interaction: evaluation of and adjustment for bias. *International Journal of Epidemiology.* 2004; 33:1014–1024. [PubMed: 15358745]
40. Piegorsch WW, Weinberg CR, Taylor JA. Non-hierarchical logistic models and case-only designs for assessing susceptibility in population-based case-control studies. *Statistics in Medicine.* 1994; 13:153–162. [PubMed: 8122051]
41. Dennis J, Hawken S, Krewski D, Birkett N, Gheorghe M, Frei J, McKeown-Eyssen G, Little J. Bias in the case-only design applied to studies of gene-environment and gene-gene interaction: a systematic review and meta-analysis. *International Journal of Epidemiology.* 2011; 40:1329–1341. [PubMed: 21729879]
42. Pande M, Amos CI, Eng C, Frazier ML. Interactions between cigarette smoking and selected polymorphisms in xenobiotic metabolizing enzymes in risk for colorectal cancer: A case-only analysis. *Molecular Carcinogenesis.* 2010; 49:974–980. [PubMed: 20886582]

43. Yang Y, Tian Y, Jin X, Yan C, Jiang F, Zhang Y, Tang J, Shen X. A case-only study of interactions between metabolic enzyme polymorphisms and industrial pollution in childhood acute leukemia. *Environmental Toxicology and Pharmacology*. 2009; 28:161–166. [PubMed: 21783998]
44. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society, Series B (Methodological)*. 1995; 57:289–300.
45. Storey JD. The positive false discovery rate: a Bayesian interpretation and the q-value. *The Annals of Statistics*. 2003; 64:2013–2035.
46. Umbach DM, Weinberg CR. The use of case-parent triads to study joint effects of genotype and exposure. *American Journal of Human Genetics*. 2000; 66:251–261. [PubMed: 10631155]
47. Vermunt, JK. LEM: A general program for the analysis of categorical data. Tilberg University; 1997.
48. Weinberg CR. Allowing for missing parents in genetic studies of case-parent triads. *Am J Hum Genet*. 1999; 64:1186–1193. [PubMed: 10090904]
49. Wang LY, Lee WC. Population stratification bias in the case-only study for gene-environment interactions. *American Journal of Epidemiology*. 2008; 168:197–201. [PubMed: 18497429]
50. Weinberg CR, Wilcox AJ, Lie RT. A log-linear approach to case-parent-triad data: assessing effects of disease genes that act either directly or through maternal effects and that may be subject to parental imprinting. *Am J Hum Genet*. 1998; 62:969–978. [PubMed: 9529360]
51. Laukkanen O, Lindstrom J, Eriksson J, Valle TT, Hamalainen H, Ilanne-Parikka P, Keinanen-Kiukaanniemi S, Tuomilehto J, Uusitupa M, Laakso M, Finnish S. Diabetes Prevention, Polymorphisms in the SLC2A2 (GLUT2) gene are associated with the conversion from impaired glucose tolerance to type 2 diabetes: the Finnish. *Diabetes Prevention Study Diabetes*. 2005; 54:2256–2260.
52. Gautron L, Elmquist JK. Sixteen years and counting: an update on leptin in energy balance. *J Clin Invest*. 2011; 121:2087–2093. [PubMed: 21633176]
53. Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JM. Positional cloning of the mouse obese gene and its human homologue. *Nature*. 1994; 372:425–432. [PubMed: 7984236]
54. Iida M, Murakami T, Ishida K, Mizuno A, Kuwajima M, Shima K. Phenotype-linked amino acid alteration in leptin receptor cDNA from Zucker fatty (fa/fa) rat. *Biochem Biophys Res Commun*. 1996; 222:19–26. [PubMed: 8630068]
55. Chua SC Jr, White DW, Wu-Peng XS, Liu SM, Okada N, Kershaw EE, Chung WK, Power-Keohoe L, Chua M, Tartaglia LA, Leibel RL. Phenotype of fatty due to Gln269Pro mutation in the leptin receptor (Lepr). *Diabetes*. 1996; 45:1141–1143. [PubMed: 8690163]
56. Shaw GM, Barber R, Todoroff K, Lammer EJ, Finnell RH. Microsatellites proximal to leptin and leptin receptor as risk factors for spina bifida. *Teratology*. 2000; 61:231–235. [PubMed: 10661913]
57. Goldfine ID, Maddux BA, Youngren JF, Reaven G, Accili D, Trischitta V, Vigneri R, Frittitta L. The role of membrane glycoprotein plasma cell antigen 1/ectonucleotide pyrophosphatase phosphodiesterase 1 in the pathogenesis of insulin resistance and related abnormalities. *Endocrine reviews*. 2008; 29:62–75. [PubMed: 18199690]
58. Rasmussen SK, Urhammer SA, Pizzuti A, Echwald SM, Ekstrom CT, Hansen L, Hansen T, Borch-Johnsen K, Frittitta L, Trischitta V, Pedersen O. The K121Q variant of the human PC-1 gene is not associated with insulin resistance or type 2 diabetes among Danish Caucasians. *Diabetes*. 2000; 49:1608–1611. [PubMed: 10969849]
59. Gonzalez-Sanchez JL, Martinez-Larrad MT, Fernandez-Perez C, Kubaszek A, Laakso M, Serrano-Rios M. K121Q PC-1 gene polymorphism is not associated with insulin resistance in a Spanish population. *Obes Res*. 2003; 11:603–605. [PubMed: 12740448]
60. Keshavarz P, Inoue H, Sakamoto Y, Kunika K, Tanahashi T, Nakamura N, Yoshikawa T, Yasui N, Shiota H, Itakura M. No evidence for association of the ENPP1 (PC-1) K121Q variant with risk of type 2 diabetes in a Japanese population. *J Hum Genet*. 2006; 51:559–566. [PubMed: 16607460]
61. Maranghi M, Prudente S, D’Erasmus L, Morini E, Ciociola E, Coletta P, Verrienti A, Arciello S, Copetti M, Pellegrini F, Santini SA, Morano S, Filetti S, Trischitta V. The ectonucleotide pyrophosphatase phosphodiesterase 1 (ENPP1) K121Q polymorphism modulates the beneficial

- effect of weight loss on fasting glucose in non-diabetic individuals. *Nutr Metab Cardiovasc Dis*. 2012
62. Pan W, Ciociola E, Saraf M, Tumurbaatar B, Tuvdendorj D, Prasad S, Chandalia M, Abate N. Metabolic consequences of ENPP1 overexpression in adipose tissue. *Am J Physiol Endocrinol Metab*. 2011; 301:E901–911. [PubMed: 21810932]
 63. Wan C, Zhang T, Wang B, Han Y, Zhang C, Zhang Y, Gong H, Jin F, Wang L. Obesity risk associated with the K121Q polymorphism of the glycoprotein PC-1 gene. *Diabetes Obes Metab*. 2006; 8:703–708. [PubMed: 17026496]
 64. Grarup N, Urhammer SA, Ek J, Albrechtsen A, Glumer C, Borch-Johnsen K, Jorgensen T, Hansen T, Pedersen O. Studies of the relationship between the ENPP1 K121Q polymorphism and type 2 diabetes, insulin resistance and obesity in 7,333 Danish white subjects. *Diabetologia*. 2006; 49:2097–2104. [PubMed: 16865358]
 65. Trocino RA, Akazawa S, Takino H, Takao Y, Matsumoto K, Maeda Y, Okuno S, Nagataki S. Cellular-tissue localization and regulation of the GLUT-1 protein in both the embryo and the visceral yolk sac from normal and experimental diabetic rats during the early postimplantation period. *Endocrinology*. 1994; 134:869–878. [PubMed: 8299581]
 66. Maeda Y, Akazawa S, Akazawa M, Takao Y, Trocino RA, Takino H, Kawasaki E, Yokota A, Okuno S, Nagataki S. Glucose transporter gene expression in rat conceptus during early organogenesis and exposure to insulin-induced hypoglycemic serum. *Acta Diabetol*. 1993; 30:73–78. [PubMed: 8219261]
 67. Hogan A, Heyner S, Charron MJ, Copeland NG, Gilbert DJ, Jenkins NA, Thorens B, Schultz GA. Glucose transporter gene expression in early mouse embryos. *Development*. 1991; 113:363–372. [PubMed: 1765007]
 68. Dempster LNAP, Rubin DB. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society Series B*. 1977; 39:1–28.
 69. Correa A, Gilboa SM, Besser LM, Botto LD, Moore CA, Hobbs CA, Cleves MA, Riehle-Colarusso TJ, Waller DK, Reece EA. Diabetes mellitus and birth defects. *Am J Obstet Gynecol*. 2008; 199:237, e231–239. [PubMed: 18674752]

Highlights

- Single-gene analyses indicate that maternal genes associated with metabolic conditions may influence the risk of neural tube defects (NTDs)
- In order to evaluate the interactions between maternal and fetal genes related to glucose homeostasis and the risk of NTDs we investigated 23 single nucleotide polymorphisms among 7 maternal metabolic genes and 2 fetal metabolic genes
- Samples were obtained from 737 NTD case-parent triads included in the National Birth Defects Prevention Study
- We found 5 statistically significant maternal-fetal gene-gene interactions
- Our findings suggest that maternal metabolic genes associated with hyperglycemia and insulin resistance and fetal metabolic genes involved in glucose homeostasis may interact to increase the risk of NTDs

Table 1

Metabolic genes and SNPs included in maternal-fetal gene-gene interaction analysis

Gene symbol	Ref SNP	Chr ^a	Position	Alleles ^b	SNP information	MAF ^c (CEU)	Selection criteria
<i>TCF7L2</i>	rs12255372	10	114808902	G/T	Intron	0.21	Diabetes-associated
<i>TCF7L2</i>	rs7903146	10	114758349	C/T	Intron	0.22	Diabetes-associated
<i>TCF7L2</i>	rs290487	10	114909731	C/T	Intron	0.27	Diabetes-associated
<i>TCF7L2</i>	rs10885390	10	114640797	T/A	Intergenic	0.24	Diabetes-associated
<i>TCF7L2</i>	rs3814573	10	114898093	C/T	Intron	0.40	Diabetes-associated
<i>UCP2</i>	rs660339	11	73689104	G/A	Missense	0.43	Obesity/diabetes
<i>ENPP1</i>	rs1044498	6	132172368	A/C	Missense	0.31	Insulin resistance
<i>FTO</i>	rs9939609	16	53820527	T/A	Intron	0.38	Obesity, BMI
<i>FTO</i>	rs8050136	16	53816275	C/A	Intron	0.37	Obesity, BMI
<i>FTO</i>	rs1421085	16	53800954	T/C	Intron	0.26	Obesity, BMI
<i>FTO</i>	rs17817449	16	53813367	T/G	Intron	0.35	Obesity, BMI
<i>ADRB3</i>	rs4994	8	37823798	T/C	Missense	0.10	Obesity, BMI
<i>PPARG</i>	rs1801282	3	12393125	C/G	Intron	0.06	Obesity-associated
<i>PPARGC1A</i>	rs8192678	4	23815662	G/A	Missense	0.30	Obesity/metabolic disorders
<i>PPARGC1A</i>	rs3736265	4	23814707	G/A	Missense	0.11	Obesity/metabolic disorders
<i>LEP</i>	rs11760956	7	127891087	G/A	Intron	0.29	tagSNP
<i>LEP</i>	rs12706831	7	127887068	T/G	Intron	0.46	tagSNP
<i>LEP</i>	rs3828942	7	127894305	G/A	Intron	0.45	tagSNP
<i>LEP</i>	rs2071045	7	127892980	T/C	Intron	0.26	tagSNP
<i>LEP</i>	rs2167270	7	127881349	G/A	5'utr	0.35	tagSNP
<i>SLC2A2</i>	rs11924032	3	170735099	G/A	Intron	0.31	tagSNP
<i>SLC2A2</i>	rs6785233	3	170756985	T/G	Intergenic	0.19	tagSNP
<i>SLC2A2</i>	rs5400	3	170732300	C/T	Missense	0.21	Diabetes-associated, cholesterol levels

^aChr (Chromosome) Genomic Build 37.1; group term GRCh37^bRefSNP alleles: reference allele/risk allele (minor allele)^cMAF (Minor Allele Frequency) source: 1000 Genomes project

Table 2

Characteristics of neural tube defect case-parent triads (n=737), National Birth Defects Prevention Study, 1999–2007

Characteristic	No.	%
Phenotype		
Spina bifida	449	60.9
Anencephaly	217	29.4
Encephalocele	71	9.6
Infant sex		
Male	337	47.9
Female	366	52.1
Maternal age		
<20	83	11.3
20–34	556	75.4
35	98	13.3
Race/ethnicity		
Non-Hispanic White	439	59.8
Non-Hispanic Black	34	4.6
Hispanic	221	30.1
Other	40	5.5
Education (years)		
<12	142	19.3
12	184	25.0
13–15	226	30.7
>15	185	25.0
Folic acid supplementation ^a		
No	351	47.6
Yes	386	52.4
Body mass index (kg/m ²)		
Underweight (<18.5)	28	4.1
Normal (18.5–24.9)	336	48.6
Overweight (25.0–29.9)	152	21.9
Obese (≥30)	176	25.4
Pre-pregnancy diabetes		
No	724	98.2
Yes	13	1.8
Gestational diabetes		
No	667	95.8
Yes	29	4.2

^aThree months before conception through the first month of pregnancy

Table 3
 Top maternal-fetal metabolic gene-gene interactions associated with neural tube defects, National Birth Defects Prevention Study, 1999–2007

Maternal SNP	Fetal SNP	Interaction OR ^a	95% CI ^b	p-value	q-value ^c	LRT p-value from step 2 ^{d,e}
<i>ENPP1</i> rs1044498	<i>SLC2A2</i> rs6785233	3.65	2.32–5.74	2.09E-08	0.001	0.00004
<i>LEP</i> rs12706831	<i>SLC2A2</i> rs6785233	0.45	0.29–0.71	0.0005	0.016	0.00001
<i>ENPP1</i> rs1044498	<i>SLC2A2</i> rs5400	1.98	1.34–2.92	0.0006	0.016	0.001
<i>LEP</i> rs2071045	<i>SLC2A2</i> rs5400	0.50	0.32–0.77	0.0016	0.03	0.008
<i>LEP</i> rs2071045	<i>SLC2A2</i> rs6785233	0.46	0.27–0.77	0.0029	0.04	0.06
<i>LEP</i> rs12706831	<i>SLC2A2</i> rs5400	0.59	0.41–0.86	0.0059	0.08	NE
<i>LEP</i> rs11760956	<i>SLC2A2</i> rs6785233	0.63	0.41–0.99	0.0450	0.44	NE
<i>LEP</i> rs3828942	<i>SLC2A2</i> rs6785233	0.64	0.42–0.99	0.0473	0.44	NE

^a Interaction odds ratio (OR) from step 1 (case-only analysis)

^b Confidence interval (CI)

^c False discovery rate q-value

^d Only interactions where $q < 0.05$ in step 1 were confirmed in step 2 (log-linear analysis in case-parent triads)

^e Not estimated (NE)

Table 4

Top maternal-fetal metabolic gene-gene interactions in non-obese mothers associated with neural tube defects, National Birth Defects Prevention Study, 1999–2007

Maternal SNP	Fetal SNP	Interaction OR ^a	95% CI ^b	p-value
<i>ENPP1</i> rs1044498	<i>SLC2A2</i> rs6785233	3.24	1.83–5.74	5.18E-05
<i>LEP</i> rs12706831	<i>SLC2A2</i> rs6785233	0.54	0.31–0.93	0.0027
<i>ENPP1</i> rs1044498	<i>SLC2A2</i> rs5400	1.78	1.12–2.86	0.0153
<i>LEP</i> rs2071045	<i>SLC2A2</i> rs5400	0.58	0.37–0.91	0.0259
<i>LEP</i> rs2071045	<i>SLC2A2</i> rs6785233	0.49	0.27–0.88	0.0172
<i>LEP</i> rs12706831	<i>SLC2A2</i> rs5400	0.58	0.37–0.91	0.0176
<i>LEP</i> rs11760956	<i>SLC2A2</i> rs6785233	0.85	0.50–1.44	0.5506
<i>LEP</i> rs3828942	<i>SLC2A2</i> rs6785233	0.72	0.43–1.25	0.2552

^aInteraction odds ratio (OR) from case-only analysis

^bConfidence interval (CI)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript