



## NIH PUBLIC ACCESS

## Author Manuscript

*Nat Med.* Author manuscript; available in PMC 2009 January 26.

Published in final edited form as:

*Nat Med.* 2008 April ; 14(4): 413–420. doi:10.1038/nm1741.**CCL3L1-CCR5 genotype influences durability of immune recovery during antiretroviral therapy of HIV-1–infected individuals****Sunil K Ahuja<sup>1,3</sup>, Hemant Kulkarni<sup>1,2,13</sup>, Gabriel Catano<sup>1,2,13</sup>, Brian K Agan<sup>4,7,13</sup>, Jose F Camargo<sup>1,2</sup>, Weijing He<sup>1,2</sup>, Robert J O'Connell<sup>4,5</sup>, Vincent C Marconi<sup>4,5,7</sup>, Judith Delmar<sup>4,5,7</sup>, Joseph Eron<sup>8</sup>, Robert A Clark<sup>1,2</sup>, Simon Frost<sup>9,10</sup>, Jeffrey Martin<sup>11</sup>, Seema S Ahuja<sup>1,2</sup>, Steven G Deeks<sup>12</sup>, Susan Little<sup>9</sup>, Douglas Richman<sup>9,10</sup>, Frederick M Hecht<sup>12</sup>, and Matthew J Dolan<sup>4,7</sup>**<sup>1</sup>*Veterans Administration Research Center for AIDS and HIV-1 Infection, South Texas Veterans Health Care System, 7400 Merton Minter, San Antonio, Texas 78229, USA.*<sup>2</sup>*Department of Medicine, University of Texas Health Science Center, 7703 Floyd Curl Drive, San Antonio, Texas 78229, USA.*<sup>3</sup>*Departments of Microbiology and Immunology and Biochemistry, University of Texas Health Science Center, 7703 Floyd Curl Drive, San Antonio, Texas 78229, USA.*<sup>4</sup>*Infectious Disease Clinical Research Program, Uniformed Services University, 4301 Jones Bridge Road, Bethesda, Maryland 20814, USA.*<sup>5</sup>*Infectious Diseases Service, Wilford Hall United States Air Force Medical Center, 2200 Bergquist Drive, Lackland Air Force Base, Texas 78236, USA.*<sup>6</sup>*Henry M. Jackson Foundation, Wilford Hall United States Air Force Medical Center, 2200 Bergquist Drive, Lackland Air Force Base, Texas 78236, USA.*<sup>7</sup>*San Antonio Military Medical Center, 3851 Roger Brooke Drive, Fort Sam Houston, Texas 78234, USA.*<sup>8</sup>*University of North Carolina, 4030 Bondurant Hall, Chapel Hill, North Carolina 27599, USA.*<sup>9</sup>*Antiviral Research Center, Department of Medicine, University of California, 150 West Washington Street, San Diego, California 92103, USA.*<sup>10</sup>*Veterans Affairs San Diego Healthcare System, 3350 La Jolla Village Drive, San Diego, California 92161, USA.*<sup>11</sup>*Department of Epidemiology and Biostatistics, University of California, 185 Berry Street, San Francisco, California 94107, USA.*<sup>12</sup>*Department of Medicine, San Francisco General Hospital, University of California–San Francisco, 995 Potrero Avenue, San Francisco, California 94110, USA.***Abstract**Correspondence should be addressed to S.K.A. ([ahujas@uthscsa.edu](mailto:ahujas@uthscsa.edu)) or M.J.D. ([mdolan@hjf.org](mailto:mdolan@hjf.org)).<sup>13</sup>These authors contributed equally to this work.**AUTHOR CONTRIBUTIONS**

S.K.A., H.K. and M.J.D. conceptualized the research, conducted statistical analyses, analyzed the data and wrote the manuscript. G.C., B.K.A., J.F.C., W.H., R.J.O., V.C.M., J.D., J.E., R.A.C., S.F., J.M., S.S.A., S.G.D., S.L., D.R., and F.M.H. made substantial contributions to conceptual ideas, experimental data, analysis of data, preparation of the manuscript and/or cohort data and development. S.K.A. and M.J.D. directed the study and obtained the funding for the genetic work.

Note: Supplementary information is available on the Nature Medicine website.

Reprints and permissions information is available online at <http://npg.nature.com/reprintsandpermissions>

The basis for the extensive variability seen in the reconstitution of CD4<sup>+</sup> T cell counts in HIV-infected individuals receiving highly active antiretroviral therapy (HAART) is not fully known. Here, we show that variations in *CCL3L1* gene dose and *CCR5* genotype, but not major histocompatibility complex HLA alleles, influence immune reconstitution, especially when HAART is initiated at <350 CD4<sup>+</sup> T cells/mm<sup>3</sup>. The *CCL3L1-CCR5* genotypes favoring CD4<sup>+</sup> T cell recovery are similar to those that blunted CD4<sup>+</sup> T cell depletion during the time before HAART became available (pre-HAART era), suggesting that a common *CCL3L1-CCR5* genetic pathway regulates the balance between pathogenic and reparative processes from early in the disease course. Hence, *CCL3L1-CCR5* variations influence HIV pathogenesis even in the presence of HAART and, therefore, may prospectively identify subjects in whom earlier initiation of therapy is more likely to mitigate immunologic failure despite viral suppression by HAART. Furthermore, as reconstitution of CD4<sup>+</sup> cells during HAART is more sensitive to *CCL3L1* dose than to *CCR5* genotypes, *CCL3L1* analogs might be efficacious in supporting immunological reconstitution.

---

HIV-1 infection is characterized by extensive variability in the degree of depletion or recovery in CD4<sup>+</sup> T cell (abbreviated as CD4<sup>+</sup>) counts before or after receipt of HAART, respectively. Variability in CD4<sup>+</sup> depletion cannot be attributed solely to the extent of viral replication (as reflected by the viral load)<sup>1,2</sup>, as host factors can influence HIV disease course independently of viral load<sup>2,3</sup>. Similarly, variability in the depth of HAART-induced suppression of viral replication cannot be the sole factor underlying intersubject differences in recovery of CD4<sup>+</sup> counts. For example, the majority of HIV-positive subjects who initiate therapy with a low CD4<sup>+</sup> nadir during untreated disease (<200 cells/mm<sup>3</sup>) do not fully recover peripheral or mucosal CD4<sup>+</sup> counts, despite HIV-suppressive HAART<sup>4-12</sup>. Conversely, some HIV-positive individuals show sustained recovery of CD4<sup>+</sup> counts in the face of incomplete viral suppression<sup>5,10,13-20</sup>. However, the host factors that mediate this variable CD4<sup>+</sup> recovery are largely unknown.

Here we tested the hypothesis that CD4<sup>+</sup> depletion and recovery reflect a dynamic balance between pathogenic and reparative processes integrally linked by specific common genetic pathways programmed during early stages of HIV disease. The importance of events occurring within the initial phases of HIV disease is highlighted by the observation that the magnitude of early immune damage determines the development and durability of virus-specific immunity, which dictates, in part, steady-state viral load, initial CD4<sup>+</sup> loss and, subsequently, HIV disease course<sup>1,3,21-23</sup>. However, once host genetic factors skew the balance toward pathogenic processes (CD4<sup>+</sup> depletion) during early untreated infection, it is unclear whether this imbalance can be fully reversed, even in the presence of HAART-induced viral suppression.

To test our hypothesis, we determined whether specific genetic factors that skew the balance toward pathogenic or reparative processes during early HIV disease also exert a negative or positive influence, respectively, on recovery of CD4<sup>+</sup> counts during HAART. We focused on polymorphisms in the genes encoding *CCR5*, the major HIV co-receptor, and *CCL3L1*, its most potent HIV-suppressive ligand<sup>24</sup>, for three reasons. First, *CCR5* genotypes and *CCL3L1* gene copy number (that is, gene dose) influence HIV-AIDS susceptibility<sup>3,25-29</sup>. Second, *CCL3L1-CCR5* genetic risk groups (GRGs), defined according to different inherited combinations of *CCL3L1* dose and *CCR5* polymorphisms, influence the extent of early immune damage<sup>3</sup>. Third, *CCL3L1-CCR5* GRGs affect HIV disease progression by affecting viral load and parameters independent of the viral load (for example, cell-mediated immunity)<sup>3</sup>.

To provide a context for our hypothesis, we considered two additional issues. First, we assessed whether the impact of *CCL3L1-CCR5* GRGs on CD4<sup>+</sup> recovery is mitigated by HAART-induced suppression of viral replication during acute or early infection. Second, HLA alleles are crucial in determining the viral load set-point, degree of CD4<sup>+</sup> depletion and HIV disease

course<sup>29,30</sup>, but whether those HLA alleles that influence early immune damage also affect CD4<sup>+</sup> recovery is unknown; thus we studied the effect of HLA alleles on recovery of CD4<sup>+</sup> counts during HAART.

## RESULTS

### ***CCL3L1*, *CCR5*, GRG and CD4<sup>+</sup> dynamics during HIV infection**

During the pre-HAART era, the overall CD4<sup>+</sup> counts within the HIV<sup>+</sup> Wilford Hall Medical Center (WHMC) cohort declined progressively, reaching ~300 cells/mm<sup>3</sup> (Fig. 1a). During the HAART era, CD4<sup>+</sup> counts increased, peaked in 2001, and remained at ~500 cells/mm<sup>3</sup> thereafter (Fig. 1a).

However, the time trends for the CD4<sup>+</sup> counts differed according to a person's *CCL3L1* and *CCR5* genetic makeup, that is, low or high *CCL3L1* dose (Fig. 1b) and detrimental or non-detrimental *CCR5* genotype<sup>27</sup> (Fig. 1c). Both *CCL3L1*<sup>low</sup> and *CCR5*<sup>det</sup> genotype conferred rapid CD4<sup>+</sup> loss and impaired CD4<sup>+</sup> recovery during the pre- and post-HAART eras, respectively (Fig. 1b,c). However, the differences in CD4<sup>+</sup> recovery associated with *CCL3L1*<sup>low</sup> and *CCL3L1*<sup>high</sup> (Fig. 1b) were more prominent than those associated with *CCR5*<sup>det</sup> and *CCR5*<sup>non-det</sup> (Fig. 1c). Concordantly, among subjects of European and African descent, a stepwise increase in *CCL3L1* copy number was associated with a stepwise increase in CD4<sup>+</sup> recovery rates during HAART (Supplementary Fig. 1a online). Analyses of the joint effects of *CCL3L1* and *CCR5* revealed that going from a low to moderate to high *CCL3L1*-*CCR5* GRG was associated with accelerating depletion during the pre-HAART era or increasingly impaired recovery during the HAART era (Fig. 1d).

### ***CCL3L1*, *CCR5*, GRG and CD4<sup>+</sup> recovery during chronic infection**

Time trends of CD4<sup>+</sup> counts indexed from the time of initiation of HAART contrasting *CCL3L1* dose, *CCR5* genotype and GRG (Fig. 1e> and Table 1) were similar to those detected by calendar-era analysis in the WHMC HIV<sup>+</sup> cohort during the HAART era (Fig. 1b–d). We next determined whether suppression of viral replication by HAART overshadowed the effects of variations in *CCL3L1* and *CCR5* on CD4<sup>+</sup> recovery. Those HIV-positive individuals who had HAART-induced viral load suppression had robust increases in CD4<sup>+</sup> counts and slower AIDS progression rates compared with those who were non-viral load suppressors (Supplementary Fig. 1b,c). Additionally, although HIV-infected individuals with transient or sustained viral load suppression did not differ in their rates of disease progression, gains in CD4<sup>+</sup> counts were significantly greater in those with sustained viral load suppression (Supplementary Fig. 1b,c). However, even among subjects with sustained viral load suppression, *CCL3L1*-*CCR5* variations affected CD4<sup>+</sup> recovery (Table 1 and Supplementary Fig. 1d). Additionally, these effects were also evident when complementary statistical methods (mixed models) were used (Supplementary Fig. 2 online).

### ***CCL3L1*, *CCR5*, GRG and durability of CD4<sup>+</sup> recovery**

Inspection of CD4<sup>+</sup> trajectories indicated that during the first 2 years of HAART, CD4<sup>+</sup> counts increased regardless of genotype, but differed thereafter according to variations in *CCL3L1* and *CCR5* (Fig. 1e). Analyses of CD4<sup>+</sup> recovery rates before and after 2 years of HAART substantiated these observations (Fig. 1f).

### **Additive effects of *CCL3L1* and *CCR5***

Although a low *CCL3L1* dose is a feature common to both high and moderate GRGs (Fig. 1d), subjects in a high GRG had significantly lower rates of CD4<sup>+</sup> gains as compared with those in a moderate GRG (Fig. 1f and Table 1). Furthermore, among all subjects who had half of the

average *CCL3L1* dose of their ethnic or racial group, those who also had a detrimental *CCR5* genotype lost CD4<sup>+</sup> cells at a faster rate than those who had a nondetrimental *CCR5* genotype (Supplementary Fig. 1e). These findings indicate that the effects of a detrimental *CCR5* genotype and a low *CCL3L1* dose on CD4<sup>+</sup> recovery are additive, and that a nondetrimental *CCR5* genotype attenuates the negative impact of a low *CCL3L1* dose on CD4<sup>+</sup> gains.

### ***CCL3L1*, *CCR5*, GRG and CD4<sup>+</sup> recovery during early infection**

HIV-positive subjects in the Acute Infection and Early Disease Research Program (AIEDRP) were stratified according to whether they received HAART during acute or early infection, and we analyzed the effects of *CCL3L1* dose and *CCR5* genotype on CD4<sup>+</sup> recovery in these two groups of subjects separately. Among subjects who initiated HAART during acute infection and who attained viral load suppression, a low GRG was associated with both a higher baseline CD4<sup>+</sup> count at time of enrollment into the cohort (619.10 cells/mm<sup>3</sup> versus 440.44 cells/mm<sup>3</sup> in those with low versus moderate-high GRG, respectively;  $P = 0.0253$ ), and an increase of ~150 cells/mm<sup>3</sup> in mean CD4<sup>+</sup> counts through 12 months of follow-up (Fig. 2a). In contrast, in subjects with a moderate or high GRG, post-HAART CD4<sup>+</sup> counts remained similar to those detected at baseline (~500 CD4<sup>+</sup> cells/mm<sup>3</sup>; Fig. 2a). The higher baseline CD4<sup>+</sup> counts observed in those with a low GRG than in those with a moderate or high GRG is consistent with our previous report showing that possession of a low GRG blunts the extent of the initial loss in CD4<sup>+</sup> cells<sup>3</sup>.

Similar findings were detected in those who received HAART during early infection. Viral load suppression during early HIV infection was associated with robust gains (19.8 CD4<sup>+</sup> cells/month) in CD4<sup>+</sup> counts (Fig. 2b). However, even among those who initiated HAART during early infection and who attained viral load suppression, the CD4<sup>+</sup> recovery rate was greatest in those with a low GRG (Fig. 2b). Of note, the individual components of the GRGs—*CCL3L1* and *CCR5*—influenced CD4<sup>+</sup> recovery (Fig. 2c).

### **GRG effects on CD4<sup>+</sup> recovery according to timing of HAART**

We next determined whether CD4<sup>+</sup> recovery was sensitive to both the *CCL3L1-CCR5* GRG (or *CCL3L1* dose alone) and the immunological status of a subject (as reflected by CD4<sup>+</sup> count at the time of initiation of HAART). Among individuals from the WHMC cohort who initiated HAART at <350 CD4<sup>+</sup> cells/mm<sup>3</sup>, a low GRG was associated with CD4<sup>+</sup> gains, whereas recovery was highly muted in those with a moderate or high GRG (Fig. 2d and Table 2). Notably, each 50-cell decrease in the CD4<sup>+</sup> count at the time at which HAART was initiated was associated with a step-wise decline in recovery rates in those possessing a moderate or high GRG and, consequently, a progressively widening difference in CD4<sup>+</sup> recovery rates for those with a low versus moderate or high GRG (Table 2). Notably, the latter associations persisted even when the high GRG (prevalence of 8%) was removed from analyses (Table 2), indicating that the associations observed are not solely due to a high GRG and that a moderate GRG (prevalence of 42%) also has a significant negative impact on CD4<sup>+</sup> recovery rates. In subjects who initiated HAART at ≥350 CD4<sup>+</sup> cells/mm<sup>3</sup>, the time trends in CD4<sup>+</sup> counts did not reveal significant differences in CD4<sup>+</sup> recovery (Fig. 2d). Similar patterns of CD4<sup>+</sup> recovery were observed in therapy-naïve and therapy-experienced subjects (Supplementary Fig. 3a online). Notably, the differential impact of genotype in subjects who differed according to the CD4<sup>+</sup> count at HAART initiation was even more prominent when subjects were categorized according to *CCL3L1* dose (Fig. 2e).

### **Specificity of GRG effects on CD4<sup>+</sup> recovery**

The influence of *CCL3L1-CCR5* GRGs on CD4<sup>+</sup> recovery rates could not be explained by significant inter-GRG differences in viral load suppression or other parameters such as the

number of subjects who initiated HAART at  $<350$  or  $\geq 350$  CD4<sup>+</sup> cells/mm<sup>3</sup> according to GRG status, the age at which HAART was initiated, or the pre-HAART CD4<sup>+</sup> cell count or viral load (Supplementary Methods and Supplementary Note online). We found that even after adjustment for these covariates (potential confounders), a moderate or high GRG remained as a significant independent determinant of impaired CD4<sup>+</sup> recovery rate after 2 years of HAART in those initiating HAART at  $<350$  CD4<sup>+</sup> cells/mm<sup>3</sup> (Table 3). A trend toward a negative effect of a moderate-high GRG on CD4<sup>+</sup> recovery rates was also observed in those who initiated HAART at  $\geq 350$  cells/mm<sup>3</sup> (Table 3). These results, together with the findings discussed below, indicate that the effect of GRG status on CD4<sup>+</sup> recovery is independent of other parameters that are also known to influence immune reconstitution during HAART.

### **Influence of disease-modifying HLA alleles on CD4<sup>+</sup> recovery**

To test further the generalizability of our hypothesis, we focused on those HLA alleles that influence early immune damage as reflected by steady-state viral load, initial CD4<sup>+</sup> loss and delayed-type hypersensitivity (DTH) skin test responses, an *in vivo* marker of cell-mediated immunity. We addressed this by adopting two unbiased approaches. First, we determined which of the HLA class I alleles at the A, B, and C loci and class II alleles at the DRB1 locus independently influenced AIDS progression rates in a multivariate model. In the combined analyses of European and African Americans from the WHMC, HLA-A\*68, HLA-B\*57 and HLA-C\*16 were the only three HLA alleles that predicted altered disease progression rates, with HLA-A\*68 and HLA-C\*16 associating with disease acceleration and HLA-B\*57 associating with disease retardation (Fig. 3a). Of these three alleles, only the HLA-B\*57 allele also influenced all three markers of early immune damage, as well as the cumulative CD4<sup>+</sup> cell count (Fig. 3b and Supplementary Fig. 3b). Although possession of the HLA-B\*57 allele was associated with a striking reduction in disease progression rates independently of other covariates such as steady-state viral load and GRGs (Fig. 3a), there were no differences in the overall CD4<sup>+</sup> recovery according to HLA-B\*57 status (Fig. 3c). Thus, an overall protective effect of the HLA-B\*57 allele was not evident during HAART, and it was noteworthy that during the first 2–3 years after starting HAART possession of this allele was associated with impaired CD4<sup>+</sup> cell recovery (Fig. 3c).

In an alternative approach, we sought to capture the contribution of all HLA alleles, rather than focusing on specific alleles. This approach takes into account the prevailing viewpoint that a common class I allele confers a detrimental on HIV disease course, whereas a rare class I allele confers a beneficial effect<sup>31</sup>. We assigned each HIV<sup>+</sup> subject an HLA allele frequency score that provided a measure of whether they possessed a common (high score) or rare (low score) HLA allele. In agreement with the prevailing viewpoint, a high score was associated with greater early immune damage and lower cumulative CD4<sup>+</sup> counts than a low score (Fig. 3d). Despite this, after initiation of HAART, the CD4<sup>+</sup> time trends according to HLA allele frequency score were similar (Fig. 3e), indicating that HLA alleles that mitigate early immune damage (for example, lower initial loss of CD4<sup>+</sup> cells) do not necessarily confer salutary effects on CD4<sup>+</sup> recovery during HAART.

### **Effects of GRGs on CD4<sup>+</sup> recovery are independent of HLA**

Although a direct effect of HLA alleles on CD4<sup>+</sup> recovery was not detected, we could not exclude the possibility of gene-gene interactions between two biological systems (that is, *CCL3L1-CCR5* and HLA) that influence early immune damage and, consequently, these interactions affecting CD4<sup>+</sup> recovery. We therefore statistically adjusted the effects of GRGs on CD4<sup>+</sup> recovery for the three HLA alleles that affected HIV disease progression and for HLA allele frequency score. These analyses showed that after 2 years of HAART, the influence of GRG on CD4<sup>+</sup> recovery rates was independent of HLA alleles (Table 3).

## DISCUSSION

By studying subjects who received HAART during chronic infection, we demonstrate that although both *CCL3LI-CCR5* genotype and HLA alleles affect the extent of early immune damage and HIV disease progression independently of viral load (on the basis of results shown here and previously<sup>3</sup>), only variations in the *CCL3LI-CCR5* axis influence recovery of CD4<sup>+</sup> counts. This supports our hypothesis that particular genetic pathways (for example, the *CCL3LI-CCR5* axis) have dominant effects on both pathogenic and reparative processes during HIV disease.

We also investigated subjects who received HAART during acute or early infection with the following supposition: if there is no relationship between *CCL3LI-CCR5* genotype-dependent differences in early immune damage and the subsequent recovery of CD4<sup>+</sup> counts during HAART, then initiation of therapy during this early phase would overshadow the impact of *CCL3LI-CCR5* genotype on CD4<sup>+</sup> gains. However, this was not the case. Hence, collectively, the results of this study emphasize that certain *CCL3LI-CCR5* genotypes promote HIV pathogenesis despite viral load suppression by HAART during acute, early or chronic HIV disease; that is, they have an effect on pathogenesis that is independent of the viral load. This inference of a viral load-independent effect on HIV pathogenesis by *CCL3LI-CCR5* genotypes amplifies and extends the conclusions of our previous study<sup>3</sup>. However, given the known effects of *CCL3LI* dose and *CCR5* genotypes on viral entry, we cannot definitively exclude the possibility that they also influence CD4<sup>+</sup> reconstitution by affecting the viral infection-target cell component of the host-pathogen relationship, that is, by influencing the extent and rate of suppression of viral replication during HAART.

Our observations also underscore that HLA alleles that retard HIV disease course might have contrasting effects during HAART, conferring no effects on CD4<sup>+</sup> cell recovery or possibly, as observed for the HLA-B\*57 allele, a detrimental effect during the initial years of HAART. Recent findings by others concur with these inferences<sup>32,33</sup>. The reasons for why the beneficial effects of HLA alleles observed during untreated HIV disease are lost during HAART are unclear, but might relate in part to the following conceptual model. During untreated HIV disease, the magnitude of viral replication contributes to CD4<sup>+</sup> cell depletion, and, in this scenario, HLA-dependent restriction of viral replication helps mitigate CD4<sup>+</sup> cell loss. However, when viral replication is suppressed by HAART, the ability of a person's HLA alleles to recognize HIV peptides and mount an effective cytotoxic Tlymphocyte response may not be so consequential. Thus, in the latter scenario, HLA alleles that have protective effects on CD4<sup>+</sup> cells in the context of viremia during untreated HIV disease might have minimal beneficial effects on CD4<sup>+</sup> recovery during HAART. In contrast, by influencing viral replication, cell-mediated immunity, and processes that are independent of the viral load and cell-mediated immunity<sup>3</sup>, *CCL3LI-CCR5* genotypes may affect the CD4<sup>+</sup> cell loss observed before initiation of HAART and the CD4<sup>+</sup> cell gain during HAART.

The differing effects of low and moderate or high *CCL3LI-CCR5* GRGs on reconstitution of CD4<sup>+</sup> counts were evident in subjects in whom HAART-induced suppression of viral replication was sufficient to sustain robust gains in CD4<sup>+</sup> counts and was also present after accounting for potential gene-gene interactions with HLA alleles and other parameters that influence HIV disease course (for example, age, pre-HAART CD4<sup>+</sup> cell count or viral load). These divergent effects suggest that a moderate or high *CCL3LI-CCR5* GRG might serve as a genetic basis for the phenotype of incomplete immunologic reconstitution despite sustained viral suppression, whereas a low GRG might underlie the converse phenotype of partial immunologic-clinical stability or recovery in the face of incomplete or transient viral suppression. The clinical importance of identifying genetic factors that contribute to the former phenotype is underscored by the finding that impaired immune reconstitution despite HAART-

induced viral load suppression is associated with an increased risk of both AIDS-related and non-AIDS-related morbidity and mortality<sup>10,11,34-38</sup>.

The genotype-phenotype associations reported here are robust because of several features of the study. First, it includes a large number of HIV-positive subjects representative of the general US population. Second, in our study cohorts, confounding factors such as unequal access to health care and loss of subjects to follow-up were minimized. Third, we found agreement between *CCL3LI-CCR5* genotype and phenotype at multiple levels: similar genotype-dependent effects were observed whether HAART was given during acute, early or chronic infection; genotypes linked with robust recovery of CD4<sup>+</sup> cell numbers also associate with improved CD4<sup>+</sup> cell function during HAART<sup>3,27</sup>; and there is concordance in the genotypes that mitigate or promote CD4<sup>+</sup> depletion during the pre-HAART era and those genotypes that associate with robust or impaired CD4<sup>+</sup> recovery during the HAART era. Substantiating the last point, previous studies have found that an HIV disease-retarding genotype (heterozygosity for the *CCR5-Δ32*-containing HHG\*2 haplotype<sup>25</sup>) has beneficial effects on CD4<sup>+</sup> recovery, whereas an HIV disease-accelerating *CCR5* genotype (homozygosity for *CCR5* HHE/HHE<sup>25</sup>) has detrimental effects on CD4<sup>+</sup> recovery during HAART<sup>39-42</sup>. However there is an important caveat: as the findings in these reports and our study are derived from cohorts from the US, the genotype-phenotype relationships observed herein might pertain specifically to clade B HIV-1–host interactions, and thus it is possible that similar relationships might not be detected within the context of other host–non-B clade HIV settings.

Although *CCL3LI* and *CCR5* genotype affect CD4<sup>+</sup> recovery across a spectrum of both high and low CD4<sup>+</sup> counts at which HAART was initiated, it is remarkable that the greatest impact of these genotypes appears to be on sustaining the initial CD4<sup>+</sup> gains after 2 years of HAART, especially in those individuals initiating HAART at <350 CD4<sup>+</sup> cells/mm<sup>3</sup>. This is noteworthy because the GRG-dependent inflections in CD4<sup>+</sup> gains between 2 and 3 years could be missed in clinical trials, which are typically of a shorter duration. The prominent beneficial effects associated with a low *CCL3LI-CCR5* GRG on CD4<sup>+</sup> recovery in HIV-positive individuals initiating HAART at <350 CD4<sup>+</sup> cells/mm<sup>3</sup> suggests that the magnitude of immune reconstitution may be more sensitive to the genetically determined (that is, *CCL3LI-CCR5* genetic status) immunologic, rather than virologic, status of the infected host.

On the basis of these findings and those from our previous studies<sup>3,25,26</sup>, we propose that very soon after exposure to virus, a person's *CCL3LI-CCR5* genotype serves as a rheostat, setting the balance between pathogenic and reparative processes that contribute, in part, to the observed wide heterogeneity in the dynamics of CD4<sup>+</sup> cell counts and function not just during untreated HIV infection, but also during therapy. As this phenotypic pathway is genetically programmed during early stages of the disease<sup>3,25,26</sup> and can persist during HAART, *CCL3LI-CCR5* genotype-dependent HIV pathogenesis may continue in spite of therapy, resulting in impaired immune reconstitution. Thus, the current debate about when to initiate antiretroviral therapy<sup>43</sup> might need to be redirected toward first assessing for whom therapy should be considered, on the basis of the host genotype. For HIV-positive individuals with ≥350 CD4<sup>+</sup> cells/mm<sup>3</sup>, the current treatment guidelines<sup>44</sup> recommend 'consideration' of therapy, and these guidelines provide some general criteria regarding which people might benefit the most from earlier initiation of HAART. We suggest that, given the poor immunologic response in individuals with a moderate or high *CCL3LI-CCR5* GRG, this genetic parameter may also be a useful guide in deciding whether to initiate therapy in those with ≥350 CD4<sup>+</sup> cells/mm<sup>3</sup>. Furthermore, our findings suggest that, for subjects with a moderate or high GRG in whom therapy is delayed, in addition to HAART, immunologically based adjunctive therapies (for example, recombinant interleukin-2 or recombinant interleukin-7)<sup>45</sup> might need to be developed to diminish the risk of impaired durability of CD4<sup>+</sup> responses. The observation that recovery of CD4<sup>+</sup> counts during HAART is more

sensitive to the copy number of *CCL3L1* than to *CCR5* genotype raises the possibility that *CCL3L1* mimetics might be efficacious in promoting immune reconstitution. Finally, it may be useful to evaluate whether a subject's *CCL3L1* gene dose and *CCR5* genotype will affect their response to *CCR5* blockers used in HIV therapy.

## METHODS

### Study cohorts

The characteristics of the WHMC cohort, a prospective observational cohort that is a component of the United States Military's Tri-Service AIDS Clinical Consortium Natural History Study, and the AIEDRP cohort are described previously<sup>3,27,46</sup> and in the Supplementary Methods. Subjects within the WHMC and AIEDRP cohorts received HAART during chronic and acute-early infection, respectively. Subjects gave informed consent for these studies, and the study was approved by the Institutional Review Boards at the participating institutions (WHMC, University of California at San Diego and University of California at San Francisco).

### Genotyping

Methods used to assess *CCL3L1* copy number, *CCR5* genotype and assign a low, moderate and high *CCL3L1-CCR5* GRG are described previously<sup>3,27</sup> and in the Supplementary Methods. Briefly, *CCL3L1*<sup>low</sup> indicates a *CCL3L1* gene copy below the population-specific median and *CCL3L1*<sup>high</sup> indicates a gene copy number greater than or equal to the population-specific median<sup>27</sup>. The average gene dose in European Americans is two copies of *CCL3L1*, whereas it is three or four copies in HIV-positive and HIV-negative African Americans, respectively<sup>27</sup>. *CCR5*<sup>det</sup> indicates a genotype associated with a rapid rate of disease progression to AIDS and death, and *CCR5*<sup>nondet</sup> indicates all other *CCR5* genotypes<sup>27</sup>. A low GRG is defined by possession of a high *CCL3L1* gene dose plus nondetrimental *CCR5* genotype (*CCR5*<sup>nondet</sup>), a moderate GRG is defined by a low *CCL3L1* gene dose and nondetrimental *CCR5* genotype or a high *CCL3L1* gene dose and detrimental *CCR5* genotype (*CCR5*<sup>det</sup>), and a high GRG is defined by a low *CCL3L1* dose plus detrimental *CCR5* genotype<sup>27</sup> (schematic shown in Fig. 1d). The prevalence of a high and moderate GRG was 8% and 42%, respectively, in the WHMC HIV<sup>+</sup> cohort<sup>3,27</sup>. Methods for genotyping of HLA alleles are described in the Supplementary Methods. Although several HLA allele scoring systems have been described<sup>33,47-50</sup>, we adapted one<sup>33</sup>, and the details are provided in the Supplementary Methods.

### Outcomes

The primary outcome variables were changes in CD4<sup>+</sup> T cell counts, plasma HIV-1 viral loads and time trends for these two laboratory parameters. Viral load suppression was defined as attainment of at least two viral load measurements that were below the assay-specific limits of detection on HAART. Sustained viral load suppression was defined as the attainment of at least two viral load measurements that were below the level of detection, and all subsequent viral load measurements were less than 500 copies/ml. Transient viral load suppression was defined as the attainment of at least two viral load measurements below the level of detection and at least one subsequent viral load measurement that exceeded 500 copies/ml on HAART. Other endpoints analyzed were parameters of early immune damage reflected by the baseline CD4<sup>+</sup> T cell count, steady-state viral load and DTH skin test responses, as well as a parameter designated as the cumulative CD4<sup>+</sup> T cell count, which reflects the extent of CD4<sup>+</sup> T cell count changes during HIV disease; these parameters are described previously<sup>3</sup> and in the Supplementary Methods.



## Statistical analyses

Detailed statistical methods and the number of subjects and CD4<sup>+</sup> T cell count or viral load measurements for the data shown herein are in the Supplementary Methods and Supplementary Note. The basis for conducting the linear regressions in two segments, that is, during the first 2 years after initiation of HAART and in the years thereafter, are also described in the Supplementary Methods.

## Supplemental Material

Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGMENTS


The people and funding agencies that made this work possible are listed in the Acknowledgements section in the Supplementary information online.

## References

1. Douek DC, Picker LJ, Koup RA. T cell dynamics in HIV-1 infection. *Annu. Rev. Immunol* 2003;21:265–304. [PubMed: 12524385]
2. Rodriguez B, et al. Predictive value of plasma HIV RNA level on rate of CD4 T-cell decline in untreated HIV infection. *J. Am. Med. Assoc* 2006;296:1498–1506.
3. Dolan MJ, et al. CCL3L1 and CCR5 influence cell-mediated immunity and affect HIV-AIDS pathogenesis via viral entry-independent mechanisms. *Nat. Immunol* 2007;8:1324–1336. [PubMed: 17952079]
4. Moore RD, Keruly JC. CD4<sup>+</sup> cell count 6 years after commencement of highly active antiretroviral therapy in persons with sustained virologic suppression. *Clin. Infect. Dis* 2007;44:441–446. [PubMed: 17205456]
5. Grabar S, et al. Clinical outcome of patients with HIV-1 infection according to immunologic and virologic response after 6 months of highly active antiretroviral therapy. *Ann. Intern. Med* 2000;133:401–410. [PubMed: 10975957]
6. Tarwater PM, et al. Increase and plateau of CD4 T-cell counts in the 3(1/2) years after initiation of potent antiretroviral therapy. *J. Acquir. Immune Defic. Syndr* 2001;27:168–175. [PubMed: 11404539]
7. Hunt PW, et al. Continued CD4 cell count increases in HIV-infected adults experiencing 4 years of viral suppression on antiretroviral therapy. *AIDS* 2003;17:1907–1915. [PubMed: 12960823]
8. Moore DM, et al. Discordant immunologic and virologic responses to highly active antiretroviral therapy are associated with increased mortality and poor adherence to therapy. *J. Acquir. Immune Defic. Syndr* 2005;40:288–293. [PubMed: 16249702]
9. Valdez H, et al. Limited immune restoration after 3 years' suppression of HIV-1 replication in patients with moderately advanced disease. *AIDS* 2002;16:1859–1866. [PubMed: 12351945]
10. Kaufmann GR, et al. Characteristics, determinants, and clinical relevance of CD4 T cell recovery to <500 cells/μl in HIV type 1-infected individuals receiving potent antiretroviral therapy. *Clin. Infect. Dis* 2005;41:361–372. [PubMed: 16007534]
11. Podlekareva D, et al. Factors associated with the development of opportunistic infections in HIV-1-infected adults with high CD4<sup>+</sup> cell counts: a EuroSIDA study. *J. Infect. Dis* 2006;194:633–641. [PubMed: 16897662]
12. Mehandru S, et al. Lack of mucosal immune reconstitution during prolonged treatment of acute and early HIV-1 infection. *PLoS Med* 2006;3:e484. [PubMed: 17147468]
13. Deeks SG, Barbour JD, Martin JN, Swanson MS, Grant RM. Sustained CD4<sup>+</sup> T cell response after virologic failure of protease inhibitor-based regimens in patients with human immunodeficiency virus infection. *J. Infect. Dis* 2000;181:946–953. [PubMed: 10720517]
14. Ledergerber B, et al. Predictors of trend in CD4-positive T-cell count and mortality among HIV-1-infected individuals with virological failure to all three antiretroviral drug classes. *Lancet* 2004;364:51–62. [PubMed: 15234856]


15. Deeks SG. Durable HIV treatment benefit despite low-level viremia: reassessing definitions of success or failure. *J. Am. Med. Assoc* 2001;286:224–226.
16. Kaufmann GR, et al. CD4 T-lymphocyte recovery in individuals with advanced HIV-1 infection receiving potent antiretroviral therapy for 4 years: the Swiss HIV Cohort Study. *Arch. Intern. Med* 2003;163:2187–2195. [PubMed: 14557216]
17. Garcia F, et al. Long-term CD4<sup>+</sup> T-cell response to highly active antiretroviral therapy according to baseline CD4<sup>+</sup> T-cell count. *J. Acquir. Immune Defic. Syndr* 2004;36:702–713. [PubMed: 15167289]
18. Tenorio AR, et al. HIV-1–infected antiretroviral-treated patients with prolonged partial viral suppression: clinical, virologic, and immunologic course. *J. Acquir. Immune Defic. Syndr* 2003;34:491–496. [PubMed: 14657759]
19. Anastos K, et al. The prognostic importance of changes in CD4<sup>+</sup> cell count and HIV-1 RNA level in women after initiating highly active antiretroviral therapy. *Ann. Intern. Med* 2004;140:256–264. [PubMed: 14970148]
20. Grabar S, et al. Response to highly active antiretroviral therapy at 6 months and long-term disease progression in HIV-1 infection. *J. Acquir. Immune Defic. Syndr* 2005;39:284–292. [PubMed: 15980687]
21. Zhang ZQ, et al. The impact of early immune destruction on the kinetics of postacute viral replication in rhesus monkey infected with the simian-human immunodeficiency virus 89.6P. *Virology* 2004;320:75–84. [PubMed: 15003864]
22. Mellors JW, et al. Plasma viral load and CD4<sup>+</sup> lymphocytes as prognostic markers of HIV-1 infection. *Ann. Intern. Med* 1997;126:946–954. [PubMed: 9182471]
23. Lifson JD, et al. Containment of simian immunodeficiency virus infection: cellular immune responses and protection from rechallenge following transient postinoculation antiretroviral treatment. *J. Virol* 2000;74:2584–2593. [PubMed: 10684272]
24. Nibbs RJ, Yang J, Landau NR, Mao JH, Graham GJ. LD78β, a non-allelic variant of human MIP-1α (LD78α), has enhanced receptor interactions and potent HIV suppressive activity. *J. Biol. Chem* 1999;274:17478–17483. [PubMed: 10364178]
25. Gonzalez E, et al. Race-specific HIV-1 disease–modifying effects associated with CCR5 haplotypes. *Proc. Natl. Acad. Sci. USA* 1999;96:12004–12009. [PubMed: 10518566]
26. Mangano A, et al. Concordance between the CC chemokine receptor 5 genetic determinants that alter risks of transmission and disease progression in children exposed perinatally to human immunodeficiency virus. *J. Infect. Dis* 2001;183:1574–1585. [PubMed: 11335892]
27. Gonzalez E, et al. The influence of *CCL3L1* gene–containing segmental duplications on HIV-1/AIDS susceptibility. *Science* 2005;307:1434–1440. [PubMed: 15637236]
28. Martin MP, et al. Genetic acceleration of AIDS progression by a promoter variant of CCR5. *Science* 1998;282:1907–1911. [PubMed: 9836644]
29. Kaslow RA, Dorak T, Tang JJ. Influence of host genetic variation on susceptibility to HIV type 1 infection. *J. Infect. Dis* 2005;191(Suppl 1):S68–S77. [PubMed: 15630678]
30. Carrington M, O'Brien SJ. The influence of HLA genotype on AIDS. *Annu. Rev. Med* 2003;54:535–551. [PubMed: 12525683]
31. Nguyen L, et al. Frequent human leukocyte antigen class I alleles are associated with higher viral load among HIV type 1 seroconverters in Thailand. *J. Acquir. Immune Defic. Syndr* 2004;37:1318–1323. [PubMed: 15385741]
32. Rauch A, et al. HLA-Bw4 homozygosity is associated with an impaired CD4-T-cell recovery after initiation of antiretroviral therapy. *Clin. Infect. Dis.* in the press
33. Brumme ZL, et al. Effects of human leukocyte antigen class I genetic parameters on clinical outcomes and survival after initiation of highly active antiretroviral therapy. *J. Infect. Dis* 2007;195:1694–1704. [PubMed: 17471440]
34. Moore DM, et al. Disease progression in patients with virological suppression in response to HAART is associated with the degree of immunological response. *AIDS* 2006;20:371–377. [PubMed: 16439870]
35. Weber R, et al. Liver-related deaths in persons infected with the human immunodeficiency virus: the D:A:D study. *Arch. Intern. Med* 2006;166:1632–1641. [PubMed: 16908797]

36. Gutierrez F, et al. Clinical Outcome of HIV-infected patients with sustained virologic response to antiretroviral therapy: long-term follow-up of a multicenter cohort. *PLoS ONE* 2006;1:e89. [PubMed: 17183720]
37. El-Sadr WM, et al. CD4<sup>+</sup> count-guided interruption of antiretroviral treatment. *N. Engl. J. Med* 2006;355:2283–2296. [PubMed: 17135583]
38. Lewden C, et al. HIV-infected adults with a CD4 cell count greater than 500 cells/mm<sup>3</sup> on long-term combination antiretroviral therapy reach same mortality rates as the general population. *J. Acquir. Immune Defic. Syndr* 2007;46:72–77. [PubMed: 17621240]
39. Valdez H, Purvis SF, Lederman MM, Fillingame M, Zimmerman PA. Association of the CCR5Δ32 mutation with improved response to antiretroviral therapy. *J. Am. Med. Assoc* 1999;282:734.
40. Yamashita TE, et al. Immunologic and virologic response to highly active antiretroviral therapy in the Multicenter AIDS Cohort Study. *AIDS* 2001;15:735–746. [PubMed: 11371688]
41. Kasten S, et al. Positive influence of the Δ32CCR5 allele on response to highly active antiretroviral therapy (HAART) in HIV-1-infected patients. *Eur. J. Med. Res* 2000;5:323–328. [PubMed: 10958764]
42. O'Brien TR, et al. Effect of chemokine receptor gene polymorphisms on the response to potent antiretroviral therapy. *AIDS* 2000;14:821–826. [PubMed: 10839590]
43. Holmberg SD, Palella FJ Jr, Lichtenstein KA, Havlir DV. The case for earlier treatment of HIV infection. *Clin. Infect. Dis* 2004;39:1699–1704. [PubMed: 15578373]
44. Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents. Department of Health and Human Services; Jan 29. 2008 Panel on Antiretroviral Guidelines for Adult and Adolescents; p. 1-128. <<http://www.aidsinfo.nih.gov/ContentFiles/AdultandAdolescentGl.pdf>>, accessed 29 February 2008
45. Pahwa S. Role of common γ chain-utilizing cytokines for immune reconstitution in HIV infection. *Immunol. Res* 2007;38:373–386. [PubMed: 17917047]
46. Hecht FM, et al. A multicenter observational study of the potential benefits of initiating combination antiretroviral therapy during acute HIV infection. *J. Infect. Dis* 2006;194:725–733. [PubMed: 16941337]
47. Saah AJ, et al. Association of HLA profiles with early plasma viral load, CD4<sup>+</sup> cell count and rate of progression to AIDS following acute HIV-1 infection. Multicenter AIDS Cohort Study. *AIDS* 1998;12:2107–2113. [PubMed: 9833851]
48. Kaslow RA, et al. Influence of combinations of human major histocompatibility complex genes on the course of HIV-1 infection. *Nat. Med* 1996;2:405–411. [PubMed: 8597949]
49. Tang J, et al. Host genetic profiles predict virological and immunological control of HIV-1 infection in adolescents. *AIDS* 2002;16:2275–2284. [PubMed: 12441799]
50. Carrington M, Nelson G, O'Brien SJ. Considering genetic profiles in functional studies of immune responsiveness to HIV-1. *Immunol. Lett* 2001;79:131–140. [PubMed: 11595300]




**Figure 1.** Influence of *CCL3L1-CCR5* genotypes on CD4<sup>+</sup> T cell dynamics in the WHMC HIV<sup>+</sup> cohort. (a–d) Trajectories of CD4<sup>+</sup> counts in the WHMC cohort across calendar years of cohort membership computed before (a) and after accounting for *CCL3L1* dose (b), *CCR5* genotype (c) and *CCL3L1-CCR5* GRGs (d). Dotted lines in a demarcate the indicated therapy eras, and mono and dual reflect therapy with one or two antiretroviral agents that were prevalent during the pre-HAART era. The time trends shown were estimated by nonlinear generalized estimating equations (GEE) modeling of CD4<sup>+</sup> counts and depicted as mean (thick lines) and 95% confidence bands (shaded regions). The genetic components that comprise *CCL3L1-CCR5* GRGs are shown above d. (e) Trajectories of CD4<sup>+</sup> counts (by nonlinear GEE) from time of initiation of HAART according to (from left to right) *CCL3L1* dose, *CCR5* genotype and *CCL3L1-CCR5* GRG status. (f) Trajectories of CD4<sup>+</sup> counts during HAART according to *CCL3L1* dose, *CCR5* genotype and GRG status during the first 2 years after initiation of HAART and in the years thereafter (modeled by linear GEE in two segments demarcated by the dashed lines). *P* values indicate differences in rate of change in CD4<sup>+</sup> counts for the indicated groups (derived by the Student's *t*-test) during the first 2 years after initiation of HAART and in the years thereafter. Rate of change in CD4<sup>+</sup> cell counts per month (with s.e.m. in parentheses) during the first 2 years of HAART in those with a high or low *CCL3L1* dose, nondetrimental or detrimental *CCR5* genotype, and low, moderate or high GRG were 5.23

(0.72), 4.73 (1.12), 5.29 (0.57), 4.18 (2.11), 6.36 (0.57), 6.20 (0.98), and 3.12 (2.86), respectively. Rate of change in CD4<sup>+</sup> cell counts/month (s.e.m.) after 2 years of HAART in those with a high or low *CCL3L1* dose, nondetrimental or detrimental *CCR5* genotype, and low, moderate or high GRG were 1.27 (0.23), -0.04 (0.39), 1.20 (0.21), -0.05 (0.61), 0.96 (0.17), -0.09 (0.17), and -3.54 (1.22), respectively. The color code for plots shown in **e** and **f** are shown above **b-d**.



**Figure 2.** Influence of *CCL3L1-CCR5* on CD4<sup>+</sup> recovery in subjects initiating HAART during acute or early infection (AIEDRP cohorts, **a-c**) and impact of the timing of HAART on effects of *CCL3L1-CCR5* on CD4<sup>+</sup> recovery in the WHMC cohort (**d,e**). **(a)** Loess curves depict trajectories of CD4<sup>+</sup> counts in those who initiated HAART during acute infection and who attained viral load suppression according to GRG status (top). Difference in average CD4<sup>+</sup> counts by GRG in the first year of HAART is also shown (bottom). **(b)** Loess curves showing CD4<sup>+</sup> count changes in subjects according to those who did or did not achieve viral load suppression during early infection (top) and according to GRG status in those who achieved HAART-induced viral load suppression during early infection (bottom). For those initiating HAART during early infection, rate of change in CD4<sup>+</sup> cell counts/month (s.e.m.) in those with low or moderate-high GRG were 21.62 (1.09) and 12.52 (1.59), respectively. **(c)** Loess curves for CD4<sup>+</sup> count changes according to *CCL3L1* dose (top) or *CCR5* genotype (bottom) in subjects with HAART-induced viral load suppression during early infection. Rate of change in CD4<sup>+</sup> counts/month (s.e.m.) in those with a high or low *CCL3L1* dose and nondetrimental or detrimental *CCR5* genotypes were 20.30 (1.02), 12.71 (1.79), 20.18 (0.97), and 12.56 (2.63), respectively. **(d)** Trajectories of CD4<sup>+</sup> counts (mean (bold lines) and 95% confidence bands by nonlinear GEE) according to GRG status for subjects in WHMC who initiated HAART at <350 (top) or ≥350 (bottom) CD4<sup>+</sup> cells/mm<sup>3</sup>. The data are for all subjects who received HAART. **(e)** Trajectories of CD4<sup>+</sup> counts according to *CCL3L1* dose in subjects who were therapy-naïve and initiated HAART at <350 (top) or ≥350 CD4<sup>+</sup> cells/mm<sup>3</sup> (bottom). The HAART-initiating CD4<sup>+</sup> T cell count is the average CD4<sup>+</sup> T cell count during the year preceding initiation of therapy. Where indicated, the moderate and high GRGs were combined into a single category (brown).



**Figure 3.** Influence of HLA alleles on early events and CD4<sup>+</sup> T cell dynamics in subjects from the WHMC HIV<sup>+</sup> cohort. **(a)** Independent disease-modifying effects of HLA alleles. The plot shows the relative hazards (RH, diamonds) and 95% confidence interval (CI, error bars) for rates of progression to AIDS for the variables listed along the x-axis. Results are derived from the final model using stepwise multivariate Cox proportional hazards regression analyses for time to AIDS (1987 Centers for Disease Control criteria) and are based on data from 719 European- and African-American subjects from the WHMC HIV<sup>+</sup> cohort. bCD4, baseline CD4<sup>+</sup> T cell count; VL, steady-state viral load; nCD4, nadir CD4<sup>+</sup> T cell count during HIV disease course; %CD4, percentage of baseline CD4<sup>+</sup> T cell counts; these parameters as well as the therapy era and DTH skin test reactions are as described previously<sup>3</sup>. The RH (95% CI) for bCD4, nCD4 and %CD4 are 1.00 (0.99 – 1.00), 0.99 (0.99 – 0.99) and 0.97 (0.96 – 0.99), respectively. **(b)** Association of the HLA-B\*57 allele (pink, present; black, absent) with parameters that reflect early immune damage (steady-state VL, baseline CD4, DTH responses) and cumulative CD4<sup>+</sup> T cell counts. Diamonds and error bars represent the mean and 95% CI, and the P values were obtained using the Student's *t*-test. **(c)** Recovery of CD4<sup>+</sup> T cell counts after initiation of HAART on the basis of possession of a HLA-B\*57 allele (by nonlinear GEE). **(d)** Association of HLA class I allele frequency score groups (low score representing rare (orange) and high score representing common (green) class I HLA alleles) with the indicated parameters. **(e)** Recovery of CD4<sup>+</sup> T cell counts after initiation of HAART on the basis of the HLA class I allele frequency score groups representative of rare and common HLA class I alleles (by nonlinear GEE).

**Table 1**  
Influence of variations in *CCL3L1* and *CCR5* on CD4<sup>+</sup> recovery rates

Genotype	Overall	Viral load suppression	
		Suppressed	Transient
<i>CCL3L1</i>			
<i>CCL3L1</i> <sup>high</sup>	2.80 (0.14)	3.43 (0.17)	5.41 (0.32)
<i>CCL3L1</i> <sup>low</sup>	0.71 (0.24)	0.73 (0.24)	-0.24 (0.29)
<i>P</i> <sub>low versus high</sub>	$6.9 \times 10^{-13}$	$2.0 \times 10^{-17}$	$8.1 \times 10^{-13}$
<i>CCR5</i>			
<i>CCR5</i> <sup>nondefect</sup>	2.38 (0.37)	2.76 (0.14)	5.18 (0.98)
<i>CCR5</i> <sup>defect</sup>	1.77 (0.13)	2.29 (0.48)	4.40 (0.26)
<i>P</i> <sub>defect versus nondefect</sub>	0.1229	0.3495	0.1786
GRG status			
Low	2.89 (0.13)	3.45 (0.15)	5.08 (0.28)
Moderate	1.42 (0.20)	1.67 (0.22)	3.68 (0.35)
High	-0.85 (0.76)	-0.77 (0.70)	-1.74 (0.69)
<i>P</i> <sub>low versus moderate</sub>	$1.7 \times 10^{-9}$	$2.8 \times 10^{-10}$	$4.4 \times 10^{-9}$
<i>P</i> <sub>moderate versus high</sub>	$4.3 \times 10^{-3}$	$1.2 \times 10^{-3}$	$2.3 \times 10^{-3}$
<i>P</i> <sub>low versus high</sub>	$3.1 \times 10^{-6}$	$6.5 \times 10^{-8}$	$7.0 \times 10^{-8}$

Data are from subjects from the WHMC HIV<sup>+</sup> cohort who received HAART during chronic infection. Values indicate the mean (s.e.m.) rate of change of CD4<sup>+</sup> cells per month after initiation of HAART, as estimated by regression coefficients computed by linear GEE. A negative number indicates a decrease in the rate of CD4<sup>+</sup> cell counts. Data for analyses conducted by using square-root-transformed CD4<sup>+</sup> T cell counts, as well as segmented regressions to determine effects of genotypes on CD4<sup>+</sup> recovery during and after the first 2 years of HAART, are shown in Supplementary Table 1 online.



**Table 2**  
GRG effects on CD4<sup>+</sup> recovery in those initiating HAARTat <350 CD4<sup>+</sup> cells/mm<sup>3</sup>

Pre-HAART CD4 count (cells/mm <sup>3</sup> ) <sup>a</sup>	GRG status			Low versus moderate GRG CD4 difference (s.e.m.), P
	Low	Moderate-high	Moderate	
<350	3.23 (0.21) <sup>b</sup>	0.83 (0.22)	0.80 (0.24)	2.40 (0.29), 5.8 × 10 <sup>-13</sup>
<300	3.52 (0.21)	0.28 (0.26)	0.23 (0.28)	3.24 (0.33), 5.0 × 10 <sup>-16</sup>
<250	3.47 (0.25)	-0.26 (0.27)	-0.33 (0.30)	3.73 (0.37), 1.9 × 10 <sup>-15</sup>
<200	4.51 (0.24)	-0.23 (0.33)	-0.22 (0.33)	4.74 (0.41), 3.6 × 10 <sup>-16</sup>
<150	4.60 (0.30)	-0.16 (0.37)	-0.16 (0.37)	4.76 (0.47), 1.1 × 10 <sup>-11</sup>
<100	4.82 (0.34)	-0.06 (0.41)	-0.06 (0.41)	4.88 (0.54), 1.5 × 10 <sup>-9</sup>
<50	4.59 (0.52)	-0.47 (0.90)	-0.47 (0.90)	5.06 (1.04), 2.0 × 10 <sup>-4</sup>

Data are for subjects from the WHMC HIV<sup>+</sup> cohort who received HAART during chronic infection.

<sup>a</sup> Pre-HAART CD4<sup>+</sup> T cell count was the average of all CD4<sup>+</sup> T cell counts during the 3 months before initiation of HAART.

<sup>b</sup> Values are defined as in Table 1. Similar analyses were also conducted by using square-root-transformed CD4<sup>+</sup> T cell counts (Supplementary Fig. 1f).

**Table 3**  
Independent effects of *CCL3L1-CCR5* GRGs on CD4<sup>+</sup> recovery

Pre-HAART CD4	Low GRG	Moderate-high GRG	P
Model 1: unadjusted			
During first 2 years of HAART			
≥ 350	7.07 (1.00) <sup>d</sup>	6.87 (1.44)	0.9093
< 350	6.05 (0.76)	5.13 (0.89)	0.4332
After 2 years of HAART			
≥ 350	0.65 (0.40)	-0.65 (0.62)	0.0808
< 350	1.90 (0.35)	0.12 (0.34)	0.0004
Model 2: adjusted for covariates <sup>b</sup>			
During first 2 years of HAART			
≥ 350	5.16 (1.19)	4.58 (1.49)	0.7611
< 350	6.03 (0.69)	4.49 (0.62)	0.1008
After 2 years of HAART			
≥ 350	1.01 (0.50)	-0.45 (0.59)	0.0626
< 350	2.07 (0.40)	0.11 (0.38)	0.0007
Model 3: adjusted for covariates <sup>b</sup> and HLA-A*68, HLA-B*57 and HLA-C*16 alleles			
During first 2 years of HAART			
≥ 350	4.77 (1.21)	5.15 (1.71)	0.8565
< 350	6.36 (0.76)	4.97 (0.33)	0.0971
After 2 years of HAART			
≥ 350	1.02 (0.48)	-1.55 (0.68)	0.0028
< 350	2.44 (0.46)	0.20 (0.36)	0.0003
Model 4: adjusted for covariates <sup>b</sup> and HLA allele frequency score			
During first 2 years of HAART			
≥ 350	4.72 (0.17)	5.15 (1.57)	0.8258
< 350	6.38 (0.78)	4.97 (0.43)	0.1206
After 2 years of HAART			
≥ 350	1.01 (0.49)	-1.56 (0.90)	0.0033
< 350	2.45 (0.46)	0.19 (0.36)	0.0003

Data are for subjects from the WHMC HIV<sup>+</sup> cohort and indicate the mean (s.e.m.) rate of change of CD4<sup>+</sup> cell counts per month after initiation of HAART, as estimated by regression coefficients computed by linear GEE.

<sup>a</sup> Values are defined as in Table 1.

<sup>b</sup> Covariates were calendar year of HAART initiation, age at HAART initiation, pre-HAART viral load, achievement of viral load suppression and prior receipt of antiretroviral therapy. Pre-HAART CD4<sup>+</sup> T cell count is as defined in Table 2. Similar results for models 1 and 2 were obtained in subjects who were naive for HAART (Supplementary Table 2 online).