

Use of Antihypotensive Therapies in Extremely Preterm Infants

AUTHORS: Beau Batton, MD,^{a,b} Lei Li, PhD,^c Nancy S. Newman, RN,^a Abhik Das, PhD,^d Kristi L. Watterberg, MD,^e Bradley A. Yoder, MD,^f Roger G. Faix, MD,^f Matthew M. Laughon, MD, MPH,^g Barbara J. Stoll, MD,^h Krisa P. Van Meurs, MD,ⁱ Waldemar A. Carlo, MD,^j Brenda B. Poindexter, MD, MS,^k Edward F. Bell, MD,^l Pablo J. Sánchez, MD,^m Richard A. Ehrenkranz, MD,ⁿ Ronald N. Goldberg, MD,^o Abbot R. Laptook, MD,^p Kathleen A. Kennedy, MD, MPH,^q Ivan D. Frantz III, MD,^r Seetha Shankaran, MD,^s Kurt Schibler, MD,^t Rosemary D. Higgins, MD,^u and Michele G. Walsh, MD, MS^a for the Eunice Kennedy Shriver National Institute of Child Health & Human Development Neonatal Research Network

^aDepartment of Pediatrics, Rainbow Babies & Children's Hospital, Case Western Reserve University, Cleveland, Ohio; ^bDepartment of Pediatrics, Southern Illinois University School of Medicine, Springfield, Illinois; ^cStatistics & Epidemiology Unit, RTI International, Research Triangle Park, North Carolina; ^dStatistics & Epidemiology Unit, RTI International, Rockville, Maryland; ^eUniversity of New Mexico Health Sciences Center, Albuquerque, New Mexico; ^fDivision of Neonatology, Department of Pediatrics, University of Utah School of Medicine, Salt Lake City, Utah; ^gDepartment of Pediatrics, University of North Carolina, Chapel Hill, North Carolina; ^hDepartment of Pediatrics, Emory University School of Medicine & Children's Healthcare of Atlanta, Atlanta, Georgia; ⁱDepartment of Pediatrics, Stanford University School of Medicine, Palo Alto, California; ^jDivision of Neonatology, University of Alabama at Birmingham, Birmingham, Alabama; ^kDepartment of Pediatrics, Indiana University School of Medicine, Indianapolis, Indiana; ^lDepartment of Pediatrics, University of Iowa, Iowa City, Iowa; ^mDepartment of Pediatrics, University of Texas Southwestern Medical Center, Dallas, Texas; ⁿDepartment of Pediatrics, Yale University School of Medicine, New Haven, Connecticut; ^oDepartment of Pediatrics, Duke University, Durham, North Carolina; ^pDepartment of Pediatrics, Women & Infants' Hospital, Brown University, Providence, Rhode Island; ^qDepartment of Pediatrics, University of Texas Medical School at Houston, Houston, Texas; ^rDivision of Newborn Medicine, Department of Pediatrics, Floating Hospital for Children, Tufts Medical Center, Boston, Massachusetts; ^sDepartment of Pediatrics, Wayne State University, Detroit, Michigan; ^tDepartment of Pediatrics, Cincinnati Children's Hospital Medical Center & University of Cincinnati College of Medicine, Cincinnati, Ohio; and ^uEunice Kennedy Shriver National Institute of Child Health & Human Development, National Institutes of Health, Bethesda, Maryland

KEY WORDS

extremely preterm infant, antihypotensive therapy, blood pressure, hypotension

ABBREVIATIONS

BP—blood pressure
GA—gestational age
IVH—intraventricular hemorrhage
MAP—mean arterial pressure
NRN—Neonatal Research Network
ROP—retinopathy of prematurity

(Continued on last page)



WHAT'S KNOWN ON THIS SUBJECT: Extremely preterm infants who receive antihypotensive therapy have worse outcomes than untreated infants. The reasons for this are not clear. High-quality randomized trials have not been performed to date because of logistical challenges, thereby necessitating alternative methods of investigation.



WHAT THIS STUDY ADDS: Antihypotensive therapy administration was not associated with improved in-hospital outcomes for any of the 15 definitions of low blood pressure investigated. Alternative methods of deciding who to treat are needed to maximize patient benefit and minimize harm.

abstract



OBJECTIVE: To investigate the relationships among blood pressure (BP) values, antihypotensive therapies, and in-hospital outcomes to identify a BP threshold below which antihypotensive therapies may be beneficial.

METHODS: Prospective observational study of infants 23^{0/7} to 26^{6/7} weeks' gestational age. Hourly BP values and antihypotensive therapy use in the first 24 hours were recorded. Low BP was investigated by using 15 definitions. Outcomes were examined by using regression analysis controlling for gestational age, the number of low BP values, and illness severity.

RESULTS: Of 367 infants enrolled, 203 (55%) received at least 1 antihypotensive therapy. Treated infants were more likely to have low BP by any definition ($P < .001$), but for the 15 definitions of low BP investigated, therapy was not prescribed to 3% to 49% of infants with low BP and, paradoxically, was administered to 28% to 41% of infants without low BP. Treated infants were more likely than untreated infants to develop severe retinopathy of prematurity (15% vs 8%, $P = .03$) or severe intraventricular hemorrhage (22% vs 11%, $P < .01$) and less likely to survive (67% vs 78%, $P = .02$). However, with regression analysis, there were no significant differences between groups in survival or in-hospital morbidity rates.

CONCLUSIONS: Factors other than BP contributed to the decision to use antihypotensive therapies. Infant outcomes were not improved with antihypotensive therapy for any of the 15 definitions of low BP investigated. *Pediatrics* 2013;131:e1865–e1873

The intrinsically abnormal condition of extremely preterm infants and their evolving complex physiology make it difficult to identify an acceptable range of blood pressure (BP) values in the immediate postnatal period. Currently, there is not a validated or widely accepted definition of hypotension in this population.¹ Difficulty with assessing organ perfusion, multiple disease processes, and unpredictable adaptation to extrauterine life also make deciding when to institute antihypotensive therapy challenging. Consequently, BP management is highly variable.^{2,3} The frequency of antihypotensive therapy use during the transition from intrauterine to postnatal life ranges from 29% to 82% across NICUs.^{3,4} Extremely preterm infants who receive these therapies have higher mortality and morbidity rates versus untreated gestational age (GA) matched infants,^{4–12} but it is unclear whether these worse outcomes are due to the cause of low BP, associated organ hypoperfusion, therapy for low BP, or a combination of these and other factors. Interpretation of BP management data is also complicated by methodologic limitations and confounding factors, both known and unknown.¹

Randomized placebo controlled trials investigating BP management in this population are lacking.¹ This is at least partly due to the many challenges of studying critical therapeutic interventions shortly after birth in such a vulnerable patient population. These include the inability to obtain timely and ethically valid informed consent, insufficient physician equipoise, identification of appropriate inclusion and exclusion criteria, and enrollment or selection biases.^{13–16}

Difficulties with randomized placebo controlled trials have led to alternative methods of investigating BP management in preterm infants.^{1,7–9,17}

However, these studies are limited by their retrospective design, small sample sizes, and inconsistent definition of low BP. Currently, there is no known BP threshold below which extremely preterm infants are at an increased risk for a poor outcome and little evidence that antihypotensive therapy improves outcomes for infants with low BP, however defined. The objectives of this study were to prospectively examine BP management in the first 24 hours for extremely preterm infants and to investigate the relationship between recorded BP values, antihypotensive therapy use, and in-hospital infant outcomes in an effort to identify a BP threshold below which therapy may be beneficial.

METHODS

This was a prospective observational study of inborn extremely preterm infants 23^{0/7} to 26^{6/7} weeks' GA born at 1 of 16 academic centers of the Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network (NRN). Infants were excluded if they died in the delivery room, had a major birth defect, or had intensive care withheld or withdrawn shortly after birth because the clinical care team felt the situation was hopeless. Research personnel used study-specific data forms to record hourly BP values and the administration of all antihypotensive therapies in the first 24 hours. BP values were obtained from an arterial catheter when available or by oscillography. Antihypotensive therapy was defined as receipt of a fluid bolus (at least 10 mL/kg of crystalloid), dopamine, dobutamine, epinephrine, hydrocortisone, vasopressin, or any blood product. Therapies were administered at the discretion of the clinical care team. Data were recorded on maternal demographics, the infant's

initial condition, and in-hospital outcomes. Standard definitions were applied for intraventricular hemorrhage (IVH),¹⁸ necrotizing enterocolitis,¹⁹ retinopathy of prematurity (ROP),²⁰ and bronchopulmonary dysplasia.²¹ This study was approved by the institutional review board at each participating center and was conducted either with informed parent consent for each infant before enrollment or with a waiver of consent from the center's institutional review board.

For all analyses, 15 definitions of low BP were investigated: 1, 2, or ≥ 3 systolic, diastolic, or mean BP values less than or equal to the fifth percentile; 1, 2, or ≥ 3 mean arterial pressure (MAP; in mm Hg) values less than or equal to the infant's GA equivalent (in weeks); and 1, 2, or ≥ 3 MAP values ≤ 25 mm Hg. Low BP values were not necessarily consecutive. At each postnatal hour, BP percentiles were constructed for different populations (all infants, only infants who did not receive therapy, and at each specific GA) by using 2 sets of BP values (all BP values versus only invasive BP values). The fifth percentile was numerically similar (within 2 mm Hg) for all populations analyzed, and results were statistically similar irrespective of which construct was used to define the fifth percentile. Only results from the entire study population are reported. Data analyses were performed at the NRN Data Coordinating Center (RTI International, Research Triangle Park, NC). Data were entered remotely by electronic submission and periodically reviewed for quality control. Statistical analysis was performed by using SAS 9.2 software (SAS Institute, Inc, Cary, NC). The *t* test was used for continuous variables, and the χ^2 test was used for categorical variables to compare differences between infants who did versus did not receive antihypotensive therapy. Associations between antihypotensive

therapy and in-hospital outcomes were examined by using logistic models with a random intercept for NRN center while controlling for GA, illness severity, and the number of low BP values. Illness severity was defined a priori as the cumulative number of the following: a positive initial blood culture, an initial hematocrit $\leq 30\%$, a 1-minute Apgar score of ≤ 3 , a pH < 7.10 in the first 24 hours, or need for delivery room chest compressions. Regression analysis was used to investigate the relationship between NRN center variability in the frequency of low BP values and the rate of antihypotensive therapy administration and to investigate the impact of NRN center variability in the rate of antihypotensive therapy administration on patient outcomes.

RESULTS

From July 21, 2010, to January 21, 2011, there were 367 infants enrolled, including 203 (55%) infants who received ≥ 1 antihypotensive therapy and 104 (28%) who received a vasoactive drug (Fig 1). Fifteen enrolled infants (8 untreated, 7 treated) died in the first

24 hours. Of the 203 treated infants, 135 (67%) received a fluid bolus, 102 (50%) received a blood product, and 92 (45%), 25 (12%), and 18 (9%) received dopamine, hydrocortisone, and dobutamine, respectively. One patient received vasopressin. Many infants given a vasoactive drug also received a fluid bolus or blood product. The frequency of antihypotensive therapy use was inversely related to birth weight ($P < .001$) and GA ($P < .001$); 130 (64%) infants with a birth weight ≤ 750 g received therapy versus 73 (45%) infants with a birth weight of 750 to 1000 g and 88 (66%) infants born at 23 to 24 weeks' GA received therapy versus 115 (49%) infants born at 25 to 26 weeks' GA.

There were 18 709 BP values recorded (6236 systolic, 6227 diastolic, and 6246 MAP). An umbilical arterial catheter was placed for 298 (81.2%) infants, invasive BP values were obtained from 306 (83.4%) infants, and 14 593 (78%) BP values were obtained from an arterial catheter. Antihypotensive therapies were administered at similar rates for infants with low BP based on

invasive versus noninvasive BP values. For the definitions of low BP investigated, the likelihood of treatment increased with the number of low BP values (Fig 2; $P < .001$); antihypotensive therapy was administered to 28% to 41% of infants without any low BP values, 51% to 77% of infants with 1 or 2 low BP values, and 71% to 97% of infants with ≥ 3 low BP values. Antihypotensive therapy use was lowest in infants who never had an MAP ≤ 25 mm Hg or the infant's GA equivalent (28% for both definitions of low BP). Three or more low BP values less than or equal to the fifth percentile of the systolic (83%), diastolic (93%), or MAP (97%) BP was associated with the highest antihypotensive therapy rates. Most treated infants had multiple low BP values. For example, 110 (54%) treated infants had ≥ 3 MAP values less than or equal to the infant's GA equivalent, including 81 (78%) infants who received a vasoactive drug.

Some, but not all, baseline characteristics differed between the 164 untreated infants and the 203 infants who received therapy (Table 1). In-hospital outcomes for these groups are presented in Table 2. The significant difference in the rate of survival to hospital discharge between the 2 groups was primarily due to a higher mortality rate after the first postnatal week for infants who received therapy (19% vs 11%) as the rate of survival to 1 week was not significantly different between groups. Logistic regression analysis with a random intercept for center controlling for GA, severity of illness, and the number of low BP values did not show any significant effects of antihypotensive therapy on rates of IVH or grade III/IV IVH, ROP, morbidity-free survival to hospital discharge (morbidity: necrotizing enterocolitis, ROP, bronchopulmonary dysplasia, grade III/IV IVH, or cystic periventricular




FIGURE 1
Extremely preterm infant study enrollment including classification by study group.




FIGURE 2 Percentage of infants with stated definition of low BP who received at ≥ 1 antihypertensive therapy.

leukomalacia), or survival through postnatal day 7 or until hospital discharge ($P > .05$ for all analyses, data not shown). Results were unchanged for each of the 15 definitions of low BP, when analysis was restricted to only infants with ≥ 1 low BP value, and when only invasive BP values were used (data not shown). There was not a definition of low BP identified for which infants who received therapy had improved rates of IVH, ROP, survival, or morbidity-free survival.

There was significant variation across NRN centers in the rate of antihypertensive

therapy administration ($P < .01$), incidence of low BP ($P < .01$ for all definitions of low BP), and rate of survival to hospital discharge ($P < .01$; Fig 3). With regression analysis, the center frequency of antihypertensive therapy administration was not significantly associated with the center incidence of low BP for any of the definitions of low BP investigated ($P > .05$ for each of the 15 definitions of low BP [range in P values: .29–.89]). In addition, for each definition of low BP investigated, the center frequency of antihypertensive therapy administration was not

significantly associated with the center incidences of IVH, grade III/IV IVH, other morbidities, or survival to hospital discharge ($P > .05$ for each of the 15 definitions of low BP [range in P values: .30–.84], data not shown).

DISCUSSION

In this prospective study of 367 extremely preterm infants, 55% received an antihypertensive therapy and 28% received a vasoactive drug. For each definition of low BP investigated, the likelihood of receiving therapy increased with the number of low BP values recorded. Antihypertensive therapy was often provided to infants without low BP and, paradoxically, not prescribed to infants with low BP. The observation that the NRN center rate of therapy was not significantly related to the center incidence of low BP is additional evidence that factors other than BP values contributed to the decision to provide antihypertensive therapy. Degree of prematurity and infant size appeared to influence this decision as the likelihood of receiving treatment was inversely related to GA and birth weight. In-hospital outcomes were not improved with therapy for any of the 15 definitions of low BP investigated, including those with ≥ 3 low BP values.

Results from the current study are consistent with previous investigations.^{3–5,10,17,22} In a study by Laughon et al, 82% of infants 23 to 27 weeks' GA received an antihypertensive therapy, including 34% who received a vasopressor.³ Although that study had some limitations that do not apply to the current one, the findings for each were similar in that the rate of therapy use varied across NICUs; smaller, less mature infants were more likely to receive treatment; and the decision to provide treatment was strongly influenced by which center provided care. Other studies compared outcomes

TABLE 1 Baseline Characteristics for Infants Who Did or Did Not Receive Antihypertensive Therapy in the First 24 Hours

Initial Characteristic	No Therapy ($n = 164$)	Administered Therapy ($n = 203$)	P Value
Received maternal antibiotics, n (%)	126 (77)	160 (79)	.58
Received (any) prenatal steroids, n (%)	147 (90)	188 (93)	.31
Vaginal delivery, n (%)	53 (32)	68 (33)	.84
Multiple gestation, n (%)	39 (24)	63 (31)	.12
Male gender, n (%)	73 (45)	100 (49)	.36
Birth wt, g, mean \pm SD	764 \pm 161	698 \pm 156	<.01
GA, weeks, mean \pm SD	25.5 \pm 0.9	25.1 \pm 1.1	<.01
1-min Ap \dot{g} ar ≤ 3 , n (%)	70 (43)	122 (60)	<.01
5-min Ap \dot{g} ar ≤ 5 , n (%)	47 (29)	80 (39)	.03
DR chest compressions, n (%)	13 (8)	25 (12)	.17
First hematocrit $< 30\%$, n (%)	8 (5)	38 (19)	<.01
Positive initial blood culture, n (%)	—	8 (4)	.01
(Any) pH < 7.10 , n (%)	5 (3)	27 (13)	<.01

DR, delivery room.

TABLE 2 In-hospital Outcomes for Infants Who Did or Did Not Receive Antihypotensive Therapy in the First 24 Hours

In-hospital Outcomes	No Therapy (n = 164)	Administered Therapy (n = 203)	P Value
Necrotizing enterocolitis requiring surgery, n (%)	11 (7)	16 (8)	.92
Bronchopulmonary dysplasia, n (%)	75 (46)	92 (45)	.26
Cystic periventricular leukomalacia, n (%)	7 (4)	11 (5)	.60
Intervention for ROP, n (%)	13 (8)	31 (15)	.03
(Any) IVH, n (%)	43 (26)	83 (41)	<.01
Grade 3/4 IVH, n (%)	18 (11)	44 (22)	<.01
Survived 24 h, n (%)	156 (95)	186 (92)	.19
Survived ≥1 week, n (%)	146 (89)	174 (86)	.20
Survived to hospital discharge, n (%)	128 (78)	137 (67)	.02
Morbidity-free survival, ^a n (%)	24 (15)	11 (5)	<.01

^a Morbidities: necrotizing enterocolitis, ROP, bronchopulmonary dysplasia, grade 3 or 4 IVH, or periventricular leukomalacia.

between infants with low BP who received an antihypotensive therapy and those who did not.^{4,17,22,23} In those studies, treatment was associated with similar or worse infant outcomes when compared with untreated infants, but no study identified a definition of low BP for which treatment improved outcomes. Neither the current study nor others support the routine use of any antihypotensive therapy for any of the current definitions of low BP in extremely preterm infants.^{1,3–5,11,17,22–25}


BP values were recorded hourly for all infants because the relationship between low BP and antihypotensive

therapy in previous studies has been influenced by a disproportionately higher number of BP values obtained from treated infants. Infants for whom intensive care was withheld or withdrawn in the first 24 hours (n = 22) were excluded so that analyses were not influenced by infants whose death was imminent irrespective of which therapies were administered or withheld. Analysis was limited to the first 24 hours because this is when the majority of infants who receive therapy are treated,^{3,4} and the evaluation, etiology, and management of low BP that occurs later may be different. Severity of illness based on factors considered

a priori as likely to affect the decision to administer therapy for low BP was controlled for with regression analysis because previous studies have suggested infants who receive antihypotensive therapy have worse outcomes because they are initially more ill.^{1,23,24,26} Multiple definitions of low BP were investigated because there is not an accepted definition of hypotension in this population. Although an MAP less than or equal to the infant's GA is the most common definition used,²⁴ it is not evidence based and was first suggested in a policy statement on the management of respiratory distress syndrome.²⁷ Additional strengths of this study are the prospective data collection by experienced research personnel using a uniform approach and analysis conducted by well-trained experts. Study limitations include the lack of information regarding some variables that may have contributed to the decision to administer antihypotensive therapies, variability in infant enrollment across NRN centers, and inconsistency in how noninvasive BP values were obtained.

This study was conducted because there is a lack of information to guide BP management in extremely preterm infants.^{1,2,26,28} Large placebo-controlled trials have not been completed to date, and the limitations of previous studies make interpretation of their results difficult.^{1–5,10,13,15,17,22,23} Two prospective interventional studies are in their early phases (Clinicaltrials.gov NCT01482559 and NCT01434251), but results are not expected from either before 2016. The prospective data collection, number of infants enrolled, and detailed data analysis plan of this study provide some of the strongest data to date regarding BP management in extremely preterm infants.

Despite the lack of evidence supporting the routine use of any antihypotensive

**FIGURE 3**

Center variation in the rate of antihypotensive therapy administration, frequency of low BP, and incidence of hospital survival.

therapy, there may be some benefit from such therapies for some extremely preterm infants. In the current study, 12% of infants were anemic at birth, and increasing the blood volume may be appropriate in such situations. In addition, 2% of infants had early-onset sepsis, and the high risk of death or neurodevelopmental impairment in such cases^{29,30} may outweigh the risks associated with antihypotensive therapy. Some extremely preterm infants with perceived low BP also have strong clinical or biochemical evidence of poor perfusion. These infants appear to be at greater risk of a poor outcome,^{4,24,25,31,32} and in this scenario, the benefits of therapy may outweigh the risks even though neither can be accurately predicted.^{1,2,22,33} However, infants with perceived low BP usually have adequate perfusion,^{23,25,34,35} and the benefit of treatment has not been established for these infants. In this situation, therapies to increase BP appear also to be used to try to prevent or improve undocumented organ hypoperfusion, primarily cerebral blood flow.^{34,35} This approach is challenging because BP may not correlate with perfusion^{34–37}; infants with low BP may have adequate cerebral blood flow,^{25,35,38,39} vasoactive drugs do not always increase cerebral perfusion^{25,34} and have not improved outcomes,⁴⁰ and treatment of low BP has been associated with similar or worse rates of intracranial abnormalities and impaired neurodevelopment versus matched untreated infants.^{3–5,17,22,23} These factors make it difficult to determine if an extremely preterm infant with perceived low BP but clinically adequate perfusion would benefit from or be harmed by therapy.

CONCLUSIONS

This prospective multicenter study of extremely preterm infants examined the relationship among 15 definitions of

low BP, antihypotensive therapy, and in-hospital outcomes. Therapy was not associated with better in-hospital outcomes for any definition of low BP investigated. A numeric cutoff for deciding when to administer antihypotensive therapies, such as an MAP less than or equal to the infant's GA, is not evidence based and cannot be recommended. Until there are data to suggest otherwise, antihypotensive therapy should be used cautiously for these infants because treatment of low BP is associated with similar or worse infant outcomes without evidence of benefit.^{4,17,22,23} Large, high-quality studies are needed to support evidence based recommendations for BP management in this population.

ACKNOWLEDGMENTS

Data collected at participating sites of the Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network were transmitted to RTI International, the data coordinating center (DCC) for the network, which stored, managed, and analyzed the data for this study. On behalf of the Neonatal Research Network, Drs Abhik Das (DCC principal investigator) and Lei Li (DCC statistician) had full access to all the data in the study and take responsibility for the integrity of the data and accuracy of the data analysis. We are indebted to our medical and nursing colleagues and the infants and their parents who agreed to take part in this study. The following investigators, in addition to those listed as authors, participated in this study:

NRN Steering Committee Chair: Michael S. Caplan, MD, University of Chicago, Pritzker School of Medicine (2006–2011); Richard A. Polin, MD, Division of Neonatology, College of Physicians and Surgeons, Columbia University (2011–present)

Alpert Medical School of Brown University and Women & Infants Hospital of Rhode Island (grant U10 HD27904): William Oh, MD; Angelita M. Hensman, RNC-NIC, BSN; Dawn Andrews, RN; Kristen Angela, RN; Susan Barnett, RRT; Kim Francis, RN; Dan Gingras, RRT; Sarah Lillie, RRT; Suzy Ventura

Case Western Reserve University, Rainbow Babies & Children's Hospital (grants U10 HD21364, M01 RR80): Avroy A. Fanaroff, MD; Bonnie S. Siner, RN; Deanne E. Wilson-Costello, MD

Cincinnati Children's Hospital Medical Center, University Hospital, and Good Samaritan Hospital (grants U10 HD27853, M01 RR8084): Barbara Alexander, RN; Cathy Grisby, BSN, CCRC; Lenora Jackson; Kristin Kirker; Estelle Fischer, MHA, MBA

Duke University School of Medicine, University Hospital, Alamance Regional Medical Center, and Durham Regional Hospital (grants U10 HD40492, M01 RR30): Ronald N. Goldberg, MD; C. Michael Cotten, MD, MHS; Kimberley A. Fisher, PhD, FNP-BC, IBCLC; Sandy Grimes, RN, BSN; University of North Carolina at Chapel Hill (grant UL1 RR25747): Carl L. Bose, MD; Gennie Bose, RN; Cindy Clark, RN

Emory University, Children's Healthcare of Atlanta, Grady Memorial Hospital, and Emory University Hospital Midtown (grants U10 HD27851, UL1 RR25008, M01 RR39): David P. Carlton, MD; Ellen C. Hale, RN, BS, CCRC; Amy K. Hutchinson, MD

Eunice Kennedy Shriver National Institute of Child Health and Human Development: Stephanie Wilson Archer, MA

Indiana University, University Hospital, Methodist Hospital, Riley Hospital for Children, and Wishard Health Services (grants U10 HD27856, M01 RR750): Leslie D. Wilson, BSN CCRC; Dianne E. Herron, RN

RTI International (grant U10 HD36790): Dennis Wallace, PhD; Jeanette O'Donnell Auman, BS; Margaret Cunningham, BS;

Carolyn M. Petrie Huitema, MS; James W. Pickett II, BS; Kristin M. Zaterka-Baxter, RN, BSN

Stanford University, Lucile Packard Children's Hospital (grants U10 HD27880, M01 RR70): David K. Stevenson, MD; M. Bethany Ball, BS, CCRC; Melinda S. Proud, RCP

Tufts Medical Center, Floating Hospital for Children (grants U10 HD53119, M01 RR54): John M. Fiascone, MD; Anne Furey, MPH; Brenda L. MacKinnon, RNC; Ellen Nylen, RN, BSN

University of Alabama at Birmingham Health System and Children's Hospital of Alabama (grants U10 HD34216, M01 RR32): Namasivayam Ambalavanan, MD; Monica V. Collins, RN, BSN, MaEd; Shirley S. Cosby, RN, BSN

University of Iowa Children's Hospital and Mercy Medical Center (grants U10 HD53109, UL1 RR24979, M01 RR59): Edward F. Bell, MD; Dan L.

Ellsbury, MD; Karen J. Johnson, RN, BSN; Donia D. Campbell, RNC-NIC; Rachael M. Hyland, BA

University of New Mexico Health Sciences Center (grants U10 HD53089, M01 RR997): Robin K. Ohls, MD; Conra Backstrom Lacy, RN

University of Texas Southwestern Medical Center at Dallas, Parkland Health and Hospital System, and Children's Medical Center Dallas (grants U10 HD40689, M01 RR633): Laura Grau, RN; Melissa Martin, RN; Araceli Solis, RRT; Lizette E. Torres, RN; Diana M. Vasil, RNC-NIC; Kerry Wilder, RN

University of Texas Health Science Center at Houston Medical School, Children's Memorial Hermann Hospital (U10 HD21373): Jon E. Tyson, MD, MPH; Beverly Foley Harris, RN, BSN; Anna E. Lis, RN, BSN; Sarah Martin, RN, BSN; Georgia E. McDavid, RN; Patti

L. Pierce Tate, RCP; Sharon L. Wright, MT (ASCP)

University of Utah, University Hospital, Intermountain Medical Center, and Primary Children's Medical Center (grants U10 HD53124, M01 RR64, UL1 RR25764): Karen A. Osborne, RN, BSN, CCRC; Jill Burnett, RNC; Cynthia Spencer, RNC; Kimberlee Weaver-Lewis, RN, BSN; Karie Bird, RN; Karen Zanetti, RN; Laura Cole, RN

Wayne State University, University of Michigan, Hutzel Women's Hospital, and Children's Hospital of Michigan (grant U10 HD21385): Beena G. Sood, MD, MS; John Barks, MD; Rebecca Bara, RN, BSN; Mary Johnson, RN, BSN

Yale University and Yale-New Haven Children's Hospital, and Bridgeport Hospital (grants U10 HD27871, UL1 RR24139): Monica Konstantino, RN, BSN; JoAnn Poulsen, RN

REFERENCES

- Dempsey EM, Barrington KJ. Treating hypotension in the preterm infant: when and with what: a critical and systematic review. *J Perinatol*. 2007;27(8):469–478
- Evans N. Which inotrope for which baby? *Arch Dis Child Fetal Neonatal Ed*. 2006;91(3):F213–F220
- Laughon M, Bose C, Allred E, et al; ELGAN Study Investigators. Factors associated with treatment for hypotension in extremely low gestational age newborns during the first postnatal week. *Pediatrics*. 2007;119(2):273–280
- Batton B, Batton D, Riggs T. Blood pressure during the first 7 days in premature infants born at postmenstrual age 23 to 25 weeks. *Am J Perinatol*. 2007;24(2):107–115
- Fanaroff JM, Wilson-Costello DE, Newman NS, Montpetite MM, Fanaroff AA. Treated hypotension is associated with neonatal morbidity and hearing loss in extremely low birth weight infants. *Pediatrics*. 2006;117(4):1131–1135
- Goldstein RF, Thompson RJ, Oehler JM, et al. Influence of acidosis, hypoxemia, and hypotension on neurodevelopmental outcome in very low birth weight infants. *Pediatrics*. 1995;95(2):238–243
- Low JA, Froese AB, Galbraith RS, Smith JT, Sauerbrei EE, Derrick EJ. The association between preterm newborn hypotension and hypoxemia and outcome during the first year. *Acta Paediatr*. 1993;82(5):433–437
- Martens SE, Rijken M, Stoelhorst GM, et al; Leiden Follow-Up Project on Prematurity, The Netherlands. Is hypotension a major risk factor for neurological morbidity at term age in very preterm infants? *Early Hum Dev*. 2003;75(1–2):79–89
- Hall RW, Kronsberg SS, Barton BA, Kaiser JR, Anand KJ; NEOPAIN Trial Investigators Group. Morphine, hypotension, and adverse outcomes among preterm neonates: who's to blame? Secondary results from the NEOPAIN trial. *Pediatrics*. 2005;115(5):1351–1359
- Finer NN, Powers RJ, Ou CH, Durand D, Wirtschafter D, Gould JB; California Perinatal Quality Care Collaborative Executive Committee. Prospective evaluation of postnatal steroid administration: a 1-year experience from the California Perinatal Quality Care Collaborative. *Pediatrics*. 2006;117(3):704–713
- Ewer AK, Tyler W, Francis A, Drinkall D, Gardosi JO. Excessive volume expansion and neonatal death in preterm infants born at 27–28 weeks gestation. *Paediatr Perinat Epidemiol*. 2003;17(2):180–186
- Clark CE, Clyman RI, Roth RS, Sniderman SH, Lane B, Ballard RA. Risk factor analysis of intraventricular hemorrhage in low-birth-weight infants. *J Pediatr*. 1981;99(4):625–628
- Vain NE, Barrington KJ. Feasibility of evaluating treatment of early hypotension in extremely low birth weight infants. *J Pediatr*. 2012;161(1):4–7
- Rich WD, Auten KJ, Gantz MG, et al; National Institute of Child Health and Human Development Neonatal Research Network. Antenatal consent in the SUPPORT trial: challenges, costs, and representative enrollment. *Pediatrics*. 2010;126(1). Available at: www.pediatrics.org/cgi/content/full/126/1/e215
- Batton BJ, Li L, Newman NS, et al. Feasibility study of early blood pressure management in extremely preterm infants. *J Pediatr*. 2012;161:65–69.e1
- Rich W, Finer NN, Gantz MG, et al; SUPPORT and Generic Database Subcommittees of the Eunice Kennedy Shriver National Institute of Child Health and Human

- Development Neonatal Research Network. Enrollment of extremely low birth weight infants in a clinical research study may not be representative. *Pediatrics*. 2012;129(3):480–484
17. Logan JW, O'Shea TM, Allred EN, et al; ELGAN Study Investigators. Early postnatal hypotension and developmental delay at 24 months of age among extremely low gestational age newborns. *Arch Dis Child Fetal Neonatal Ed*. 2011;96(5):F321–F328
 18. Papile LA, Burstein J, Burstein R, Koffler H. Incidence and evolution of subependymal and intraventricular hemorrhage: a study of infants with birth weights less than 1,500 gm. *J Pediatr*. 1978;92(4):529–534
 19. Bell MJ, Ternberg JL, Feigin RD, et al. Neonatal necrotizing enterocolitis. Therapeutic decisions based upon clinical staging. *Ann Surg*. 1978;187(1):1–7
 20. International Committee for the Classification of Retinopathy of Prematurity. The International Classification of Retinopathy of Prematurity revisited. *Arch Ophthalmol*. 2005;123(7):991–999
 21. Walsh MC, Wilson-Costello D, Zadell A, Newman N, Fanaroff A. Safety, reliability, and validity of a physiologic definition of bronchopulmonary dysplasia. *J Perinatol*. 2003;23(6):451–456
 22. Batton B, Zhu X, Fanaroff J, et al. Blood pressure, anti-hypotensive therapy, and neurodevelopment in extremely preterm infants. *J Pediatr*. 2009;154(3):351–357, e1
 23. Dempsey EM, Al Hazzani F, Barrington KJ. Permissive hypotension is associated with good neurodevelopmental outcome in extremely low birthweight infants. *Arch Dis Child Fetal Neonatal Ed*. 2009;94(4):F241–F244
 24. Dempsey EM, Barrington KJ. Diagnostic criteria and therapeutic interventions for the hypotensive very low birth weight infant. *J Perinatol*. 2006;26(11):677–681
 25. Bonestroo H, Lemmers P, Baerts W, van Bel F. Effect of antihypotensive treatment on cerebral oxygenation of preterm infants without PDA. *Pediatrics*. 2011;128(6). Available at: www.pediatrics.org/cgi/content/full/128/6/e1502
 26. Short BL, Van Meurs K, Evans JR; Cardiology Group. Summary proceedings from the cardiology group on cardiovascular instability in preterm infants. *Pediatrics*. 2006;117(3 pt 2):S34–S39
 27. Joint Working Party of British Association of Perinatal Medicine and the Research Unit of the Royal College of Physicians. Development of audit measures and guidelines for good practice in the management of neonatal respiratory distress syndrome. Report of a Joint Working Group of the British Association of Perinatal Medicine and the Research Unit of the Royal College of Physicians. *Arch Dis Child*. 1992;67(spec no 10):1221–1227
 28. Food and Drug Administration. Studies of drugs in neonates challenging but necessary. *AAP News*. 2012;33:7
 29. Schlapbach LJ, Aebischer M, Adams M, et al; Swiss Neonatal Network and Follow-Up Group. Impact of sepsis on neurodevelopmental outcome in a Swiss National Cohort of extremely premature infants. *Pediatrics*. 2011;128(2). Available at: www.pediatrics.org/cgi/content/full/128/2/e348
 30. Stoll BJ, Hansen NI, Sánchez PJ, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Early onset neonatal sepsis: the burden of group B Streptococcal and *E. coli* disease continues. *Pediatrics*. 2011;127(5):817–826
 31. Pöschl JM, Weiss T, Fallahi F, Linderkamp O. Reactive hyperemia of skin microcirculation in septic neonates. *Acta Paediatr*. 1994; 83(8):808–811
 32. Tibby SM, Hatherill M, Murdoch IA. Capillary refill and core-peripheral temperature gap as indicators of haemodynamic status in paediatric intensive care patients. *Arch Dis Child*. 1999;80(2):163–166
 33. Noori S, Seri I. Neonatal blood pressure support: the use of inotropes, lusitropes, and other vasopressor agents. *Clin Perinatol*. 2012;39(1):221–238
 34. Garner R, Burchfield D. Treatment of presumed hypotension in very low birthweight neonates: effects on regional cerebral oxygenation. *Arch Dis Child Fetal Neonatal Ed*. 2013;98(2):F117–F121
 35. Giliberti P, Giordano L, Chello G, De Leonibus C, Giliberti P. The scenarios of shock in newborn infants. *J Matern Fetal Neonatal Med*. 2010;23(suppl 3):27–29
 36. El-Khuffash AF, Walsh K, Molloy EJ. Blood pressure correlates poorly with left ventricular output and celiac artery blood flow in preterm infants in the first 48 hours of life. *J Neonatal Perinatal Med*. 2008;1(1):37–41
 37. Cayabyab R, McLean CW, Seri I. Definition of hypotension and assessment of hemodynamics in the preterm neonate. *J Perinatol*. 2009;29(suppl 2):S58–S62
 38. Gilmore MM, Stone BS, Shepard JA, Czosnyka M, Easley RB, Brady KM. Relationship between cerebrovascular dysautoregulation and arterial blood pressure in the premature infant. *J Perinatol*. 2011;31(11):722–729
 39. Tyszczyk L, Meek J, Elwell C, Wyatt JS. Cerebral blood flow is independent of mean arterial blood pressure in preterm infants undergoing intensive care. *Pediatrics*. 1998;102(2 pt 1):337–341
 40. Osborn DA, Paradisis M, Evans N. The effect of inotropes on morbidity and mortality in preterm infants with low systemic or organ blood flow. *Cochrane Database Syst Rev*. 2007;(1):CD005090

(Continued from first page)

Dr Batton is the lead study investigator for the Eunice Kennedy Shriver National Institute of Child Health & Human Development Neonatal Research Network (NRN) Early Blood Pressure study and the chair of the Early Blood Pressure Protocol Subcommittee. He developed the protocol with the assistance of the subcommittee, trained the study coordinators and principal investigators (PIs), and managed study implementation. As the Study PI at Case Western Reserve University, he oversaw study implementation at that site, which enrolled 25 infants in this study. Dr Batton drafted the manuscript and received input from the other authors as part of manuscript revision. Dr Walsh is the PI at Case Western Reserve University and the vice chair of the Early Blood Pressure Protocol Subcommittee. She mentored Dr Batton through protocol development and implementation, and, as the PI, oversaw subject recruitment and study implementation at her site, which enrolled 25 infants. She provided critical revisions to the manuscript and approved the final version of the manuscript. Dr Li served as the primary statistician for the study, providing statistical input for protocol development and completing the statistical analyses for the manuscript. He developed the tables and figures for the manuscript, provided critical revision to the manuscript, and approved the final version of the manuscript. Ms Newman is the NRN coordinator at Case Western Reserve University and a member of the Early Blood Pressure Protocol Subcommittee. She provided day-to-day assistance with protocol and procedural questions both at her site and for PIs and coordinators at other NRN sites. Ms Newman provided critical revision of the manuscript as well as approval of the final manuscript. Dr Das is the PI for the NRN Data Coordinating Center and a member of the Early Blood Pressure Protocol Subcommittee. Dr Das oversaw all aspects of the statistical analysis, provided critical revisions to the manuscript, and approved the final version of the manuscript. Dr Watterberg is the PI at the University of New Mexico and a member of the Early Blood Pressure Protocol Subcommittee. As the PI, she oversaw subject recruitment and study implementation at her site, which enrolled 13 infants in this study. She contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr Yoder is the

Alternate PI at the University of Utah and a member of the Early Blood Pressure Protocol Subcommittee. As the Alternate PI, he helped oversee subject recruitment and study implementation at his site, which enrolled 27 infants. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Faix is the PI at the University of Utah and a member of the Early Blood Pressure Protocol Subcommittee. As the PI, he oversaw subject recruitment and study implementation at his site, which enrolled 27 infants in this study. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Laughon is the site investigator at the University of North Carolina, a satellite site of Duke University in the NRN and a member of the Early Blood Pressure Protocol Subcommittee. As the investigator, he oversaw subject recruitment and study implementation at the satellite site. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Stoll is the PI at Emory University School of Medicine and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, she oversaw subject recruitment and study implementation at her site, which enrolled 19 infants in this study. She contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Van Meurs is the PI at Stanford University and a member of the Early Blood Pressure Protocol Subcommittee. As the PI, she oversaw subject recruitment and study implementation at her site, which enrolled 18 infants in this study. She contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Carlo is the PI at the University of Alabama and a member of the Early Blood Pressure Protocol Subcommittee. As the PI, he oversaw subject recruitment and study implementation at his site, which enrolled 36 infants in this study. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Poindexter is the PI at the Indiana University School of Medicine and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, she oversaw subject recruitment and study implementation at her site, which enrolled 37 infants in this study. She contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Bell is the PI at the University of Iowa and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, he oversaw subject recruitment and study implementation at his site, which enrolled 30 infants in this study. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Sánchez is the PI at the University of Texas Southwestern Medical Center and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, he oversaw subject recruitment and study implementation at his site, which enrolled 29 infants in this study. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Ehrenkranz is the PI at Yale University School of Medicine and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, he oversaw subject recruitment and study implementation at his site, which enrolled 31 infants in this study. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Goldberg is the PI at Duke University and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, he oversaw subject recruitment and study implementation at his site, which enrolled 31 infants in this study. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Laptook is the PI at the Women & Infants' Hospital at Brown University and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, he oversaw subject recruitment and study implementation at his site, which enrolled 21 infants in this study. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Kennedy is the PI at the University of Texas Medical School at Houston and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, she oversaw subject recruitment and study implementation at her site, which enrolled 16 infants in this study. She contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Frantz was the PI at the Floating Hospital for Children, Tufts Medical Center and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, he oversaw subject recruitment and study implementation at his site, which enrolled 14 infants in this study. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Shankaran is the PI at Wayne State University and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, she oversaw subject recruitment and study implementation at her site, which enrolled 13 infants in this study. She contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Schibler is the PI at Cincinnati Children's Hospital Medical Center and a member of the NRN Steering Committee, which oversees and approves all investigations within the NRN. As the PI, he oversaw subject recruitment and study implementation at his site, which enrolled 7 infants in this study. He contributed critical revisions of the manuscript and approved the final manuscript for submission. Dr. Higgins served as the Program Scientist for the National Institute of Child Health & Human Development NRN and a member of the Early Blood Pressure Protocol Subcommittee. Dr. Higgins helped develop the protocol, oversaw study implementation, and assisted with data edits from the sites. She also provided critical revision to the manuscript and approved the final version of the manuscript.

This trial has been registered at www.clinicaltrials.gov (identifier NCT00874393).

www.pediatrics.org/cgi/doi/10.1542/peds.2012-2779

doi:10.1542/peds.2012-2779

Accepted for publication Feb 7, 2013

Address correspondence to Beau Batton, MD, Southern Illinois University School of Medicine, Department of Pediatrics, Division of Neonatology, PO Box 19676, Springfield, IL 62794. E-mail: bbatton@siumed.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2013 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: *Dr Sanchez reports receiving grants from Glaxosmithkline and Abbott Laboratories for studies unrelated to this investigation; the other authors have indicated they have no financial relationships relevant to this article to disclose.*

FUNDING: The National Institutes of Health and the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) provided grant support, including funding from the Best Pharmaceuticals for Children Act, for the Neonatal Research Network's Early Blood Pressure Observational Study. Funded by the National Institutes of Health (NIH).