

Cerebrospinal Fluid Pressure Decreases with Older Age

David Fleischman¹, John P. Berdahl², Jana Zaydlarova³, Sandra Stinnett⁴, Michael P. Fautsch³, R. Rand Allingham^{4*}

1 Department of Ophthalmology, University of North Carolina, Chapel Hill, North Carolina, United States of America, **2** Vance Thompson Vision, University of South Dakota Medical Center, Sioux Falls, South Dakota, United States of America, **3** Department of Ophthalmology, Mayo Clinic, Rochester, Minnesota, United States of America, **4** Duke University Eye Center, Durham, North Carolina, United States of America

Abstract

Purpose: Clinical studies implicate low cerebrospinal fluid pressure (CSFP) or a high translaminar pressure difference in the pathogenesis of primary open angle glaucoma (POAG) and normal tension glaucoma (NTG). This study was performed to examine the effect of age, sex, race and body mass index (BMI) on CSFP.

Methods: Electronic medical records from all patients who had a lumbar puncture (LP) performed at the Mayo Clinic from 1996–2009 were reviewed. Information including age, sex, race, height and weight, ocular and medical diagnoses, intraocular pressure (IOP) and LP opening pressure was obtained. Patients using medications or with medical diagnoses known to affect CSFP, and those who underwent neurosurgical procedures or where more than one LP was performed were excluded from analysis.

Results: Electronic medical records of 33,922 patients with a history of having an LP during a 13-year period (1996–2009) were extracted. Of these, 12,118 patients met all entry criteria. Relative to mean CSFP at age group 20–49 (mean 11.5 ± 2.8 mmHg), mean CSFP declined steadily after age 50, with percent reduction of 2.5% for the 50–54 age group (mean 11.2 ± 2.7 mmHg, $p < 0.002$) to 26.9% for the 90–95 group (mean 8.4 ± 2.4 mmHg, $p < 0.001$). Females had lower CSFP than males throughout all age groups. BMI was positively and independently associated with CSFP within all age groups.

Conclusion: There is a sustained and significant reduction of CSFP with age that begins in the 6th decade. CSFP is consistently lower in females. BMI is positively and independently associated with CSFP in all age groups. The age where CSFP begins to decline coincides with the age where the prevalence of POAG increases. These data support the hypothesis that reduced CSFP may be a risk factor for POAG and may provide an explanation for the mechanism that underlies the age-related increase in the prevalence of POAG and NTG.

Citation: Fleischman D, Berdahl JP, Zaydlarova J, Stinnett S, Fautsch MP, et al. (2012) Cerebrospinal Fluid Pressure Decreases with Older Age. *PLoS ONE* 7(12): e52664. doi:10.1371/journal.pone.0052664

Editor: Yingfeng Zheng, Zhongshan Ophthalmic Center, China

Received: August 30, 2012; **Accepted:** November 19, 2012; **Published:** December 26, 2012

Copyright: © 2012 Fleischman et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by National Institutes of Health research grants EY 15736 (MPF), EY 07065 (MPF); Mayo Foundation, Rochester, MN (MPF); and Research to Prevent Blindness, New York, NY (MPF is a recipient of a Lew R. Wasserman Merit Award and the Department of Ophthalmology, Mayo Clinic; Department of Ophthalmology, University of North Carolina Hospitals, and Duke University Eye Center are recipients of unrestricted grants). The funding agencies had no role in the design or conduct of this research.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: rand.allingham@duke.edu

Introduction

Primary open angle glaucoma (POAG) is the most common form of glaucoma, a family of disorders with varying biological and environmental risk factors. Well-described risk factors for POAG include elevated intraocular pressure (IOP), advancing age, African and Hispanic ancestry [1], reduced central corneal thickness [1–5], and a positive family history. Recent studies suggest that lower cerebrospinal fluid pressure (CSFP) may contribute to the risk of developing open angle glaucoma, including both POAG and normal tension glaucoma [6–8]. The lamina cribrosa located at the anterior face of the optic nerve is situated between two pressurized compartments, the intraocular and subarachnoid spaces, and is the site of retinal ganglion cell axon loss in glaucoma. Both elevated IOP and reduced CSFP increase translaminar pressure. It is hypothesized that lower cerebrospinal fluid pressure acts similarly to elevated IOP at the

optic nerve head, increasing risk for glaucomatous neuropathy. Retrospective and prospective clinical studies have found that CSFP is lower in patients with POAG and increased in patients with ocular hypertension (OHT) [6–9]. Importantly, this hypothesis offers an explanation for the mechanism of disease in patients with normal tension glaucoma (NTG), which occurs in the absence of statistically elevated IOP. Conversely, as elevated CSFP would theoretically reduce the translaminar pressure difference, higher CSFP could function as a protective factor for glaucoma in patients with OHT.

Even though the cerebrospinal fluid (CSF) has been studied for many decades, there is remarkably little published about what constitutes a “normal” CSFP. The primary method to determine CSFP in patients is to perform a lumbar puncture (LP). Since performing an LP carries some risk of adverse events to the patient, it has generally been reserved for patients suspected of harboring serious diseases and is rarely performed in otherwise

healthy subjects. Consequently, current CSFP reference ranges have been generated from either small groups of volunteers or from larger, less well-characterized groups [10–12].

The purpose of the current study was to investigate the effect of age, sex, race, and body mass index on CSFP in a large dataset based on a long-standing electronic medical record system.

Methods

Patient Selection

This was a retrospective chart analysis from the Mayo Clinic's electronic medical records system of patients over age 20 years who underwent a diagnostic lumbar puncture from 1996 to 2009. Following Institutional Review Board approval granted from the Mayo Clinic in Rochester, Minnesota, a list of patients from December 1, 1996 to December 31, 2009 was generated by searching for diagnostic lumbar puncture by CPT code 62270. All subjects were de-identified according to Mayo Clinic protocol.

Lumbar Puncture

At the Mayo Clinic (Rochester, MN) trained teams perform lumbar punctures in most cases. These teams use a standardized method that is performed similarly for all patients. Using this approach, patients are placed in the lateral decubitus position and either the L3 to L4 or L4 to L5 interspace is identified and anesthetized. A 3.5-inch 20-g spinal needle with a 3-way stopcock is inserted into the subarachnoid space. A 550-mm manometer is attached to the stopcock and the column of CSF fluid is allowed to equilibrate. The patient is asked to remain still and not to speak. For some patients, a bedside lumbar puncture may be difficult due to body habitus or due to other anatomical issues. In those cases, the lumbar puncture was performed by Radiology under fluoroscopic guidance. The electronic database extraction does not indicate which method was used for LP. In either method, the meniscus of the CSF fluid is read and reported in millimeters of water. For this study we converted millimeters of water to millimeters of mercury in order to simplify comparison with IOP for data analysis and discussion (1 mmHg = 13.6 mm H₂O).

Extracted Information

Information abstracted from this search included time and date of lumbar puncture, as well as the time and date of the CSF analysis. Patient's age, sex, race, ethnicity, height, and weight were obtained at the time of LP. A list of up to 15 separate current and past diagnoses was obtained as well as all medications. Blood pressures measured at the various visits of the patients who underwent LP were available for review. Only blood pressures recorded within one month of the lumbar puncture (either before or after) were included for analysis. The blood pressure measured closest to the time of lumbar puncture was selected. If blood pressure was measured one day before and one day following LP we used the earlier measurement for analysis.

All admissions or procedures performed at the Mayo Clinic were reviewed for each patient. BMI for each patient was calculated manually [$BMI = (\text{weight in kilograms}) / (\text{height in meters})^2$] and rounded to the nearest whole number. BMI less than 10 and greater than 50 were excluded as these values were felt to be the result of data entry errors. The height and weight used for calculation of BMI was performed from measurements recorded in the hospital stay associated with the LP.

IOP Data

Patients who met initial screening criteria and had an ophthalmologic exam including IOP measurements within one-

year of the LP were identified. The method of measurement of IOP was not recorded in the dataset. Patients with POAG or other eye disease were not excluded from this study (with the exception of idiopathic intracranial hypertension since it is associated with altered CSFP). If repeat ophthalmologic exams were performed, the IOP measurement closest to LP data was used for evaluation. Right eye, left eye, and the average of both eyes IOP were regressed against age.

Screening

Information from specific patients was given a unique, randomly generated identifier and entered into a database. Patients with medical conditions, head trauma, taking medications known to alter CSFP, or having a neurosurgical procedure were excluded. A total of 134 different medications and diagnostic codes of medical conditions that might affect CSFP were used for manual screening of all electronic medical records. Filters were applied regardless of time sequence leading to the LP. For example, if a patient developed a condition known to alter CSFP at any time point, they were excluded from the study. All patients with multiple LPs were excluded. Patients with CSFP values <60 or >250 mmH₂O (<4.41 mmHg or >18.38 mmHg), considered outside the accepted normal range, were excluded from analysis [13].

Analysis

Subjects who met all inclusion criteria were stratified into 1, 5 and 10-year age groups. For the purpose of this study, we used World Health Organization criterion for the age of adulthood, which is ≥ 20 years [14]. Mean CSFP and standard deviation was calculated for each age group. A two-tailed student's t-test was used to measure the difference between each group and to determine significance. A p-value < 0.05 was used for statistical significance.

Multivariate linear regression models, both segmented and non-segmented by age, were created for multiple sets of variables. Age, BMI, sex, race (Black, Caucasian, Asian, Other), presence of diabetes mellitus (DM), mean arterial pressure (MAP), and time of LP were regressed for the purpose of determining independent association with CSFP.

Results

The CPT database extraction protocol identified electronic medical records for 33,922 patients who had diagnostic lumbar punctures performed between December 1, 1996 and December 31, 2009. The number meeting initial inclusion criteria was 13,715, which is 40.4% of the original extraction group. The final dataset used for analysis contained 12,118 patients after exclusion for CSFP values out of the normal range, age less than 20, and BMI < 10 and > 50. The distribution and demographic data of the analysis dataset are listed in **Table 1 and 2**. The age distribution of the study group was evenly distributed with mean age of 54.4 years \pm 15.2, with a median age of 55. Caucasians were the largest group in this population (79.1%), followed by "Unknown" (17.1%), Black (1.4%), and "Other" (1.0%). Regarding ethnicity, 0.6% self-identified as Hispanic, 17.4% as Not-Hispanic, and 81.9% were documented as "Unknown." Sex was evenly distributed between males and females, 49% and 51% respectively. While indications for LP were not included in the extracted data, the primary diagnosis was available for the medical visit where the LP was performed. There were a total of 1,404 unique diagnoses recorded among the patients included in the final dataset. Of all the patients included in the final dataset, over

Table 1. Derivation of the final study population after adoption of all entry criteria.

Disposition	N	Percent
Extracted in data base	13715	100
Excluded*	1597	11.6
- Out of range CSFP values (<4.4 or >18.4)	1152	8.4
- Age ≤19	532	3.9
- BMI ≤10 or ≥50	27	0.2
Included in overall study	12118	88.4
Included in analysis of BMI out of overall included in study	4314	35.6

*114 were in two exclusion categories.

doi:10.1371/journal.pone.0052664.t001

69,041 diagnoses were listed. The top ten primary diagnoses for the analysis dataset are listed in **Table 3**.

There was no difference in CSFP values measured by 1, 5, or 10 year age groupings between ages 20 to 49. Therefore we used mean CSFP for the 20–49 age group as a baseline to compare to subsequent five-year age groups. Compared to the mean CSFP for age group 20–49, a statistically significant difference in mean CSFP was evident in the 50–54 age group (2.5%, $p<0.002$). (**Table 4 and Figure 1**) Mean CSFP progressively decreased for each subsequent 5-year age group. CSFP reduction was 13.3% (1.53 mmHg) at age 80, 19.4% (2.23 mmHg) at age 85, and

Table 3. Top ten primary diagnoses at visit of lumbar puncture for patients ages 20–95.

	Primary Diagnosis	N	%
1	IDIOPATHIC PERIPHERAL NEUROPATHY, NOS	887	7.69%
2	SKIN SENSATION DISTURBANCES	424	3.67%
3	MULTIPLE SCLEROSIS	397	3.44%
4	HEADACHE	370	3.21%
5	AMYOTROPHIC SCLEROSIS	367	3.18%
6	ABNORMALITY OF GAIT	286	2.48%
7	SPINAL CORD DISEASE NOS	250	2.17%
8	OTHER MALAISE & FATIGUE	195	1.69%
9	PERSISTENT MENTATION DISORDER NEC	164	1.42%
10	HYPERTENSION NOS	161	1.40%

N: Number of Patients; % = Percent of all primary diagnoses.

doi:10.1371/journal.pone.0052664.t003

26.9% (3.09 mmHg) at age 90–95. A similar analysis, which grouped the study population as young adults (20–49 years), late-middle age (50–69), and older adults (greater than 70 years), was

Table 2. Demographics and disposition of study patients following final screening.

Demographic	Statistic	All Study Patients	Patients with Recorded BMI	
	N	12118	4314	
Age	Mean (SD)	54.4 (15.2)	55.0 (15.2)	
	Min, Median, Max	20, 55, 95	20, 55, 91	
BMI	N	4314		
	Mean (SD)	26.7 (5.2)	26.7 (5.2)	
	Min, Median, Max	10.1, 26.1, 49.1	10.1, 26.1, 49.1	
Gender	N (%)			
	- Female	6224 (51)	2250 (52)	
	- Male	5894 (49)	2064 (48)	
Race	N (%)			
		- American Indian/Eskimo	44 (0.4)	20 (0.5)
		- Black	171 (1.4)	73 (1.7)
		- Caucasian	9586 (79.1)	3409 (79.0)
		- Native Hawaiian/Pacific Islander	4 (0.03)	4 (0.1)
		- Other	121 (1.00)	43 (1.0)
		- Asian	81 (0.7)	23 (0.5)
		- Unknown	2073 (17.1)	726 (17.9)
		- Choose not to disclose	38 (0.3)	16 (0.4)
		Ethnicity	N (%)	
- Hispanic	73 (0.6)			29 (0.7)
- Not Hispanic	2107 (17.4)			1036 (24.0)
- Unknown	9919 (81.8)			3238 (75.1)
- Choose not to disclose	19 (0.2)			11 (0.2)

doi:10.1371/journal.pone.0052664.t002

Cerebrospinal Fluid Pressure and Age




Figure 1. Mean CSFP within specific age groups. Bars represent one standard deviation.
doi:10.1371/journal.pone.0052664.g001

performed. The 20–49 group had a mean CSFP of 11.5 mm Hg \pm 2.7, which was 5.3% (10.9 mmHg \pm 2.7; $p < 0.0001$) higher than the 50–69 group, and 13.1% (10.0 mm Hg \pm 2.6; $p < 0.0001$) higher than the group of older adults, or those greater than 70 years.

To determine the relationship of age and BMI on CSFP, a two-way analysis of variance was fit to the data. For the subset of subjects with recorded BMI, stratifications were created by evenly distributing the N of the subgroups. Both age ($p < 0.001$) and BMI (< 0.001) were significant predictors of CSFP. There was no difference in the trend of CSFP in the BMI categories over the age categories ($p = 0.611$): overall, CSFP was negatively correlated with advancing age after age 50 and was positively correlated with BMI regardless of age and within all age groups. (Table 5) This finding was consistent using either segmented (correlation: 0.223, standard error [SE]: 0.01; $p < 0.001$) or non-segmented (correlation: 0.23, SE: 0.01; $p < 0.001$) linear regression models, both of which confirmed that BMI and age are independently associated with CSFP.

CSFP was significantly different between sexes. On average, males have a higher CSFP than females by approximately 0.9 mmHg. The rate of change of CSFP with age for males and females was the same (ages 20–50, $p = 0.369$; ages 50–95, $p = 0.067$). Although this difference was significant, sex as a variable explains only a small portion of the variability of CSFP in this dataset ($r^2 = 0.07$, $p < 0.0001$). (Figure 2)

We attempted to look at the effect of race on CSFP with age. Overall, mean CSFP was higher in Black patients compared to Caucasians (12.01 mmHg \pm 2.9 and 10.9 mmHg \pm 2.7, respectively). The N for all Black patients within this study was 171, or 1.4% of the dataset. The N was further reduced by incorporation of other independent factors such as BMI for regression analyses. The lack of statistical power precluded our ability to determine the role, if any, between race and ethnicity with CSFP.

Other variables such as blood pressure, diabetes, and the time of lumbar puncture were examined. Mean arterial pressure (MAP) within one month of LP was positively, but weakly, correlated with CSFP (correlation: 0.016, SE: 0.00; $p < 0.0001$). A statistically significant, but weakly positive correlation exists between MAP

and age in this study group. ($r^2 = 0.02$, $p < 0.0001$). Blood pressure data was available for approximately one third of patients ($N = 4197$ or 34.6%), and the respective model explains only a small amount of the observed variability in CSFP. CSFP was not correlated with time of lumbar puncture (correlation: 0.012, SE: 0.01; $p = 0.3881$) or diabetes mellitus status (correlation: -0.213 , SE: 0.15; $p = 0.149$).

A subgroup of patients ($N = 441$) who had IOP and CSFP recorded were selected for IOP and CSFP analysis. Whereas CSFP changes significantly with age, there was no statistically significant difference in mean IOP with age in this group. (Table 6, Figure 3).

Discussion

There is a growing body of evidence that indicates CSFP is lower in patients with primary open angle glaucoma compared to non-glaucomatous subjects and additionally, is lower in normal tension versus the high tension form of POAG.^{6–8} If CSFP is related to the pathogenesis of POAG then it is important to explore factors that affect CSFP. In this study we report the effect of age on CSFP. We have found that mean CSFP is stable for the first 50 years of life after which there is a steady decline. The observed change in CSFP is substantial, with the mean reduction in measured CSFP being 3.1 mm Hg (26.9%) by age 90. We also determined that the effect of age on CSFP is independent of BMI, and that BMI is positively and independently associated with CSFP.

The few studies that have examined CSFP and age failed to find a relationship [11,12,15,16]. In a study of 200 mostly syphilitic subjects, performed by Masserman in 1935, no significant correlation was found between age and resting cerebrospinal fluid pressure in those aged 16 to 48 [7]. Our study corroborates CSFP stability in this age group. Studies by Ekstedt and, more recently, Malm were conducted on 100 and 40 subjects respectively [11,16]. No correlation was found between CSFP and age in these studies. However, after stratification by age, it is likely that these studies were underpowered to observe this phenomenon. In contrast, the current study was based on measurements obtained on 12,118

Table 4. Mean cerebrospinal fluid pressure (mm Hg) by age categories.

CSF Pressure by Age Category			
Age Category	Statistic	P-Value*	Percent Lower**
20–49	N	4592	
	Mean (SD)	11.49 (2.75)	
	Min/Median/Max	4.41, 11.47, 18.38	
50–54	N	1402	
	Mean (SD)	11.20 (2.67)	0.001 2.5
	Min/Median/Max	4.41, 11.03, 18.38	
55–59	N	1435	
	Mean (SD)	11.09 (2.71)	<0.001 3.5
	Min/Median/Max	4.41, 11.03, 18.38	
60–64	N	1150	
	Mean (SD)	10.68 (2.60)	<0.001 7.0
	Min/Median/Max	4.41, 10.59, 18.38	
65–69	N	1269	
	Mean (SD)	10.46 (2.60)	<0.001 9.0
	Min/Median/Max	4.41, 10.29, 18.38	
70–74	N	1084	
	Mean (SD)	10.08 (2.51)	<0.001 12.3
	Min/Median/Max	4.41, 10, 18.38	
75–79	N	781	
	Mean (SD)	9.93 (2.62)	<0.001 13.6
	Min/Median/Max	4.41, 9.85, 18.24	
80–84	N	318	
	Mean (SD)	9.96 (2.72)	<0.001 13.3
	Min/Median/Max	4.56, 9.71, 18.38	
85–89	N	79	
	Mean (SD)	9.26 (3.05)	<0.001 19.4
	Min/Median/Max	4.71, 8.68, 18.38	
90–95	N	11	
	Mean (SD)	8.40 (2.35)	<0.001 26.9
	Min/Median/Max	5.29, 8.24, 13.24	

Percent difference is compared against the 20–49 year age group.
 *P-value based on F-test of difference in CSF means between each older age category and age category 20–49.
 **Percentage lower of mean in age category than mean in 20–49 age category.
 doi:10.1371/journal.pone.0052664.t004

subjects, a dataset at least two orders of magnitude larger than previous reports.

Prospective and retrospective studies have shown that BMI is positively associated with CSFP in different populations [17–19]. In a clinical interventional study of 71 patients in China, Ren and colleagues identified a positive correlation between BMI and CSFP [18]. These investigators did not find a significant

Table 5. Bivariate analysis of mean cerebrospinal fluid pressure (mm Hg) as a function of BMI within specific age groups.

Age	N	Mean CSFP	BMI	N	Mean CSFP
20–42	861	11.56	10.1–22.3	241	9.92
			22.3–24.8	167	11.01
			24.8–27.5	148	11.78
			27.5–30.8	148	12.59
42–51	767	11.31	30.8–49.1	157	13.49
			10.1–22.3	156	9.37
			22.3–24.8	154	10.76
			24.8–27.5	143	11.16
51–60	953	11.04	27.5–30.8	127	12.28
			30.8–49.1	187	12.82
			10.1–22.3	154	9.09
			22.3–24.8	197	10.00
60–69	796	10.61	24.8–27.5	170	10.74
			27.5–30.8	200	11.78
			30.8–49.1	232	12.81
			10.1–22.3	139	8.60
69–91	937	9.84	22.3–24.8	145	9.60
			24.8–27.5	172	10.52
			27.5–30.8	181	11.26
			30.8–49.1	159	12.63
69–91	937	9.84	10.1–22.3	169	8.18
			22.3–24.8	202	9.19
			24.8–27.5	227	10.07
			27.5–30.8	210	10.50
69–91	937	9.84	30.8–49.1	129	11.57

doi:10.1371/journal.pone.0052664.t005

relationship between lower CSFP with age. In the present study, we found that BMI and age independently influence CSFP. However, whereas CSFP declines beginning in the 6th decade of life, BMI is positively associated with CSFP in every age group we studied. A biomechanical explanation for the relationship between cerebrospinal fluid pressure and body mass index has been suggested for patients with idiopathic intracranial hypertension (IIH). Studies of IIH suggest that obesity, in particular central obesity, increases intra-abdominal pressure, which ultimately causes an increase in venous pressure and consequently intracranial pressure [20,21]. It is plausible that a similar mechanism is in play in the entire spectrum of BMI. Interestingly, it has also been reported that lower BMI is a possible risk factor for glaucoma, or conversely, increased BMI may be protective [22–24].

We have found that CSFP is higher in males compared to females within all age groups after controlling for BMI and age. Although males consistently have higher CSFP values, the model was unable to pronounce the relationship between advancing age and sex. If lower CSFP is a risk factor for POAG, this would suggest that women are at increased risk for POAG. However most population-based studies have found little difference in prevalence and incidence between sexes [25–27]. In fact, Rudnicka and colleagues' meta-analysis of population-based studies concluded that, in general, men have a higher prevalence of POAG than women (OR = 1.37 [1.22, 1.53]) [28,29]. On the




Figure 2. Segmented multivariate linear regression model CSFP (mm Hg) by age and sex. For each year of increase in age, CSFP decreased 0.009 mmHg ($p=0.132$) for age <50 and by 0.056 mmHg, ($p<0.001$) for age greater than 50. Prior to age 50, the slope is not different between males and females ($p=0.369$). After age 50, the gender difference in the slope is borderline significant ($p=0.067$). The r^2 of this model is 0.071.

doi:10.1371/journal.pone.0052664.g002

other hand, Drance reported that female patients with normal tension glaucoma have a more rapid progression than males [30]. It may be that the effect of sex on CSFP, 6.5% lower mean CSFP in women in this study, is too small in existing populations-based studies. It is also possible that other variables, such as hormonal influences, exert an effect on POAG risk [23].

We examined the effect of race and CSFP. Mean CSFP appeared to be higher in Black patients compared to Caucasians. However, the number of Black patients in this dataset is small (1.4% of the study population) and is consequently inadequate to draw a meaningful conclusion on this important topic. For similar reasons, the effect of the time lumbar puncture is performed, mean arterial pressure, and diabetes mellitus on CSFP are not possible.

IOP and Age




Figure 3. Average intraocular pressure against age. Formula for the regression: $y = -0.0144x + 15.575$, $r^2 = 0.003$.

doi:10.1371/journal.pone.0052664.g003

Table 6. Intraocular pressure, cerebrospinal fluid pressure, and translaminar pressure difference by age groups.

Age	IOP AVG	CSFP	N	TLPd	Difference*	P value
20–49	14.9	10.5	178	4.4		
50–59	14.9	11.3	106	3.6	0.41%	0.884
60–69	14.7	9.9	75	4.8	1.39%	0.746
70–79	14.7	8.3	65	6.4	1.43%	0.726
>80	13.8	7.6	17	6.2	7.36%	0.193

The percentage difference is calculated by mean IOP average of each age group against mean IOP average at age 20–49; significance measured by two-tailed Student's t-test. IOP: intraocular pressure; CSFP: cerebrospinal fluid pressure; TLPd: translaminar pressure difference.

doi:10.1371/journal.pone.0052664.t006

Well-powered studies of this type, with a particular focus on the effect of race would be of great interest.

We have found that CSFP progressively decreases after age 50. This suggests that either the resistance to CSFP outflow is reduced or that CSF production decreases with age. Cerebrospinal fluid is produced in the choroid plexus within the third, fourth, and inferior horn of the lateral ventricles, and drains in the arachnoid villi into the cerebral venous system, as well as through lymphatic channels [31–35]. There is no reported evidence that CSF outflow resistance decreases with age, rather most studies report CSF flow resistance increases [34,36]. However, there is evidence that the choroid plexus undergoes aging changes leading to decreased CSF production [37–39].

Levels of vasopressin, a hormone that regulates the choroid plexus, is affected by aging [40]. Vasopressin receptor (V1) activation in the choroid plexus has been shown to decrease blood flow and reduce CSF secretion. Vasopressin-secreting neurons show increased activity with aging. Vasopressin levels are elevated in the CSF of old rats and in plasma of older humans [41]. May and colleagues examined the rate of CSF production in 7 young and 7 elderly healthy normal volunteers [37]. Although opening CSF pressures were similar between the two groups, the mean CSF production was reduced by over 50% in the older age group (0.19 ± 0.07 versus 0.41 ± 0.24 ml/min, $p < 0.02$). These data support the role of reduced CSF production rather than increased drainage as the most likely cause of lower CSFP with age.

We hypothesize that the decline in CSFP with age in conjunction with essentially stable IOP would produce an age-related increase in the translaminar pressure gradient. These findings carry potentially major implications for risk of POAG with age. A study by Morgan et al found that the mechanical effect of altering CSFP on the lamina cribrosa is equivalent to or greater than altering IOP [42,43]. Assuming equivalence for IOP and CSFP on the translaminar pressure gradient, we can examine the effect of a similar rise in IOP on POAG risk using data provided by population-based studies. In the Baltimore Eye Survey the relative risk of POAG increased from 2.8 for an IOP group of 19–21 mmHg to 12.8 for IOPs of 22–24 mmHg, a 4 fold increase in risk produced by an average 3 mmHg increase in IOP [44]. The Beaver Dam Eye Study reported that patients with glaucoma had an average IOP of 20.1 mmHg compared to 15.3 mmHg in unaffected controls, which represents a mean difference of approximately 5 mmHg [45]. The Tajimi Eye Study [46] is of particular interest since over 90% of untreated POAG cases had an IOP at diagnosis within the normal range. These investigators found that a rise of 4 mmHg increased POAG risk 2 fold even

within the normal IOP range. These studies support the notion that relatively small increases in IOP and therefore the translaminar pressure gradient, similar to those induced by age on CSFP observed in this study, may be associated with substantial changes in relative risk for POAG.

The time frame for age-dependent reduction in CSFP mirrors that seen for the rise in POAG prevalence in population-based studies performed in the United States and globally. In the United States, the Baltimore Eye Survey, Beaver Dam Eye Study, and the Los Angeles Latino Eye Study have all found that POAG prevalence begins to rise in the 5th and 6th decades and progressively increases through the 8th or 9th decades [1,44,45,47–49]. Studies conducted in Australia, Europe, China, and Japan describe similar trends [50–59]. Taken together the hypothesized effect of a reduction of CSFP with age is compatible with observed increases in the prevalence of POAG that are consistently observed in populations throughout the world. This hypothesis may be particularly applicable in the Hispanic and Japanese populations where the great majority, 80% and 92%, respectively, of those with POAG have a normal IOP at the time of diagnosis. Interestingly, a genome-wide association study of POAG identified a genetic locus on chromosome 8q22 that is strongly associated with normal pressure glaucoma and contains putative regulatory sites that are active in choroid plexus epithelial cells, the cell type responsible for CSF production [60].

We recognize several limitations in this study. Although the dataset is robust in size this is still subject to the challenges encompassed by a retrospective analysis. While diagnoses were available for each of these patients, it was not possible to review the actual individual medical records for accuracy, although studies of accuracy between ICD-9 codes and actual medical records have demonstrated acceptable congruency [61,62]. Some information from the CSF analysis, including cell counts and differentials, were not available so it is possible that subjects with abnormal CSF parameters were inadvertently included in the analysis. Similarly, although data were carefully reviewed to exclude conditions that affect CSFP, it is possible that some subjects developed conditions for which they would have been excluded at a later date or some conditions may exert a currently unknown influence on CSFP.

Another important consideration is the interpretation of the values reported in this study. Lenfeldt found that CSFP measured by lumbar puncture accurately represents intracranial pressure as long as the patient is in the lateral decubitus position [63]. However, the intracranial pressure differs from that within the orbital CSF space due to differences in fluid dynamics. Furthermore, altering position, for example upright to supine posture, induces substantial changes in CSFP [64,65]. These positional CSFP changes occur routinely, as do changes from diurnal variation.

This study utilized a “snapshot” in time for a single individual so these values do not reflect CSFP as a function of time for any particular patient or group of patients. Additionally, this study did not discriminate between bedside lumbar punctures in the lateral decubitus position and fluoroscopy-guided lumbar punctures which may account for minor differences in pressures. Despite these limitations, we feel that the robust patient dataset compensates for many of these limitations.

Finally, the coefficient of determination (r^2) in the majority of the regression models indicates that the group of predictors chosen (age, sex, BMI, race) together are only modestly predictive of CSFP. This highlights the need to investigate and identify other variables that may impact CSFP.

In conclusion, these data support the hypothesis that CSFP may play a role in the pathogenesis of POAG through its contribution to translaminal pressure. CSFP decreases significantly and steadily after age 50. The observed change in CSFP with age parallels the rise in prevalence of POAG and is of a magnitude that is sufficient to produce a significant increased risk for glaucoma. This finding offers a previously unrecognized mechanism to help explain age-related risk for POAG in general, and for the large number of patients with statistically normal IOP in particular. If cerebrospinal fluid pressure is in fact an important factor in the pathogenesis of glaucoma, this could possibly introduce new avenues for therapeutics, assessment of glaucoma risk and estimation of progression.

References

- Francis BA, Varma R, Vigen C, Lai MY, Winarko J, et al. (2011) Population and High-Risk group screening for glaucoma: The Los Angeles Latino Eye Study. *Invest Ophthalmol Vis Sci* 52(9):6257–6264
- Herndon LW, Weizer JS, Stinnett SS (2004) Central corneal thickness as a risk factor for advanced glaucoma damage. *Arch Ophthalmol* 122:17–21
- Gordon MO, Beiser JA, Brandt JD, Heuer DK, Higginbotham EJ, et al. (2002) The Ocular Hypertension Treatment Study: baseline factors that predict the onset of primary open-angle glaucoma. *Arch Ophthalmol* 120:714–720
- Wu RY, Zheng YF, Wong TY, Cheung CY, Loon SC, et al. (2011) Relationship of central corneal thickness with optic disc parameters: The Singapore Malay Eye Study. *Invest Ophthalmol Vis Sci* 52(3):1320–1324
- Saenz-Frances F, Garcia-Feijo J, Janez L, Borrego-Sanz L, Martinez de la Casa JM, et al. (2011) Comparing corneal variables in healthy subjects and patients with primary open-angle glaucoma. *Invest Ophthalmol Vis Sci* 52(6):3683–3688
- Berdahl JP, Allingham RR, Johnson DH (2008) Cerebrospinal fluid pressure is decreased in primary open-angle glaucoma. *Ophthalmology* 115:763–8
- Berdahl JP, Fautsch MP, Stinnett SS, Allingham RR (2008) Intracranial pressure in primary open angle glaucoma, normal tension glaucoma, and ocular hypertension: a case-control study. *Invest Ophthalmol Vis Sci* 49:5412–8
- Ren R, Jonas JB, Tian G, Zhen Y, Ma K, et al. (2010) Cerebrospinal Fluid Pressure in Glaucoma: A Prospective Study. *Ophthalmology* 117:259–266
- Ren R, Zhang X, Wang N, Li B, Tian G, Jonas JB (2011) Cerebrospinal fluid pressure in ocular hypertension. *Acta Ophthalmol* 89(2):e142–8
- Gilland O, Tourtellote WW, O'Tauma L, Henderson WG (1974) Normal cerebrospinal fluid pressure. *J Neurosurg* 40(5):587–93.
- Malm J, Jacobsson J, Birgander R, Eklund A (2011) Reference values for CSF outflow resistance and intracranial pressure in healthy elderly. *Neurology* 76(10):903–9.
- Masserman JH (1935) Cerebrospinal hydrodynamics. VI. Correlations of the pressure of the cerebrospinal fluid with age, blood pressure and the pressure index. *Arch Neurol Psychiatry* 34(3):564–566.
- Fishman R (1992) "Pseudotumor cerebri and related disorders." *Cerebrospinal Fluid in Disease of the Nervous System*. 2nd edition. Philadelphia, PA: W. B. Saunders Company: 71–151.
- World Health Organization (1995) "Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee." *WHO Technical Report Series: 854*. Geneva, World Health Organization.
- Czosnyka M, Czosnyka ZH, Whitfield PC, Donovan T, Pickard JD (2001) Age dependence of cerebrospinal pressure-volume compensation in patients with hydrocephalus. *J Neurosurg* 94(3):482–6.
- Ekstedt J (1978) CSF hydrodynamic studies in man. 2. Normal hydrodynamic variables related to CSF pressure and flow. *J Neurol Neurosurg Psychiatry* 41(4):345–53.
- Fleischman D, Allingham RR, Berdahl J, Fautsch M (2011) Body mass, spinal fluid, and glaucoma. *Ophthalmology* 118(6):1225–6; author reply 1226.
- Ren R, Wang N, Zhang X, Tian G, Jonas JB (2012) Cerebrospinal fluid pressure correlated with body mass index. *Graefes Arch Clin Exp Ophthalmol* 250(3):445–6
- Berdahl JP, Fleischman D, Zaydlarova J, Stinnett SS, Allingham RR, Fautsch MP (2012) Body Mass Index has a linear relationship with Cerebrospinal Fluid Pressure. *Invest Ophthalmol Vis Sci* 53(3):1422–7.
- Karahalios DG, ReKate HL, Khayata MH, Apostolides PJ (1996) Elevated intracranial venous pressure as a universal mechanism in pseudotumor cerebri of varying etiologies. *Neurology* 46(1):198–202
- Sugerman HJ, DeMaria EJ, Felton WL, Nakatsuka M, Sismanis A (1997) Increased intra-abdominal pressure and cardiac filling pressures in obesity-associated pseudotumor cerebri. *Neurology* 49(2):507–11
- Leske MC, Connell AM, Wu SY, Hyman LG, Schachat AP (1995) Risk factors for open-angle glaucoma. The Barbados Eye Study. *Arch Ophthalmol* 113:918–24.
- Pasquale LR, Willett WC, Rosner BA, Kang JH (2010) Anthropometric Measures and their Relation to Incident Primary Open-Angle Glaucoma. *Ophthalmology* 117:1521–9.
- Ramdas WD, Wolfs RC, Hofman A, de Jong PT, Vingerling JR, et al. (2011) Lifestyle and risk of developing open-angle glaucoma: The Rotterdam Study. *Arch Ophthalmology* 129(6):767–72
- Tielsch JM, Sommer A, Katz J, Royall RM, Quigley HA, et al. (1991) Racial variations in the prevalence of primary open-angle glaucoma: The Baltimore Eye Survey. *JAMA* 266(3):369–374.
- Nizankowska MH, Kaczmarek R (2005) Prevalence of glaucoma in the Wroclaw population. The Wroclaw epidemiological study. *Ophthalmic Epidemiol* 12(6):363–71.
- Nizankowska MH, Kaczmarek R (2004) Prevalence of open angle glaucoma and ocular hypertension as a risk factor for primary open angle glaucoma in Wroclaw population: Wroclaw Epidemiology Study. *Klin Oczna* 106 (1–2 Suppl):147–52
- Rudnicka AR, Mt-Isa S, Owen CG, Cook DG, Ashby D (2006) Variations in Primary Open-Angle Glaucoma Prevalence by Age, Gender, and Race: A Bayesian Meta-Analysis. *Invest Ophthalmol Vis Sci* 47(10):4254–4261
- Leske MC (2007) Open-Angle Glaucoma-An epidemiologic overview. *Ophthalmic Epidemiology* 14:166–172
- Drance S, Anderson DR, Schulzer M (2001) Risk factors for progression of visual field abnormalities in normal-tension glaucoma. *Am J Ophthalmol* 131:699–708
- McComb GJ, Hyman S (1984) Lymphatic drainage of cerebrospinal fluid in the cat. In: *Hydrocephalus*. Eds: Shapiro K, Marmarou A, Portnoy H. New York: Raven Press: 83–97
- McComb GJ, Hyman S (1990) Lymphatic drainage of cerebrospinal fluid in the primate. In: *Pathophysiology of the Blood-Brain Barrier*. Eds: Johansson BB, Widner H. New York: Elsevier Science Publishers: 421–37
- Zakharov A, Papaiconomou C, Koh L, Djencic J, Bozanic-Sosic R, et al. (2004) Integrating the roles of extracranial lymphatics and intracranial veins in cerebrospinal fluid absorption in sheep. *Microvasc Res* 67: 96–104
- Nagra G, Johnston MG (2007) Impact of ageing on lymphatic cerebrospinal fluid absorption in the rat. *Neuropathology and Applied Neurobiology* 33: 684–691
- Papaiconomou C, Zakharov A, Azizi N, Djencic J, Johnston M (2004) Reassessment of the pathways responsible for cerebrospinal fluid absorption in the neonate. *Childs Nerv Syst* 20: 29–36
- Albeck MJ, Skak C, Nielsen PR, Olsen KS, Borgesen SE, et al. (1998) Age dependency of resistance to cerebrospinal fluid outflow. *J Neurosurg* 89: 275–8
- May C, Kaye JA, Atack JR, Schapiro MB, Friedland RP, et al. (1990) Cerebrospinal fluid production is reduced in healthy aging. *Neurology* 40(3 Pt 1):500–3.
- Redzic ZB, Preston JE, Duncan JA, Chodobski A, Szymdynger-Chodobska J (2005) The choroid plexus cerebrospinal fluid system: from development to aging. *Curr Top Dev Biol* 71: 1–52
- Serot JM, Bene MC, Faure GC (2003) Choroid plexus, aging of the brain, and Alzheimer's disease. *Front Biosci* 8:s1515–21
- Faraci FM, Mayhan WG, Heisted DD (1990) Effect of vasopressin on cerebrospinal fluid production: possible role of vasopressin V1 receptors. *Am J Physiol* 258:R94–99.
- Frolkis VV, Kvmititskaia-Ryzhova T, Dubiley TA (1999) Vasopressin, hypothalamo-neurohypophyseal system and ageing. *Arch Geront Ger* 29:109–214.
- Morgan WH, Yu D, Alder VA, Cringle SJ, Cooper RL, et al. (1998) The Correlation between Cerebrospinal Fluid Pressure and Retrolaminar Tissue Pressure. *Invest Ophthalmol Vis Sci* 39:1419–1428
- Morgan WH, Chauhan BC, Yu DY, Cringle SJ, Alder VA, et al. (2002) Optic disc movement with variations in intraocular and cerebrospinal fluid pressure. *Invest Ophthalmol Vis Sci* 43(10):3236–42

Acknowledgments

We would like to thank Mr. Francis Aning for his help in extracting data from the medical records system at Mayo Clinic.

Author Contributions

Conceived and designed the experiments: DF JPB MPF RRA. Performed the experiments: DF JZ. Analyzed the data: DF JPB JZ SSS MPF RRA. Contributed reagents/materials/analysis tools: DF SSS. Wrote the paper: DF JPB MPF RRA.

44. Sommer A, Tielsch JM, Katz J, Quigley HA, Gottsch JD, et al. (1991) Relationship between intraocular pressure and primary open angle glaucoma among White and Black Americans. *Arch Ophthalmol* 109:1090–1095
45. Klein BE, Klein R, Linton KL (1992) Intraocular Pressure in an American Community. The Beaver Dam Eye Study. *Invest Ophthalmol Vis Sci* 33(7):2224–8
46. Iwase A, Suzuki Y, Araie M, Yamamoto T, Abe H, et al. (2004) The prevalence of primary open-angle glaucoma in Japanese. The Tajimi Study. *Ophthalmology* 111:1641–8.
47. Klein BE, Klein R, Sponsel WE, Franke T, Cantor LB, et al. (1992) Prevalence of glaucoma: the Beaver Dam Eye Study. *Ophthalmology*. 99:1499–1504.
48. Francis BA, Varma R, Chopra V, Lai MY, Shtir C, et al. (2008) Intraocular pressure, central corneal thickness, and prevalence of open-angle glaucoma: the Los Angeles Latino Eye Study. *Am J Ophthalmol*. 146(5):741–6.
49. Friedman DS, Eye Diseases Prevalence Research Group (2004) Prevalence of open-angle glaucoma among adults in the United States. *Arch Ophthalmol*. 122(4):532–8.
50. Quigley HA, Broman AT (2006) The number of people with glaucoma worldwide in 2010 and 2020. *Br J Ophthalmol*. 90(3):262–7
51. Mitchell P, Smith W, Attebo K, Healey PR (1996) Prevalence of open-angle glaucoma in Australia. The Blue Mountains Eye Study. *Ophthalmology* 103:1661–9.
52. Wensor MD, McCarty CA, Stanislavsky YL, Livingston PM, Taylor HR (1998) The prevalence of glaucoma in the Melbourne Visual Impairment Project. *Ophthalmology* 105:733–9.
53. Anton A, Andrada MT, Mujica V, Calle MA, Portela J, et al. (2004) Prevalence of primary open-angle glaucoma in a Spanish population. The Segovia Study. *J Glaucoma* 13:371–6.
54. Bonomi L, Marchini G, Marraffa M, Bernardi P, De Franco I, et al. (1998) Prevalence of glaucoma and intraocular pressure distribution in a defined population. The Egna-Neumarkt Study. *Ophthalmology* 105:209–15.
55. Cedrone C, Culasso F, Cesareo M, Zapelloni A, Cedrone P, et al. (1997) Prevalence of glaucoma in Ponza, Italy: a comparison with other studies. *Ophthalmic Epidemiology* 4:59–72.
56. Dielemans I, Vingerling JR, Wolfs RC, Hofman A, Grobbee DE, et al. (1994) The prevalence of primary open-angle glaucoma in a population-based study in The Netherlands. *Ophthalmology* 11:1851–5.
57. Foster PJ, Oen FT, Machin D, Ng TP, Devereux JG, et al. (2000) The prevalence of glaucoma in Chinese residents of Singapore. A cross-sectional population survey of the Tanjong Pagar District. *Arch Ophthalmol* 118:1105–11.
58. Shiose Y, Kitazawa Y, Tsukahara S, Akamatsu T, Mizokami K, et al. (1991) Epidemiology of glaucoma in Japan—a nationwide glaucoma survey. *Jap J Ophthalmol* 35:133–55.
59. Yoshida M, Okada E, Mizuki N, Kokaze A, Sekine Y, et al. (2001) Age-specific prevalence of open-angle glaucoma and its relationship to refraction among more than 60,000 asymptomatic Japanese subjects. *J Clin Epidemiol* 54:1151–8.
60. Wiggs JL, Yaspan BL, Hauser MA, Kang JH, Allingham RR, et al. (2012) Common variants at 9p21 and 8q22 are associated with increased susceptibility to optic nerve degeneration in glaucoma. *PLoS Genet*. 8(4):e1002654.
61. Dixon J, Sanderson C, Elliot P, Walls P, Jones J, et al. (1998) Assessment of the reproducibility of clinical coding in routinely collected hospital activity data: a study in two hospitals. *J Public Health Med*. 20(1):63–9
62. Campbell SE, Campbell MK, Grimshaw JM, Walker AE (2001) A systematic review of discharge coding accuracy. *J Public Health Med* 23(3):205–11
63. Lenfeldt N, Koskinen LOD, Bergenheim AT, Malm J, Eklund A (2007) CSF pressure assessed by lumbar puncture agrees with intracranial pressure. *Neurology*. 68:155–158
64. Magnaes B (1976) Body position and cerebrospinal fluid pressure. Part 1: clinical studies on the effect of rapid postural changes. *J Neurosurg* Jun;44(6):687–97
65. Magnaes B (1976) Body position and cerebrospinal fluid pressure. Part 2: clinical studies on orthostatic pressure and the hydrostatic indifferent point. *J Neurosurg* Jun;44(6):698–705