



NIH PUBLIC ACCESS

Author Manuscript

J Org Chem. Author manuscript; available in PMC 2013 February 3.


Published in final edited form as:

J Org Chem. 2012 February 3; 77(3): 1449–1456. doi:10.1021/jo202322k.

Chemoenzymatic synthesis of Uridine Diphosphate-GlcNAc and Uridine Diphosphate-GalNAc analogs for the preparation of unnatural glycosaminoglycans

Sayaka Masuko[†], Smritilekha Bera[†], Dixy E. Green[‡], Michel Weïwer[§], Jian Liu[□], Paul L. DeAngelis[‡], and Robert J. Linhardt^{†,*}[†]Department of Chemistry and Chemical Biology, Center for Biotechnology and Interdisciplinary Studies, Rensselaer Polytechnic Institute, Troy, NY 12180, USA[‡]Department of Biochemistry and Molecular Biology, University of Oklahoma Health Sciences Center, Oklahoma Center for Medical Glycobiology, 940 Stanton L. Young Blvd., Oklahoma City, OK 73126, USA[□]Division of Medicinal Chemistry and Natural Products, Eshelman School of Pharmacy, University of North Carolina, Chapel Hill, NC 27599[§]Broad Institute of Harvard and MIT, Cambridge, MA 02142, USA

Abstract



Eight *N*-acetylglucosamine-1-phosphate and *N*-acetylgalactosamine-1-phosphate analogs have been synthesized chemically and were tested for their recognition by the GlmU uridylyltransferase enzyme. Among these, only substrates that have an amide linkage to the C-2 nitrogen were transferred by GlmU to afford their corresponding uridine diphosphate(UDP)-sugar nucleotides. Resin-immobilized GlmU showed comparable activity to non-immobilized GlmU and provides a more facile final step in the synthesis of an unnatural UDP-donor. The synthesized unnatural UDP-donors were tested for their activity as substrates for glycosyltransferases in the preparation of unnatural glycosaminoglycans *in vitro*. A subset of these analogs was useful as donors, increasing the synthetic repertoire for these medically important polysaccharides.

1. Introduction

Glycosaminoglycans (GAGs) are a family of polysaccharides that are prevalent both as free glycans and in the core structures of glycans in glycoproteins¹ and glycolipids,² affecting cell–cell interactions during various metabolic processes.³ They are composed of repeating disaccharide units containing hexosamine residues; the other part of the repeat is often an uronic acid residue. *N*-Acetylglucosamine (GlcNAc) is found in heparosan, heparin, heparan sulfate, and hyaluronan. *N*-acetylgalactosamine (GalNAc) is found in chondroitin sulfate and dermatan sulfate.

^{*}Corresponding author linhar@rpi.edu.

Supporting Information

¹H and ¹³C NMR spectra for all compounds are available in the Supporting Information. This material is available free of charge via the Internet at <http://pubs.acs.org/>.

Monosaccharides in the form of UDP-nucleotides act as donors being transferred to a glycosyl acceptor by a glycosyltransferase or a synthase in the polysaccharide biosynthetic pathway.⁴ Natural and unnatural GAGs can be prepared in the laboratory⁵ by using recombinant GAG synthases and chemically prepared UDP-sugar nucleotides. With the advent of recombinant technology, glycosyltransferases and synthases have now become readily available. A major limitation to the enzymatic preparation of GAG polysaccharides and oligosaccharides is the limited availability and cost of sugar nucleotides and of sugar nucleotide analogs and derivatives due to the difficulty of their syntheses.⁶ Though they can be prepared in several ways, a purely chemical approach requires a long and tedious synthesis and yields of UDP-donor are generally low. Enzymatic synthesis offers an alternative to the use of protection and deprotection steps required for chemical synthesis⁷ and circumvents the difficulties inherent to the formation of a pyrophosphate bond.

Wang *et al.* reported the enzymatic synthesis of UDP-GlcNAc/UDP-GalNAc analogs using recombinant *Escherichia coli* *N*-acetylglucosamine-1-phosphate uridyltransferase (GlmU).⁸ However they reported low yields, especially for *N*-modified GalNAc-1-phosphate analogs and later turned to recombinant human UDP-GalNAc pyrophosphorylase (AGX1) for the synthesis of UDP-GalNAc analogs.⁹ More recently, Chen *et al.* reported a one-pot three-enzyme synthesis of UDP-GlcNAc derivatives using an *N*-acetylhexosamine 1-kinase and GlmU but only a few derivatives were successfully synthesized.¹⁰ Here we report the chemical synthesis of eight different GlcNAc-1-phosphate and GalNAc-1-phosphate analogs and their recognition by the GlmU enzyme (Figure 1).

2. Results and Discussion

2.1 Synthesis of *N*-acetylglucosamine-1-phosphate and *N*-acetylgalactosamine-1-phosphate analogs

Eight natural and unnatural *N*-acetylglucosamine-1-phosphates and *N*-acetylgalactosamine-1-phosphates were chemically synthesized from commercially available monosaccharides. Our first attempt aimed at synthesizing GlcNH₂-1-phosphate, since the amine group can be easily functionalized at a later step. As shown in Scheme 1, reactions of glucosamine hydrochloride (**1**) with *in situ* generated TfN₃ afforded 2-azido sugar **2** in 89% yield, which was subsequently peracetylated.¹¹ MacDonald phosphorylation¹² of the protected 2-azido sugar gave monophosphate **4** in a modest 46% yield. Subjecting **4** to Zemplén conditions removed its acetyl groups, affording GlcN₃-1-phosphate (**5**). Finally, reduction of **5** by hydrogenation using the Lindlar's catalyst¹³ afforded GlcNH₂-1-phosphate (**6**) in quantitative yield.

Next, GlcNAc-1-phosphate (**7**), the natural substrate of GlmU, was chemically synthesized from **6** through *N*-acetylation using Ac₂O in MeOH/H₂O (1/3) followed by treatment with MeOH/H₂O/Et₃N (2/2/1) to remove any *O*-acetyl groups that had been introduced (Scheme 2). Next, GalNAc-1-phosphate (**9**), an unnatural substrate of GlmU, was synthesized to investigate the impact of the C4 configuration on GlmU recognition. Compound **9**, a C4 epimer of **7**, was prepared from the commercially available galactosamine (**8**).

Unnatural UDP-GlcNAc and UDP-GalNAc analogs having a 'clickable'¹⁴ alkyne group were also synthesized. In our initial approach, we attempted to functionalize glucosamine hydrochloride in the first step with propiolic acid under several different coupling conditions, but this led to low yields and separation difficulties. Instead, the stepwise synthesis of GlcN-alkyne-1-phosphate was undertaken as shown in Scheme 3. 2-Azido-*D*-glucose tetraacetate (**3**) was reduced by hydrogenation with Lindlar's catalyst to afford **10**. The amino sugar **10** was then acylated using propiolic anhydride, prepared by treating propiolic acid with DCC in DCM, and affording the alkyne derivative in 70% yield.

MacDonald phosphorylation¹² followed by de-*O*-acetylation provided GlcN-alkyne-1-phosphate (**13**). GalN-alkyne-1-phosphate (**18**) was similarly synthesized using CBz protection of the amino group (Scheme 4):

Next, a bioisostere of GlcNAc-1-phosphate bearing an *N*-trifluoroacetamide was synthesized (Scheme 5). The *N*-trifluoroacetyl (TFA) group represents an excellent protecting group in the chemoenzymatic synthesis of GAGs,¹⁵ as it can undergo selective deprotection by mild base treatment and then can be readily *N*-sulfonated to afford heparin and heparan sulfate type glycans. First, *D*-glucosamine hydrochloride (**1**) was protected as an *N*-trifluoroacetyl by treatment of CF₃CO₂Et¹⁶ followed by peracetylation. Phosphorylation¹² followed by Zemplén deprotection afforded GlcN-TFA-1-phosphate (**22**).

A ‘thiol-clickable’¹⁷ analog, with an *N*-ene, was next synthesized from commercially available tetra-*O*-acetyl glucosamine (**23**). The per-*O*-acetylated sugar was chosen as the starting material, as the unprotected monosaccharide was prone to polymerization upon treatment with acryloyl chloride. Thus, treatment of the tetraacetate with acryloyl chloride introduced an ene group to afford compound **24**. Phosphorylation proceeding in low yield was followed by deacetylation to give monophosphate **26**.

2.2 Preparation of the UDP-Sugar Nucleotides

After the successful synthesis of a small library of GlcNAc-1-phosphate and GalNAc-1-phosphate analogs, each substrate was tested for its recognition by GlmU from *Escherichia coli*.¹⁵ The enzymatic reaction was performed in Tris buffer with MgCl₂, DTT, UTP, and inorganic pyrophosphatase at 30 °C for 3 h (2 days for UDP-GalNAc analogs).¹⁸ The results are shown in Table 1.

Out of the eight analogs tested, six were accepted as substrates by GlmU, affording the corresponding UDP-sugar nucleotides in high yields. Neither GlcN₃-1-phosphate (**5**) nor GlcNH₂-1-phosphate (**6**) was accepted by GlmU. GlcNAc-1-phosphate (**7**) and GlcN-TFA-1-phosphate (**22**) were excellent substrates, yielding more than 70% conversion. Since GalNAc-1-phosphate (**9**) was accepted by GlmU, the 4-OH group appears not to play a critical role in enzyme recognition. Both GlcN-alkyne-1-phosphate (**13**) and GalN-alkyne-1-phosphate (**18**) gave moderate yields of UDP-sugars using GlmU. Based on these observations, it appears that the amide bond on the 2-amino functional group is essential for substrate recognition by GlmU but the configuration at C4 is not important.

Immobilization of (His)₆-tagged enzymes on Ni beads has been useful in carbohydrate synthesis.^{19, 20} Since the reactions above using the soluble enzyme showed promising yields, we examined the immobilization of GlmU for future applications in large-scale synthesis. Taking advantage of the C-terminal (His)₆-tag, GlmU was immobilized on Ni Sepharose beads that were then added to the reaction mixture using **22** as the substrate. Immobilized GlmU gave high yields of UDP-GlcN-TFA, comparable to that of the solution-based reaction. The same beads could be recycled and reused for a second reaction, and retained high enzymatic activity. Recycling the enzyme through its immobilization avoids the requirement of repeated enzyme expression, and can allow for continuous processes and eases the purification of UDP-sugar nucleotide product.

Although UDP-GlcN₃ and UDP-GlcNH₂ could not be synthesized using GlmU, they were chemically synthesized from the aforementioned GlcN₃-1-phosphate (**5**) (Scheme 7). Compound **5** was converted to its pyridinium salt, then stirred with UMP-morpholidate²¹ in pyridine for 5 days to afford UDP-GlcN₃ (**33**). The azide group was then reduced to an amine by hydrogenation using Lindlar’s catalyst, affording UDP-GlcNH₂ (**34**). This free

amino group was then functionalized to afford the following unnatural UDP-GlcNAc analogs: UDP-GlcN-biotin (**35**), UDP-GlcN-Fmoc (**36**), and UDP-GlcN-*t*Boc (**37**).

These ten UDP-sugar nucleotides were then tested for incorporation by various GAG synthases *in vitro*. PmHAS and PmCS produce hyaluronan and chondroitin polysaccharides, respectively.²² PmHS1 and PmHS2 are two homologous isozymes that both can polymerize heparosan, the precursor of heparan sulfate and heparin.⁵ The hexosamine analogs were tested in a chain polymerization assay using radioactively labeled UDP-[³H]GlcUA; a radioactive signal results only if the hexosamine is co-polymerized into the disaccharide repeats of a GAG product.^{5, 23}

The results shown in Table 2 demonstrate that some analogs function well with one or more synthases, but a few analogs did not incorporate into polymer under our conditions. In addition to increasing the chemical functionality of synthetic GAGs, the substrate specificity information yields insights into the synthases' recognition of their donor precursors.

Conclusions

In conclusion, we synthesized a small library of GlcNAc/GalNAc-1-phosphate analogs for their recognition by the GlmU uridyltransferase to synthesize UDP-GlcNAc/GalNAc analogs. Only analogs with an amide linkage at the C-2 nitrogen were successfully accepted by GlmU and gave the product in moderate to high yields. Analogues without an amide linkage were converted to UDP-sugar donors by chemical synthesis. The UDP-sugars were then tested for incorporation by GAG synthases using a radioactive assay, which showed that natural and unnatural UDP-sugars were incorporated into the polysaccharide. These results demonstrate that unnatural GAGs with a desired functionality can be synthesized *in vitro* and has promising therapeutic applications.

Experimental Section

General

¹H and ¹³C NMR spectra were recorded at 500 MHz for ¹H NMR, 125 MHz for ¹³C NMR, or 600 MHz for ¹H NMR, 150 MHz for ¹³C NMR, or 800 MHz (800 MHz for ¹H NMR, 200 MHz for ¹³C NMR) with Topsin 2.1 software. Mass data were acquired by high-resolution ESI-MS. Thin-layer chromatography (TLC) was carried out using plates of silica gel 60 with fluorescent indicator and revealed with UV light (254 nm) when possible and Von's reagent or ninhydrin solution in ethanol. Flash chromatography was performed using silica gel 230–400 mesh. Yields are given after purification, unless otherwise noted. When reactions were performed under anhydrous conditions, the mixtures were maintained under argon. Compounds were named following IUPAC rules as applied by Beilstein-Institute AutoNom (version 2.1) software for systematic names in organic chemistry.

Expression of GlmU enzyme

The procedures for the expression and purification of GlmU were described previously.²⁴ Briefly, GlmU was cloned into a PET 21b vector (Novagen) to form a C-terminal (His)₆-tagged fusion protein. The expression was carried out in BL21 star cells (Invitrogen). The protein was purified by a Ni-Sepharose 6 Fast Flow column (GE Health) following a standard procedure.

General procedure for the MacDonald phosphorylation

Crystalline phosphoric acid (8.1 mmol) was dried *in vacuo* over phosphorous pentoxide for 12 h. The 1-*O*-acetyl sugar derivative (1 mmol) was added and the mixture was heated at 60

°C *in vacuo*. After 2 h, heating was ceased and the resulting dark black mixture was dissolved in anhydrous THF (5 mL). The solution was cooled to 0 °C and concentrated ammonium hydroxide (0.5 mL) was added until ~pH 7. The precipitate of ammonium phosphate was filtered off and washed with THF (20 mL). The combined filtrate and washings were evaporated to give a syrupy residue that was purified by flash column chromatography (EtOAc 100% to EtOAc/MeOH 50/50).

General procedure for Zemplén deacetylation

Sodium methoxide (3 mmol) in MeOH (0.5 M) was added to the peracetylated phosphate (1 mmol) in MeOH (7 mL) at 0 °C. The reaction was stirred at 0 °C for 1 h and quenched with Amberlite IR-120 (H⁺). Amberlite was removed through filtration and the filtrate was concentrated *in vacuo* to afford the fully deprotected phosphate in quantitative yield.

General procedure for the enzymatic preparation of the UDP-sugar nucleotides

A reaction mixture containing the monophosphate, glucosamine-1-phosphate acetyltransferase/N-acetyl glucosamine-1-phosphate uridylyltransferase (GlmU), Tris-HCl (46 mM, pH 7.0), MgCl₂ (5 mM), dithiothreitol (200 μM), UTP (2.5 mM), inorganic pyrophosphatase (0.012 units/μL), from Sigma-Aldrich, was incubated at 30 °C for 3 h. For GalNAc substrates, a second portion of GlmU was added to the reaction mixture on the second day. The completion of the reaction was monitored by TLC (*n*-butanol/formic acid/methanol=4/8/1) or through MS. Then 3 reaction volumes of ethanol was added to quench the reaction and stored overnight at -20 °C. Centrifugation at 3000 rcf for 30 min afforded the crude product in the supernatant which was then concentrated and further purified on a BioGel (Bio-Rad) P2 column (2.5 × 65cm) and eluted with H₂O. Fractions were collected and those containing the product as determined by UV absorbance at 262 nm were combined and freeze-dried to afford the pure product.

2-azido-2-deoxy-D-glucopyranose-3,4,6-triacetate-1-dihydrogen phosphate (4)

—Brown oil; ¹H NMR (500 MHz, MeOD): δ (ppm) 5.73 (s, 1H), 5.49 (t, *J* = 9.5 Hz, 1H), 5.12 (t, *J* = 9.74 Hz, 1H), 4.36 (d, *J* = 10 Hz, 2H), 4.14 (m, 1H), 3.65 (d, *J* = 9 Hz, 1H), 2.08 (s, 3H), 2.04 (s, 6H); ¹³C NMR (125 MHz, MeOD): δ (ppm) 171.1, 170.2, 170.0, 93.4, 70.7, 68.4, 68.0, 61.4, 61.2, 61.1; HRMS-FAB: [M-H] calcd. for C₁₂H₁₇N₃O₁₁P⁻ 410.0606 Found: 410.0601.

2-azido-2-deoxy-D-glucopyranose-1-dihydrogen phosphate (5)

—Light brown amorphous solid; ¹H NMR (500 MHz, D₂O): δ (ppm) 5.54 (dd, 1H, *J* = 7.4 Hz, 3.1 Hz), 3.88-3.72 (m, 4H), 3.48 (t, 1H, *J* = 9.5 Hz), 3.29 (d, 1H, *J* = 2.5 Hz); ¹³C NMR (125 MHz, D₂O): δ (ppm) 94.0, 72.8, 70.7, 69.5, 62.8, 60.3; HRMS-FAB: [M-H] calcd. for C₆H₁₁N₃O₈P⁻ 284.0289 Found: 284.0289.

2-amino-2-deoxy-D-glucopyranose-1-dihydrogen phosphate (6)

—Lindlar's catalyst (20 mg) was added to a stirred solution of **5** (75 mg, 0.228 mmol) in MeOH/H₂O, 1/1 (10 mL). A hydrogen balloon was attached to the reaction flask and the reaction was stirred overnight at room temperature. The reaction mixture was filtered through Celite and rinsed with water. The filtrate and washings were combined and concentrated to afford **6** in quantitative yield.

¹H NMR (800 MHz, D₂O): δ (ppm) 5.59 (1H, dd, *J* = 7.4 Hz, 3.1 Hz), 3.90-3.84 (3H, m), 3.73 (1H, m), 3.45 (1H, m), 3.26 (1H, m); ¹³C NMR (200 MHz, D₂O): δ (ppm) 90.8, 72.1, 69.6, 69.4, 60.1, 54.4; HRMS-FAB: [M-H] calcd. for C₆H₁₃NO₈P⁻ 258.0384 Found: 258.0391.

2-deoxy-2-[(1-oxo-2-propyn-1-yl)amino]- β -glucopyranose-1,3,4,6-tetraacetate (11)—To a stirred solution of DCC (740 mg, 3.59 mmol) in dry CH_2Cl_2 (19.5 mL), propiolic acid (0.18 mL, 2.92 mmol) was slowly added at 0 °C. After 1h, **10** (1.267 g, 2.924 mmol), dissolved in dry CH_2Cl_2 (7 mL), and DIPEA was added dropwise. The reaction mixture was stirred for 1 h at 0 °C, then warmed to room temperature and then stirred for another 2 h before the solvents were removed *in vacuo*. The crude product was purified by silica gel chromatography (MeOH/ CH_2Cl_2 , 1/20) to yield **11** as a colorless amorphous solid (1.20 g, 70%). ^1H NMR (500 MHz, CDCl_3): δ (ppm) 6.01 (d, 1 H, $J = 9.4$ Hz), 5.69 (d, 1 H, $J = 8.7$ Hz), 5.14 (m, 2 H), 4.34 (m, 1 H), 4.27 (dd, 1 H, $J = 4.2, 12.5$ Hz), 4.12 (d, 1 H, $J = 12.5$ Hz), 3.81 (m, 1 H), 2.13-2.04 (4s, 12 H); ^{13}C NMR (125 MHz, CDCl_3): δ (ppm) 171.2, 170.7, 169.5, 169.4, 152.1, 92.0, 76.5, 74.7, 72.5, 72.2, 68.0, 61.6, 52.6, 20.8-20.5; HRMS-FAB: $[\text{M}+\text{Na}]$ calcd. for $\text{C}_{17}\text{H}_{21}\text{NNaO}_{10}^+$ 422.1063 Found: 422.1065.

2-deoxy-2-[(1-oxo-2-propyn-1-yl)amino]- β -glucopyranose-3,4,6-triacetate-1-dihydrogen phosphate (12)—Brown oil; ^1H NMR (500 MHz, CDCl_3): δ (ppm) 6.01 (d, $J = 9.4$ Hz, 1 H), 5.69 (d, $J = 8.7$ Hz, 1 H), 5.14 (m, 2 H), 4.34 (m, 1 H), 4.27 (dd, $J = 4.2, 12.5$ Hz, 1 H), 4.12 (d, $J = 12.5$ Hz, 1 H), 3.81 (m, 1 H), 2.13-2.04 (4s, 12 H); ^{13}C NMR (125 MHz, CDCl_3): δ (ppm) 171.2, 170.7, 169.5, 169.4, 152.1, 92.0, 76.5, 74.7, 72.5, 72.2, 68.0, 61.6, 52.6, 20.8-20.5; HRMS-FAB: $[\text{M}+\text{Na}]$ calcd. for $\text{C}_{17}\text{H}_{21}\text{NNaO}_{10}^+$ 422.1063 Found: 422.1065.

2-deoxy-2-[(1-oxo-2-propyn-1-yl)amino]- β -glucopyranose-1-dihydrogen phosphate (13)—Light brown amorphous solid; ^1H NMR (500 MHz, D_2O): δ (ppm) 5.32 (dd, $J = 3.3, 7.0$ Hz, 1 H), 3.87 (dt, $J = 2.6, 10.5$ Hz, 1 H), 3.72 (dd, $J = 2.4, 4.2$ Hz, 1 H), 3.70-3.64 (m, 3 H) 3.38 (dd, $J = 9.1, 10.0$ Hz, 1 H), 3.35 (s, 1 H); ^{13}C NMR (125 MHz, D_2O): δ (ppm) 152.5, 95.5, 77.8, 75.0, 73.2, 63.9, 62.9, 56.7, 56.6; HRMS-FAB: $[\text{M}-\text{H}]$ calcd. for $\text{C}_9\text{H}_{13}\text{NO}_9\text{P}^-$ 310.0328 Found: 310.0323.

2-deoxy-2-[(phenylmethoxy)carbonyl]amino]- β -galactopyranose-1,3,4,6-tetraacetate (15)—Galactosamine-HCl (5.0 g, 23 mmol) was dissolved in 130 mL of 0.5 M NaOMe in MeOH and stirred for 35 min. The reaction mixture was cooled to 0 °C, 3.23 mL (23.2 mmol) of Et_3N and 5.0 g (23 mmol) of benzyl chloroformate were added, and the mixture was stirred for 5 h. The thick suspension was concentrated and dissolved in 25 mL of Ac_2O and 50 mL of pyridine. A catalytic amount of *N,N*-(dimethylamino)pyridine (DMAP) was added, and the solution was stirred overnight. It was concentrated and coevaporated with toluene (6×25 mL). The resulting syrup was dissolved in 40 mL of CHCl_3 and washed with saturated CuSO_4 (2×20 mL), saturated NaHCO_3 (2×25 mL), and brine (1×25 mL). The combined aqueous layers were extracted with 40 mL of CHCl_3 , and the combined organic layers were dried over Na_2SO_4 . The crude product was concentrated and purified by silica gel chromatography eluting with a gradient of 200/1 to 20/1, $\text{CHCl}_3/\text{MeOH}$, to yield 6.3 g (52%) of a yellow syrup comprising a mixture of anomers. ^1H NMR (500 MHz, CDCl_3): δ 7.32 (m, 5 H), 6.22 (dd, 1 H, $J = 3.65$ Hz), 5.39 (m, 1 H), 5.12-4.99 (m, 3 H), 4.43 (m, 1 H), 4.12-3.98 (m, 3 H), 2.12 (s, 3 H), 2.09 (s, 3 H), 1.98 (s, 3 H), 1.89 (s, 3 H); ^{13}C NMR (125 MHz, CDCl_3): δ 170.4, 170.3, 170.3, 168.8, 155.4, 136.7, 128.5, 91.4, 77.3, 68.4, 68.0, 67.1, 66.7, 61.2, 48.6, 20.8, 20.5, 20.4. HRMS-FAB: $[\text{M}+\text{Na}]$ calcd. for $\text{C}_{22}\text{H}_{25}\text{NNaO}_{10}^+$ 486.1376 Found: 486.1379.

2-amino-2-deoxy- β -galactopyranose-1,3,4,6-tetraacetate hydrochloride (16)— ^1H NMR (500 MHz, CD_3OD): δ 6.37 (d, 1 H, $J = 3.6$ Hz), 5.50 (dd, 1 H, $J = 1.3, 3.2$ Hz), 5.36 (dd, 1 H, $J = 3.2, 8.3$ Hz), 4.43 (m, 1 H), 4.12 (d, 1 H, $J = 6.6$ Hz), 4.09 (d, 1 H, $J = 6.6$ Hz), 3.96 (dd, 1 H, $J = 3.6, 11.4$ Hz), 2.21 (s, 3 H), 2.14 (s, 3 H), 2.06 (s, 3 H), 1.89 (s, 3 H);

δ 169.8, 169.4, 168.8, 168.5, 91.6, 67.6, 65.6, 61.2, 55.4, 42.4, 19.1, 17.3, 15.8, 11.2.
HRMS-FAB: $[M+Na]$ calcd. for $C_{14}H_{21}NNaO_9^+$ 370.1114 Found: 370.1117.

2-deoxy-2-[(1-oxo-2-propyn-1-yl)amino]- β -galactopyranose-1,3,4,6-tetraacetate (17)—Colorless oil; 1H NMR (500 MHz, $CDCl_3$): δ 6.10 (d, 1 H, $J = 3.6$ Hz), 5.32 (m, 1 H), 5.09 (dd, 1 H, $J = 3.2, 11.6$ Hz), 4.56 (dd, 1 H, $J = 3.2, 11.6$ Hz), 4.16 (t, 1 H, $J = 6.6$ Hz), 3.97 (m, 2 H), 3.57 (m, 1 H), 2.93 (s, 1 H), 2.05 (s, 3 H), 2.04 (s, 3 H), 1.90 (s, 3 H), 1.89 (s, 3 H); ^{13}C NMR (125 MHz, $CDCl_3$): δ 170.5, 170.2, 169.4, 168.8, 152.3, 92.5, 77.2, 74.4, 71.5, 70.1, 66.2, 61.3, 49.4, 24.8-22.0; HRMS-FAB: $[M+Na]$ calcd. for $C_{17}H_{21}NNaO_{10}^+$ 422.1063 Found: 422.1068.

2-deoxy-2-[(1-oxo-2-propyn-1-yl)amino]- α - β -galactopyranose-1-(dihydrogen phosphate) (18)—Light brown amorphous solid; 1H NMR (500 MHz, CD_3OD): δ 5.32 (dd, 1 H, $J = 3.4, 7.2$ Hz), 4.17 (dt, 1 H, $J = 2.6, 10.8$ Hz), 4.00 (dd, 1 H, $J = 5.7, 6.3$ Hz), 3.86 (m, 1 H), 3.80 (dd, 1 H, $J = 3.1, 10.0$ Hz), 3.59 (m, 1 H), 3.34 (s, 1 H); ^{13}C NMR (125 MHz, CD_3OD): δ (ppm) 152.6, 91.4, 68.0, 67.9, 67.5, 67.2, 66.0, 61.9, 61.3; HRMS-FAB: $[M-H]$ calcd. for $C_9H_{13}NO_9P^-$ 310.0328 Found: 310.0321.

2-deoxy-2-[(trifluoroacetyl)amino]- β -glucopyranose (19)— Et_3N (3.50 mL, 25 mmol) and CF_3CO_2Et (3.8 mL, 32 mmol) were added to β -glucosamine hydrochloride (5.375 g, 25 mmol) in MeOH (15 mL). The reaction mixture was stirred at room temperature overnight. The solution was concentrated *in vacuo* and recrystallized in MeOH to afford *N*-trifluoroacetylated **19** ($\alpha/\beta = 1:0.8$) (5.26 g, 19.2 mmol, 71%). 1H NMR (500 MHz, D_2O): δ (ppm) 8.28 (d, $J = 0.76$ Hz, 1H), 5.11 (d, $J = 3.49$ Hz, 1H), 4.63 (d, $J = 12.5$ Hz, 1H), 3.84 (dd, $J = 10.73, 3.53$ Hz, 1H), 3.77 (d, $J = 1.44$ Hz, 1H), 3.75-3.73 (m, 1H), 3.73-3.71 (m, 1H), 3.71-3.69 (m, 1H), 3.68 (d, $J = 2.31$ Hz, 1H), 3.64 (d, $J = 5.20$ Hz, 1H), 3.63 (d, $J = 2.43$ Hz, 1H), 3.62-3.60 (m, 1H), 3.59 (dd, $J = 4.87, 2.14$ Hz, 1H), 3.49-3.45 (m, 1H), 3.34 (ddd, $J = 10.06, 2H$); ^{13}C NMR (125 MHz, D_2O): δ (ppm) 94.8, 90.5, 76.7, 73.8, 71.7, 70.9, 70.7, 70.4, 61.4, 61.2, 57.7, 55.1; HRMS-FAB: $[M-H]$ calcd. for $C_8H_{11}F_3NO_6^-$ 274.0544 Found: 274.0541.

2-deoxy-2-[(trifluoroacetyl)amino]- β -glucopyranose-1,3,4,6-tetraacetate (20)—Pyridine (56.3 mL) and Ac_2O (32.5 mL) were added to **19** (5.26 g, 19.2 mmol). The reaction mixture was stirred for 3 hours at room temperature. The solution was concentrated *in vacuo* and co-evaporated with toluene (10 mL) three times. The resulting residue was recrystallized in Et_2O (15 mL) and petroleum ether (5 mL) to afford peracetylated **20** ($\alpha/\beta = 1:0.33$, 8.51 g, 19.2 mmol, quant.). 1H NMR (500 MHz, $CDCl_3$): δ (ppm) 6.60 (d, $J = 9.28$ Hz, 1H), 6.51 (d, $J = 8.50$ Hz, 1H), 6.27 (d, $J = 3.66$ Hz, 1H), 5.76 (d, $J = 8.71$ Hz, 1H), 5.35-5.28 (m, 1H), 5.25 (t, $J = 9.74$, 1H), 5.22-5.16 (m, 1H), 4.45 (ddd, $J = 10.84, 8.69, 3.66$ Hz, 1H), 4.33-4.27 (m, 2H), 4.15 (dd, $J = 12.56, 2.28$ Hz, 1H), 4.09 (dd, $J = 12.50, 2.33$ Hz, 1H), 4.04 (ddd, $J = 9.95, 3.97, 2.32$ Hz, 1H), 3.85 (ddd, $J = 9.50, 4.50, 2.18$ Hz, 1H), 2.23 (s, 3H), 2.13 (s, 6H), 2.11 (s, 6H), 2.08 (s, 3H), 2.08 (s, 3H), 2.07 (s, 3H), 2.06 (s, 3H); ^{13}C NMR (500 MHz, $CDCl_3$): δ (ppm) 171.7, 170.7, 169.2, 168.6, 157.5, 157.2, 116.5, 114.2, 91.7, 89.6, 72.8, 71.9, 70.1, 69.7, 67.9, 67.2, 61.6, 61.3, 53.1, 51.7; HRMS-FAB: $[M-H]$ calcd. for $C_{16}H_{19}F_3NO_{10}^-$ 442.0967 Found: 442.0961.

2-deoxy-2-[(trifluoroacetyl)amino]- β -glucopyranose-3,4,6-triacetate-1-dihydrogen phosphate (21)— 1H NMR (500 MHz, D_2O): δ (ppm) 5.70-5.55 (m, 1H), 5.43 (s, 1H), 5.14 (s, 1H), 4.36 (s, 2H), 4.24-4.09 (m, 1H), 3.34 (d, $J = 19.11$ Hz, 1H), 2.08 (s, 3H), 2.01 (s, 3H), 1.94 (s, 3H); ^{13}C NMR (125 MHz, D_2O): δ (ppm) 171.3, 170.4, 169.8, 157.2, 114.7, 92.7, 71.2, 68.2, 68.1, 61.4, 52.8; HRMS-FAB: $[M-H]$ calcd. for $C_{14}H_{18}F_3NO_{12}P^-$ 480.0524 Found: 480.0524.

2-deoxy-2-[(trifluoroacetyl)amino]- β -glucopyranose-1-dihydrogen phosphate (22)— ^1H NMR (500 MHz, D_2O): δ (ppm) 5.44 (dd, $J = 7.32, 3.23$ Hz, 1H), 4.03 (ddd, $J = 10.57, 3.17, 1.84$ Hz, 1H), 3.94 (ddd, $J = 10.12, 4.89, 2.22$ Hz, 1H), 3.77 (dd, $J = 12.38, 4.94$ Hz, 1H), 3.87 (ddd, $J = 12.31, 9.99, 5.66$ Hz, 1H), 3.51 (dd, $J = 10.06, 9.14$ Hz, 1H); ^{13}C NMR (125 MHz, D_2O): δ (ppm) 159.4, 116.7, 92.6, 72.6, 69.9, 69.5, 60.2, 54.3; HRMS-FAB: $[\text{M}-\text{H}]$ calcd. for $\text{C}_8\text{H}_{12}\text{F}_3\text{NO}_9\text{P}^-$ 354.0207 Found: 354.0206.

2-deoxy-2-[(1-oxo-2-ethen-1-yl)amino]- α - β -glucopyranose-1,3,4,6-tetraacetate (24)— Et_3N (145 μL , 1.04 mmol) was added to a stirred solution of 1,3,4,6-*O*-acetyl-2-amino-2-deoxy- β -D-glucopyranose-HCl (200 mg, 0.521 mmol). The solution was cooled to 0 $^\circ\text{C}$ and acryloyl chloride (50 μL , 0.625 mmol) was added dropwise. The reaction mixture was stirred overnight at room temperature. The solution was concentrated in vacuo and purified on silica gel chromatography (hexanes 100% to hexanes/EtOAc = 1/1) to afford **24** as a clear syrup (170 mg, 81%).

^1H NMR (500 MHz, CDCl_3): δ (ppm) 6.96 (d, $J = 9.59$ Hz, 1H), 6.20 (d, $J = 17.1$ Hz, 1H), 6.00 (dd, $J = 17.1, 10.4$ Hz, 1H), 5.68 (d, $J = 8.8$ Hz, 1H), 5.61 (d, $J = 9$ Hz, 1H), 5.17 (t, $J = 11$ Hz, 1H), 5.07 (t, $J = 9.65$ Hz, 1H), 4.34 (dd, $J = 19.42, 9.61$ Hz, 1H), 4.21 (td, $J = 9.10, 4.57$ Hz, 1H), 4.08 (dd, $J = 12.44, 2.15$ Hz, 1H), 3.81 (ddd, $J = 9.85, 4.55, 2.19$ Hz, 1H), 2.04-1.95 (s, 12H); ^{13}C NMR (125 MHz, CDCl_3): δ (ppm) 171.2, 170.9, 169.7, 169.4, 166.0, 130.1, 127.4, 92.4, 72.6, 72.6, 68.0, 61.7, 61.7, 52.5, 50.3; HRMS-FAB: $[\text{M}+\text{H}]$ calcd. for $\text{C}_{17}\text{H}_{24}\text{NO}_{10}^+$ 402.1400 Found: 402.1395.

2-deoxy-2-[(1-oxo-2-ethen-1-yl)amino]- α - β -glucopyranose-3,4,6-triacetate-1-(dihydrogen phosphate) (25)—Brown oil; ^1H NMR (500 MHz, MeOD): δ (ppm) 6.28 (d, $J = 9.92$ Hz, 17 Hz, 1H), 6.22 (dd, $J = 17.5, 2.05$ Hz, 1H), 5.66 (dd, $J = 9.93, 2.05$ Hz, 1H), 5.52 (dd, $J = 6.63, 3.26$ Hz, 1H), 5.36 (dd, $J = 10.49, 9.59$ Hz, 1H), 5.11 (t, $J = 9.75, 9.75$ Hz, 1H), 4.39-4.34 (m, 1H), 4.31 (dd, $J = 9.20, 6.05$ Hz, 2H), 4.29-4.26 (m, 1H), 4.17-4.11 (m, 1H), 2.00 (s, 6H), 1.94 (s, 3H); ^{13}C NMR (125 MHz, MeOD): δ (ppm) 171.1, 170.5, 169.9, 166.7, 130.3, 126.1, 93.7, 71.4, 68.3, 68.2, 61.5, 51.9, 50.3; HRMS-FAB: $[\text{M}-\text{H}]$ calcd. for $\text{C}_{15}\text{H}_{21}\text{NO}_{12}\text{P}^-$ 438.0807 Found: 438.0800.

2-deoxy-2-[(1-oxo-2-ethen-1-yl)amino]- α - β -glucopyranose-3,4,6-1-(dihydrogen phosphate) (26)—Light brown amorphous solid; ^1H NMR (500 MHz, D_2O): δ (ppm) 6.14 (dd, $J = 17.14, 10.20$ Hz, 1H), 6.04 (dd, $J = 17.14, 1.35$ Hz, 1H), 5.60 (dd, $J = 10.23, 1.34$ Hz, 1H), 5.28 (dd, $J = 6.63, 3.26$ Hz, 1H), 3.88-3.83 (m, 1H), 3.68 (dd, $J = 6.66, 3.67$ Hz, 1H), 3.65-3.62 (m, 3H), 3.59 (m, 1H); ^{13}C NMR (125 MHz, D_2O): δ (ppm) 165.8, 129.4, 128.0, 93.5, 72.6, 70.4, 69.4, 60.1, 53.7; HRMS-FAB: $[\text{M}-\text{H}]$ calcd. for $\text{C}_9\text{H}_{15}\text{NO}_9\text{P}^-$ 312.0490 Found: 312.0486.

P'-[2-(acetylamino)-2-deoxy- α - β -glucopyranosyl] ester-uridine 5'-(trihydrogen diphosphate), (UDP-GlcNAc, 27)— ^1H NMR (500 MHz, D_2O): δ (ppm) 7.87 (dd, $J = 8.08, 2.41$ Hz, 1H), 5.93-5.86 (m, 2H), 5.44 (d, $J = 3.28$ Hz, 1H), 4.32-4.25 (m, 2H), 4.24-4.08 (m, 3H), 3.94-3.88 (m, 1H), 3.86 (ddd, $J = 8.03, 5.20, 2.66$ Hz, 1H), 3.80 (dd, $J = 17.63, 9.17$ Hz, 1H), 3.73 (ddd, $J = 18.49, 9.53, 5.39$ Hz, 2H), 3.47 (ddd, $J = 9.52, 5.51, 3.38$ Hz, 1H), 2.02-1.95 (s, 3H); ^{13}C NMR (125 MHz, D_2O): δ (ppm) 174.7, 166.7, 152.1, 141.4, 102.6, 94.3, 88.4, 83.0, 73.7, 72.9, 70.8, 69.5, 69.4, 64.9, 60.2, 53.5, 22.0; HRMS-FAB: $[\text{M}-\text{H}]$ calcd. for $\text{C}_{17}\text{H}_{26}\text{N}_3\text{O}_{17}\text{P}_2^-$ 606.0743 Found: 606.0734.

P'-[2-(1-oxo-2-propyn-1-yl)amino-2-deoxy- α - β -glucopyranosyl] ester-uridine 5'-(trihydrogen diphosphate) (UDP-GlcN-alkyne, 28)—White fluffy powder; ^1H NMR (600 MHz, D_2O): δ (ppm) 7.89 (d, $J = 8.1$ Hz, 1 H), 5.89 (d, $J = 4.6$ Hz, 1 H), 5.88 (d, $J = 8.9$ Hz, 1 H), 5.47 (dd, $J = 3.2, 6.9$ Hz, 1 H), 4.28 (m, 2 H), 4.20 (m, 1 H), 4.18-4.16 (m, 1

H), 4.12-4.09 (m, 1 H), 3.99 (dt, $J = 3.1, 10.5$ Hz, 1 H), 3.87-8.83 (m, 1 H), 3.79-3.77 (m, 1 H), 3.76-3.70 (m, 2 H), 3.49-3.44 (m, 2 H); ^{13}C NMR (150 MHz, D_2O): δ (ppm) 166.2, 154.8, 151.8, 141.6, 102.6, 94.1, 88.4, 88.3, 83.2, 73.8, 72.0, 69.6, 68.3, 67.4, 64.9, 60.2, 50.3, 50.2; HRMS-FAB: [M] calcd. for $\text{C}_{18}\text{H}_{25}\text{N}_3\text{O}_{17}\text{P}_2^+$ 617.0659 Found: 617.0656.

P'-[2-deoxy-2-[(2,2,2-trifluoroacetyl)amino]- α -D-glucopyranosyl] ester-uridine 5'-(trihydrogen diphosphate) (UDP-GlcN-TFA, 29)—White fluffy powder; ^1H NMR (600 MHz, D_2O): δ (ppm) 7.88 (d, $J = 8.11$ Hz, 1H), 5.90 (m, 2H), 5.58-5.51 (dd, 1H, $J = 7.0$ Hz, 3.2 Hz), 4.29 (m, 2H), 4.21 (s, 1H), 4.20-4.14 (m, 1H), 4.11 (ddd, $J = 11.68, 5.39, 3.12$ Hz, 1H), 4.04 (d, $J = 10.57$ Hz, 1H), 3.88 (dd, $J = 20.10, 9.82$ Hz, 2H), 3.81 (d, $J = 11.83$ Hz, 1H), 3.78-3.73 (m, 1H), 3.52 (dd, $J = 9.53, 5.39$ Hz, 1H); ^{13}C NMR (150 MHz, D_2O): δ (ppm) 166.4, 160.0, 151.9, 141.8, 117.0, 102.7, 93.9, 88.5, 83.2, 73.9, 73.1, 70.3, 69.7, 69.5, 64.9, 60.2, 54.4; HRMS-FAB: [M-H] calcd. for $\text{C}_{17}\text{H}_{23}\text{F}_3\text{N}_3\text{O}_{17}\text{P}_2^-$ 660.0460 Found: 660.0463.

P'-[2-deoxy-2-[(1-oxo-2-ethen-1-yl)amino]-2-deoxy- α -D-glucopyranosyl] ester-uridine 5'-(trihydrogen diphosphate) (UDP-GlcN-ene, 30)—White fluffy powder; ^1H NMR (800 MHz, D_2O): δ (ppm) 7.88 (d, $J = 8.09$ Hz, 1H), 6.35-6.29 (dd, $J = 10.4, 18.4$ Hz, 1H), 6.19-6.13 (d, $J = 16.8$ Hz, 1H), 5.92-5.86 (m, 2H), 5.74-5.68 (d, $J = 10.4$ Hz, 1H), 5.46-5.42 (m, 1H), 4.31-4.25 (m, 2H), 4.21-4.18 (m, 1H), 4.18-4.08 (m, 2H), 3.97-3.93 (m, 1H), 3.88-3.84 (m, 1H), 3.78-3.73 (m, 3H), 3.50-3.46 (m, 1H); ^{13}C NMR (200 MHz, D_2O): δ (ppm) 168.7, 166.1, 151.7, 141.4, 129.6, 127.9, 102.5, 94.4, 88.3, 83.1, 73.7, 72.9, 70.9, 69.5, 69.3, 64.8, 60.1, 53.6; HRMS-FAB: [M-H] calcd. for $\text{C}_{18}\text{H}_{26}\text{N}_3\text{O}_{17}\text{P}_2^-$ 618.0743 Found: 618.0741.

P'-[2-(acetylamino)-2-deoxy- α -D-galactopyranosyl] ester-uridine 5'-(trihydrogen diphosphate) (UDP-GalNAc, 31)— ^1H NMR (500 MHz, D_2O): δ (ppm) 7.79 (d, $J = 8.2$ Hz, 1 H), 5.81 (br d, $J = 4.2$ Hz, 1 H); 5.79 (br d, $J = 8.3$, 1 H); 5.39 (dd, $J = 6.9, 3.3$ Hz, 1 H); 4.22-4.19 (m, 2 H), 4.13-4.08 (m, 3 H), 4.05-4.01 (m, 2 H); 3.88 (d, $J = 3.0$ Hz, 1 H), 3.78 (dd, $J = 11.8, 3.2$ Hz, 1 H); 3.73 (m, 1 H), 3.62 (m, 1H), 1.92 (s, 3 H); HRMS-FAB: [M] calcd. for $\text{C}_{17}\text{H}_{27}\text{N}_3\text{O}_{17}\text{P}_2^+$ 607.0816 Found: 607.0818.

P'-[2-(1-oxo-2-propyn-1-yl)amino]-2-deoxy- α -D-galactopyranosyl] ester-uridine 5'-(trihydrogen diphosphate) (UDP-GalN-alkyne, 32)—White fluffy powder; ^1H NMR (600 MHz, D_2O): δ (ppm) 7.87 (d, $J = 8.1$ Hz, 1 H), 5.89 (d, $J = 4.8$ Hz, 1 H), 5.86 (d, $J = 8.2$ Hz, 1 H), 5.50 (dd, $J = 3.4, 6.9$ Hz, 1 H), 4.28 (m, 2 H), 4.22 (dt, $J = 3.1, 10.6$ Hz, 1 H), 4.20 (m, 1 H), 4.17 (m, 1 H), 4.11-4.09 (m, 2 H), 3.97 (br d, $J = 3.0$ Hz, 1 H), 3.91 (dd, $J = 3.1, 10.9$ Hz, 1 H), 3.70-3.64 (m, 3 H); ^{13}C NMR (150 MHz, D_2O): δ (ppm) 166.2, 155.0, 151.8, 141.6, 102.6, 94.2, 88.4, 83.2, 83.1, 73.7, 72.0, 69.7, 68.3, 67.5, 65.0, 60.9, 50.3, 50.2; HRMS-FAB: [M] calcd. for $\text{C}_{18}\text{H}_{25}\text{N}_3\text{O}_{17}\text{P}_2^+$ 617.0659 Found: 617.0661.

P'-[2-azido-2-deoxy- α -D-glucopyranosyl] ester-uridine 5'-(trihydrogen diphosphate) (UDP-GlcN₃, 33)—UMP-morpholidate (211 mg, 0.303 mmol) was dried by dissolving in anhydrous pyridine (5 mL) and concentrating *in vacuo* three times. Fully deprotected phosphate **5** (77 mg, 0.217 mmol) was converted to its PyH^+ form by passing through an Amberlite (PyH^+) column. UMP-Morpholidate was dissolved in pyridine (5 mL) and added to phosphate **5** (PyH^+). The reaction mixture was stirred for 5 days at room temperature. The solvent was removed by concentration and the resulting residue was dissolved in H_2O (10 mL). Insoluble material was removed through filtration and the soluble filtrate was concentrated *in vacuo*. The residue was loaded onto a BioGel P2 column (2.5 × 65 cm) and eluted with H_2O . Fractions were collected and those containing the product as

determined by TLC (*n*BuOH/HCO₂H/H₂O = 4/8/1) were combined and freeze-dried to afford a white powder (50 mg, 0.0756 mmol, 35%).

¹H NMR (500 MHz, D₂O): δ (ppm) 7.90 (d, *J* = 8.12 Hz, 1H), 5.91 (m, 2H), 5.61 (dd, *J* = 7.48, 3.32 Hz, 1H), 4.30 (m, 2H), 4.23-4.18 (m, 1H), 4.15 (m, 2H), 3.89-3.83 (m, 2H), 3.78 (dd, *J* = 12.47, 2.34 Hz, 1H), 3.71 (dd, *J* = 12.51, 4.28 Hz, 1H), 3.46 (dd, *J* = 10.12, 9.16 Hz, 1H), 3.30 (m, 1H); ¹³C NMR (125 MHz, D₂O): δ (ppm) 166.6, 152.2, 141.9, 102.9, 94.6, 88.5, 83.5, 74.0, 73.0, 70.9, 69.9, 69.6, 63.1, 63.0, 60.3; HRMS-FAB: [M-H] calcd. for C₁₅H₂₂N₅O₁₆P₂⁻ 590.0542 Found: 590.0539.

P'-[2-amino-2-deoxy-α-D-glucopyranosyl] ester-uridine 5'-(trihydrogen diphosphate) (UDP-GlcNH₂, 34)—¹H NMR (800 MHz, D₂O): δ (ppm) 7.87 (d, *J* = 8.09 Hz, 1H), 5.93-5.85 (m, 2H), 5.64-5.56 (m, 1H), 4.29 (d, *J* = 2.66 Hz, 2H), 4.23-4.19 (m, 1H), 4.19-4.09 (m, 2H), 3.85-3.81 (m, 1H), 3.80-3.72 (m, 2H), 3.72-3.66 (m, 1H), 3.45-3.39 (m, 1H), 3.01-2.91 (m, 1H); ¹³C NMR (200 MHz, D₂O): δ (ppm) 166.6, 152.2, 141.9, 102.7, 94.7, 88.5, 83.1, 73.8, 73.2, 71.8, 69.6, 69.1, 65.0, 60.1, 54.7; HRMS-FAB: [M-H] calcd. for C₁₅H₂₄N₃O₁₆P₂⁻ 564.0637 Found: 564.0638.

P'-[2-deoxy-2-[[5-[(3a*S*,4*S*,6a*R*)-hexahydro-2-oxo-1*H*-thieno[3,4-*d*]imidazol-4-yl]-1-oxopentyl]amino]-α-D-glucopyranosyl] ester uridine 5'-(trihydrogen diphosphate) (UDP-GlcN-biotin, 35)—To a stirred solution of **34** (5 mg, 8.2 μmol) in HEPES buffer (2.5 mL, pH 8) was added 2.5 mL of a 10mM biotin-NHS solution in DMF at 0 °C. The reaction mixture was stirred for 2 h at 0 °C then concentrated. The crude mixture was purified on a BioGel P2 column (2.5 × 65cm) and eluted with H₂O. Fractions were collected and those containing the product as determined by TLC (*n*BuOH/HCO₂H/H₂O = 4/8/1) were combined and freeze-dried to afford **35** as a white powder.

¹H NMR (600 MHz, D₂O): δ (ppm) 7.92-7.87 (d, *J* = 7.8 Hz, 1H), 5.92-5.83 (m, 2H), 5.44-5.39 (m, 1H), 4.54-4.49 (m, 1H), 4.37-4.33 (m, 1H), 4.31-4.24 (m, 2H), 4.21-4.18 (m, 1H), 4.18-4.08 (m, 2H), 3.95-3.90 (m, 1H), 3.88-3.83 (m, 1H), 3.82-3.77 (m, 1H), 3.76-3.69 (m, 2H), 3.49-3.43 (m, 1H), 2.95-2.88 (m, 1H), 2.72-2.65 (m, 1H), 2.32-2.25 (m, 1H), 1.70-1.29 (m, 8H); ¹³C NMR (150 MHz, D₂O): δ (ppm) 177.5, 166.2, 158.1, 153.3, 141.7, 102.6, 94.6, 84.5, 83.1, 73.8, 73.0, 70.9, 69.6, 64.9, 62.0, 60.2, 55.1, 48.1, 39.7, 35.4, 27.9, 27.5, 25.0; HRMS-FAB: [M-H] calcd. for C₂₅H₃₈N₅O₁₈P₂S⁻ 790.1413 Found: 790.1401.

P'-[2-deoxy-2-[[Fluorenylmethoxycarbonyl]amino]-α-D-glucopyranosyl] ester uridine 5'-(trihydrogen diphosphate) (UDP-GlcN-Fmoc, 36)—To a stirred solution of **34** (5 mg, 8.2 μmol) in 1,4-dioxane/H₂O, 1/1 (0.8 mL) was added NaHCO₃ (6.9 mg, 8.2 μmol) and FmocCl (5.3 mg, 20.5 μmol). The reaction was stirred overnight at room temperature then concentrated. The crude mixture was purified on a BioGel P2 column (2.5 × 65 cm) and eluted with H₂O. Fractions were collected and those containing the product as determined by TLC (*n*BuOH/HCO₂H/H₂O, 4/8/1) were combined and freeze-dried to afford **36** as a white powder.

¹H NMR (600 MHz, D₂O): δ (ppm) 7.92-7.87 (d, *J* = 7.8 Hz, 1H), 5.92-5.83 (m, 2H), 5.44-5.39 (m, 1H), 4.54-4.49 (m, 1H), 4.37-4.33 (m, 1H), 4.31-4.24 (m, 2H), 4.21-4.18 (m, 1H), 4.18-4.08 (m, 2H), 3.95-3.90 (m, 1H), 3.88-3.83 (m, 1H), 3.82-3.77 (m, 1H), 3.76-3.69 (m, 2H), 3.49-3.43 (m, 1H), 2.95-2.88 (m, 1H), 2.72-2.65 (m, 1H), 2.32-2.25 (m, 1H), 1.70-1.29 (m, 8H); ¹³C NMR (150 MHz, D₂O): δ (ppm) 163.1, 162.9, 143.5, 141.0, 140.8, 128.0, 127.5, 125.5, 125.4, 120.1, 117.3, 102.3, 94.9, 88.5, 82.8, 73.6, 73.0, 71.7, 71.2, 64.9, 60.4, 46.8; HRMS-FAB: [M-H] calcd. for C₃₀H₃₄N₃O₁₈P₂⁻ 786.1318 Found: 786.1318.

P'-[2-deoxy-2-[(1,1-dimethylethoxy)carbonyl]amino]- α -D-glucopyranosyl] ester uridine 5'-(trihydrogen diphosphate) (UDP-GlcN-tBoc, **37)**—Boc₂O (4.53 μ L, 0.0197 mmol) was added to a stirred solution of **34** (8 mg, 0.0131 mmol) in 1,4-dioxane/H₂O, 1/1 (1 mL). The pH of the solution was adjusted to pH 10 using 10% NaOH (aq.) and the reaction was stirred overnight at room temperature. A second portion of Boc₂O (4.53 μ L, 0.0197 mmol) was added and the reaction was again stirred overnight at room temperature. Ethyl acetate (1 mL) was added to the reaction mixture and the organic phase was removed from the resulting biphasic mixture. The aqueous phase was loaded onto a BioGel P2 column (2.5 \times 65cm) and eluted with H₂O. Fractions were collected and those containing the product as determined by TLC (*n*BuOH/HCO₂H/H₂O = 4/8/1) were combined and freeze-dried to afford **37** as a white powder (5.0 mg, 0.007 mmol, 30%).

¹H NMR (500 MHz, D₂O): δ (ppm) 7.96 (d, *J* = 8.00 Hz, 1H), 6.00 (d, *J* = 4.00 Hz, 1H), 5.98-5.95 (m, 2H), 5.54-5.49 (m, 1H), 4.37 (t, *J* = 4.33 Hz, 2H), 4.29 (s, 1H), 4.24 (s, 1H), 4.20 (dd, *J* = 6.04, 2.82 Hz, 1H), 3.93 (dd, *J* = 10.00, 3.16 Hz, 1H), 3.86-3.82 (m, 1H), 3.72 (d, *J* = 5.77 Hz, 1H), 3.68 (s, 1H), 3.54 (d, *J* = 9.43 Hz, 1H), 1.45 (s, 9H); ¹³C NMR (125 MHz, D₂O): δ (ppm) 178.2, 167.4, 157.9, 141.5, 102.7, 95.0, 88.0, 85.2, 81.3, 73.8, 72.9, 69.6, 69.5, 65.9, 64.8, 60.2, 55.0, 27.6; HRMS-FAB: [M-H] calcd. for C₂₀H₃₂N₃O₁₈P₂⁻ 664.1162 Found: 664.1166.

Radioactive assay

A given UDP-hexosamine analog (1–4 mM) and radioactive ³H-UDP-GlcUA (0.6–1 mM, 0.1 μ Ci) were co-incubated in 20–50 μ L reactions containing 50 mM Tris pH 7.2, 1 mM MnCl₂, (as well as 1 M ethylene glycol for PmHAS and PmCS) with 0.5–45 μ g of purified recombinant enzyme at 30°C for 3–16 hrs. The reaction mixture was then quenched with detergent and analyzed by descending paper chromatography.²³ As a negative control for assay background, a reaction with no UDP-hexosamine was tested in parallel (chain polymerization can only occur when UDP-GlcUA and a functional UDP-hexosamine are present simultaneously).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank Dr. Dmitri Zagorevski for MS data acquisition. The authors are also grateful for funding from the National Institutes of Health in the form of grants HL62244 and HL094463.

References

1. van Heijenoort J. *Glycobiology*. 2001; 11(3):25R–36R.
2. Wu AM. *Neurochem. Res.* 2002; 27(7–8):593–600. [PubMed: 12374194]
3. Capila I, Linhardt RJ. *Angew. Chem. Int. Edit.* 2002; 41(3):391–412.
4. Watkins WM. *Carbohydr. Res.* 1986; 149(1):1–12.
5. Sismey-Ragatz AE, Green DE, Otto NJ, Rejzek M, Field RA, DeAngelis PL. *J. Biol. Chem.* 2007; 282(39):28321–28327. [PubMed: 17627940]
6. Karst NA, Linhardt RJ. *Curr Med Chem.* 2003; 10(19):1993–2031. [PubMed: 12871100]
7. Xu Y, Masuko S, Takiuddin M, Xu H, Liu R, Jing J, Mousa SA, Linhardt RJ, Liu J. *Science*. 2011; 334(6055):498–501. [PubMed: 22034431]
8. Guan W, Cai L, Fang J, Wu B, Wang PG. *Chem. Commun.* 2009; 45:6976–6978.
9. Guan W, Cai L, Wang PG. *Chem.-Eur. J.* 2010; 16(45):13343–13345. [PubMed: 21031374]

10. Chen Y, Thon V, Li Y, Yu H, Ding L, Lau K, Qu J, Hie L, Chen X. *Chem. Commun.* 2011; 47(38):10815–10817.
11. Yan RB, Yang F, Wu YF, Zhang LH, Ye XS. *Tetrahedron Lett.* 2005; 46(52):8993–8995.
12. MacDonald DL. *J. Org. Chem.* 1966; 31(2):513–516. [PubMed: 5902475]
13. Ikeda S, Shibuya M, Iwabuchi Y. *Chem. Commun.* 2007; 5:504–506.
14. Rostovtsev VV, Green LG, Fokin VV, Sharpless KB. *Angew. Chem. Int. Edit.* 2002; 41(14):2596–2599.
15. Liu T, Xu Y, Chen M, Weïwer M, Zhou X, Bridges AS, DeAngelis PL, Zhang Q, Linhardt RJ, Liu J. *J Biol Chem.* 2010; 285(44):34240–34249. [PubMed: 20729556]
16. Curphey TJ. *J. Org. Chem.* 1979; 44(15):2805–2807.
17. Dondoni A. *Angew. Chem. Int. Edit.* 2008; 47(47):8995–8997.
18. Mengin-Lecreulx D, van Heijenoort J. *J. Bacteriol.* 1994; 176(18):5788–5795. [PubMed: 8083170]
19. Bintein F, Auge C, Lubineau A. *Carbohydr. Res.* 2003; 338(11):1163–1173.
20. Faijes M, Saura-Valls M, Perez X, Conti M, Planas A. *Carbohydr. Res.* 2006; 341(12):2055–2065.
21. Sala RF, MacKinnon SL, Palcic MM, Tanner ME. *Carbohydr. Res.* 1998; 306(1–2):127–136.
22. Jing W, DeAngelis PL. *Glycobiology.* 2003; 13(10):661–671. [PubMed: 12799342]
23. Jing W, DeAngelis PL. *Glycobiology.* 2000; 10(9):883–889. [PubMed: 10988250]
24. Chen M, Bridges A, Liu J. *Biochemistry.* 2006; 45(40):12358–12365. [PubMed: 17014088]







Figure 1.
Enzymatic synthesis of UDP-GlcNAc and UDP-GalNAc analogs




Scheme 1.
Synthesis of GlcNH₂-1-phosphate




Scheme 2.
Synthesis of GlcNAc/GalNAc-1-phosphate




Scheme 3.
Synthesis of GlcN-alkyne-1-phosphate




Scheme 4.
Synthesis of GalN-alkyne-1-phosphate



Scheme 5.
GlcN-TFA-1-phosphate




Scheme 6.
Synthesis of GlcN-ene-1-phosphate









Scheme 7.
Chemical synthesis of UDP-GlcN derivatives

Table 1

GlcNAc/GalNAc-1-phosphate analog acceptance by GlmU



| | Substrate | R | Yield (%) | Product |
|------------------|-----------|--|-----------|----------------------------|
| GlcNAc & analogs | 5 | N ₃ | n.r. | - |
| | 6 | NH ₂ | n.r. | - |
| | 7* |  | 85 | UDP-GlcNAc, 27 |
| | 13 |  | 50 | UDP-GlcN-alkyne, 28 |
| | 22 |  | 78 | UDP-GlcN-TFA, 29 |
| | 26 |  | 65 | UDP-GlcN-ene, 30 |
| GalNAc & analogs | 9* |  | 70 | UDP-GalNAc, 31 |
| | 18 |  | 60 | UDP-GalN-alkyne, 32 |

* natural precursor

Table 2

Analog usage by GAG synthases as monitored by co-polymerization assay.^{5, 23} The incorporation of [³H]GlcUA radioactivity into polysaccharide is shown.

| UDP-donor | ³ H]GlcUA incorporation (disintegrations/min × 10 ⁻³) | | | |
|-----------------|--|------------------|-----------------|-----------------|
| | PmHAS | PmCS | PmHS1 | PmHS2 |
| 27 ^a | 70–120 | na ^b | 25–80 | 25–70 |
| 28 | 1 | na | 12 | 16 |
| 29 | 34 | na | NS ^c | 12 |
| 30 | 7 | na | 12 | 14 |
| 33 | NS ^d | na | NS ^d | NS ^d |
| 35 | NS ^d | na | NS ^d | NS ^d |
| 36 | NS | na | NS | 1.2 |
| 37 | 6 | na | 13 | 13 |
| 31 ^a | na | 110 ^d | na | na |
| 32 | na | 82 ^d | na | na |

^a natural precursor positive control (PmCS only utilizes UDP-GalNAc derivatives while rest of synthases only employ UDP-GlcNAc derivatives)

^b na, not applicable

^c NS, not significant (≤5-fold over background of the negative control without any UDP-hexosamine present)

^d Values for overnight reaction time