



University Medical Center Groningen

University of Groningen

Screening for coronary artery calcium in a high-risk population

Denissen, Sabine Jam; van der Aalst, Carlijn M; Vonder, Marleen; Gratama, Jan Willem C; Adriaansen, Henk J; Kuijpers, Dirkjan; Roeters van Lennep, Jeanine E; Vliegenthart, Rozemarijn; van der Harst, Pim; Braam, Richard L

Published in:
European journal of preventive cardiology

DOI:
[10.1177/2047487320932263](https://doi.org/10.1177/2047487320932263)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Denissen, S. J., van der Aalst, C. M., Vonder, M., Gratama, J. W. C., Adriaansen, H. J., Kuijpers, D., Roeters van Lennep, J. E., Vliegenthart, R., van der Harst, P., Braam, R. L., van Dijkman, P. R., Oudkerk, M., & de Koning, H. J. (2020). Screening for coronary artery calcium in a high-risk population: the ROBINSICA trial. *European journal of preventive cardiology*, [2047487320932263]. <https://doi.org/10.1177/2047487320932263>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Screening for coronary artery calcium in a high-risk population: the ROBINSICA trial

Sabine JAM Denissen¹, Carlijn M van der Aalst¹, Marleen Vonder², Jan Willem C Gratama³, Henk J Adriaansen⁴, Dirkjan Kuijpers^{5,6}, Jeanine E Roeters van Lennep⁷, Rozemarijn Vliegenthart², Pim van der Harst^{2,8,9}, Richard L Braam¹⁰, Paul RM van Dijkman^{11,12}, Matthijs Oudkerk^{13,14} and Harry J de Koning¹

European Journal of Preventive
Cardiology
0(0) 1–5

© The Author(s) 2020



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/2047487320932263

journals.sagepub.com/home/cpr



Cardiovascular disease (CVD) remains the main cause of death worldwide, accounting for 44% of all non-communicable disease deaths, of which most are attributable to coronary heart disease (CHD).¹ Coronary artery calcification (CAC) has a strong association with major cardiovascular events and mortality, and has a high risk-predictive value of CHD in asymptomatic individuals.^{2,3} It has been argued that the amount of CAC, expressed in the CAC score, can be used in population-based screening.

The Dutch Risk Or Benefit IN Screening for Cardiovascular disease (ROBINSICA) trial is the first large-scale population-based randomised controlled trial (RCT) to investigate whether CAC screening followed by preventive treatment is effective in reducing CHD-related morbidity and mortality in asymptomatic individuals.^{4,5} The aim of this study was to investigate the CAC prevalence and predictors in the ROBINSICA trial, which included an asymptomatic high-risk potential target population from the general population.

The rationale and design of the ROBINSICA trial have been described before.⁵ Briefly, 43,447 potentially high-risk women (55–74 years) and men (45–74 years) from the national population registry who completed a baseline questionnaire to assess sociodemographic and health characteristics and gave informed consent were randomly allocated (1:1:1) to either the control arm, intervention arm A (screening according to traditional risk factors) or intervention arm B (CAC screening). The current study focuses on the CAC screening arm (Figure 1). The Minister of Health authorised the ROBINSICA trial in 2013.

CAC screening was performed using computed tomography scanning to identify CVD risk according to the CAC score, which represents the total amount of

any CAC.⁶ CAC scores were categorised into low (Agatston 0–99), high (Agatston 100–399) and very high (Agatston ≥ 400) risk.²

The effects of baseline characteristics on CAC score were analysed using a two-step approach regression analyses for modelling presence, both any CAC

¹Department of Public Health, Erasmus Medical Centre, The Netherlands

²Centre for Medical Imaging North-East Netherlands (CMI-NEN), University Medical Centre Groningen, The Netherlands

³Department of Radiology and Nuclear Medicine, Gelre Hospitals, The Netherlands

⁴Clinical Chemistry and Hematology Laboratory, Gelre Hospitals, The Netherlands

⁵Department of Radiology, University Medical Center Groningen, The Netherlands

⁶Department of Radiology, Haaglanden Medical Centre Bronovo, The Netherlands

⁷Department of Internal Medicine, Erasmus Medical Centre, The Netherlands

⁸Department of Cardiology, University Medical Centre Groningen, The Netherlands

⁹Department of Cardiology, University Medical Centre Utrecht, The Netherlands

¹⁰Department of Cardiology, Gelre Hospitals, The Netherlands

¹¹Department of Cardiology, Leids University Medical Centre, The Netherlands

¹²Department of Cardiology, Haaglanden Medical Centre Bronovo, The Netherlands

¹³University of Groningen, University Medical Centre Groningen, The Netherlands

¹⁴Institute for Diagnostic Accuracy – iDNA, The Netherlands

The first two authors contributed equally.

Trial registration number: NTR6471.

Corresponding author:

Sabine Denissen, Department of Public Health, Erasmus MC, P.O. Box 2040, 3000 CA Rotterdam, The Netherlands.

Email: s.denissen@erasmusmc.nl




Figure 1. Flowchart of the ROBINSICA trial study design in which CAC scoring is performed in intervention arm B. BMI: body mass index; CAC: coronary artery calcium; CHD: coronary heart disease; CT: computed tomography; CVD: cardiovascular disease; CVRM: cardiovascular risk management; SCORE: systematic coronary risk evaluation.

and CAC score of 400 or higher (multivariable backward logistic regression), and extent (multivariable backward linear regression of the log-transformed CAC score) of CAC in women and men separately. Variables included in the models were age, educational level, waist circumference cut-off (88 cm for women and 102 cm for men), body mass index (BMI) cut-off (30 kg/m^2), family history of CHD, smoking, diabetes mellitus, hypertension and/or hypercholesterolemia in the past year, and baseline use of either antihypertensive or lipid-lowering medication (according to self-reported data from the baseline questionnaire). A *P* value of less than 0.05 was considered statistically significant. All analyses were performed using IBM SPSS Statistics version 24.0.

Of the 12,950 screened participants, 48.1% were women and 94.2% were born in The Netherlands. The median age was 64 years in women and 62 years in men. Regarding CVD risk factors, 20.0% were current smokers at baseline, 3.4% reported diabetes mellitus, 16.4% and 15.3% reported being diagnosed with hypertension and hypercholesterolemia, respectively, in the year before baseline, and 44.9% reported a family history of CHD.

CAC was absent in 39.2% of the total study population. Overall, 48% of women had a zero CAC score compared to 20.7% of men in the same age category and 31.2% of all men. Furthermore, 16.8% of women had a CAC score of 100 or higher compared to 40.0% of men in the same age category and 30.7% of all men. The CAC distribution in the ROBINSICA trial is compared to the German Heinz Nixdorf Recall Study and the American Multi Ethnic Study of Atherosclerosis in the Supplementary material.

Age, high waist circumference, family history of CHD, smoking at baseline, diabetes mellitus, self-reported hypertension or hypercholesterolemia at baseline and baseline use of either antihypertensive or lipid-lowering medication were all selected as predictors in the backward regression analysis of the presence of CAC and CAC of 400 or greater, and in the linear regression for predicting the log-transformed CAC extent in women (Table 1). Age, educational level, high BMI, family history of CHD, smoking at baseline, diabetes mellitus, self-reported hypercholesterolemia at baseline and baseline use of either antihypertensive or lipid-lowering medication were selected as predictors in the analyses for men (Table 1). A higher educational level predicted a lower CAC score in men. The composition of the predictors differed moderately in the models for women and men.

The associations of age, male sex, diabetes mellitus and smoking with higher CAC scores are well known.⁷ A lower socioeconomic status, indicated by educational level, significantly predicted a higher extent of CAC in

men. This association is possibly a result of a less favourable lifestyle in terms of smoking, diet and physical activity.⁸ Diabetes mellitus was one of the strongest predictors of CAC presence in women. This is in line with previous research in which diabetes mellitus was identified to have a greater impact in women compared to men.⁹ Moreover, diabetes mellitus was a strong predictor for CAC extent in both sexes, suggesting that it is the most important risk factor for CAC development after sex and age. Regarding BMI and waist circumference, our results confirm earlier findings that BMI is not a strong predictor for the presence of CAC, while waist circumference is more predictive of CAC presence.¹⁰ The predictive value of the baseline use of either antihypertensive or lipid-lowering medication in CAC development was also seen in previous research. However, statins have been associated with increased CAC scores, but not with more CVD events. It is suggested that statins induce CAC progression and, at the same time, plaque repair.¹¹

This study contributes to evidence on identifying the optimal target population for screening from the general population that will gain most healthy life-years from screening and subsequent treatment. All inclusion criteria for the ROBINSICA trial (smoking, waist circumference, BMI and a family history of CHD) were statistically significant predictors of CAC. Future analyses should provide evidence on whether the study population includes individuals who benefit most.

A main limitation is that the ROBINSICA population is not representative of all ethnic groups as a result of a homogeneous distribution, although ethnicity is known to affect CAC prevalence and severity. Another possible limitation is that study participants tend to be generally healthier than similar individuals not responding to the participation invitation (healthy volunteer effect). However, the inclusion and exclusion criteria should have minimised this effect. Furthermore, participants using both cholesterol-lowering and antihypertensive medication were excluded from the trial, which might have affected the found associations of CAC with CVD medication. Finally, baseline data were obtained using a self-reported questionnaire, rather than diagnostic test measures, and might entail some inaccuracies.

In conclusion, this currently largest population-based RCT for CAC screening in asymptomatic middle-aged Caucasian individuals showed that 30.7% of men and 16.8% of women with a CAC score of 100 or greater urgently require preventive treatment. To a large extent, male sex and increasing age, followed by diabetes mellitus and smoking, influence CAC distribution. These results can therefore help determine the best risk prediction and prevention

Table 1. Baseline predictors for the presence and extent of coronary artery calcium and for a coronary artery calcium score of 400 or higher.

	Logistic regression for CAC presence (CAC score = 0 vs. > 0)		Linear regression for log-transformed CAC extent ^a		Logistic regression for CAC ≥ 400	
	Odds ratio (95% CI)	P value	Coefficient (95% CI)	P value	Odds ratio (95% CI)	P value
Women						
Age, per 10 years	2.74 (2.46–3.05)	<0.001***	0.48 (0.36–0.59)	<0.001***	3.58 (2.79–4.64)	<0.001***
Waist circumference cut-off ^b	1.22 (1.03–1.46)	0.024*				
Family history of CHD ^c	1.63 (1.45–1.84)	<0.001***	0.20 (0.08–0.31)	0.001**	1.70 (1.31–2.21)	<0.001***
Smoker at baseline	2.00 (1.69–2.37)	<0.001***	0.35 (0.19–0.51)	<0.001***	2.18 (1.55–3.05)	<0.001***
Diabetes mellitus	2.26 (1.51–3.39)	<0.001***	0.33 (0.04–0.63)	0.026*		
Hypertension in past year (self-reported)	1.27 (1.06–1.51)	0.008**				
Hypercholesterolemia in past year (self-reported)	1.15 (0.98–1.36)	0.097				
Antihypertensive medication	1.28 (1.09–1.49)	0.002**	0.28 (0.14–0.42)	<0.001***	1.93 (1.46–2.57)	<0.001***
Lipid-lowering medication	1.81 (1.41–2.32)	<0.001***	0.22 (0.02–0.42)	0.033*	2.25 (1.54–3.30)	<0.001***
	Area under the curve: 0.685		Adjusted R²: 0.043		Area under the curve: 0.744	
Men						
Age, per 10 years	2.97 (2.52–3.46)	<0.001***	0.70 (0.60–0.80)	<0.001***	<0.001***	<0.001***
Educational level ^d			-0.15 (-0.30–-0.002)		0.046*	0.85 (0.68–1.05)
Low			-0.15 (-0.29–-0.02)		0.026*	0.77 (0.63–0.93)
Medium						
High						
Body Mass Index cut-off ^e			0.23 (0.07–0.38)		0.004**	1.33 (1.07–1.66)
Family history of CHD ^c	1.38 (1.18–1.62)	<0.001***	0.27 (0.15–0.38)	<0.001***	<0.001***	<0.001***
Smoker at baseline	1.35 (1.12–1.64)	0.002**	0.30 (0.16–0.44)	<0.001***	<0.001***	<0.001***
Diabetes mellitus			0.39 (0.13–0.66)		0.004**	1.39 (0.96–2.03)
Hypercholesterolemia in past year (self-reported)	1.25 (0.99–1.59)	0.063				0.085
Antihypertensive medication	1.94 (1.55–2.43)	<0.001***	0.32 (0.18–0.46)	<0.001***	<0.001***	<0.001***
Lipid-lowering medication	1.48 (1.10–2.00)	0.011*	0.40 (0.21–0.58)	<0.001***	<0.001***	0.004**
	Area under the curve: 0.686		Adjusted R²: 0.077		Area under the curve: 0.698	

CAC: coronary artery calcium; CHD: coronary heart disease; CI: confidence interval.

^aLog-transformation in individuals with CAC score > 2 because transforming CAC score > 0 did not result in a normal distribution.^bTrial inclusion criteria cut-off for waist circumference: ≥ 88 cm for women and ≥ 102 cm for men.^cFamily history of myocardial infarction or sudden death before the age of 65 years in first and second-degree relatives.^dEducational levels: low; primary, lower secondary general or lower vocational education, medium; intermediate vocational or higher secondary education, high; higher vocational education or university.^eTrial inclusion criteria cut-off for body mass index: ≥ 30 kg/m².

*P < 0.05, **P < 0.01, ***P < 0.001.

strategy should screening for a high risk of developing CVD be (cost)-effective.

Acknowledgements

The author(s) would like to thank the European Union for funding the ROBINSCA trial and the Ministry of Health, Welfare and Sports for the ethical approval to perform the trial. Naturally, The author(s) thank all participants for their participation. Furthermore, they would like to thank M Quak for the extensive research assistance, RADventure for developing the data management system and the IVA group for handling all questionnaires and letters. Finally, they thank all employees of the radiology departments of the screening centres (Gelre Hospital Apeldoorn, Bronovo Hospital The Hague and University Medical Centre Groningen) for scanning participants.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The ROBINSCA trial was funded by an advanced grant (agreement no. 294604) of the European Research Council.

Supplemental material

Supplemental material for this article is available online.

References

1. Wilkins E, Wilson L, Wickramasinghe K, et al. *European Cardiovascular Disease Statistics 2017*. Brussels: European Heart Network, 2017.
2. Greenland P, Blaha MJ, Budoff MJ et al. Coronary calcium score and cardiovascular risk. *J Am Coll Cardiol* 2018; 72: 434–447.
3. Budoff MJ, Young R, Burke G, et al. Ten-year association of coronary artery calcium with atherosclerotic cardiovascular disease (ASCVD) events: the multi-ethnic study of atherosclerosis (MESA). *Eur Heart J* 2018; 39: 2401–2408.
4. McEvoy JW, Martin SS, Blaha MJ, et al. The case for and against a coronary artery calcium trial: means, motive, and opportunity. *JACC Cardiovasc Imaging* 2016; 9: 994–1002.
5. Van der Aalst CM, Vonder M, Gratama JM, et al. Risk or Benefit in Screening for Cardiovascular Disease (ROBINSCA): the rationale and study design of a population-based randomized-controlled screening trial for cardiovascular disease. *J Clin Trials* 2019; 9: 1000361.
6. Vonder M, van der Aalst CM, Vliegenthart R, et al. Coronary artery calcium imaging in the ROBINSCA Trial: rationale, design, and technical background. *Acad Radiol* 2018; 25: 118–128.
7. Pletcher MJ, Sibley CT, Pignone M, et al. Interpretation of the coronary artery calcium score in combination with conventional cardiovascular risk factors: the Multi-Ethnic Study of Atherosclerosis (MESA). *Circulation* 2013; 128: 1076–1084.
8. Djekic D, Angeras O, Lappas G, et al. Impact of socio-economic status on coronary artery calcification. *Eur J Prev Cardiol* 2018; 25: 1756–1764.
9. Nicoll R, Wiklund U, Zhao Y, et al. Gender and age effects on risk factor-based prediction of coronary artery calcium in symptomatic patients: a Euro-CCAD study. *Atherosclerosis* 2016; 252: 32–39.
10. Roy SK, Zeb I, Kadakia J, et al. Body surface area is a predictor of coronary artery calcium, whereas body mass index is not. *Coron Artery Dis* 2012; 23: 113–117.
11. Henein M, Granasen G, Wiklund U, et al. High dose and long-term statin therapy accelerate coronary artery calcification. *Int J Cardiol* 2015; 184: 581–586.