



## LJMU Research Online

**Kelly, R and Roberts, JW**

**Investigation into the dynamic visual acuity of skilled cricketers using a continuous motion task**

<http://researchonline.ljmu.ac.uk/id/eprint/14038/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Kelly, R and Roberts, JW (2020) Investigation into the dynamic visual acuity of skilled cricketers using a continuous motion task. Journal of Sports Medicine and Physical Fitness. ISSN 0022-4707**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

**Investigation into the dynamic visual acuity of skilled cricketers using a continuous  
motion task**

Robert Kelly & James W. Roberts<sup>1†</sup>

<sup>1</sup>: Liverpool Hope University

Psychology, Action and Learning of Movement (PALM) Laboratory

School of Health Sciences

Hope Park, Liverpool

L16 9JD

**Running head:** Dynamic visual acuity in cricketers

†Author JWR is now affiliated with Liverpool John Moores University, Brain and Behaviour Laboratory, Research Institute of Sport and Exercise Sciences (RISES), Byrom Street, Tom Reilly Building, Liverpool, L3 5AF

Corresponding author:

James W. Roberts

Liverpool John Moores University

Brain & Behaviour Laboratory

Research Institute of Sport & Exercise Sciences (RISES)

Byrom Street, Tom Reilly Building, Liverpool, L3 5AF

E-mail: J.W.Roberts@ljmu.ac.uk

1 **Abstract**

2 BACKGROUND: Great demands are imposed upon the perceptual-motor system when  
3 undertaking ball-throwing and -hitting tasks including cricket. That is, performers must detect  
4 and resolve object details while on the move – something referred to as dynamic visual acuity  
5 (DVA). The present study aimed to investigate DVA in skilled cricketers and non-cricketers  
6 using a more immediate or real-time assessment.

7 METHODS: Skilled cricketers and non-cricketers had to detect the presence of the gap  
8 within a Landolt-C ring as it moved horizontally or vertically, while progressively increasing  
9 the size until the participants registered a response. Measures were taken as the mean  
10 (dynamic) minimum angle of resolution of the object size at the moment that participants  
11 correctly responded to the gap. Objects would move at either a high, medium or low velocity.

12 RESULTS: There was greater dynamic visual acuity in the skilled cricketers compared to  
13 non-cricketers ( $p < .05$ ). There was a reduced negative influence of object velocity on  
14 dynamic visual acuity in the skilled cricketers compared to non-cricketers ( $p < .05$ ).

15 CONCLUSIONS: We suggest these findings contribute to the growing evidence surrounding  
16 DVA within ball-throwing and -hitting sports, while making some assertions as to the  
17 implications for the cricket performance setting.

18

19 **Key words:** ball sports; perceptual-cognitive; ocular pursuit; Landolt-C

## 1 **Introduction**

2           In the sport of cricket, players have to contend with the substantial demands placed  
3 upon the perceptual-motor system. For example, cricket batsmen are exposed to bowling  
4 velocities that can reach in excess of 40 m/s (approximately 144 km/h), which assumes a ball  
5 flight time of near 600 ms with a 17-m distance between the creases<sup>1,2</sup>. Moreover, fielders  
6 have to judge and intercept balls that travel at an extremely high velocity following contact  
7 with the bat (e.g., 92-126 km/h; <sup>3</sup>). Thus, it is of great interest to explore the perceptual  
8 abilities of skilled cricketers who are able to perform under such constraints.

9           With this in mind, it is worthwhile evaluating the possible sources of visual  
10 information that promote skilled cricket performance (e.g., catching a small-sized ball within  
11 a short period of time). The key cues include the expansion of objects within the sagittal  
12 plane (monocular), relative retinal velocities or motion parallax (monocular), and retinal  
13 disparities between the two eyes (binocular). A common characteristic for each of these cues  
14 surrounds the ability to resolve objects within a dynamic or relative moving environment –  
15 something that is referred to as dynamic visual acuity (DVA)<sup>4,5,6</sup>.

16           Since its inception, there have been a growing number of studies showing superior  
17 DVA in skilled athletes compared to novices or non-athletes<sup>7,8,9,10,11</sup> (for alternative findings,  
18 see <sup>12</sup> and <sup>13</sup>). Within the context of ball-throwing and -hitting sports (e.g., baseball, cricket),  
19 this finding has been primarily attributed to the advanced oculomotor abilities in skilled  
20 athletes – they exhibit a low-latency rapid eye movement (i.e., saccade) prior to the object  
21 reaching the “hitting zone”<sup>1,10,14</sup>. Along these lines, skilled athletes appear to demonstrate a  
22 reduced decline in DVA following an increasing object velocity (“velocity resistant”; <sup>15</sup>; see  
23 also, <sup>16</sup>).

24           Nevertheless, the previously used DVA tasks have typically featured the sudden  
25 appearance and disappearance of visual targets, where performers are afforded the

1 opportunity to deliberate over their choice of response. For example, performers are  
2 presented a Landolt-C ring at varying eccentricities for a brief temporal window (~100 ms)  
3 before taking their time to decide on what direction the gap in the ring was facing (e.g., <sup>6</sup>). In  
4 the context of cricket, these types of tasks are not entirely suitable for reflecting the spatial  
5 and temporal dynamics of cricket performance, where precise visual information must be  
6 processed within a comparatively short period of time (<1 s). Thus, it could be informative to  
7 alternatively incorporate a continually moving target that demands an immediate response.  
8 With this in mind, the present study adopts a previously designed DVA task that uniquely  
9 features a continually moving object (<15 deg/s), which progressively increases in size until  
10 the performer can positively resolve it – an increasing target size coincides with an increasing  
11 presentation time (see <sup>16,17</sup>). Indeed, these newly introduced parameters may more closely  
12 reflect the common performance setting of tracking a ball in preparation for catching or  
13 intercepting it. Specifically, the presence of a continual object motion promotes the retinal  
14 velocities and object-tracking eye movements (i.e., smooth pursuit) (~60 deg/s;<sup>18</sup>) that are  
15 also required to perceive the ball in-flight. Likewise, the presence of a response-time  
16 contingency resembles the selection and initiation processes that enable performers to  
17 physically interact with the ball.

18         At the same time, it is not entirely accepted that DVA, and other related visual  
19 abilities, can positively discriminate skill levels within sport. Indeed, prior assessments of  
20 generalizable visual abilities have alternatively indicated limited differences in skilled  
21 compared to less-skilled athletes<sup>19,20,21</sup>. Likewise, visual training interventions that have been  
22 designed to enhance these generalizable visual abilities have failed to benefit athletes in both  
23 their visual- and sport-specific skills<sup>22</sup>. In the context of cricket, recent findings have shown  
24 that the ability to anticipate bowls and subsequently hit the ball when under degraded vision  
25 (courtesy of wearing plus dioptré lenses) can remain relatively unaffected<sup>23,24,25</sup>. Taken

1 together, these lines of research appear to contest the influence of generalizable visual  
2 abilities within sport, while advocating a primary role of specialised perceptual-cognitive  
3 skills – it is not how information is seen, but how it is used that is essential.

4 To this end, the present study seeks to broadly expand upon evidence of DVA in  
5 cricketers compared to non-cricketers, including their responses to varying object velocities.  
6 More specifically, we aimed to examine these issues using a previously adopted DVA task  
7 (see <sup>16,17</sup>) that potentially encapsulates many aspects of the perceptual, oculomotor and  
8 response demands in cricket. In so doing, we can advance the ecological validity of DVA  
9 within the context of interceptive ball sports, while further advancing our understanding of  
10 the role of generalizable dynamic visual abilities. We hypothesised that there would be a  
11 generally superior DVA within cricketers compared to non-cricketers. Additionally, we  
12 hypothesised that there would be a smaller decline in DVA following an increase in object  
13 velocity (i.e., “velocity resistant”) for cricketers compared to non-cricketers.

14

## 15 **Materials and Methods**

### 16 *Participants*

17 Sixteen male participants volunteered for the study (8 skilled cricketers, 8 non-  
18 cricketers; age range = 18-24 years).<sup>1</sup> All participants reported normal or corrected-to-normal  
19 static vision and no known neurological conditions. The skilled group comprised of sub-elite  
20 varsity-standard cricket players that reported at least 8.5 years of competitive experience<sup>5</sup>.  
21 While the unskilled group were comparatively young and active, they reported no  
22 competitive or extended recreational experience within cricket, nor competed within any  
23 other interceptive ball sports around the time of testing. The study was approved by the local  
24 ethics committee, and conducted in accordance with the Declaration of Helsinki (1964,  
25 2013).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

*Apparatus*

Stimuli were generated and controlled via Matlab (2018b) (The Mathworks Inc., Natick, MA) running Psychtoolbox (version 3.0.11)<sup>28</sup>. A Samsung UHD TV (screen size = 109.5 x 62 cm, screen resolution = 1360 x 768, temporal resolution = 75 Hz) was used to display the stimuli. The display was vertically oriented and adjusted so that the centre of the screen could be aligned with the participants' line of gaze. Participants were stood 2-m from the display, and provided a keypad that was connected to a universal serial bus extension so they could freely respond to the stimulus.

*Stimulus and Procedure*

The stimulus consisted of a standard black Landolt-C ring on a white background. The gap within the ring occupied a 1/5 of the entire diameter (equivalent to a single leg of an "E" optotype). Thereafter, the stimulus parameters were closely adapted from work conducted by Muiños and Ballesteros<sup>16</sup>. That is, the initial size of the ring was 3.02 mm, which was progressively increased by 1 pixel (.026°) every 2.3 secs. The ring was oriented so that the gap could face directly up, down, left or right, and translated across the horizontal or vertical mid-line of the screen (see Figure 1). The ring was moved at a constant velocity of either 15°/s (.536 m/s; high), 9.15°/s (.322 m/s; medium), or 3.06°/s (.107 m/s; low). If the ring reached the outer edges of the display without a keyed response being made, then it was simply reversed so that it could be moved in the opposite direction with the same gap orientation and velocity.

The task required participants to detect the presence of the ring and respond to the direction of the gap by quickly pressing a key on the keypad. Arrows were placed over the keys 2, 4, 6 and 8, which corresponded to the directions down, left, right and up, respectively.

1 The target object became increasingly larger in size over the course of each trial, which  
2 unfolded indefinitely until participants issued a response. Therein, the movement of the  
3 object and changes in size would momentarily cease until the participants self-selected  
4 another key in order to commence the next trial. There were a total of 120 trials comprising  
5 of 10 trials per variation of object velocity (high, medium, low) and movement direction  
6 (horizontal, vertical), which were randomly presented throughout the experiment. There was  
7 a further prompt to undertake a two-minute break at half way (60 trials). Participants were  
8 provided 25 trials of practice before any formal data collection.

9

10 [Insert Figure 1 about here]

11

## 12 *Data Management and Analysis*

13 The size of the ring following each response was stored for further analysis. The trials  
14 featuring a correct response (i.e., selected key corresponded with the direction of the gap)  
15 were adapted to calculate a logMAR acuity score.<sup>2</sup> This measure was based on a logarithmic  
16 transformation of the ratio between the test and standard minimum angle of resolution  
17 (MAR):

18

$$19 \text{ test MAR} = \text{object size (in pixels)} \times .026^\circ$$

$$20 \text{ logMAR} = \log_{10}(\text{test MAR} / .0833^\circ)$$

21

22 The analysis involved entering the participant mean logMAR acuity scores into a  
23 three-way mixed design ANOVA with group (cricketers, non-cricketers) as the between-  
24 measures factor, and velocity (high, medium, slow) and direction (horizontal, vertical) as the  
25 repeated-measures factors. The equal variance of differences (Sphericity) assumption was



1 evaluated using Mauchly's test. In the event of a violation, then the Huynh-Feldt adjusted  
2 value was adopted when Epsilon was  $>.75$ , while the Greenhouse-Geisser value was adopted  
3 when Epsilon was  $<.75$ . For the ANOVA at least, effect sizes were indicated by partial eta-  
4 squared ( $\eta^2$ ). In the event of a statistically significant effect involving more than two means,  
5 then a Tukey HSD post hoc procedure was undertaken.

6 In order to capture the potential "velocity resistant" characteristic of DVA within  
7 highly skilled athletes<sup>15</sup>, while corroborating the effects from our main omnibus ANOVA, we  
8 additionally analysed the within-participant slope coefficients that pertained to the linear  
9 relation between logMAR scores and object velocities (15°/s, 9.15°/s, 3.06°/s). Indeed, a  
10 more deleterious effect of velocity on DVA should manifest in a steeper gradient. The  
11 assumptions of parametric data, including a normal distribution and homogeneity of variance,  
12 were evaluated using the Shapiro-Wilk and Levene's tests, respectively. Therein, the  
13 cricketers and non-cricketers were compared using an independent samples t-test. In this  
14 instance, the effect size was indicated by Cohen's  $d_s$ <sup>29</sup>. All inferential statistical analyses were  
15 declared as significant at  $p < .05$ .

16

## 17 **Results**

18 LogMAR is interpreted as lower scores representing better acuity. Thus, the dynamic  
19 logMAR acuity scores (grand  $M = .15$  logMAR,  $SD = .07$ ) were generally worse than  
20 standard levels of static acuity (.00 logMAR or 20/20), which reflects the ubiquitous finding  
21 of a decline in object resolution during relative moving conditions.

22 ANOVA revealed a significant main effect of group,  $F(1, 14) = 5.58, p = .033, partial$   
23  $\eta^2 = .29$ , as the skilled cricketers were significantly lower than the non-cricketers.<sup>2</sup> There was  
24 also a significant main effect of velocity,  $F(2, 28) = 44.42, p = .00, partial \eta^2 = .76$ , as an  
25 increasing object velocity proved detrimental to dynamic acuity. However, these effects were

1 superseded by a significant group x velocity interaction,  $F(2, 28) = 4.07, p = .028, \text{partial } \eta^2$   
2  $= .23$ , which indicated that the detrimental effect of object velocity on dynamic acuity was  
3 less apparent for skilled cricketers (see Figure 2). Indeed, the post hoc analysis revealed that  
4 there was a significantly higher logMAR score in the high compared to the medium velocity  
5 condition, which was also higher than the low velocity condition, for the non-cricketers ( $ps <$   
6  $.05$ ). However, there was only a significant difference between the extremely high and low  
7 velocity conditions ( $p < .05$ ), and no significant differences surrounding comparisons with  
8 the medium velocity condition ( $ps > .05$ ), for the skilled cricketers. There were no further  
9 statistically significant main, or interaction effects (direction, group x direction:  $F_s < 1$ ;  
10 velocity x direction:  $F(2, 28) = 1.50, p = .24, \text{partial } \eta^2 = .10$ ; group x velocity x direction:  
11  $F(2, 28) = 1.09, p = .35, \text{partial } \eta^2 = .07$ ).

12         Meanwhile, the independent t-test on individual participant slope coefficients  
13 indicated a significantly larger slope for non-cricketers compared to cricketers for the  
14 horizontal stimuli ( $t(14) = 2.41, p < .05, d_s = 1.21$ ), and a similar trend for the vertical stimuli  
15 ( $t(14) = 1.89, p = .08, d_s = .95$ ) (see Table 1).

16

17                                 [Insert Figure 2 and Table 1 about here]

18

## 19 **Discussion**

20         The present study aimed to examine the DVA underlying skilled cricketers compared  
21 to non-cricketers, as well as the potential modulation of DVA under varying object velocities.  
22 Importantly, we adopted a DVA task<sup>16,17</sup> that uniquely featured a continual object motion  
23 (incorporating retinal velocities and object-tracking eye movements; e.g., watching the ball  
24 trajectory) and response time-contingency (response initiation being coincident with visual  
25 perception; e.g., initiating the response to catch during perception of the ball). Specifically,

1 the task required participants to respond to the direction of a gap within a Landolt-C ring,  
2 which continuously moved in the horizontal or vertical direction while progressively  
3 increasing in size. In addition, the velocity of the target object was varied across trials (slow,  
4 medium, fast). The findings generally showed that the skilled cricketers were better than non-  
5 cricketers. While the skilled cricketers demonstrated a decline in DVA for the fast compared  
6 to slow object velocities, there was a limited difference for the comparisons involving the  
7 medium object velocity. Meanwhile, the non-cricketers demonstrated an incremental decline  
8 in DVA from the fast to medium object velocities, and medium to slow object velocities.  
9 These findings were corroborated by the much smaller positive linear relations between  
10 logMAR scores and object velocities for the cricketers compared to non-cricketers. These  
11 skill-level differences also indicated a medium-to-large effect size.

12         The presence of a decline in DVA following the increasing object velocities indicates  
13 a deleterious effect of velocity across all skill levels. However, there appeared to be a much  
14 smaller decline within the skilled cricketers compared to non-cricketers (see Figure 2). This  
15 outcome closely reflects the “velocity resistant” characteristics of skilled athletic  
16 performance<sup>16</sup>. This characteristic feature may be attributed to the specialised oculomotor  
17 abilities of skilled compared to less-skilled performers. That is, skilled performers within  
18 ball-throwing and -hitting sports indicate enhanced low-latency rapid saccades<sup>1,10</sup> and smooth  
19 pursuits<sup>30,31</sup> in anticipation of time-to-contact. To elucidate, skilled perceptual-motor  
20 performance within cricket assumes an initial rapid detection of the ball flight, which is  
21 impossible to continuously track due to its extremely high angular velocity (ball velocity  
22  $>500^\circ/\text{s}$  vs. pursuit eye-tracking  $\sim 60^\circ/\text{s}$ ). Thus, the performer tends to generate anticipatory  
23 eye movements that accommodate the resolution of object details as the object/performer  
24 moves<sup>15</sup>. While the present object velocities ( $<15^\circ/\text{s}$ ) were substantially lower than the  
25 previously evidenced angular velocities (e.g., <sup>10,11</sup>), we have extended upon the notion of

1 velocity resistance in sport athletes compared to novices or non-athletes by introducing an  
2 object-tracking task feature.

3 Unlike the influence of object velocity, there was a limited influence of motion  
4 direction as it failed to discriminate DVA between skilled cricketers and non-cricketers.  
5 Previous evidence has rendered at least some influence of motion direction when the stimulus  
6 is deemed to be partially similar to the characteristics of the sport performance-setting (<sup>16</sup>; see  
7 also, <sup>20</sup>). For example, cricketers may have alternatively benefited from motion within the  
8 sagittal plane (i.e., depth) as opposed to the current fronto-parallel plane (i.e.,  
9 horizontal/vertical) because it more closely resembles the approach of ball for catching or  
10 hitting. Likewise, it is relevant to consider the implications of motion direction on the  
11 recruitment of unique neural pathways that are specialised for the processing of particular  
12 visual characteristics (e.g., translational vs. radial optic flow<sup>32,33,34</sup>; upper vs. lower visual  
13 fields<sup>35</sup>; low- vs high-spatial frequencies<sup>25</sup>). Thus, it is important to realise that in the absence  
14 of an influence of motion direction, there is still a strong possibility that introducing a further  
15 direction (e.g., radial) will discriminate DVA across skill levels.

16 When reflecting on the skill-level differences in DVA, we may attempt to relate these  
17 findings to the real-life performance setting. That is, the mean DVA scores for the skilled  
18 cricketers and non-cricketers were synonymous with target sizes that ranged from 3.5-4 mm  
19 and 4-5.2 mm, respectively. This difference equated to at least a single step in the  
20 modification of the stimulus object size, which was ramped up by 1 pixel (~.81 mm) every  
21 2.3 s. Along these lines, the minimum angle of resolution for the skilled cricketers and non-  
22 cricketers assumes that a moving object equivalent to the size of a cricket ball (approximately  
23 72 mm diameter) could be resolved at distances of 36-42 m and 27-37 m, respectively. While  
24 these metrics can seemingly translate abstract optometric data into real-life cricket  
25 performance, it is important to stress that they are theoretical in nature. Thus, future studies

1 should contend with the challenge of directly implicating sport performance outcomes based  
2 on lab-based controlled measures (e.g., <sup>36</sup>).

3         At the same time, it is relevant to consider that the present study failed to feature  
4 expert/professional athletes. Indeed, the obvious difficulty in recruiting such a cohort can be  
5 observed throughout the perceptual-cognitive sport literature (e.g., <sup>7,10,37,38</sup>). Thus, while the  
6 present study cannot extend the present findings to the elite/professional domain, they  
7 nevertheless reflect robust skill-level differences (i.e., large statistical effect sizes and  
8 approximately 38% mean decrease from the non-cricketers to skilled cricketers) that  
9 positively indicate dynamic visual acuity as a discriminating feature of cricket performance.  
10 Further research may elaborate on the present skill-level differences by additionally  
11 incorporating the assessment of dynamic visual abilities in expert/professional athletes.

12

### 13 **Conclusions**

14         We have strongly corroborated previous evidence of an advantage in DVA for skilled  
15 athletes compared to novices or non-athletes within rapid ball-throwing and -hitting sports,  
16 including cricket. These findings contribute to the body of literature that supports the role of  
17 generalizable dynamic visual abilities (e.g., <sup>39</sup>) as opposed to the independent role of  
18 specialised perceptual-cognitive skills (e.g., <sup>19,21,22</sup>). Furthermore, we showed that the  
19 tendency for DVA to exponentially decline with an increasing velocity is less apparent for  
20 skilled athletes. That is, they tend to be less susceptible to the deleterious effects of object  
21 velocity. While the current evidence may be considered derivative with respect to the existing  
22 DVA literature, we have additionally expanded this evidence-base to a DVA task where there  
23 was a need to immediately resolve and respond to the stimulus object, while displaying  
24 angular velocities that accommodated pursuit eye movements.

1 **References**

- 2 1) Land MF, McLeod P. From eye movements to actions: how batsmen hit the ball. *Nat*  
3 *Neurosci* 2000;3:1340-1345.
- 4
- 5 2) Müller, S., Abernethy, B., Reece, J., Rose, M., Eid, M., McBean, R., Hart, T., & Abreu,  
6 C. An in-situ examination of the timing of information pick-up for interception by cricket  
7 batsmen of different skill levels. *Psychol Sport Exerc* 2009;10:644-652.
- 8
- 9 3) Peplow, C., McErlain-Naylor, S. A., Harland, A. R., & King, M. A. Relationships  
10 between technique and bat speed, post- impact ball speed, and carry distance during  
11 arrange hitting task in cricket. *Hum Movement Sci* 2019;63:34-44.
- 12
- 13 4) Miller JW. Study of visual acuity during the ocular pursuit of moving test objects. II.  
14 Effects of direction of movement, relative movement, and illumination. *J Opt Soc Am*  
15 1958;48:803-808.
- 16
- 17 5) Miller JW, Ludvigh E. The effect of relative motion on visual acuity. *Surv Ophthalmol*  
18 1962;7:83-116.
- 19
- 20 6) Erickson GB, Citek K, Cove M, Wilczek J, Linster C, Bjarnason B, Langemo N.  
21 Reliability of a computer-based system for measuring visual performance skills.  
22 *Optometry* 2011;82:528-542.
- 23
- 24 7) Gao Y, Chen L, Yang S, Wang H, Yao J, Dai Q, et al. Contributions of visuo-oculomotor  
25 abilities to interceptive skills in sports. *Optometry Vision Sci* 2015;92:679-689.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

8) Ishigaki H, Miyao H. Differences in dynamic visual acuity between athletes and nonathletes. *Percept Motor Skill* 1993;77:835–839.

9) Rouse MW, DeLand P, Christian R, Hawley J A comparison study of dynamic visual acuity between athletes and nonathletes. *J Am Optom Assoc* 1988;59:946-950.

10) Uchida Y, Kudoh D, Higuchi T, Honda M, Kanosue K. Dynamic visual acuity in baseball players is due to superior tracking abilities. *Med Sci Sport Exer* 2013;45:319-325.

11) Uchida Y, Kudoh D, Murakami A, Honda M, Kitazawa S. Origins of superior dynamic visual acuity in baseball players: superior eye movements or superior image processing. *PLoS One* 2012;7:e31530.

12) Poltavski D, Biberdorf D. The role of visual perception measures used in sports vision programmes in predicting actual game performance in Division I collegiate hockey players. *J Sport Sci* 2015;33:597-608.

13) Wimshurst ZL, Sowden PT, Cardinale M. Visual skills and playing positions of Olympic field hockey players. *Percept Motor Skill* 2012;114:204-216.

14) Bahill AT, Laritz T. Why can't batters keep their eyes on the ball? *Am Sci* 1984;72:249–253.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

15) Sanderson FH. Visual acuity and sports performance. In: Cockerill IM, MacGillivray WW (editors). Vision and sport. Cheltenham: Stanley Thomas; 1981. p. 64-79.

16) Muiños M, Ballesteros S. Sports can protect dynamic visual acuity from aging: a study with young and older judo and karate martial arts athletes. *Atten Percept Psycho* 2015;77:2061-2073.

17) Quevedo L, Aznar-Casanova JA, Merindano MD, Solé J. A task to assess dynamic visual acuity and a valuation of the stability of its measurements. *Psicológica* 2010;31:109–128.

18) Bennett SJ, Barnes GR. Prediction in ocular pursuit. In: Elliott D, Khan MA (editors). Vision and goal-directed movement: neurobehavioral perspectives. Champaign: Human Kinetics; 2010. p.149-176.

19) Helsen WF, Starkes JL. A multidimensional approach to skilled perception and performance in sport. *Appl Cognitive Psych* 1999;13:1-27.

20) Roberts JW, Strudwick AJ, Bennett SJ. Visual function of English Premier League soccer. *Medicine and Science in Football* 2017;1:178-182.

21) Ward P, Williams, AM. Perceptual and cognitive skill development in soccer: the multidimensional nature of expert performance. *J Sport Exercise Psy* 2003;25:93-111.



- 1 22) Abernethy B, Wood JM. Do generalized visual training programmes for sport really  
2 work? An experimental investigation. *J Sport Sci* 2001;19:203-222.  
3
- 4 23) Mann DL, Abernethy B, Farrow D. The resilience of natural interceptive actions to  
5 refractive blur. *Hum Movement Sci* 2010a;29:368-400.  
6
- 7 24) Mann DL, Abernethy B, Farrow D. Visual information underpinning skilled anticipation:  
8 the effect of blur on a coupled and uncoupled in situ anticipatory response. *Atten Percept*  
9 *Psycho* 2010b;72:1317-1326.  
10
- 11 25) Ryu D, Abernethy B, Park SH, Mann DL. The perception of deceptive information can  
12 be enhanced by training that removes superficial visual information. *Front Psychol*  
13 2018;9:1132.  
14
- 15 26) Faul F, Erdfelder E, Lang AG, Buchner A. G\*Power 3: A flexible statistical power  
16 analysis program for the social, behavioral, and biomedical sciences. *Behav Res Meth*  
17 2007;39:175–191.  
18
- 19 27) Colquhoun D. An investigation of the false discovery rate and the misinterpretation of p-  
20 values. *Roy Soc Open Sci* 2014;1:140216.  
21
- 22 28) Pelli DG. The VideoToolbox software for visual psychophysics: Transforming numbers  
23 into movies. *Spat Vis* 1997;10:437–442.  
24

- 1 29) Lakens, D. Calculating and reporting effect sizes to facilitate cumulative science: a  
2 practical primer for t-tests and ANOVAs. *Front Psychol* 2013;4:863  
3
- 4 30) Bennett SJ, Barnes GR. Smooth ocular pursuit during the transient disappearance of an  
5 accelerating visual target: the role of reflexive and voluntary control. *Exp Brain Res*  
6 2006;175:1-10.  
7
- 8 31) Santos EM, Kowler E. Anticipatory smooth pursuit eye movements evoked by  
9 probabilistic cues. *J Vision* 2017;17:1-16.  
10
- 11 32) Aaen-Stockdale C, Ledgeway T, Hess RF. Second-order optic flow processing. *Vision*  
12 *Res* 2007;47:1798-1808.  
13
- 14 33) Duffy CJ, Wurtz RH. Sensitivity of MST neurons to optic flow stimuli. I. A continuum  
15 of response selectivity to large-field stimuli. *J Neurophysiol* 1991a;65:1329-1345.  
16
- 17 34) Duffy CJ, Wurtz RH. Sensitivity of MST neurons to optic flow stimuli. II. Mechanisms  
18 of response selectivity revealed by small-field stimuli. *J Neurophysiol* 1991b;65:1346-  
19 1359.  
20
- 21 35) Danckert J, Goodale MA. Superior performance for visually guided pointing in the lower  
22 visual field. *Exp Brain Res* 2001;137:303-308.  
23

- 1 36) Clark JF, Ellis JK, Bench J, Khoury J, Graman P. High-performance vision training  
2 improves batting statistics for University of Cincinnati baseball players. PLoS One  
3 2015;7:e29109.  
4
- 5 37) Camponogara I, Rodger M, Craig C, Cessari P. Expert players accurately detect an  
6 opponent's movement intentions through sound alone. J Exp Psychol Human  
7 2017;43:348-359.  
8
- 9 38) Ward P, Williams AM, Bennett SJ. Visual search and biological motion in tennis. Res Q  
10 Exercise Sport 2002;73:107-112.  
11
- 12 39) Appelbaum GL, Erickson G Sports vision training: a review of the state-of-the-art in  
13 digital training techniques. Int Rev Sport Exer P 2018;11:160-189.

1 **Notes**

2 Conflicts of interest: The authors certify that there is no conflict of interest with any financial  
3 organization regarding the material discussed in the manuscript.

4

5 Authors' contributions: Robert KELLY contributed to the study conceptualisation, data  
6 collection, data analysis, and writing. James W. ROBERTS contributed to the study  
7 conceptualisation, experimental set-up, data analysis, and writing. All authors read and  
8 approved the final version of the manuscript.

1 **Figure Captions**

2 Figure 1. Representative illustration of the screen display and visual stimuli. Landolt-C ring  
3 continuously moved in the horizontal or vertical direction across the mid-line. In this  
4 example, the progressively shaded rings imply a left-to-right motion (not present within  
5 reality). Image is drawn to scale with the current ring size being equivalent to 1.00 logMAR  
6 ( $.833^\circ$ ) relative to the screen.

7

8 Figure 2. Mean logMAR ( $\pm$ SE) acuity scores from the dynamic task as a function of group  
9 (skilled cricketers, non-cricketers), velocity (high, medium, low) and direction (horizontal,  
10 vertical). Error bars represent standard error of the mean.

1 **Tables**

2 Table 1. Mean (95% CI) individual participant slope coefficients pertaining to the relation  
3 between logMAR scores and object velocities (15°/s, 9.15°/s, 3.06°/s) for both horizontal and  
4 vertical stimuli. Values may be interpreted as the amount of increase in logMAR following a  
5 1°/s increase in object velocity.

	Skilled Cricketers	Non-Cricketers
Horizontal	.0051 (95% CI .0021 .0081)	.0099 (95% CI .0063 .0134)
Vertical	.0044 (95% CI .0011 .0078)	.0077 (95% CI .0053 .0102)

6

## 1 Footnotes

- 2 1) Posteriori power analysis was conducted using G\*Power software (version 3.1.9.4)<sup>27</sup>,  
3 including the following input parameters:  $\alpha = .05$ ,  $partial \eta^2 = .23$  (see *Results*; group x  
4 velocity interaction),  $n = 16$  (2 groups). Power ( $1-\beta$ ) was reported at .76. Combined with  
5 the knowledge that the statistical outcomes were consistent with our hypothesis and  
6 previous literature (e.g., <sup>7,8,10</sup>)<sup>28</sup>, it would suggest that there were no such false negative  
7 (Type II) or positive (Type I) errors.
- 8 2) The proportion of response errors indicated that there were no significant main, or  
9 interaction effects featuring the factor of group (group, group x velocity, group x velocity  
10 x direction:  $F_s < 1$ ; group x direction interaction approached significance,  $F(1, 14) =$   
11  $4.40$ ,  $p = .055$ ,  $partial \eta^2 = .24$ ). There was a significant main effect of velocity,  $F(2, 28)$   
12  $= 3.44$ ,  $p = .046$ ,  $partial \eta^2 = .20$ , and direction,  $F(1, 14) = 16.99$ ,  $p = .001$ ,  $partial \eta^2 =$   
13  $.55$ , but no velocity x direction interaction,  $F(2, 28) = 1.28$ ,  $p = .29$ ,  $partial \eta^2 = .08$ .
- 14 When reviewing our raw data, we determined that our stimulus programme failed to  
15 accurately evaluate the responses to the gap facing downward by not positively  
16 discriminating the correct and erred responses. This programing fault likely caused the  
17 record of response errors to appear inflated (grand  $M = 27.66\%$ ,  $SD = 3.34$ ). That said, a  
18 series of one-sample t-tests that compared the recorded response errors and the error rate  
19 assumed by chance alone (75%) showed a significantly lower-than-chance outcome for  
20 all of the groups and stimulus conditions (range  $ts = 17.49-28.32$ ,  $ps < .001$ ).