



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Influence of Aridity on Carbon Isotope Discrimination in Leaves of Stipa and Other C3 Species in Central Asian Grassland

M. Wittmer

Technische Universität München, Germany

Karl Auerswald

Technische Universität München, Germany

R. Tungalag

National University of Mongolia, Mongolia

Chunhua Bai

Chinese Academy of Sciences, China

Yongfei Bai

Chinese Academy of Sciences, China

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/1-3/43>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

M. Wittmer, Karl Auerswald, R. Tungalag, Chunhua Bai, Yongfei Bai, and Hans Schnyder

Influence of aridity on carbon isotope discrimination in leaves of *Stipa* and other C3 species in central Asian grassland

M. Wittmer¹, K. Auerswald¹, R. Tunglag², Chunhua Bai³, Yongfei Bai³, H. Schnyder¹

¹ Technische Universität München, Lehrstuhl für Grünlandlehre, Am Hochanger 1, D-85350 Freising-Weihenstephan. E-mail: auerswald@wzw.tum.de. ² National University of Mongolia, Ikh surguulin gudalmj-1, Baga toiruu, Sukhbataar district, Ulaanbaatar, Mongolia. ³ Institute of Botany, Chinese Academy of Sciences 20 Nanxincun, Xiangshan 10009, Beijing, China.

Key words: carbon isotope discrimination, *Stipa*, C3 plants, aridity, central Asian grassland

Introduction The carbon isotope discrimination (Δ) of C3 ecosystems is sensitive to water availability. Mean annual precipitation (MAP) is the most convenient and in some cases even the only known property to quantify water availability. The study reports the effects of MAP on Δ of *Stipa* species and other C3 plants.

Material and methods *Stipa* species were sampled along aridity transects in Inner Mongolia (China) and in the Republic of Mongolia in 2005 and 2006. Δ of *Stipa* was compared with published data of *Stipa* and of other C3 species (including samples collected along the 2005 and 2006 transects and published data) covering several years and regions around Mongolia. Weather data were taken from Climate Source Inc. and the NOAA NNDC network and geostatistically interpolated to obtain temporally and spatially resolved information for the sampling sites.

Results and discussion In 2005 and in 2006 Δ for *Stipa* species increased linearly with MAP. The slope (Figure 1 a) was the same in both years there was an offset of 1.1‰. The latter was caused by a difference of about 0.7 mm day⁻¹ of precipitation during the growing season. The Δ response was the same in both years when Δ was related to the year-specific mean daily precipitation during the growing season (Figure 1 b). An unbiased generalized relation of Δ with MAP was derived for *Stipa* species in Central Asian grassland. It has a slope of 0.0063‰ mm⁻¹ and predicts Δ for average growing seasons. The effect on Δ of deviations of actual from mean precipitation can be accounted for. The generalized relation of Δ with MAP was validated with published *Stipa* data (Figure 2 a). The same relationship held true for other C3 species (Figure 2 b).

Conclusions The relationships established in this study can be used to estimate the mean Δ of C3 communities from MAP or annual precipitation during the growing period and *vice versa*.




Figure 1 (a) Relation between Δ and MAP for *Stipa* species for the years 2005 (●, dashed regression line) and 2006 (□, solid regression line). (b) Relation between Δ of *Stipa* sp. and year-specific mean daily precipitation during the growing period (April–August) for the years 2005 (●) and 2006 (□).




Figure 2 Relation between Δ and mean daily precipitation (a) *Stipa* sp. from this study and from literature. (b) C3 species from this study and from literature.

Bold lines denote 95% confidence interval of the individual values, dashed lines 95% confidence interval of the regression and thin line the regression, all calculated from *Stipa* collected in 2005 and 2006.

Reference

NOAA NNDC Climate Data Online (2006). <http://cdo.ncdc.noaa.gov/CDO/cdo>.