



University of Kentucky  
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

## Citric Acid Accumulation of *Puccinellia tenuiflorais* a Specific Adaptive Response to Alkaline Stress

Hong Yan

*Northeast Normal University, China*

Shangjun Yin

*Zhejiang Wanli University, China*

Wei Zhao

*Air Force Avigation University, China*

Decheng Shi

*Northeast Normal University, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/1-4/50>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Citric acid accumulation of *Puccinellia tenuiflora* is a specific adaptive response to alkaline stress

Hong Yan<sup>1</sup>, Shang-jun Yin<sup>2</sup>, Wei Zhao<sup>3</sup>, De-cheng Shi<sup>1\*</sup>


<sup>1</sup> College of Life Sciences, Northeast Normal University, Changchun, PR China. <sup>2</sup> Department of Life Sciences, Zhejiang Wanli University, Ningbo, PR China. <sup>3</sup> The Department of Avigation Lifesaving, Air Force Avigation University, Changchun, PR China. E-mail address shidc274@nenu.edu.cn

**Key words :** *Puccinellia tenuiflora*, saline stress, alkaline stress, organic acid, citric acid, proline


**Introduction** *Puccinellia tenuiflora* is typical forages found in saline-alkaline habitats of Songnen Grassland. Owing to the special osmregulation, the species could survive in high pH conditions. In order to understand the physiologic characteristic of *P. tenuiflora* alkali tolerance, citric acid accumulation under the alkaline stress was studied.

**Materials and methods** These treatments were divided randomly into 15 groups (3 replicates per group). One group was used as the control (CK). Five groups were treated with Na<sub>2</sub>CO<sub>3</sub> solutions. Six groups were treated with NaCl solutions with concentrations. The remaining 3 groups were treated with 100 mmol l<sup>-1</sup> Na<sub>2</sub>CO<sub>3</sub> solutions, which were neutralized pH7, pH8 and pH9 using 85% H<sub>3</sub>PO<sub>4</sub>. Proline concentration was determined according to Bates *et al.* (1973). The concentration of total organic acid was determined according to Jing and Ding (1981), and citric acid was determined using the pentabromoacetone method adapted from Shi and Yin (1993).

**Results** Citric acid concentrations of *P. tenuiflora* increased with increasing stress strength by Na<sub>2</sub>CO<sub>3</sub> ( $p < 0.05$ ,  $R = 0.967$ ). Under NaCl stress, citric acid concentration decreased slightly ( $p > 0.05$ ) with increasing stress strength (Figure 1). Changes in the concentration of citric acid and total organic acid with increasing stress strength were almost parallel in the alkaline stressed seedlings of *P. tenuiflora* (data not shown). The proline concentration of *P. tenuiflora* also increased with the increases of NaCl and Na<sub>2</sub>CO<sub>3</sub>. In neutralization treatments, citric acid concentrations of *P. tenuiflora* decreased with decreasing pH value ( $p < 0.05$ ,  $R = 0.963$ ). Proline concentration of changed irregularly with pH value decreasing.



**Figure 1** Comparison of citric acid and proline accumulation between salt stress and alkaline stress in *P. tenuiflora*.



**Figure 2** Effects of neutralization on the concentrations of proline and citric acid in *P. tenuiflora*.

**Conclusions** Under alkaline stress, the organic acid metabolism of *P. tenuiflora* was changed, mainly due to the accumulation of citric acid. The citric acid accumulation was a specific physiological response to alkaline stress. The proline accumulation was physiological response to osmotic stress and the primary physiological significance of proline accumulation was osmotic adjustment.

### Reference

Shi, D. C., and Yin, L. J. (1993). Difference between salt (NaCl) and alkaline (Na<sub>2</sub>CO<sub>3</sub>) stresses on *Puccinellia tenuiflora* (Griseb.) Scribn. et Merr. plants. *Acta Botanica Sinica*, 35: 144-149. (in Chinese).