



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Comparison of Aggregates Content and Carbon Proportion of Top Soil in Perennial Forage

Lu Chen

Lanzhou University, China

Yuying Shen

Lanzhou University, China

Jing Yang

Lanzhou University, China

Yining Niu

Lanzhou University, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/2-1/24>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Comparison of aggregates content and carbon proportion of top soil in perennial forage

Lu Chen, Yuying Shen, Jing Yang, Yining Niu

College of Pastoral Agriculture Science and Technology, Lanzhou University 730020 China; E-mail: yy_she@lzu.edu.cn

Key words: total carbon (TC), total organic carbon (TOC), water-stable aggregates

Introduction Carbon storage in cropland soil, a potentially huge carbon pool, accounts for approximately 10% of global terrestrial carbon storage. In croplands, tillage is one of the key drivers behind the degradation of soil stability and leads to the accelerate decomposition of soil organic carbon. Perennial forages are widely accepted as being beneficial to the storage of soil organic matter and the formation of the soil aggregates, which can effectively improve and enhance soil fertility. In this study the carbon sequestration rate was estimated to be on average $0.332 \text{ Mg} \cdot \text{ha}^{-1} \cdot \text{a}^{-1}$ after cropland was converted to forage land, this outcome has positive effects in terms of improving sustainable agriculture, including a reduction in CO₂ emissions. The purpose of this study was to compare the carbon fixation rates of various perennial forage at different topsoil depths.

Materials and methods This study was carried out in the western Loess plateau ($35^{\circ}40' \text{ N}$, $107^{\circ}51' \text{ E}$), which has an annual rainfall of 562 mm. In 2002, seven different species of forage were sown. In August of 2007, soil samples from 0-5cm and 5-10cm depths were collected, with six subsamples collected using a cutting ring (volume of 200 g cm^{-3}) to determine bulk density. The proportion of water-stable aggregation ($>0.25 \text{ mm}$) was measured by wet sieving. Total organic carbon (TOC), total carbon (TC) were determined through a combustion method using liquiTOC (elementar, Germany).

Results For the 0-5cm layer, water-stable aggregates ($>0.25 \text{ mm}$) was the highest under bromegrass. The TC ranged between $19.687\text{--}25.75 \text{ g/kg}$, with the highest under clover. Every index measured was significant lower in the 5-10cm layer compared to 0-5cm layer, with water-stable aggregates reduced by 50%, and TOC and TC declining by approximately 20% (see table). A significant positive correlation was found between water-stable aggregates ($>0.25 \text{ mm}$) and TOC/TC ($R=0.340^*$ and 0.448^{**} , respectively).

Table 1 Water-stable aggregates ($>0.25 \text{ mm}$) carbon content and bulk density of 7 different perennial forages.

Sample	Layer	Water-stable aggregate ($>0.25 \text{ mm}$)%	Bulk density (g/cm^3)	TC (g/kg)	TOC (g/kg)	TOC/TC
AM sweet pea	0-5 cm	16.873 a	1.066 ab	19.687 a	10.163 a	0.514 a
Bromegrass		33.988 b	0.986 a	22.623 b	13.259 bcd	0.585 b
Clover		17.536 a	0.983 a	25.750 c	15.062 d	0.584 b
Crow toe		21.447 a	1.152 b	20.438 ab	10.354 a	0.506 a
Lucerne cv. longdong		19.526 a	1.155 b	21.036 ab	11.421 abc	0.541 a
Lucerne cv. saditi		18.038 a	1.119 ab	19.774 a	10.893 ab	0.547 ab
Sainfoin		17.376 a	1.149 b	23.125 bc	13.778 cd	0.585 b
AM sweet pea	5-10 cm	10.599 a	1.139 a	17.460 bc	7.931 ab	0.454 bc
Bromegrass		15.046 b	1.191 ab	17.664 bc	8.213 bc	0.465 bc
Clover		8.282 a	1.272 bc	18.518 d	8.304 bc	0.448 d
Crow toe		10.272 a	1.247 bc	17.754 bc	7.578 a	0.427 bc
Lucerne cv. longdong		8.894 b	1.197 ab	18.084 cd	8.573 c	0.474 cd
Lucerne cv. saditi		8.147 a	1.250 bc	16.347 a	7.491 a	0.458 a
Sainfoin		10.89 a	1.325 c	17.339 b	8.232 bc	0.475 b

Conclusions In this study the organic carbon of perennial forage in the topsoil accounted for over 50% of the total carbon, with water-stable aggregates and TOC showing significant differences between 0-5cm and 5-10cm layer. This finding indicates that tillage disturbs the soil causing the spatial distribution of organic matter to become more homogenous. Further, tillage has no beneficial effect in terms of both carbon fixation and maintaining soil health. Lastly this study indicated that different perennial forages result in significant differences in organic carbon fixation.