



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Effects of Fertilizer and Plant Density on the Fresh Biomass of Sudan Grass

Zhongsheng Zhi

*Inner Mongolia Agricultural University, China*

Guiping Yao

*Inner Mongolia Agricultural University, China*

Jianguang Wang

*Inner Mongolia Agricultural University, China*

Meihua Gao

*Inner Mongolia Agricultural University, China*

Meihua Li

*Inner Mongolia Agricultural University, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/2-3/8>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Effects of fertilizer and plant density on the fresh biomass of Sudan grass

Zhongsheng Zhi,\* Guiping Yao, Jianguang Wang, Meihua Gao, Meihua Li  
College of Agronomy, Inner Mongolia Agricultural University, Huhhot, China, 010018, Phone: 0471-4301317,  
E-mail: mrzszs@126.com

**Key words:** Sudan grass, Neinong No.1, mathematic model, yield, planting density, fertilization

**Introduction** *Sorghum sudanense* Stapf, cv. Neinong No.1 was a hybrid of Sorghum and Sudan grass bred by our group. The objective of this study was to evaluate the effects of planting density and fertilizing level on fresh matter (FM) yield of Neinong No.1. The optimum cultivation condition was obtained.

**Material and methods** Sudan grass No.1 was seeded with four randomized replications for 11 treatment plots and one control plot in a greenhouse in 2006. The area of each plot was 14m<sup>2</sup>. The experimental field located in Huhhot. Data were analyzed with SAS 9.0 software.

### Results and analysis

**Regression analysis of FM yield of Sudan grass** The relation between FM yield and other factors were analyzed with the regression simulation method at different harvest time. One regression equation was obtained with stepwise regression:

$$y^2 \text{ (FM weight (kg) / 2 m}^2\text{)} = -3.35695 + 0.03885x_1 + 3.87533x_2 + 0.06993x_3 - 0.36936x_4$$

It is clear that the grass height ( $x_2$ ) plays a major role in Sudan grass production. However, germination rate ( $x_1$ ) had a minor influence on the FM yield.

**Single-factor analysis** For two harvest times: July 19 and September 2, the differences of FM yield between 11 treatment plots and one control plot were not significant ( $p > 0.05$ ). If analyzed with single-factor analysis of variance, however, the FM yield of Sudan grass in No.9 treatment plot was highest ( $p < 0.15$ ), which was 14613.97kg/667m<sup>2</sup>.

**Conclusion** In current study, the results showed that cultivating measurement combination of No.9 plot, in which planting density was 45022 plants/ha and 20kg additional urea was applied to the plot during the growth, was optimum for the maximum yield of FM of tested Sudan grass, which was harvested for four times.