



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Research on Rejuvenation and Replacement Cultivated Technology of *Salix psammophila* in the Hobq Desert

Zhiying Mi  
*Inner Mongolia Agricultural University, China*

Yong Gao  
*Inner Mongolia Agricultural University, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/4-2/10>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Research on rejuvenation and replacement cultivated technology of *Salix Psammophila* in the Hobq desert

Mi Zhiying ,Gao Yong


College of Ecol. and Env. Sci ,Inner Mongolia Agric. Univ. , Huhhot , Inner Mongolia 010018 P .R .China .E-mail : nmgmizhiying@163 .com

**Key words :** the Hobq desert ,*Salix psammophila* ,rejuvenation and replacement ,height growth ,stem growth


**Introduction** *Salix psammophila* is an important species of windbreak , sand-fixation and industrial timber forest in arid and semi-arid area . Recently , Ecological Environment of natural *Salix psammophila* forest was destroyed severely . In this study , We tried to find the effect on height growth and stem growth of *Salix psammophila* .for four kinds of different treatments including Installation of Artificial Sand-break ,Leveling Stubble Along the Surface of the Ground ,Usual Leveling Stubble , Fertilization .

**Materials and methods** The research was about declining *salix psammophila* bushes in the east of the Hobq Desert . Technical methods :(1) Installation of Artificial Sand-break (Specification : 20cm high , checker board , porosity 20% -25% ) .(2) Leveling Stubble (Along the Surface of the Ground) (3) Usual Leveling Stubble (remains 20cm Stubble) . (4) Fertilization (dug a ditch of 20 cm deep filling 2 kg organic fertilizers around each *salix psammophila* bushes) .The data of four kinds of different treatments were analysed by using SAS9 .0 .

**Results** The results showed that four kinds of different treatments were all beneficial to height growth of *salix psammophila* bushes ( $P < 0.05$ ) . Compored with check sample , height growth of *salix psammophila* bushes which were treated with Artificial Sand-break ,Leveling Stubble Along the Surface of the Ground ,Usual Leveling Stubble ,Fertilization had increased by 10 .07% ,6 .06% ,12 .91% ,23 .18% ,respectively (Figure 1) . It is noteworthy that four kinds of different treatments were helpful to stem growth of *salix psammophila* bushes ( $P < 0.05$ ) . Compored with check sample , stem growth of *salix psammophila* bushes which were treated with Artificial Sand-break ,Leveling Stubble Along the Surface of the Ground ,Usual Leveling Stubble ,Fertilization had increased by 5 .20% ,3 .28% ,4 .90% ,3 .80% ,respectively (Figureure , 2) .



**Figure 1** Height growth of *Salix psammophila* by different treatment in three years .



**Figure 2** Stem growth of *Salix psammophila* by different treatment in three years .

**Conclusions** The results showed that four kinds of different treatments were all beneficial to rejuvenation and replacement of *salix psammophila* bushes ( $P < 0.05$ ) . Comprehensive effect of Fertilization was helpful to height growth of *salix psammophila* bushes . On the other hand , Leveling Stubble Along the Surface of the Ground had much more effects on stem growth .