



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Role of Grazing Cattle on Seed Dispersal of Plants in a Hill Pasture 1. Effect of Sward Structure on Grass Seed Ingestion by Cattle

M. Obara
Tohoku University, Japan

Shin-ichiro Ogura
Tohoku University, Japan

T. Shishido
Tohoku University, Japan

K. Sugawara
Tohoku University, Japan

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/3-1/33>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Role of grazing cattle on seed dispersal of plants in a hill pasture 1 . Effect of sward structure on grass seed ingestion by cattle

M . Obara , S . Ogura , T . Shishido and K . Sugawara

Graduate School of Agricultural Science , Tohoku University , Kawatabi , Naruko-onsen , Osaki , 989-6711 Japan . E-mail : s-ogura@bios.tohoku.ac.jp .

Key words : germinability , grazing cattle , seed dispersal , seed ingestion , sward structure

Introduction Seed dispersal by defecation of grazing herbivores is one of the significant factors in vegetational change (e.g . , Janzen , 1984 ; Archer and Pyke , 1991) . It is well known that the seeds of certain plant species are ingested by grazing herbivores and disseminated through the digestive tract . Recent studies conducted in Japanese hill pastures have shown that grazing cattle dispersed majority of seeds of monocotyledons such as *Carex albata* (Watanabe *et al.* , 2000 ; Obara *et al.* , 2005) . In this paper , we examined factors affecting seed ingestion by cattle using a sward-based measurement to learn timing of seed ingestion by cattle relative to seed maturity .

Materials and methods

Relationship between seed maturity and the timing of seed ingestion by cattle A hill pasture grazed by 10 beef cattle was used in our study . Fifty seedheads of three grass species (*Anthoxanthum odoratum* , *Poa pratensis* and *C. albata*) were tagged during late spring and the frequency of defoliation by cattle were recorded at 2-5 day intervals until late June . Seed germinability (30 days incubation in 15°C/12 hr light and 25°C/12 hr light) of the 3 species was also investigated by sampling of 30 seeds on each day measurements were obtained .

Sward canopy structure of 4 monocotyledon species in their reproductive stage

Plots (20 cm×20 cm each) were located in vegetation patches where *A. odoratum* , *Agrostis alba* , *C. albata* and *P. pratensis* were dominant in the pasture , on the first day of the study . Mean height of leaf mass and seedheads were measured at 10-37 day intervals during the grazing season .

Results and discussion *A. odoratum* seeds were defoliated by cattle in mid-to late May , whereas in June when the seeds matured no seed consumption by cattle was observed . In contrast , the seeds of *P. pratensis* and *C. albata* were grazed after seeds were mature .

Measurements in sward canopy structure showed that , while the height of seedheads in *A. odoratum* and *A. alba* was significantly higher than that of leaves , the height of seedheads in *C. albata* and *P. pratensis* was almost same as that of leaves (Figure 1) . The results suggest that it is difficult for grazing cattle to choose leaf mass over seedheads from the sward canopy structure in which both leaves and seeds are contained within a same grazing horizon , particularly if the density of tillers having seedheads is high .

Conclusions Plant architecture with the height of seedheads equivalent to vegetative tillers is an effective strategy for *C. albata* and *P. pratensis* to disperse the seeds via grazing animals , because the frequency of defoliation of matured seeds by animals increases .

References

- Archer , S . , Pyke , D . A . , (1991) . Plant-animal interactions affecting plant establishment and persistence on revegetated rangeland . *Journal of Range Management* , 44 : 558-565 .
 Janzen , D . H . , (1984) . Dispersal of small seeds by big herbivores : foliage is the fruit . *American Naturalist* , 123 : 338-353 .
 Obara , M . , Ogura , S . , Shishido , T . , Sugawara , K . , (2005) . Seed dispersal by defecation of cattle rotationally grazed in sown and native pastures . *Japanese Journal of Grassland Science* , 52 : 6-16 . (in Japanese with English abstract) .
 Watanabe , N . , Nishiwaki , A . , Sugawara , K . , (2002) . Dissemination of *Carex albata* Boott seeds by grazing cattle . *Grassland Science* , 48 : 142-145 .




Figure 1 Mean height of leaves and seedheads in the 4 monocotyledons . Values with different letter in each date represent significant difference (Tukey's test , $P < 0.05$) . n : not significant . * : n = 1 , ** : n = 2 (seedhead data) .