



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Bush Encroachment: A Major Threat to Pastoralists Livelihood in Ethiopia

Ebro Abule

*Adami Tulu Agricultural Research Center, Ethiopia*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/6-1/12>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Bush encroachment : a major threat to pastoralists livelihood in Ethiopia

Abule , Ebro

A dami Tulu Agricultural Research Center , P .O .Box , 35 , Zeway , Ethiopia , E-mail : abule \_ ebro@ yahoo .com

**Key words** : Bush encroachment , Pastoralists livelihood , rangeland

**Introduction** Ethiopia has a vast area of rangeland (61 to 67% of the land area) located around the periphery and they are found below 1,500 m a.s.l. They support pastoral and agro-pastoral communities (mostly Somalia , Afar and Borana) of about 9.8 million people belonging to 29 ethnic groups . The rangelands are not only important for livestock raising but also for wildlife habitats , ecotourism , irrigated agriculture , medicinal plants , conservation of biodiversity , and exploitation of natural resources (Abule *et al.* , 2005) . However , bush encroachment , the process of open grassland savanna being transformed into bushes & shrubs , is a major problem . Therefore , the objective of this paper is to review the status of bush encroachment and its effect on the livelihood of the pastoralists in Ethiopia .

**Materials and methods** Ethiopia , with a land area of 1,104,300 km<sup>2</sup> , is located in the horn of Africa bordered in the north by Eritrea and Djibouti , in the south by Kenya , in the east by Somalia and in the west by Sudan . The rangelands have unreliable and erratic rainfall , with regularly high temperature . This paper is based on reviewing literatures and documents .

**Results** According to the pastoralists compared to the past , the rangelands are bush encroached and the major triggering factors were the lack of prescribed fire , severe overgrazing , drought and the expansion of farming into the rangelands . Accordingly , there is a change in the species composition of livestock they keep from grazer (cattle and sheep) to browsers (camel and goats) . Furthermore , the pastoralists have noticed that bush encroachment is leading to agricultural and biodiversity problems . It has created a decrease in grass production , difficulty in livestock herding , damage by wildlife , and problems of bloating . The encroaching species and their density also varied from rangeland to rangeland . The major encroaching species in the Afar region are *Prosopis juliflora* , *Acacia seyal* , *Acacia mellifera* and *Acacia senegal* while in the Borana rangeland species of *Acacia* and *Commiphora* and in parts of the Somalian rangeland , *Acacia nubica* and *A . Mellifera* . The communal grazing lands are more bush encroached (greater than 2,500 woody plants/ha) than the other grazing types and *Parthineum hysterophores* is becoming a major concern . In some rangelands , the bush encroached area is about 52% of the total land area . There is no visible bush control method being undertaken by the pastoral communities on their own effort except in the Borana rangeland which was supported by different organizations (Oba *et al.* , 2000 ; Gemedo , 2004 ; Abule *et al.* , 2005 ; Admasu , 2006 ; Amaha , 2006 ; Belaynesh , 2006 ; Teshome , 2007) .

**Conclusions** Bush encroachment has created a difficulty for the pastoralists to plan effective resource management and utilization strategies . Accordingly , the control of bush encroachment must be given due attention which requires a proper understanding of causal factors , invasive species , the degree and extent of encroachment , the mechanism and the population dynamics of invasive species and long-term community based control programs .

### References

- Abule Ebro , Snyman HA and Smit GN . 2005 . Comparisons of pastoralists perceptions about rangeland resource utilization in the middle Awash valley of Ethiopia . *J . Envi Manage* . 75 : 21-35 .
- Admasu Terefe , 2006 . Pastoralists perceptions on range-livestock management practices and rangeland assessment in Hamer and Benna-Tsemay districts of South Omo Zone . An MSc Thesis Presented to the School of Graduate Studies of Alemaya University , Ethiopia . 159p .
- Amaha Kassahun , 2006 . Characterization of rangeland resources and dynamics of the pastoral Production system in the Somali region of Estern Ethiopia . A PhD Thesis Presented to the University of the Free State , Bloemfontein , South Africa . 232p .
- Belaynesh Debalkie , 2006 . Floristic composition and diversity of the vegetation , soil seed bank flora and condition of the rangelands of the Jijiga Zone , Somali Regional State , Ethiopia . An MSc Thesis Presented to the School of Graduate Studies of Alemaya University , Ethiopia . 144p .
- Gemedo Dalle , 2004 . Vegetation Ecology , Rangeland Condition and Forage Resources Evaluation in the Borana Lowlands , Southern Oromia , Ethiopia . A PhD Thesis Presented to the University of the Göttingen , Germany . 253 pp .
- Oba G , Post E , Syvertsen PO and Stenseth NC . 2000 . Bush cover and range condition assessments in relation to landscape and grazing in southern Ethiopia . *L .scape Ecol* . 15 : 535-546 .
- Teshome , Abate . 2007 . Traditional utilization practices and condition assessment of the rangelands in Rayitu district of Bale zone , Ethiopia . An MSc Thesis Presented to the School of Graduate Studies of Haramaya University , Ethiopia . 129p .