



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Carbon Dynamics and Mitigation of Methane and Nitrous Oxide Emissions in Agroecosystems with *Pinus ponderosa* (Dougl. Ex Laws) and Native Pastures Established on Degraded Volcanic Soils in the Chilean Patagonia

Francis Dube

University of Concepción, Chile

Neal B. Stolpe

University of Concepción, Chile

Erick Zagal Venegas

University of Concepción, Chile

Miguel Espinosa Bancalari

University of Concepción, Chile

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/8-1/13>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Carbon dynamics and mitigation of methane and nitrous oxide emissions in agroecosystems with *Pinus ponderosa* (Dougl . Ex Laws) and native pastures established on degraded volcanic soils in the Chilean Patagonia

Francis Dube¹ , Neal Stolpe² , Erick Zagal Venegas² and Miguel Espinosa Bancalari¹

¹ Facultad de Ciencias Forestales , Universidad de Concepción , Victoria 631 , Casilla 160-C , Chile

² Departamento de Suelos , Facultad de Agronomía , Universidad de Concepción , Vicente Méndez 595 , Casilla 537 , Chillan , Chile . Corresponding author : fdube@udec.cl

Key words : carbon sequestration , silvopastoral systems , volcanic soils , CO₂ FIX

Introduction During the 1990s , terrestrial ecosystems captured approximately 36% of the total carbon liberated to the atmosphere by combustion of fossil fuels . Consequently , there is a growing interest to study the potential of carbon sequestration in presently non-sustainable agro-ecosystems worldwide , including remote regions such as Patagonia , where large areas of degraded pastures and eroded soils are also encountered , and ranchers face increasing pressures to maintain the cattle-raising productivity of their land . They are encouraged to adopt silvopastoral systems as more sustainable practices that satisfy their socioeconomic necessities and contribute to mitigate atmospheric CO₂ . According to the 3rd IPCC report (2001) , the use of agroforestry systems in degraded lands constitute effective C sinks , especially with the use of perennial pasture and fast growing trees . The objective of this study is to investigate and model the potential to sequester C in an exotic short rotation forest plantation and degraded grasslands of the Chilean Patagonia .

Materials and methods The site was located at 730 m altitude at S 45°25' W 72°00' near Coyhaique , Chile . The soil had low bulk density (< 0.9 g cm⁻³) and high P fixation values (65-89%) . It is classified as medial , amorphic , mesic Typic Hapludands . Treatments were imposed in a randomized complete design with three replicates . Soil samples were analyzed to determine soil organic carbon (SOC) , microbial C and N , and soil respiration (C-CO₂ evolution) . Measured parameters included tree and pasture biomass . Measured SOC values were compared with those predicted by CO₂ FIX , previously calibrated to the site conditions . Mean monthly temperature and rainfall during tree growth period , current annual increment of trees , and proportions of C in stems , leaves , branches and roots were used to calibrate the model .

Results Preliminary results show that the contents of SOC are greater than those predicted by the model (Figure 1) , which indicate the need to adjust the soil parameters so that simulations better reflect reality . The unexpected result may be due to the presence of volcanic soils which have distinctive properties including allophanic clays , and higher C contents in surface soil compared to non-volcanic soils . Additionally , soil respiration was highest in the pine plantation (Figure 2) and microbial biomass was highest in the prairie (Figure 3) . Between agroecosystems , soil respiration was not correlated to SOC nor microbial biomass , perhaps due to the larger amount of roots and the presence of mycorrhizae in the plantation . Estimates of CH₄ and N₂O emissions were highest in the prairie . However , pine-based silvopastoral systems would permit addressing cattle-raising and timber production needs and allow for the sequestration of larger amounts of C in above and belowground components of plants and soil of the agroecosystem .




Figure 1 SOC measured and simulated at 0 to 40 cm depth , Chilean Patagonia .




Figure 2 Mean soil respiration at 0-40 cm depth , Chilean Patagonia .




Figure 3 Mean C and N microbial biomass at 0 to 40 cm depth .

Conclusions Once the model is recalibrated to soil conditions , it will be used with an adjacent six-year-old pine-based silvopastoral system arranged in strips to estimate total carbon balance , and the results will be used to construct C cycling models . Given the more efficient utilization of site resources and the presence of favorable microclimate , the silvopastoral system will permit a higher annual total biomass production and C capture in comparison with the prairie and the plantation ; also , the presence of highly active aerial and subterranean C cycles will result in a large increase in C capture . The silvopastoral system will have the better potential to mitigate global warming through increased sequestration of greenhouse gases , while also being a more sustainable form of land use in the long term .

Reference

IPCC . 2001 . Climate change 2001 : the scientific basis . Cambridge Univ . Press , Cambridge , UK , 881p .