



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Seed Protein Variability and Abiotic Stress in Naturalised Italian Ryegrass (*Lolium multiflorum* Lam.)

Mónica B. Sacido

Universidad Nacional del Centro de la Provincia de Buenos Aires, Argentina

Ezequiel M. Latorre

Universidad Nacional del Centro de la Provincia de Buenos Aires, Argentina

W. John Rogers

Universidad Nacional del Centro de la Provincia de Buenos Aires, Argentina

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/6-2/9>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Seed protein variability and abiotic stress in naturalised Italian ryegrass (*Lolium multiflorum* Lam.)

Sacido Mónica B.¹, Ezequiel M. Latorre¹, ex aequo and W. John Rogers^{1,2}

¹ Facultad de Agronomía, Universidad Nacional del Centro de la Provincia de Buenos Aires, C.C. 47, (B7300) Azul, República Argentina, ² Researcher of the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), República Argentina. E-mail msacido@faa.unicen.edu.ar

Abstract : Electrophoretic (SDS-PAGE) analysis of seed proteins was employed to reveal genetic variability in naturalized populations of *Lolium multiflorum* from unmanaged land in a region of the Province of Buenos Aires located in the Flooding Pampas of Argentina. The populations came from the following four site types, defined according to the species composition of the plant communities found at each: (i) a community designated 'loam' that was the least edaphologically and topographically limiting; (ii) 'half-loam' that presented intermediate limitations of drainage and alkalinity; (iii) 'alkaline lowland' that presented the most severe limits of alkalinity; and (iv) 'humid lowland' that only presented hydro-morphological limits. SDS-PAGE applied to samples obtained from each population revealed multiple banding patterns that could be divided into six electrophoretic zones, A to F. Individual bands were further sub-classified as follows: A1, A2, A3, B1, B2, B3, B4, B5, C1, C2, C3, C4, C5, D1, E1, E2 y F1. The banding patterns of each population sample were compared, once the minimum sample size required to represent the variability of the each population was derived by the Method of Pielou. In some comparisons, discrimination between populations could be made by observing only one band; for example, band C1 discriminated 'alkaline lowland' from 'humid lowland' and band C4 'half-loam' from 'alkaline lowland'. In other comparisons, it was necessary to observe several bands simultaneously. By this means, all the six pair-wise comparisons between the four populations provided sufficient information for discrimination to be made. If the divergence between populations is related to the conditions found at each topographical site, these populations may comprise a useful source of genetic variability for stress tolerance and might offer the potential for the development of new cultivars adapted to local conditions. We believe this to be the first report of the use of this methodology to characterise and distinguish natural populations of *L. multiflorum* found in different topographical positions in the grasslands of the region under study.

Key words : electrophoresis ; method of Pielou ; alkaline lowland ; humid lowland ; topographical position ; grassland




Figure 1 Example gel of observed protein profiles. Lanes 1-3 : standard cultivar Nui ; Lane 4 : half-loam ; Lanes 5-6 : loam ; Lanes 7-8 : alkaline lowland ; Lanes 9 : humid lowland . Arrows indicate the reference bands of cultivar Nui , originally designated in cultivar Dalita (ISTA , 1992) , while the lines on the right hand side represent the range of bands covered by each denominated zone A to F (modified classification of Galussi et al . , 1997) . Included is a plot produced by the programme Scion Image , where each trace corresponds to a gel lane and shows the peak area and relative position of each band .




Figure 2 Frequency profile for the bands significant in differentiating populations from distinct topographical positions ($t < 0.5\%$).