



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Carbon Dynamics (2000-2006) over the Northern Great Plains Grasslands

Li Zhang
U.S. Geological Survey

Bruce Wylie
U.S. Geological Survey

Lei Ji
U.S. Geological Survey

Tagir Gilmanov
South Dakota State University

Larry L. Tieszen
U.S. Geological Survey

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/8-1/45>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Carbon dynamics (2000—2006) over the northern Great Plains grasslands

Li Zhang¹, Bruce Wylie¹, Lei Ji¹, Tagir Gilmanov², and Larry L. Tieszen³

¹SAIC, contractor to U.S. Geological Survey (USGS) Center for Earth Resources Observation and Science (EROS), Sioux Falls, SD, 57198, USA. E-mail: lizhang@usgs.gov Work performed under USGS contract 03CRCN0001.

²South Dakota State University, Brookings, SD 57007, USA, ³USGS/EROS, Sioux Falls, SD 57198, USA

Key words : eddy covariance, flux tower, gross primary production, net ecosystem exchange

Introduction The grassland ecosystem in the U.S. Great Plains occupies was about 1.5 million km². However, the contribution of the grasslands to local and regional carbon budgets remains uncertain due to the lack of carbon flux data for the expansive grassland ecosystems under various land managements, land uses, and climate variability. An understanding of carbon fluxes across the ecosystem is essential for developing carbon budget models at regional, national, and global scales.

Materials and methods A remote sensing-based empirical model, piecewise regression model (Wylie et al., 2007; Zhang et al., 2007), was modified to estimate the grassland carbon fluxes in the U.S. northern Great Plains. The model explored the empirical relationship between environmental variables and tower-based measurements collected from six grassland flux towers and extrapolated the tower-measured data to the entire grassland ecosystem. We used this model to estimate the spatio-temporal carbon fluxes across the study area. Net ecosystem exchange (NEE) between land and atmosphere were measured using eddy covariance and Bowen-Ratio techniques. We partitioned 30-minute CO₂ fluxes into total ecosystem respiration and gross primary production (GPP) using the light response curve analysis (Gilmanov et al., 2005). The predictors include the MODIS normalized difference vegetation index (NDVI), precipitation, temperature, photosynthetic active radiation, and phenological metrics. We also incorporated the actual vegetation evapotranspiration data derived from a vegetation evapotranspiration (VegET) model (Senay and Henebry, 2007), which takes into account soil moisture and land surface phenology.

Results and discussion In the northern Great Plains grasslands, we estimated a 13 percent reduction of GPP in the dry year 2002 compared to the average GPP for years 2000-2006, which resulted in an anomalous net source of carbon dioxide (-19 gC/m²/year) to the atmosphere. The carbon budgets depend, to a great extent, upon precipitation and its distribution in this region. Specifically, the 2006 NEE map (Figure 1) shows carbon sources in drier western regions and carbon sinks in wetter eastern and southern regions, which was in line with the east-west gradient of annual precipitation across this region. The annual NEE varied from -28 g C/m²/year (source) in 2006 to 37 g C/m²/year (sink) in 2001, with sources in 2002, 2004, and 2006, and sinks in other years (Figure 2).




Figure 1 Spatial distribution of annual NEE for 2006.




Figure 2 Estimated annual NEE for 2000-2006.

Conclusions The CO₂ exchanges over the northern Great Plains grasslands are highly variable in space and time. The annual NEE transits from sinks to sources in the dry years of 2002, 2004, and 2006. If the climate becomes drier in the future, the ecosystems may change to larger carbon sources.

References

- Gilmanov, T.G., Tieszen, L.L., Wylie, B.K., Flanagan, L.B., Frank, A.B., Haferkamp, M.R., Meyers, T.P., Morgan, J.A., 2005. Integration of CO₂ flux and remotely sensed data for primary production and ecosystem respiration analyses in the Northern Great Plains: Potential for quantitative spatial extrapolation. *Global Ecology and Biogeography* 14 (3): 271-292.
- Senay, G.B., Henebry, G., 2007. Evaluating vegetation evapotranspiration (VegET) modeling results in South Dakota. In: Proceedings of the Eastern South Dakota Water Conference. October 29-31, Sioux Falls, South Dakota.
- Wylie, B.K., Fosnight, E.A., Gilmanov, T.G., Frank, A.B., Morgan, J.A., Haferkamp, M.R., Meyers, T.P., 2007. Adaptive data-driven models for estimating carbon fluxes in the Northern Great Plains. *Remote Sensing of Environment* 106: 399-413.
- Zhang, L., Wylie, B.K., Loveland, T., Fosnight, E.A., Tieszen, L.L., Ji, L., 2007. Evaluation and comparison of gross primary production estimates for the northern Great Plains grasslands. *Remote Sensing of Environment* 106: 173-189.