

University of Kentucky UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th International Rangeland Congress

Carbon Sequestration Potential of Organic Pasture in Thailand

Thumrongsakd Phonbumrung Mahidol University, Thailand

Suriyapong Watanasak Mahidol University, Thailand

Chirapol Sintunawa Mahidol University, Thailand

Charlie Navanukraha Mahidol University, Thailand

Follow this and additional works at: https://uknowledge.uky.edu/igc

Part of the Plant Sciences Commons, and the Soil Science Commons

This document is available at https://uknowledge.uky.edu/igc/21/8-1/49

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Carbon sequestration potential of organic pasture in Thailand


Thumrongsakd Phonbumrung¹, Suriyapong Watanasak¹, Chirapol Sintunawa¹, Charlie Navanukraha¹ ¹ Faculty of Environment and Resource Studies, Mahidol University, Salaya, Nakon Pathom, Thailand Correspondence: thumrongsakd[@] hotmail.com

Key words : carbon sequestration , organic pasture , environmental model


Introduction This paper provides the feature of a generic environmental model to demonstrate environmental impacts of the organic pasture used for beef production as compared with the intensive conventional pasture system. Both pastures had been utilized through the rotational grazing for 3 consecutive years. Carbon sequestration potential which is inversely associated with the Global Warming Potential was simulated.

The model The model was developed following the general stages of modeling strategies described in Forrester (1968). STELLA II version 9 software package (HPS, 1994) has been used in the model construction. The model described the interrelationships between the pasture production, cattle production, and soil organic carbon (SOC) accumulation in the farm boundary. The study focused on Northeastern Thailand . Two types of pasture systems selected for this investigation are the intensive conventional system and the organic system . The pastures were grass-legume mixed of Guinea grass (*Panicum maximum* TD 58) and Taphra stylo (*Stylosanthes guianensis* CIAT 184). The cattle were Thai Indigenous beef cattle . The system in this study involved the pasture production process , the cattle production process , the returning of waste from pasture and cattle into the soil organic carbon (SOC) , and soil nutrient and uptake by plants .

Results The organic pasture showed greater carbon sequestration potential (Figure 2) compared to the conventional pasture (Figure 1). Without external fertilization, the soil carbon under organic pasture would gradually increase from 10.0 to 12.75 t ha⁻¹ for the 20 year period. With 50 and 100 kg N ha⁻¹ yr⁻¹ fertilization, the carbon content in the pasture soils would increase to 26.01 and 35.57 t ha⁻¹, respectively. The carbon sequestration potential of the organic pastures with 0,50 and ,100 kg N ha⁻¹ yr⁻¹, were estimated to be 0.14, 0.80 and 1.28 t ha⁻¹ yr⁻¹, respectively.



1=0N 2=50 kg N ha⁻¹ yr⁻¹ 3=100 kg N ha⁻¹ yr⁻¹ Figure 1 SOC (t ha⁻¹ yr⁻¹) of an intensive pasture.



1=0N; $2=50 \text{ kg N ha}^{-1} \text{ yr}^{-1}$; $3=100 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ Figure 2 SOC (t ha⁻¹ yr⁻¹) of organic pasture.

References

Forrester, J. W. (1968) Principle of Systems. Cambridge. Wright-Allen Press Inc., Cambridge.

HPS-High Performance Systems (1994) An introduction to system thinking. High Performance Work System Inc., Hanover.