



University of Nebraska at Omaha
DigitalCommons@UNO

Student Work

12-1-1999

The effect of dredging on the plant community of a Missouri-River floodplain wetland.

Lisa Ann Peterson

Follow this and additional works at: <https://digitalcommons.unomaha.edu/studentwork>

Recommended Citation

Peterson, Lisa Ann, "The effect of dredging on the plant community of a Missouri-River floodplain wetland." (1999). *Student Work*. 3360.

<https://digitalcommons.unomaha.edu/studentwork/3360>

This Thesis is brought to you for free and open access by DigitalCommons@UNO. It has been accepted for inclusion in Student Work by an authorized administrator of DigitalCommons@UNO. For more information, please contact unodigitalcommons@unomaha.edu.



THE EFFECT OF DREDGING ON THE PLANT COMMUNITY OF A MISSOURI-
RIVER FLOODPLAIN WETLAND

A Thesis

Presented to the

Department of Biology

and the

Faculty of the Graduate College

University of Nebraska

In Partial Fulfillment

Of the Requirements for the Degree

Master of Arts

University of Nebraska at Omaha

by

Lisa A. Peterson

December 1999

UMI Number: EP74962

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI EP74962

Published by ProQuest LLC (2015). Copyright in the Dissertation held by the Author.

Microform Edition © ProQuest LLC.

All rights reserved. This work is protected against unauthorized copying under Title 17, United States Code




ProQuest LLC.
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106 - 1346

THESIS ACCEPTANCE


Acceptance for the faculty of the Graduate College,
University of Nebraska, in partial fulfillment of the
Requirements for the degree Master of Arts in Biology,
University of Nebraska at Omaha.

Committee





Chairperson



Date



ABSTRACT

A plant community gradient, consisting of Open-water, Bulrush, Grass, and Forest Zones, was evaluated both before (1995) and one-year after (1997) restoration-dredging of a wetland along the Missouri River in east-central Nebraska. Species diversity declined significantly ($P \geq 0.05$) in both the Bulrush and Grass Zones (-24 and -30 species) but not elsewhere. The Open-water Zone, which increased the most with dredging (+40 meters), was dominated by duckweed (*Lemna minor*) (35% canopy cover in 1995 and 45% in 1997) and watermeal (*Wolffia columbiana*) (53% and 61%) both before and after dredging. Coontail (*Ceratophyllum demersum*) increased significantly (7% to 25%). The greatest species decline occurred with river bulrush (*Scirpus fluviatilis*) (50% to 28%) both due to a substantial reduction in the areal extent of the Bulrush Zone (-14 m width) and a significant decline in canopy cover of bulrush within the zone (55% in 1995 to 29% in 1997). Reed canary grass (*Phalaris arundinacea*) declined significantly in all terrestrial zones (average decline of 31%). Despite declines in cover in these and other native wetland species, most survived dredging, suggesting that, at least in this one instance, dredging to restore the backwater habitat can be accomplished without a substantive loss of associated plant communities.

ACKNOWLEDGMENTS

I would like to express my appreciation to Dr. David Sutherland, whose plant identifications and editing of species lists proved invaluable; Dr. Jeffrey Peake and Michael Gilbert for their critical review of the manuscript; and to Dr. Thomas Bragg who is greatly responsible for getting this manuscript written.

I would like to thank both the University Committee on Research and the Department of Biology, University of Nebraska at Omaha, for providing financial support of this research. Also, my sincere gratitude is extended to the U.S. Army Corps of Engineers, whose technical and financial assistance made this study possible.

Finally, I would like extend a special thank you to Karen Lawrence, who provided professional assistance both in the field and in the analysis of data. However, it was her constant encouragement and support as a friend that helped me get through this process. Because of that, I dedicate this thesis to her.

TABLE OF CONTENTS

Title page	i
Thesis Acceptance page.....	ii
Abstract	iii
Acknowledgements.....	iv
List of Tables and Figures.....	vii
Introduction.....	1
Methods.....	2
Study Site	2
Background.....	4
Field Methods	5
Analysis.....	8
Results and Discussion	8
Conclusion	17
Literature Cited.....	18
Appendix.....	19
Appendix Table 1.....	20
Appendix Table 2.....	23
Appendix Table 3.....	24
Appendix Table 4.....	25
Appendix Table 5.....	27
Appendix Table 6.....	29

Appendix Table 7.....	31
Appendix Figure 1	32

LIST OF TABLES AND FIGURES

TABLES

- Table 1. Zone descriptions and characteristics.
- Table 2. Description of plant categories.
- Table 3. Average zone width and changes between 1995 and 1997.
- Table 4. Mean canopy cover by zone and year.
- Table 5. Significant differences in species diversity between pre- and post-dredging based on a *t*-test of the Shannon-Weiner diversity indices (H') for each year (Zar, 1984).

FIGURES

- Fig. 1. Great Marsh study site.
- Fig. 2. Mean canopy cover of bulrush (*Scirpus fluviatilis*) in 1995 and 1997 along a marsh-to-forest gradient.
- Fig. 3. Mean canopy cover of reed canary grass (*Phalaris arundinacea*) in 1995 and 1997 along a marsh-to-forest gradient.

APPENDIX TABLES

- Appendix Table 1. List of species identified in study quadrats at the Great Marsh.
- Appendix Table 2. Zone widths (m) for 1995 and 1997.
- Appendix Table 3. Mean canopy cover and standard deviation (SD) for the Open-water Zone (Zone 1) for 1995 and 1997.
- Appendix Table 4. Mean canopy cover and standard deviation (SD) for the Bulrush Zone (Zone 2) for 1995 and 1997.

Appendix Table 5. Mean canopy cover and standard deviation (SD) for the Grass Zone (Zone 3) for 1995 and 1997.

Appendix Table 6. Mean canopy cover and standard deviation (SD) for the Forest Zone (Zone 4) for 1995 and 1997.

Appendix Table 7. Latitude and Longitude recordings for the four corners of each study station at the Great Marsh.

APPENDIX FIGURES

Appendix Fig. 1. A: Predredge photo (1990) with study stations. B: Postdredge photo (1997) with study stations.

INTRODUCTION

The Missouri River extends 3,767 km from Montana to its mouth north of St. Louis, Missouri and drains one-sixth of the contiguous United States. Historically, the river's floodplain contained multiple braided main channels with numerous islands and sandbars and smaller side channels that provided a wide variety of water depths and flow rates (Faber and Smith, 1999). This complex river system was in a constant state of change, affected by the interaction of floods, bank and bed erosion, and subsequent deposition. Flooding was particularly important to the biological components of the ecosystem, for example, maintaining backwater areas where conditions were particularly suitable for spawning fish, insects, and waterfowl (Hesse, 1996).

The plant communities found along the Missouri River varied considerably, depending on the stage of ecological succession. Among the communities closest to the river were lakes, ponds and marshes (Weaver, 1960). Marshes were often surrounded by extensive stands of prairie cordgrass (*Spartina pectinata*) and reed canary grass (*Phalaris arundinacea*) while swamps were dominated by such species as river bulrush (*Scirpus fluviatilis*), common reed (*Phragmites australis*) and arrowhead (*Sagittaria* spp). The adjacent floodplain forest was dominated by willows (*Salix* spp) and cottonwoods (*Populus* spp).

Today, the Missouri River is manipulated through the "Mainstem Reservoir System," a series of six dams and reservoirs constructed by the U.S. Army Corps of Engineers beginning in the 1930's. Channelization along the "Lower River," from below Sioux City, Iowa to St. Louis, Missouri, was intended to reduce flooding, support barge

traffic, and generate power (Faber and Smith, 1999). In the process of channelization, however, most of the sand bars, islands and side channels were eliminated. The result of this manipulation in the Lower River has been both a marked change in the river dynamics and a loss of wetland and riparian habitat including oxbows and other backwater areas where floodplain wetlands are often concentrated (U.S. Army Corps of Engineers, 1998).

In the absence of historic river dynamics, disconnected oxbows are affected by sedimentation so that only active efforts at wetland restoration, such as through dredging, can maintain their physical character and the plant and animal communities that they support. One such restoration was initiated along the Missouri River in east-central Nebraska in 1995. This study was initiated in order to take advantage of the opportunity to conduct a pre- and post-dredging evaluation of marsh vegetation (1) to assess the response of plant communities to dredging and (2) to provide baseline data for longer-term study on recovery of the marsh from dredging.

MATERIALS AND METHODS

Study Site

The study was conducted at a detached oxbow lake situated within the 526 ha Fontenelle Forest Preserve along the eastern boundary of Nebraska (North 41°10'22" West 95°53'03") (Fig. 1). The study site, referred to as the Great Marsh, but previously known as Heron Lake, Horseshoe Lake or “the marsh,” covers approximately 14 ha of




Fig 1. Great Marsh study site

wetland and an adjoining 6 ha of seasonably wet lowland forest (U.S. Army Corps of Engineers, 1995).

Background

Between 1850 and 1900, the Missouri River shifted its channel, leaving behind the oxbow that forms the present Great Marsh. By 1995, siltation had decreased the depth of the marsh to 0.3 - 0.9 meters (Eco-Centrics, 1997).

In 1995, the U.S. Army Corps of Engineers, in consultation with the Fontenelle Forest Nature Center and the Papio-Missouri River Natural Resources District, initiated an effort to deepen the Great Marsh in order to extend its function as a backwater habitat for native biota. Dredging specifically focused on (1) removal of sediment from three arms of the marsh, (2) deepening the marsh to a depth of 0.9 –1.2 meters and (3) grading the marsh's littoral zone to avoid sharp changes in depth along its margin (Eco-Centrics, 1997). Ultimately, 85,680 cubic meters of sediment were removed from Great Marsh, most of which had accumulated as a result of erosion from residential development near Fontenelle Forest over the last 15 years.

Descriptions of the plant communities of the Great Marsh are not known prior to 1900. The earliest plant survey of Fontenelle Forest was conducted in 1959 by the Omaha Botany Club (1959) with a more recent survey conducted by Garabrandt (1988). Neither of these surveys described the Great Marsh specifically, however, based on their plant species lists, it is likely that the open water was dominated by lotus (*Nelumbo lutea*) with the surrounding plant communities dominated by bulrush (*Scirpus* spp) and reed

canary grass (*Phalaris arundinacea*). The adjacent forest would have been dominated by cottonwood (*Populus deltoides*) with a dogwood (*Cornus spp.*) and poison ivy (*Toxicodendron spp.*) understory.

Field Methods

Plant community sampling was conducted during the fall of 1995 and again in the fall of 1997. The 1995 evaluation was conducted at twelve study stations systematically located around the Great Marsh (Appendix Fig. 1.). Each study station was stratified into four distinct vegetation zones that together encompassed the principal plant community types that surrounded the Great Marsh (Table 1). Each station was geo-referenced using a Global Positioning System (GPS) unit (Appendix Table 7). Data for 1997 were collected from the same study stations used in 1995, although sampling was adjusted so as to be centered in vegetation zones that became established after dredging. Hydraulic dredging continued from May through September of 1996.

Within each vegetation zone, five, 1 x 1-meter quadrats were systematically located at 5-meter intervals along a 25-meter transect aligned parallel to the marsh edge. When zone edges were not clearly defined, the boundary was determined to be the point at which the predominant species from each abutting zone comprised 50% of the plant composition. Within each 1 x 1-meter quadrat, the canopy cover of five plant groupings (Table 2), as well as that of each species, was estimated following procedures modified from Daubenmire (1959). Cover categories used were: 0 = 0%, 1 = <1%, 2 = 1-5%, 3 = 5-25%, 4 = 25-50%, 5 = 50-75%, 6 = 75-95%, 7 = 95-99%, 8 = >99%. Botanical

Table 1. Zone descriptions and characteristics.

Zone	Descriptive Term	Characteristics
1	Open-Water Zone	Open water area and adjacent mudflat fringe dominated by floating or submerged vegetation, but including some emergent species.
2	Bulrush Zone	Emergent species. Primarily bulrush (<i>Scirpus</i> spp).
3	Grass Zone	Primarily dominated by grasses, such as reed canary grass (<i>Phalaris arundinacea</i>), and graminoid species, such as common reed (<i>Phragmites australis</i>).
4	Forest Zone	Overstory canopy greater than 5 m.

Table 2. Description of plant categories.

Category	Description
TOTCOV	Total plant cover including overstory trees and submerged aquatic plants.
SUBMUD	Floating unattached plants that move about with winds and currents. Floating attached plants with floating leaves but with roots anchored in the substrate. Submerged plants whose life cycle, with the exception of flowering, is completed beneath the surface of the water. Most are anchored to the substrate, but the vegetative portion either does not reach the surface or the terminal end lies in a horizontal position just beneath the surface.
BULRUSH	Comprised primarily of bulrush (<i>Scirpus spp.</i>) and other emergent plants whose roots and basal portion grow beneath the surface of shallow water, but whose leaves and stem are borne primarily above the surface.
GRASS	Comprised principally of graminoid species.
FORBS	Forbs, including non-woody vines.
WOODY	Non-herbaceous plants including canopy trees and shrubs.

nomenclature is based on the Great Plains Flora Association (1986). Species that could not be identified in the field were collected and identified at the University of Nebraska at Omaha Herbarium (OMA).

In addition to plant species composition, the width of each zone was measured in 1995 and estimated in 1997, the latter procedure necessitated by the substantial changes in the marsh edge resulting from dredging. For example, hip waders were required to measure plant canopy cover in the Open-water Zone in 1995, but a canoe was required in 1997.

Analysis

A *t*-test (SAS Institute, 1985) was used to test for significant differences in canopy cover between 1995 and 1997 for those species with a cover value greater than 3% in at least one zone. Between-year differences in species diversity were tested by zone using procedures based on the Shannon Weiner Index (H') as described in Zar (1984). Species Richness (S) of each zone is the total number of species recorded.

RESULTS AND DISCUSSION

Eighty-four plant species were identified in this study (Appendix Table 1). Thirty-eight were found both in 1995 and 1997. Of the remaining forty-six species, thirty, such as swamp milkweed (*Asclepias incarnata*) and path rush (*Juncus tenuis*), were found only in 1995 and sixteen were found only in 1997, including bladderwort

(*Utricularia vulgaris*) and nettle-leaved vervain (*Verbena urticifolia*) (Appendix Tables 3-6). In general, dredging reduced the canopy cover of all plant categories significantly in at least one or more zones with the exception of the submergent category, which increased in the Open-water, Bulrush and Grass Zones (Table 3, Appendix Tables 3-6).

Zone width increased only for the Open-water Zone, the intended consequence of dredging (Table 4). Both the year before and the year after dredging, the dominant herbaceous plants of this zone were duckweed (*Lemna minor*) (35% and 45% mean cover respectively) and watermeal (*Wolffia columbiana*) (53% and 61%). Coontail (*Ceratophyllum demersum*), however, increased significantly from 7% in 1995 to 25% in 1997 (Table 3, Appendix Table 3). The increase in coontail is probably a consequence of the increased water depth of the Open-water Zone. Overall, there were no significant between-year differences in species diversity for this zone (Table 5).

In the Bulrush Zone, river bulrush (*Scirpus fluviatilis*) cover decreased significantly from 47% in 1995 to 27% in 1997 (Fig. 2, Table 3, Appendix Table 4). This decrease in cover within the zone, combined with a substantial decrease in the total amount of the Bulrush Zone (Table 4), resulted in a substantial short-term decline in this species as a consequence of dredging. Reed canary grass was the only other species to occur in relatively high cover in this zone in 1995 (12% cover), although it was eliminated as a consequence of dredging (0% in 1997) (Fig. 3). High cover values for duckweed (8%) and watermeal (6%) in this terrestrial zone before dredging were largely a consequence of these floating plants being deposited on the surface during high water

Table 3. Mean canopy cover by zone and year. Only species averaging > 3% cover in either year are included. S = significant difference between years, NS = no significant difference. tr = < 0.1% cover.

Plant Category and Species by Zone	Year		Statistically Significant Difference (P>0.05)
	1995	1997	
OPEN-WATER ZONE			
TOTCOV	68.8	96.9	S
SUBMUD	66.4	96.4	S
BULRUSH	0.6	3.0	S
FORBS	1.8	1.4	NS
GRASS	7.0	0	S
WOODY	0	2.6	NS
<i>Wolffia columbiana</i> Karst. (watermeal)	52.6	61.4	NS
<i>Lemna minor</i> L. (duckweed)	35.1	45.5	NS
<i>Ceratophyllum demersum</i> L. (coontail)	7.3	25.4	S
<i>Nelumbo lutea</i> (Willd.) Pers.	6.0	8.3	NS
<i>Echinochloa crusgalli</i> (L.) Beauv.	6.1	0	S
<i>Morus alba</i> L.	0	3.1	NS
BULRUSH ZONE			
TOTCOV	82.9	38.3	S
SUBMUD	12.0	28.3	S
BULRUSH	55.3	28.5	S
FORBS	4.6	0.3	S
GRASS	29.9	2.5	S
WOODY	2.0	0.1	NS
<i>Scirpus fluviatilis</i> (Torr.) Gray (river bulrush)	50.0	27.9	S
<i>Phalaris arundinacea</i> L. (reed canary grass)	14.0	0.1	S
<i>Scirpus validus</i> Vahl	9.8	0	S
<i>Lemna minor</i> L. (duckweed)	7.9	22.6	S
<i>Echinochloa crusgalli</i> (L.) Beauv.	6	0	S
<i>Wolffia columbiana</i> Karst. (watermeal)	5.9	13.7	S
<i>Phragmites australis</i> (Cav.) Trin. ex Steud	4.3	0.3	NS
<i>Typha</i> spp	tr	4.3	NS

Table 3. Mean canopy cover by zone and year. Only species averaging > 3% cover in either year are included. S = significant difference between years, NS = no significant difference. tr = < 0.1% cover.

GRASS ZONE			
TOTCOV	86.8	32.3	S
SUBMUD	0.2	12.9	S
BULRUSH	13.6	4.2	S
FORBS	12.0	0.1	S
GRASS	68.3	26.9	S
WOODY	16.9	tr	S
<i>Phalaris arundinacea</i> L. (reed canary grass)	54.0	8.0	S
<i>Scirpus fluviatilis</i> (Torr.) Gray (river bulrush)	12.9	3.2	S
<i>Phragmites australis</i> (Cav.) Trin. Ex Steud (common reed)	11.4	8.8	NS
<i>Morus alba</i> L	8.3	0	S
<i>Salix exigua</i> Nutt.	3.5	0	NS
<i>Lemna minor</i> L.	0.2	8.0	S
<i>Wolffia columbiana</i> Karst.	tr	8.9	S
FOREST ZONE			
TOTCOV	84.0	54.8	S
SUBMUD	0	8.7	S
BULRUSH	8.0	5.8	NS
FORBS	12.2	4.3	S
GRASS	65.8	32.3	S
WOODY	59.1	28.1	S
<i>Phalaris arundinacea</i> L. (reed canary grass)	38.6	4.4	S
<i>Cornus drummondii</i> C.A. Meyer (rough-leaved dogwood)	27.2	7.8	S
<i>Elymus virginicus</i> L. (Virginia wild rye)	12.9	10.9	NS
<i>Morus alba</i> L	7.3	4.4	NS
<i>Fraxinus pennsylvanica</i> Marshall	5.3	1.2	NS
<i>Ulmus americana</i> L	1.1	6.9	NS
<i>Wolffia columbiana</i> Karst.	1.1	5.2	NS
<i>Lemna minor</i> L.	0	5.5	S
<i>Scirpus fluviatilis</i> (Torr.) Gray	7.3	5.5	NS
<i>Spartina pectinata</i> Link	2.9	4.8	NS
<i>Stachys palustris</i> L.	3.4	0	NS

Table 4. Average zone width and changes between 1995 and 1997. See Table 1 for zone descriptions and Appendix Table 2 for individual zone-width measurements.

Zone	Year	Average Zone Width (meters)	Average Change in Zone Width from 1995 to 1997
OPEN-WATER	1995	3	40-m increase
	1997	43	
BULRUSH	1995	20	14-m decrease
	1997	6	
GRASS	1995	19	17-m decrease
	1997	2	
FOREST	1995	13	9-m decrease
	1997	4	

Table 5. Significant differences in species diversity between pre- and post-dredging based on a *t*-test of the Shannon-Weiner diversity indices (H') for each year (Zar 1984). Alpha = 0.05; n = the sum of species occurrences; S = species richness.

Year	Parameter	Zone			
		Open water	Bulrush	Grass	Forest
1995	H_1'	0.8282	1.296	1.338	1.524
	n	165	332	280	320
	S	12	38	44	55
1997	H_2'	0.9403	0.9315	0.995	1.544
	n	324	122	102	224
	S	16	14	14	46
	Differ Significantly	No	Yes	Yes	No




Fig. 2. Mean canopy cover of Bulrush (*Scirpus fluviatilis*) in 1995 and 1997 along an open water-to-forest gradient. * = Significant difference between years ($P \geq 0.05$).




Fig. 3. Mean canopy cover of reed canary grass (*Phalaris arundinacea*) in 1995 and 1997 along an open water-to-forest gradient. * = Significant difference between years ($P \geq 0.05$).

periods that occurred at the marsh shortly before sampling. Overall, there was a significant decline in species diversity for the Bulrush zone from 38 species to 14 species (Table 5). Species lost included bulrush (*Scirpus validus*) and five smartweed species (*Polygonum* spp.).

As with the Bulrush Zone, the decrease in width of the Grass Zone was substantial, and changes in dominant species were significant. For example, reed canary grass dominated before dredging (53% cover), but decreased significantly to 7% cover as a consequence of dredging. Both river bulrush (12% to 3%) and common reed (11% to 8%) also decreased, although only river bulrush declined significantly (Fig. 2, Table 3, Appendix Table 5). These decreases in cover are most likely due to the mechanical removal of portions of the zone (personal observation).

The Forest Zone also declined in extent following dredging, with significant declines in understory grasses and forbs. Reed canary grass, for example, the dominant understory species, declined from 38% to 4%, a response also noted in the Grass Zone and, to a lesser extent, in the Bulrush Zone as well (Fig.3, Table 3, Appendix Table 6). Other species, such as Virginia wild rye (*Elymus virginica*), a native woodland species, persisted irrespective of treatment, although at low levels (12% and 10% cover in 1995 and 1997, respectively). Declines were also noted for woody species cover (59% to 28%) with a significant decline specifically in dogwood (*Cornus drumondii*) (28% to 8%). Species diversity, however, did not decline significantly following dredging (Table 5).

CONCLUSIONS

The Bulrush and Grass Zones were the zones most affected by dredging, each with a significant decline in species richness and a substantial reduction in zone widths (Table 5). Although much of the Bulrush and Grass Zones was replaced by open water, none of the species in those two zones completely disappeared from the marsh. In contrast, the plant diversity of the Open-water Zone was not affected by dredging, although there was a substantial increase in areal extent and in water-depth. The Forest Zone, as with the Open-water zone, was not affected substantially by dredging.

In combination, the results of this study suggest that dredging to restore wetland habitat can successfully deepen and widen the standing-water portion of marsh habitat without eliminating the principal plant communities that immediately surround it, at least not the year immediately following dredging. The significant decline in diversity that accompanied dredging, however, suggests effects that may have a longer-term impact on marsh habitat, although only future study can assess this effect (Table 5). This study provides the base for such long-term assessments. The results of this study should be regarded in the context of a single effort being performed in site-specific conditions. Data from other restoration-dredging efforts would need to be compared with the data from this study in order to suggest generalities about this process as a means of extending the life of marshes where the physical conditions that established them no longer occur.

LITERATURE CITED

- Daubenmire, R. F. 1959. A Canopy-coverage method of Vegetational Analysis. *Northwest Science*. 33(1): 43-64.
- Eco-Centrics. 1997. Post-construction fish monitoring of Great Marsh in Fontenelle Forest, Sarpy County, Nebraska. *A report to the U.S. Army Corps of Engineers Omaha District Planning Office, Omaha, Nebraska*. 13 pp.
- Faber, S., and Smith, C. 1999. *Voyage of Recovery. Restoring the Rivers of Lewis and Clark*. American Rivers, Inc. 31 pp.
- Garabrant, M. M. 1988. *A Floristic Survey of Fontenelle Forest and Neale Woods Nature Centers*. M. A. Thesis, Univ. of Nebraska at Omaha. 118 pp.
- Great Plains Flora Association. 1986. *Flora of the Great Plains*. University Press of Kansas, Lawrence. 1392 pp.
- Hesse, L. W. 1996. Floral and faunal trends in the middle Missouri River. In D.L. Galat and A. G. Frazier (eds.) *Overview of river-floodplain ecology in the upper Mississippi River basin*, Vol. 3 of J. A. Kelmelis (ed) *Science for Floodplain Management into the 21st Century*. U. S. Government Printing Office. Washington, CD, USA. p. 73-90.
- Omaha Botany Club. 1959. *Plants of Fontenelle Forest*. Fontenelle Forest Association. 136 pp.
- SAS Institute Inc. 1985. *SAS user's guide: Statistics, Version 5*. Cary, North Carolina: SAS Institute Inc. 956 pp.
- U. S. Army Corps of Engineers. 1995. *Project Modification Report with Environmental Assessment: Hidden Lake/Great Marsh Restoration Project (Section 1135)*. 61 pp.
- U. S. Army Corps of Engineers. 1998. *Summary of the Preliminary RDEIS. Master Water Control Manual*. Missouri River Review and Update Study. 20 pp.
- Weaver, J. E. 1960. Flood Plain Vegetation of the Central Missouri Valley and Contacts of Woodland with Prairie. *Ecological Monographs* Vol. 30:1. 64 pp.
- Zar, J. H. 1984. *Biostatistical Analysis*, 2nd Ed. Prentice-Hall, Englewood Cliffs, New Jersey. 718 pp.

APPENDIX

Appendix Table 1. List of species identified in study quadrats at the Great Marsh. Nomenclature follows Great Plains Flora Association. (1986)

Scientific Name	Common Name
<i>Acer</i> seedlings	Maple seedlings
<i>Acer saccharinum</i> L.	Silver maple
<i>Alisma subcordata</i> Raf.	Water plantain
<i>Alliaria petiolata</i> (Bieb.) Cavara & Grande	Garlic mustard
<i>Asclepias incarnata</i> L.	Swamp milkweed
<i>Aster ontarionis</i> Wieg.	Missouri aster
<i>Bidens cernua</i> L.	Nodding beggar-ticks
<i>Bidens frondosa</i> L.	Beggar-ticks
<i>Bidens vulgata</i> Greene	Beggar-ticks
<i>Boehmeria cylindrica</i> (L.) Swartz	False-nettle
<i>Bromus latiglumis</i> (Scribn. ex Shear) Hitchc.	Brome
<i>Bromus pubescens</i> Muhl. ex Willd.	Brome
Bryophyte - Liverwort	
Bryophyte - Moss	
<i>Campanula americana</i> L.	American or tall bellflower
<i>Carex</i> spp.	Sedge
<i>Carex pensylvanica</i> Lam.	GROUP VIII
<i>Celtis occidentalis</i> L.	Hackberry
<i>Ceratophyllum demersum</i> L.	Coontail
<i>Cinna arundinacea</i> L.	Woodreed
<i>Cornus drummondii</i> C.A. Meyer	Rough-leaved dogwood
<i>Cyperus</i> spp.	Umbrella sedge
<i>Cyperus esculentus</i> L.	Umbrella sedge
<i>Echinochloa</i> spp.	Barnyard grass
<i>Echinochloa crusgalli</i> (L.) Beauv.	Barnyard grass
<i>Echinochloa muricata</i> (Beauv.) Fernald	Barnyard grass
<i>Eclipta prostrata</i> (L.)L.	Yerba de Tajo
<i>Elymus canadensis</i> L.	Canada wild rye
<i>Elymus villosus</i> Muhl. ex Willd.	Hairy wild rye
<i>Elymus virginicus</i> L.	Virginia wild rye
<i>Equisetum</i> spp.	Horsetail
<i>Equisetum arvense</i> L.	Field horsetail
<i>Eupatorium serotinum</i> Michx.	Boneset
<i>Fraxinus</i> seedlings	Green Ash seedlings
<i>Fraxinus pennsylvanica</i> Marshall	Green Ash
<i>Galium aparine</i> L.	Catchweed bedstraw
<i>Galium trifidum</i> L.	Small bedstraw
<i>Hemerocallis fulva</i> L.	Daylily
<i>Hystrix patula</i> Moench.	Bottlebrush grass
<i>Juncus tenuis</i> Willd.	Path rush
<i>Lamiaceae</i> spp.	Mint

Appendix Table 1. List of species identified in study quadrats at the Great Marsh. Nomenclature follows Great Plains Flora Association. (1986)

Scientific Name	Common Name
<i>Leersia oryzoides</i> (L.) Swartz	Rice cutgrass
<i>Leersia virginica</i> Willd.	Whitegrass
<i>Lemna minor</i> L.	Duckweed
<i>Leonurus cardiaca</i> L.	Motherwort
<i>Morus alba</i> L.	White mulberry
<i>Muhlenbergia mexicana</i> (L.) Trin.	Wirestem muhly
<i>Muhlenbergia</i> spp.	Muhly
<i>Nelumbo lutea</i> (Willd.) Pers.	Lotus, Water chinkapin
<i>Oxalis dilleni</i> Jacq.	Gray-green wood sorrel
<i>Panicum dichotomiflorum</i> Michx.	Fall panicum
<i>Phalaris arundinacea</i> L.	Reed canary grass
<i>Phalaris arundinacea</i> L. seedlings	Reed canary grass seedlings
<i>Phragmites australis</i> (Cav.) Trin. ex Steud	Common reed
<i>Pilea pumila</i> (L.) Gray	Clearweed
<i>Platanus occidentalis</i> L.	Sycamore or Plane-tree
<i>Polygonum</i> spp.	Smartweed
<i>Polygonum amphibium</i> L.	Water smartweed
<i>Polygonum hydropiper</i> L.	Waterpepper
<i>Polygonum lapathifolium</i> L.	Pale smartweed
<i>Polygonum pennsylvanicum</i> L.	Pennsylvania smartweed
<i>Polygonum persicaria</i> L.	Lady's thumb
<i>Polygonum punctatum</i> Elliott	Water smartweed
<i>Polygonum virginianum</i> L.	Virginia knotweed
<i>Populus deltoides</i> W. Bartram ex Marshall	Cottonwood
<i>Potamogeton nodosus</i> Poir.	Longleaf pondweed
<i>Potamogeton pectinatus</i> L.	Sago pondweed
<i>Ranunculus sceleratus</i> L.	Cursed crowfoot
<i>Rumex crispus</i> L.	Curly dock
<i>Rumex</i> spp.	Dock
<i>Sagittaria latifolia</i>	Arrowhead
<i>Salix</i> spp.	Willow
<i>Salix amygdaloides</i> Anderss.	Peach-leaf willow
<i>Salix eriocephala</i> Michx.	Diamond willow
<i>Salix exigua</i> Nutt.	Sandbar willow
<i>Salix nigra</i> Marshall	Black willow
<i>Sanicula gregaria</i> Bickn.	Black snakeroot
<i>Scirpus</i> spp.	Bulrush
<i>Scirpus acutus</i> Muhl. ex Bigel.	Bulrush
<i>Scirpus fluviatilis</i> (Torr.) Gray	Bulrush
<i>Scirpus validus</i> Vahl	Bulrush
<i>Scutellaria lateriflora</i> L.	Blue skullcap or mad-dog
<i>Smilax hispida</i> Muhl. ex Torr.	Bristly greenbrier

Appendix Table 1. List of species identified in study quadrats at the Great Marsh. Nomenclature follows Great Plains Flora Association. (1986)

Scientific Name	Common Name
<i>Spartina pectinata</i> Link	Prairie cordgrass
<i>Stachys palustris</i>	Hedge-nettle
<i>Toxicodendron radicans</i> (L.) Kuntze	Poison ivy
<i>Tridens flavus</i> (L.) A. Hitchc.	Redtop
<i>Typha</i> spp.	Cattail
<i>Ulmus</i> seedlings	Elm seedlings
<i>Ulmus americana</i> L.	American elm
<i>Ulmus rubra</i> Muhl.	Slippery elm
Unidentified forb spp.	
Unidentified grass spp.	
Unidentified woody spp.	
<i>Urtica dioica</i> L.	Stinging nettle
<i>Utricularia vulgaris</i> L.	Common bladderwort
<i>Verbena urticifolia</i> L.	Nettle-leaved vervain
<i>Viola</i> spp.	Violet
<i>Viola missouriensis</i> Greene	Violet
<i>Vitis riparia</i> Michx.	River-bank grape
<i>Vitis riparia</i> seedlings	River-bank grape seedlings
<i>Vitis</i> spp.	Grape seedlings
<i>Wolffia columbiana</i> Karst.	Watermeal

Appendix Table 2. Zone widths (m) for 1995 and 1997. Direct measurements were made in 1995, but estimated in 1997. See Table 1 for zone descriptions. Study stations numbering is not sequential as numbers 9, 12, and 14 were eliminated during site selection.

Study Station	Year and Zones							
	1995				1997			
	1	2	3	4	1	2	3	4
1	3	18	6	6	27	0	0	6
2	3	8	4	6	16	0	0	6
3	3	4	3	6	11	0	0	6
4	3	9	3	5	14	0	0	6
5	3	21	15	15	46	2	0	6
6	3	20	20	17	52	1	0	6
7	3	57	16	17	85	3	6	0
8	6	53	17	30	102	0	5	0
10	0	0	85	14	50	30	6	12
11	0	18	28	13	59	0	0	0
13	3	21	9	0	34	0	0	0
15	3	11	23	25	15	38	9	0
Sum	33	240	229	154	511	74	26	48
Mean	3	20	19	13	43	6	2	4
Width Change					+40	-14	-17	-9

Appendix Table 3. Mean canopy cover and standard deviation (SD) for the Open-water Zone (Zone 1) for 1995 and 1997. See Table 2 for category descriptions. (*) indicates a significant difference in the mean of that vegetative category or species between 1995 and 1997. tr = < 0.1% cover.

Species	Zone 1			
	1995		1997	
	Mean	SD	Mean	SD
TOTCOV (*)	68.8	39.0	96.9	8.2
SUBMUD (*)	66.4	40.5	96.4	8.3
BULRUSH (*)	0.6	2.7	3.0	7.7
FORBS	1.8	7.2	1.4	5.5
GRASS (*)	7.0	15.5	0.0	0.0
WOODY	0.0	0.0	2.6	13.5
<i>Alisma subcordata</i> Raf.	0.0	0.0	0.1	0.5
<i>Ceratophyllum demersum</i> L. (*)	7.3	21.7	25.4	25.5
<i>Echinochloa crusgalli</i> (L.) Beauv. (*)	6.1	15.0	0.0	0.0
<i>Fraxinus pennsylvanica</i> Marshall	0.0	0.0	0.1	0.5
<i>Fraxinus</i> seedlings	0.0	0.0	tr	0.1
<i>Lamiaceae</i> spp.	tr	0.3	0.0	0.0
<i>Lemna minor</i> L.	35.1	38.5	45.5	32.9
<i>Morus alba</i> L.	0.0	0.0	3.1	16.6
<i>Nelumbo lutea</i> (Willd.) Pers.	6.0	13.1	8.3	13.6
<i>Phalaris arundinacea</i> L.	1.6	7.0	0.0	0.0
<i>Platanus occidentalis</i> L.	0.0	0.0	0.1	0.5
<i>Polygonum amphibium</i> L.	0.0	0.0	tr	0.1
<i>Polygonum lapathifolium</i> L.	0.1	0.3	0.0	0.0
<i>Polygonum persicaria</i> L.	tr	0.3	0.0	0.0
<i>Potamogeton nodosus</i> Poir.	0.0	0.0	0.4	2.0
<i>Potamogeton pectinatus</i> L. (*)	0.0	0.0	2.0	5.7
<i>Ranunculus sceleratus</i> L.	0.1	0.3	0.0	0.0
<i>Rumex</i> spp.	0.3	1.9	0.0	0.0
<i>Sagittaria latifolia</i> Willd.	0.0	0.0	0.3	1.9
<i>Scirpus acutus</i> Muhl. ex Bigel.	0.0	0.0	tr	0.1
<i>Scirpus fluviatilis</i> (Torr.) Gray	1.7	11.1	1.4	3.3
<i>Utricularia vulgaris</i> L.	0.0	0.0	0.5	2.0
<i>Wolffia columbiana</i> Karst.	52.6	40.8	61.4	29.8

Appendix Table 4. Mean canopy cover and standard deviation (SD) for the Bulrush Zone (Zone 2) for 1995 and 1997. See Table 2 for category descriptions. (*) indicates a significant difference in the mean of that vegetative category or species between 1995 and 1997. tr = < 0.1% cover.

Species	Zone 2			
	1995		1997	
	Mean	SD	Mean	SD
TOTCOV (*)	82.9	27.2	38.3	46.8
SUBMUD (*)	12.0	22.1	28.3	39.0
BULRUSH (*)	55.3	33.6	28.5	38.4
FORBS (*)	4.6	9.1	0.3	1.9
GRASS (*)	29.9	31.3	2.5	11.4
WOODY	2.0	12.7	0.1	0.3
<i>Alisma subcordata</i> Raf.	0.0	0.0	tr	0.3
<i>Bidens cernua</i> L.	0.3	1.9	0.0	0.0
<i>Bidens vulgata</i> Greene	tr	0.1	0.0	0.0
<i>Boehmeria cylindrica</i> (L.) Swartz	0.1	0.3	0.0	0.0
Bryophyte – Liverwort (*)	0.8	2.7	0.0	0.0
Bryophyte – Moss (*)	2.4	7.3	0.0	0.0
<i>Ceratophyllum demersum</i> L.	0.0	0.0	1.2	5.5
<i>Cyperus esculentus</i> L. (*)	0.1	0.4	0.0	0.0
<i>Echinochloa crusgalli</i> (L.) Beauv. (*)	6.8	23.0	tr	0.1
<i>Echinochloa muricata</i> (Beauv.) Fernald	0.3	1.9	0.0	0.0
<i>Echinochloa</i> spp. (*)	0.2	0.6	0.0	0.0
<i>Eclipta prostrata</i> (L.)L.	0.4	2.0	0.0	0.0
<i>Fraxinus pennsylvanica</i> Marshall	0.0	0.0	tr	0.1
<i>Leersia oryzoides</i> (L.) Swartz (*)	2.1	7.4	0.0	0.0
<i>Lemna minor</i> L. (*)	7.9	19.8	22.6	32.5
<i>Morus alba</i> L.	1.1	8.1	0.0	0.0
<i>Nelumbo lutea</i> (Willd.) Pers. (*)	tr	0.1	0.3	0.6
<i>Panicum dichotomiflorum</i> Michx.	tr	0.1	0.0	0.0
<i>Phalaris arundinacea</i> L. (*)	14.0	23.2	0.1	0.3
<i>Phalaris arundinacea</i> L. seedlings	0.9	5.2	0.0	0.0
<i>Phragmites australis</i> (Cav.) Trin. ex Steud	4.3	15.5	0.3	2.0
<i>Pilea pumila</i> (L.) Gray	0.1	0.5	0.0	0.0
<i>Platanus occidentalis</i> L.	tr	0.1	0.0	0.0
<i>Polygonum amphibium</i> L.	0.7	4.9	0.0	0.0
<i>Polygonum hydropiper</i> L.	tr	0.1	0.0	0.0
<i>Polygonum lapathifolium</i> L. (*)	0.2	0.6	0.0	0.0
<i>Polygonum pensylvanicum</i> L.	0.1	0.5	0.0	0.0
<i>Polygonum persicaria</i> L. (*)	0.2	0.6	0.0	0.0
<i>Polygonum</i> spp.	0.4	2.0	0.0	0.0

Appendix Table 4. Mean canopy cover and standard deviation (SD) for the Bulrush Zone (Zone 2) for 1995 and 1997. See Table 2 for category descriptions. (*) indicates a significant difference in the mean of that vegetative category or species between 1995 and 1997. tr = < 0.1% cover.

<i>Populus deltoides</i> W. Bartram ex Marshall	0.3	1.9	0.0	0.0
<i>Ranunculus sceleratus</i> L.	tr	0.1	0.0	0.0
<i>Rumex</i> spp.	tr	0.1	0.0	0.0
<i>Salix amygdaloides</i> Anderss.	1.6	12.6	0.0	0.0
<i>Scirpus acutus</i> Muhl. ex Bigel.	1.5	8.5	tr	0.1
<i>Scirpus fluviatilis</i> (Torr.) Gray (*)	50.0	32.7	27.9	37.6
<i>Scirpus validus</i> Vahl (*)	9.8	22.6	0.0	0.0
<i>Scutellaria lateriflora</i> L.	0.8	3.3	0.0	0.0
<i>Spartina pectinata</i> Link (*)	0.0	0.0	0.2	0.6
<i>Typha</i> spp.	tr	0.1	4.3	18.7
Unidentified forbs (*)	0.1	0.5	0.0	0.0
Unidentified grasses	1.4	6.8	0.0	0.0
Unidentified wood spp.	tr	0.1	0.0	0.0
<i>Utricularia vulgaris</i> L.	0.0	0.0	0.1	0.3
<i>Wolffia columbiana</i> Karst. (*)	5.9	14.5	13.7	25.1

Appendix Table 5. Mean canopy cover and standard deviation (SD) for the Grass Zone (Zone 3) for 1995 and 1997. See Table 2 for category descriptions. (*) indicates a significant difference in the mean of that vegetative category or species between 1995 and 1997. tr = < 0.1% cover.

Species	Zone 3			
	1995		1997	
	Mean	SD	Mean	SD
TOTCOV (*)	86.8	27.2	32.3	46.2
SUBMUD (*)	0.2	0.5	12.9	28.6
BULRUSH (*)	13.6	20.0	4.2	13.7
FORBS (*)	12.0	20.7	0.1	0.5
GRASS (*)	68.3	30.4	26.9	39.6
WOODY (*)	16.9	30.7	tr	0.1
<i>Acer saccharinum</i> L.	tr	0.3	0.0	0.0
<i>Acer</i> seedlings	0.3	1.9	0.0	0.0
<i>Alisma subcordata</i> Raf.	0.0	0.0	tr	0.1
<i>Asclepias incarnata</i> L.	0.5	2.7	0.0	0.0
<i>Boehmeria cylindrica</i> (L.) Swartz	0.1	0.5	0.0	0.0
Bryophyte - Liverwort	0.3	1.9	0.0	0.0
Bryophyte - Moss	1.4	6.8	0.0	0.0
<i>Cornus drummondii</i> C.A. Meyer	0.3	2.0	0.0	0.0
<i>Cyperus</i> spp.	0.3	1.9	0.0	0.0
<i>Echinochloa crusgalli</i> (L.) Beauv.	2.3	12.0	0.1	0.3
<i>Eclipta prostrata</i> (L.)L.	0.3	1.9	0.0	0.0
<i>Elymus virginicus</i> L.	0.1	0.6	0.0	0.0
<i>Equisetum arvense</i> L.	tr	0.3	0.0	0.0
<i>Fraxinus pennsylvanica</i> Marshall	3.0	12.2	tr	0.1
<i>Fraxinus</i> seedlings	0.9	5.2	0.0	0.0
<i>Hemerocallis fulva</i> L.	tr	0.3	0.0	0.0
<i>Lemna minor</i> L. (*)	0.2	0.5	8.0	21.7
<i>Morus alba</i> L. (*)	8.3	23.4	0.0	0.0
<i>Muhlenbergia</i> spp.	tr	0.1	0.0	0.0
<i>Nelumbo lutea</i> (Willd.) Pers. (*)	0.0	0.0	0.1	0.2
<i>Phalaris arundinacea</i> L. (*)	54.0	33.6	8.0	19.5
<i>Phalaris arundinacea</i> L. seedlings	tr	0.3	0.0	0.0
<i>Phragmites australis</i> (Cav.) Trin. ex Steud	11.4	22.3	8.8	23.2
<i>Pilea pumila</i> (L.) Gray	0.1	0.5	0.0	0.0
<i>Platanus occidentalis</i> L.	1.1	8.1	0.0	0.0
<i>Polygonum amphibium</i> L. (*)	2.4	7.6	0.0	0.0
<i>Polygonum hydropiper</i> L.	0.9	5.2	0.0	0.0
<i>Polygonum persicaria</i> L.	0.3	1.9	0.0	0.0
<i>Polygonum punctatum</i> Elliott	0.3	1.9	tr	0.1
<i>Polygonum</i> spp. (*)	2.1	7.4	0.0	0.0

Appendix Table 5. Mean canopy cover and standard deviation (SD) for the Grass Zone (Zone 3) for 1995 and 1997. See Table 2 for category descriptions. (*) indicates a significant difference in the mean of that vegetative category or species between 1995 and 1997. tr = < 0.1% cover.

<i>Populus deltoides</i> W. Bartram ex Marshall	1.4	8.3	tr	0.3
<i>Salix amygdaloides</i> Anderss.	0.3	2.0	0.0	0.0
<i>Salix eriocephala</i> Michx.	1.1	8.1	0.0	0.0
<i>Salix exigua</i> Nutt.	3.5	16.0	0.0	0.0
<i>Scirpus acutus</i> Muhl. ex Bigel.	0.0	0.0	tr	0.1
<i>Scirpus fluviatilis</i> (Torr.) Gray (*)	12.9	20.2	3.2	11.3
<i>Scirpus</i> spp.	0.6	4.8	0.0	0.0
<i>Scirpus validus</i> Vahl	tr	0.3	0.0	0.0
<i>Scutellaria lateriflora</i> L.	0.7	4.9	0.0	0.0
<i>Smilax hispida</i> Muhl. ex Torr.	tr	0.1	0.0	0.0
<i>Spartina pectinata</i> Link (*)	1.6	5.8	tr	0.1
<i>Stachys palustris</i> L.	tr	0.3	0.0	0.0
<i>Typha</i> spp.	1.6	6.0	1.4	6.8
<i>Ulmus americana</i> L.	1.7	11.1	0.0	0.0
Unidentified forbs (*)	tr	0.1	0.0	0.0
Unidentified wood spp.	0.1	0.3	0.0	0.0
<i>Viola missouriensis</i> Greene	0.1	0.5	0.0	0.0
<i>Vitis riparia</i> Michx.	0.3	1.9	0.0	0.0
<i>Wolffia columbiana</i> Karst. (*)	tr	0.1	8.9	26.1

Appendix Table 6. Mean canopy cover and standard deviation (SD) for the Forest Zone (Zone 4) for 1995 and 1997. See Table 2 for category descriptions. (*) indicates a significant difference in the mean of that vegetative category or species between 1995 and 1997. tr = < 0.1% cover.


Species	Zone 4			
	1995		1997	
	Mean	SD	Mean	SD
TOTCOV (*)	84.0	27.4	54.8	47.1
SUBMUD (*)	0.0	0.0	8.7	20.7
BULRUSH	8.0	18.3	5.8	14.5
FORBS (*)	12.2	17.4	4.3	9.9
GRASS (*)	65.8	27.8	32.3	36.1
WOODY (*)	59.1	42.0	28.1	37.3
<i>Acer saccharinum</i> L.	tr	0.1	0.1	0.6
<i>Acer</i> seedlings (*)	0.3	0.8	0.0	0.0
<i>Alliaria petiolata</i> (Bieb.) Cavara & Grande	0.0	0.0	0.3	2.0
<i>Asclepias incarnata</i> L.	tr	0.3	0.1	0.3
<i>Aster ontarionis</i> Wieg.	0.0	0.0	tr	0.1
<i>Bidens frondosa</i> L.	0.0	0.0	tr	0.1
<i>Boehmeria cylindrica</i> (L.) Swartz	0.4	2.0	0.0	0.0
<i>Bromus latiglumis</i> (Scribn. ex Shear) Hitchc.	0.0	0.0	0.8	3.3
<i>Bromus pubescens</i> Muhl. ex Willd.	0.3	2.0	0.0	0.0
<i>Campanula americana</i> L.	tr	0.1	0.0	0.0
<i>Carex pensylvanica</i> Lam.	0.8	4.9	0.0	0.0
<i>Carex</i> spp.	0.8	4.9	tr	0.1
<i>Celtis occidentalis</i> L.	1.2	8.1	1.3	8.3
<i>Cinna arundinacea</i> L.	tr	0.1	0.3	2.0
<i>Cornus drummondii</i> C.A. Meyer (*)	27.2	38.6	7.8	21.8
<i>Echinochloa crusgalli</i> (L.) Beauv.	0.0	0.0	0.1	0.3
<i>Elymus canadensis</i> L.	2.3	10.4	0.0	0.0
<i>Elymus villosus</i> Muhl. ex Willd.	1.3	8.3	0.0	0.0
<i>Elymus virginicus</i> L.	12.9	23.5	10.9	23.9
<i>Equisetum arvense</i> L.	0.7	4.8	0.3	1.9
<i>Equisetum</i> spp.	0.1	0.3	0.0	0.0
<i>Eupatorium serotinum</i> Michx.	0.0	0.0	tr	0.1
<i>Fraxinus pennsylvanica</i> Marshall	5.3	15.1	1.2	5.5
<i>Galium aparine</i> L.	0.0	0.0	0.1	0.3
<i>Galium trifidum</i> L.	0.0	0.0	0.1	0.5
<i>Hystrix patula</i> Moench.	1.7	9.3	2.0	7.4
<i>Juncus tenuis</i> Willd.	0.3	1.9	0.0	0.0
<i>Leersia virginica</i> Willd.	0.0	0.0	0.1	0.3
<i>Lemna minor</i> L. (*)	0.0	0.0	5.5	13.8

Appendix Table 6. Mean canopy cover and standard deviation (SD) for the Forest Zone (Zone 4) for 1995 and 1997. See Table 2 for category descriptions. (*) indicates a significant difference in the mean of that vegetative category or species between 1995 and 1997. tr = < 0.1% cover.

<i>Leonurus cardiaca</i> L.	0.3	2.0	0.1	0.3
<i>Morus alba</i> L.	7.3	23.7	4.4	17.3
<i>Muhlenbergia mexicana</i> (L.) Trin.	0.0	0.0	0.5	2.7
<i>Muhlenbergia</i> spp.	0.1	0.3	0.0	0.0
<i>Oxalis dilleni</i> Jacq.	0.0	0.0	0.1	0.3
<i>Phalaris arundinacea</i> L. (*)	38.6	37.0	4.4	9.3
<i>Phragmites australis</i> (Cav.) Trin. ex Steud	tr	0.1	0.1	0.3
<i>Platanus occidentalis</i> L.	0.0	0.0	0.7	4.9
<i>Polygonum hydropiper</i> L.	0.5	2.7	0.8	4.9
<i>Polygonum lapathifolium</i> L.	0.3	1.9	0.0	0.0
<i>Polygonum punctatum</i> Elliott	0.0	0.0	0.1	0.3
<i>Polygonum</i> spp.	1.4	6.8	0.1	0.5
<i>Polygonum virginianum</i> L.	0.3	2.0	tr	0.1
<i>Populus deltoides</i> W. Bartram ex Marshall	3.1	15.1	0.4	2.0
<i>Rumex crispus</i> L.	tr	0.1	0.0	0.0
<i>Salix amygdaloides</i> Anderss.	0.0	0.0	0.1	0.5
<i>Salix eriocephala</i> Michx.	4.0	15.7	0.0	0.0
<i>Salix exigua</i> Nutt.	1.7	12.6	0.3	1.9
<i>Salix nigra</i> Marshall	1.5	11.0	0.3	2.0
<i>Salix</i> spp. (*)	5.2	19.9	0.0	0.0
<i>Sanicula gregaria</i> Bickn. (*)	0.2	0.7	0.0	0.0
<i>Scirpus fluviatilis</i> (Torr.) Gray	7.3	17.3	5.5	14.5
<i>Scutellaria lateriflora</i> L.	tr	0.3	0.0	0.0
<i>Smilax hispida</i> Muhl. ex Torr.	0.4	2.0	tr	0.1
<i>Spartina pectinata</i> Link	2.9	11.5	4.8	16.7
<i>Stachys palustris</i> L.	3.4	15.5	0.0	0.0
<i>Toxicodendron radicans</i> (L.) Kuntze	1.0	5.2	0.1	0.3
<i>Tridens flavus</i> (L.) A. Hitchc.	0.9	5.2	0.0	0.0
<i>Ulmus americana</i> L.	1.1	8.1	6.9	22.7
<i>Ulmus rubra</i> Muhl.	6.4	21.4	2.8	13.6
<i>Ulmus</i> seedlings	tr	0.1	0.0	0.0
Unidentified forbs	0.4	2.0	0.0	0.0
Unidentified grasses	0.6	4.8	0.6	2.7
Unidentified wood spp.	4.1	18.2	0.0	0.0
<i>Urtica dioica</i> L.	0.3	2.0	0.1	0.3
<i>Verbena urticifolia</i> L.	0.0	0.0	tr	0.1
<i>Viola missouriensis</i> Greene (*)	3.5	10.8	0.1	0.6
<i>Viola</i> spp.	0.0	0.0	0.8	4.9
<i>Vitis riparia</i> Michx.	3.1	12.3	0.0	0.0
<i>Vitis riparia</i> seedlings	tr	0.3	0.0	0.0
<i>Vitis</i> spp.	tr	0.3	0.3	1.9
<i>Wolffia columbiana</i> Karst.	1.1	5.5	5.2	17.4

Appendix Table 7. Latitude and Longitude recordings for the four corners of each study station at the Great Marsh. Study stations numbered is not sequential as numbers 9, 12, and 14 were eliminated during site selection. All GPS points were obtained through a post-process differential-correction of rover data. For relocation purposes, a real-time kinematics unit would be required. All North latitude designations are 41° and all West longitude designations are 95°.

Study Station	Latitude	Longitude	Latitude	Longitude	Latitude	Longitude	Latitude	Longitude	Latitude	Longitude
1	10°15.5932"	53°18.7705"	10°15.9619"	53°17.8901"	10°16.7373"	53°18.4826"	10°16.1556"	53°19.3695"		
2	10°18.9060"	53°13.3959"	10°19.3688"	53°12.2041"	10°19.9608"	53°12.9063"	10°19.5839"	53°13.9228"		
3	10°22.3182"	53°07.0887"	10°22.4846"	53°06.3345"	10°22.8554"	53°06.6132"	10°22.6566"	53°07.3455"		
4	10°26.8159"	52°53.0336"	10°26.8297"	52°51.9394"	10°27.3751"	52°52.1232"	10°27.3797"	52°53.3088"		
5	10°23.5143"	52°51.2124"	10°24.0894"	52°49.8780"	10°25.3708"	52°50.9845"	10°24.8593"	52°52.2119"		
6	10°18.2614"	53°04.9778"	10°18.6220"	53°03.7183"	10°20.3838"	53°04.5914"	10°20.0038"	53°05.4878"		
7	10°15.3900"	53°11.0946"	10°15.6456"	53°10.1306"	10°18.5634"	53°11.8675"	10°18.6110"	53°13.1324"		
8	10°12.9810"	53°16.3541"	10°13.0847"	53°15.3095"	10°15.7679"	53°17.6585"	10°15.4956"	53°18.7297"		
10	10°04.4923"	53°13.7590"	10°05.1268"	53°13.9772"	10°06.7931"	53°16.0007"	10°06.1721"	53°16.4363"		
11	09°56.0932"	53°10.2512"	09°56.1676"	53°09.7426"	09°58.7082"	53°12.4831"	09°58.5816"	53°13.0794"		
13	09°59.9406"	53°18.0331"	10°00.3930"	53°16.8887"	10°01.1240"	53°17.0121"	10°01.0103"	53°18.4213"		
15	10°06.6900"	53°20.7077"	10°07.3009"	53°17.6500"	10°08.5444"	53°18.8058"	10°07.5769"	53°21.0650"		



Appendix Figure1. A: Predredge photo (1990) with study stations. B: Postredge photo (1997) with study stations.