

Yale University

EliScholar – A Digital Platform for Scholarly Publishing at Yale

Cowles Foundation Discussion Papers

Cowles Foundation

8-1-2003

On the Empirical Content of Quantal Response Equilibrium

Philip A. Haile

Ali Hortaçsu

Grigory Kosenok

Follow this and additional works at: <https://elischolar.library.yale.edu/cowles-discussion-paper-series>



Part of the [Economics Commons](#)

Recommended Citation

Haile, Philip A.; Hortaçsu, Ali; and Kosenok, Grigory, "On the Empirical Content of Quantal Response Equilibrium" (2003). *Cowles Foundation Discussion Papers*. 1706.

<https://elischolar.library.yale.edu/cowles-discussion-paper-series/1706>

This Discussion Paper is brought to you for free and open access by the Cowles Foundation at EliScholar – A Digital Platform for Scholarly Publishing at Yale. It has been accepted for inclusion in Cowles Foundation Discussion Papers by an authorized administrator of EliScholar – A Digital Platform for Scholarly Publishing at Yale. For more information, please contact elischolar@yale.edu.

**ON THE EMPIRICAL CONTENT OF
QUANTAL RESPONSE EQUILIBRIUM**

By

Philip A. Haile, Ali Hortaçsu and Grigory Kosenok

August 2006

COWLES FOUNDATION DISCUSSION PAPER NO. 1432R



**COWLES FOUNDATION FOR RESEARCH IN ECONOMICS
YALE UNIVERSITY
Box 208281
New Haven, Connecticut 06520-8281**

<http://cowles.econ.yale.edu/>

On the Empirical Content of Quantal Response Equilibrium*

Philip A. Haile[†] Ali Hortaçsu[‡] Grigory Kosenok[§]

August 22, 2006

Abstract

The quantal response equilibrium (QRE) notion of McKelvey and Palfrey (1995) has recently attracted considerable attention, due in part to its widely documented ability to rationalize observed behavior in games played by experimental subjects. However, even with strong *a priori* restrictions on unobservables, QRE imposes no falsifiable restrictions: it can rationalize any distribution of behavior in any normal form game. After demonstrating this, we discuss several approaches to testing QRE under additional maintained assumptions.

Keywords: quantal response equilibrium, falsifiability, testable restrictions, regular quantal response equilibrium, rank-cumulative probabilities, Block-Marschak polynomials

*This paper subsumes material in two earlier drafts: one, first circulated widely in August 2003, with the same title, and a second with the title “On the Empirical Content of Quantal Choice Models.” We received helpful comments from a number of colleagues and seminar audiences. We are also grateful to a co-editor and referees for very helpful comments and for encouraging us to develop our ideas for testing more fully. Kun Huang and Dmitry Shapiro provided capable research assistance. Financial support from the National Science Foundation and the Alfred P. Sloan Foundation is gratefully acknowledged.

[†]Department of Economics and Cowles Foundation, Yale University, and NBER, 37 Hillhouse Avenue, Box 208264, New Haven, CT 06520, philip.haile@yale.edu.

[‡]Department of Economics, University of Chicago, and NBER, 1126 E. 59th St., Chicago, IL 60637, hortacsu@uchicago.edu.

[§]New Economic School, Nakhimovsky Prospekt 47, Suite 1721 (11), 117418 Moscow, Russia, gkosenok@nes.ru.

1 Introduction

The quantal response equilibrium (QRE) notion of McKelvey and Palfrey (1995) can be viewed as an extension of standard random utility models of discrete (“quantal”) choice to strategic settings, or as a generalization of Nash equilibrium that allows noisy optimizing behavior while maintaining the internal consistency of rational expectations. Formally, QRE is based on the introduction of random perturbations to the payoffs associated with each action a player can take.¹ The realizations of these perturbations affect which action is the best response to the equilibrium distribution of opponents’ behavior.

Both interpretations of QRE have strong intuitive appeal, and much recent work has shown that QRE can rationalize behavior in a variety of experimental settings where Nash equilibrium fails to do so. In particular, when parameters (of the distributions of payoff perturbations) are chosen so that the predicted distributions of outcomes match the data as well as possible, the fit is often very good. McKelvey and Palfrey’s original paper demonstrated the ability of QRE to explain departures from Nash equilibrium behavior in several games. Since then, the success of QRE in matching observed behavior has been demonstrated in a variety of experimental settings, including all-pay auctions (Anderson, Goeree and Holt (1998)), first-price auctions (Goeree, Holt and Palfrey (2002)), alternating-offer bargaining (Goeree and Holt (2000)), coordination games (Anderson, Goeree and Holt (2001)), and the “traveler’s dilemma” (Capra, Goeree, Gomez and Holt (1999), Goeree and Holt (2001)).² The quotation below, from Camerer, Ho and Chong (2004), suggests the impact this evidence has had:³

Quantal response equilibrium (QRE), a statistical generalization of Nash, almost always explains the direction of deviations from Nash and should replace Nash as the static benchmark to which other models are routinely compared.

Given this recent work and its influence, it is natural to ask how informative the ability of QRE to fit the data really is. Our first result provides a strong negative answer to this question for the type of data often considered in the literature: QRE is not falsifiable in any normal form game, even with significant *a priori* restrictions on payoff perturbations. In particular, any behavior can be rationalized by a QRE, even when each player’s payoff perturbations are restricted to be

¹We give a more complete discussion in the following section. The literature has considered generalizations of the QRE to extensive form games (McKelvey and Palfrey, 1998) and games with continuous strategy spaces (e.g., Anderson, Goeree and Holt (2002)). We restrict attention to normal form games for simplicity.

²Dufwenberg, Gneezy, Goeree and Nagel (2002) suggest that they find an exception proving the rule, noting “Our results are unusual in that we document a feature of the data that is impossible to reconcile with the [QRE].”

³See also, e.g., the provocatively titled paper of Goeree and Holt (1999).

independent across actions or to have identical marginal distributions for all actions. Hence, an evaluation of fit in a single game (no matter the number of replications) is uninformative without strong *a priori* restrictions.

This first result implies no critique of the QRE notion, but merely points to the challenge of developing valid approaches to testing the QRE hypothesis. Testing requires maintained hypotheses beyond those of the QRE notion itself, and the most useful approach may depend on the application. We discuss several promising approaches. Each maintains restrictions on how distributions of payoff perturbations can vary across related normal-form games, leading to falsifiable comparative statics predictions.

In the following section we define notation, review the definition of QRE, and discuss common application of the QRE in the literature. We then present our non-falsifiability result in section 3. Section 4 provides a discussion of the result, leading to our exploration of testing approaches in section 5. We conclude in section 6.

2 Quantal Response Equilibrium

2.1 Model and Definition

Here we review the definition of a QRE, loosely following McKelvey and Palfrey (1995). We refer readers to their paper for additional detail, including discussion of the relation of QRE to other solution concepts. Consider a finite n -person normal form game Γ . The set of pure strategies (actions) available to player i is denoted by $S_i = \{s_{i1}, \dots, s_{iJ_i}\}$, with $S = \times_i S_i$. Let Δ_i denote the set of all probability measures on S_i . Let $\Delta \equiv \times_i \Delta_i$ denote the set of probability measures on S , with elements $p = (p_1, \dots, p_n)$. For simplicity, let p_{ij} represent $p_i(s_{ij})$.

Payoffs of Γ are given by functions $u_i(s_i, s_{-i}) : S_i \times_{j \neq i} S_j \rightarrow \mathbb{R}$. In the usual way, these payoff functions can be extended to the probability domain by letting $u_i(p) = \sum_{s \in S} p(s) u_i(s)$. Hence, e.g., the argument s_{ij} of the payoff function $u_i(s_{ij}, s_{-i})$ is reinterpreted as shorthand for a probability measure in Δ_i placing all mass on strategy s_{ij} . Finally, for every $p_{-i} \in \times_{j \neq i} \Delta_j$ and $p = (p_i, p_{-i})$, define $\bar{u}_{ij}(p) = u_i(s_{ij}, p_{-i})$ and $\bar{u}_i(p) = (\bar{u}_{i1}(p), \dots, \bar{u}_{iJ_i}(p))$.

The QRE notion is based on the introduction of payoff perturbations associated with each action of each player. For player i let

$$\hat{u}_{ij}(p) = \bar{u}_{ij}(p) + \epsilon_{ij}$$

where the vector of perturbations $\epsilon_i \equiv (\epsilon_{i1}, \dots, \epsilon_{iJ_i})$ is drawn from a joint density f_i . For all i and j , ϵ_{ij} is assumed to have the same mean, which may be normalized to zero. Each player i is then

assumed to use action s_{ij} if and only if⁴

$$\hat{u}_{ij}(p) \geq \hat{u}_{ik}(p) \quad \forall k = 1, \dots, J_i. \quad (1)$$

Given a vector $u'_i = (u'_{i1}, \dots, u'_{iJ_i}) \in \mathbb{R}^{J_i}$, let

$$R_{ij}(u'_i) = \{\epsilon_i \in \mathbb{R}^{J_i} : u'_{ij} + \epsilon_{ij} \geq u'_{ik} + \epsilon_{ik} \quad \forall k = 1, \dots, J_i\}. \quad (2)$$

Conditional on the distribution p_{-i} characterizing the behavior of i 's opponents, $R_{ij}(\bar{u}_i(p))$ is the set of realizations of the vector ϵ_i that would lead i to choose action j . Let

$$\sigma_{ij}(u'_i) = \int_{R_{ij}(u'_i)} f_i(\epsilon_i) d\epsilon_i$$

denote the probability of realizing a vector of shocks in $R_{ij}(u'_i)$ and let $\sigma_i = (\sigma_{i1}, \dots, \sigma_{iJ_i})$.

McKelvey and Palfrey (1995) call σ_i player i 's *statistical best response function* or *quantal response function*. Given the baseline payoffs of the game Γ , a distribution of play by i 's opponents, and a joint distribution of i 's payoff perturbations, σ_i describes the probabilities with which each of i 's strategies will be chosen by i . A *quantal response equilibrium* is attained when the distribution of behavior of all players is consistent with their statistical best response functions. More precisely, letting $\sigma = (\sigma_1, \dots, \sigma_n)$ and $\bar{u} = (\bar{u}_1, \dots, \bar{u}_n)$, a QRE is a fixed point of the composite function $\sigma \circ \bar{u} : \Delta \rightarrow \Delta$, which maps joint distributions over all players' pure strategies into statistical best responses for all players.

Definition 1 A quantal response equilibrium (QRE) is any $\pi \in \Delta$ such that for all $i \in 1, \dots, n$ and all $j \in 1, \dots, J_i$, $\pi_{ij} = \sigma_{ij}(\bar{u}_i(\pi))$.

There are several possible interpretations of the QRE notion. One need not take the payoff perturbations literally. The idea that players use strategies that are merely “usually close” to optimal rather than “always fully” optimal has natural appeal, and the QRE offers a coherent formalization of this idea—one that closes the model of error-prone decisions with the assumption of rational expectations about opponents' behavior. One may also view the perturbations as a device for “smoothing out” best response functions in the hope of obtaining more robust and/or plausible

⁴This rule is consistent with rational choice by i given the payoff function \hat{u}_{ij} if the following assumptions are added: (1) ϵ_i and $\epsilon_{i'}$ are independent for $i' \neq i$; (2) the “baseline” payoff functions $u_i(s_i, s_{-i})$ and densities f_i are common knowledge; and (3) for each player i the vector ϵ_i is i 's private information. As McKelvey and Palfrey (1995) show for a particular distribution of perturbations, under these assumptions a QRE is a Bayesian Nash equilibrium of the resulting game of incomplete information. Note that in this case, given the correctly anticipated equilibrium behavior of opponents, each player faces a standard polychotomous choice problem with additive random expected utilities. This observation is useful in estimation, since the distribution of equilibrium play by opponents will typically be directly observable to the researcher. It is also used in the proof of Theorem 1 below.

predictions (cf. Rosenthal, 1989). However, as McKelvey and Palfrey (1995) suggest, the payoff perturbations can have natural economic interpretations as well.⁵ Each ϵ_{ij} could reflect the error made by player i in calculating his expected utility from strategy j , due perhaps to unmodeled costs of information processing. Alternatively, ϵ_{ij} might reflect unmodeled determinants of i 's utility from using strategy j . This interpretation is appealing in many applications since a fully specified theoretical model can, of course, only approximate a real economic environment. Furthermore, any true payoff function $\tilde{u}_i(s_{ij}, p_{-i})$ can be represented as the sum of an arbitrary “baseline” payoff $u_i(s_{ij}, p_{-i})$ and a correction $\epsilon_{ij}(p_{-i}) = \tilde{u}_i(s_{ij}, p_{-i}) - u_i(s_{ij}, p_{-i})$. If the game underlying the baseline payoffs $u_i(s_{ij}, p_{-i})$ provides a good approximation to the truth, representing $\epsilon_{ij}(p_{-i})$ by a random variable that does not depend p_{-i} (as in the QRE) might be useful for predicting behavior or as an empirical model.⁶

2.2 Application and Evaluation

Following McKelvey and Palfrey (1995), application of the QRE to data from experiments has typically proceeded by first specifying the joint densities f_i (up to a finite-dimensional parameter) for all players. In every application we are aware of, it has been assumed for simplicity that ϵ_{ij} is independently and identically distributed (i.i.d.) across all j . In most applications it is assumed that every ϵ_{ij} is an independent draw from an extreme value distribution, yielding the familiar convenient logit choice probabilities

$$p_{ij} = \frac{e^{\lambda \bar{u}_{ij}(p)}}{\sum_{k=1}^{J_i} e^{\lambda \bar{u}_{ik}(p)}}. \quad (3)$$

With p observable, the unknown parameter λ is then easily estimated by maximum-likelihood.⁷

Typically the ability of the QRE to rationalize the data is then assessed based on the match between the observed probabilities on each pure strategy and those predicted by the QRE at the estimated parameter value(s).⁸ Although formal testing is uncommon, visual inspection often suggests a very good fit. Since a QRE would simply be a Nash equilibrium if perturbations were degenerate, the fit *must* improve when one adds the freedom to choose the best fitting member of

⁵See also Chen, Friedman and Thisse (1997). Interpretations mirror those for random utility models in the discrete choice literature.

⁶Examples of empirical applications of QRE include Signorino (1999), Seim (2002), Goeree, Holt and Palfrey (2002), Bajari and Hortaçsu (2003) Sweeting (2004), and Augereau et al. (forthcoming). See also Bajari (1998).

⁷In the applications that have avoided the logit formulation, a power function specification has been used, but the approach is the same. In the logit specification, $1/\lambda$ is proportional to the variance of the payoff perturbations, with equilibrium behavior converging to a Nash equilibrium as $\lambda \rightarrow \infty$.

⁸See, e.g., Baye and Morgan (2004), Cason and Reynolds (2005), Goeree, Holt and Palfrey (2002), Guarnaschelli, McKelvey and Palfrey (2000), McKelvey and Palfrey (1995, 1998), McKelvey, Palfrey and Weber (2000), Fey, McKelvey and Palfrey (1996).

a parametric family. In fact, however, the fit is often greatly improved. The following excerpt from Fey, McKelvey and Palfrey (1996, p. 286–287), which relies on this type of comparison in centipede games, is typical of the conclusions drawn from this fit:

Among the models we evaluate, the Quantal Response Equilibrium model best explains the data. It offers a better fit than the Learning model and, as it is an equilibrium model, is internally consistent. It also accounts for the pattern of increasing take probabilities within a match. These facts lend strong support to the Quantal Response Equilibrium model.

3 How Informative is Fit?

One might expect the QRE notion to impose considerable structure on the types of behavior consistent with equilibrium. As Goeree, Holt and Palfrey (2002) have suggested, the QRE requires a “consistency condition that the probabilities which determine expected utility. . . match the choice probabilities. . . that result from probabilistic choice.” Put differently, only probabilities that form a fixed point of the composite mapping $\sigma \circ \bar{u}$ can form a QRE, and experience suggests that fixed points are special.

However, the freedom to choose the joint densities f_i to fit the data gives considerable flexibility to QRE, particularly if one is unwilling to assume *a priori* that payoff perturbations are i.i.d.. To see this, consider relaxing the assumption of i.i.d. perturbations across each player’s strategies in one of two ways. Let

$$\mathcal{I}_J = \{\text{joint pdfs for } J \text{ independent, mean-zero random variables}\}$$

$$\mathcal{S}_J = \{\text{joint pdfs for } J \text{ mean-zero random variables with identical marginal distributions}\}.$$

Joint densities f_i in the set \mathcal{I}_J imply independence of ϵ_{ij} across strategies j , without requiring that they be identically distributed. Joint densities f_i in \mathcal{S}_J allow dependence of ϵ_{ij} and ϵ_{ik} , $k \neq j$, but require ϵ_{ij} to be identically distributed for all j .

The following result shows that even when payoff perturbations are restricted to come from densities in one of these fairly restrictive classes, QRE imposes no restriction on behavior. For *any* game and *any* distribution of observed behavior on the interior of the J_i -dimensional simplex for each i , there exist densities from $\mathcal{I}_{J_i} \forall i$, as well as densities from $\mathcal{S}_{J_i} \forall i$, any of which will enable a QRE to match the distribution of behavior of each player perfectly.⁹

⁹As the proof makes clear, the results apply to the “1-player” case of an additive random utility discrete choice model. For that paradigm, Berry (1994) has shown that if utilities for each choice j are given by $\bar{u}_j + \epsilon_{ij}$ and an

Theorem 1 Take any finite n -player normal form game Γ with $j = 1, \dots, J_i$ pure strategies for each player i . For any p on the interior of Δ ,

- (i) there exist joint probability density functions $f_i \in \mathcal{I}_{J_i} \forall i$ such that p forms a QRE of Γ .
- (ii) there exist joint probability density functions $f_i \in \mathcal{S}_{J_i} \forall i$ such that p forms a QRE of Γ .

Proof. Given p_{-i} , the probability that player i plays action j in a QRE is given by

$$\sigma_{ij}(\bar{u}_i(p)) = \Pr \{ \epsilon_{ij} \geq \epsilon_{ik} + \bar{u}_{ik}(p) - \bar{u}_{ij}(p) \quad \forall k = 1, \dots, J_i \}.$$

Noting that $\bar{u}_{ij}(p)$ and $\bar{u}_{ik}(p)$ depend only on p_{-i} , let

$$H_i^{jk}(p_{-i}) = \bar{u}_{ik}(p) - \bar{u}_{ij}(p).$$

Part (i) [part (ii)] will then be proven if we can show that for each player i and any given $(p_{i1}, \dots, p_{iJ_i}) \in (0, 1)^{J_i}$, a density $f_i \in \mathcal{I}_{J_i}$ [$f_i \in \mathcal{S}_{J_i}$] can be found that implies

$$\Pr \left\{ \epsilon_{ij} \geq \epsilon_{ik} + H_i^{jk}(p_{-i}) \quad \forall k = 1, \dots, J_i \right\} = p_{ij} \quad j = 1, \dots, J_i \quad (4)$$

i.e., that the probabilities p_{ij} are in fact best responses given p_{-i} .

For simplicity, for both part (i) and part (ii), we will consider here the case of a game in which every player has two pure strategies. An Appendix shows how to generalize these results to the case of an arbitrary number of strategies for each player.

(i) Take player 1 and let p_{1j} be the (given) probability that player 1 chooses strategy s_{1j} . Let $(\epsilon_{11}, \epsilon_{12})$ be independent draws from two-point distributions such that

$$\epsilon_{1j} = \begin{cases} \alpha_j & \text{with prob. } q_j \\ -\frac{q_j}{1-q_j} \alpha_j & \text{with prob. } 1 - q_j \end{cases}$$

for some $\alpha_j > 0$ and $q_j \in (0, 1)$ to be determined.¹⁰ By construction, each ϵ_{1j} has mean zero. Suppose $H_1^{12}(p_{-1}) > 0$ (the complementary case is analogous). Figure 1 illustrates. Realizations of $(\epsilon_{11}, \epsilon_{12})$ in the shaded region lead to strategy s_{11} being chosen over s_{12} . Set $\alpha_1 > H_1^{12}(p_{-1})$ and $\alpha_2 = \gamma [\alpha_1 - H_1^{12}(p_{-1})]$ for some $\gamma \in [0, 1)$. Let $q_2 = 1/2$. To match p_{11} exactly, set $q_1 = p_{11}$. Repeating the argument for each player then proves the result.

arbitrary joint distribution of the perturbations $\{\epsilon_{ij}\}_{j=1}^J$ is given, there exists a (unique) vector of mean utilities $(\bar{u}_1, \dots, \bar{u}_J)$ that will rationalize arbitrary probabilities on choices $\{1, \dots, J\}$. This contrasts with our result where, in the discrete choice case, arbitrary mean utilities are given and we choose a distribution of mean-zero disturbances to match arbitrary data. Dagsvik (1994), McFadden and Train (2000), and Joe (2001) consider related problems of choosing distributions from particular families to approximate choice probabilities (they also consider variation in the set of choices and/or choice characteristics). See also McFadden (1978). As in Berry (1994), all of these allow mean utilities to be chosen to fit the data. None of these results implies the others, although ours is more relevant to experimental settings, where mean payoffs are given.

¹⁰The two-point support is used only to provide a simple construction. Our prior working paper, Haile et al (2003), showed that the mixtures of univariate normal densities (replacing the mixtures of Dirac-delta functions here) can be used to obtain the same result with continuously distributed perturbations.




Figure 1: Illustration for part (i).

Part (ii). Suppose $H_i^{12}(p_{-i}) > 0$ (the case $H_i^{12}(p_{-i}) \leq 0$ is analogous) and choose any $\delta_1 > \bar{u}_{i2}(p) - \bar{u}_{i1}(p)$. Let $\epsilon_{i2} = \xi$ be uniformly distributed on $[-\kappa, \kappa]$, where $\kappa > \frac{\delta_1}{2}$ will be chosen below. Let

$$\epsilon_{i1} = \begin{cases} \xi + \delta_1 & \xi + \delta_1 \leq \kappa \\ \xi + \delta_1 - 2\kappa & \xi + \delta_1 > \kappa. \end{cases}$$

or, letting \oplus represent addition on the circle $[-\kappa, \kappa]$ (see Figure 2),

$$\epsilon_{i1} = \xi \oplus \delta_1.$$

The marginal distributions of ϵ_{i1} and ϵ_{i2} are then both uniform on $[-\kappa, \kappa]$. In Figure 2, the bold arc of the circle shows the set of realizations of ξ that yield $\epsilon_{i2} > \epsilon_{i1}$ (one such realization is shown). The length of this arc (divided by 2κ) determines the probability of this event which, since $\delta_1 > \bar{u}_{i2}(p) - \bar{u}_{i1}(p)$, is also the probability that choice 1 is preferred to choice 2. Then because $\epsilon_{i1} > \epsilon_{i2}$ if and only if $\epsilon_{i2} \leq \kappa - \delta_1$, we have

$$\begin{aligned} p_{i1} &= \Pr(\epsilon_{i1} > \bar{u}_{i2} - \bar{u}_{i1} + \epsilon_{i2}) \\ &= \Pr(\epsilon_{i2} \leq \kappa - \delta_1) \\ &= 1 - \frac{\delta_1}{2\kappa}. \end{aligned}$$

Because we are free to choose any $\kappa > \frac{\delta_1}{2}$, any $p_{i1} \in (0, 1)$ can be matched. Repeating the argument for each player then proves the result. \square




Figure 2: Illustration for part (ii).

4 Discussion

Theorem 1 shows that when the assumption of i.i.d. payoff perturbations is partially relaxed, any distribution of behavior by each player is consistent with a QRE. Hence, any falsifiable implication of QRE must be derived from additional maintained hypotheses on payoff perturbations. Even if one views the perturbations only as a device for smoothing out best response functions, one must be concerned about whether the way this is done is important. Theorem 1 shows that this choice can completely determine equilibrium predictions. This raises at least three important questions.

One question is how relevant our result is, given the literature's focus on i.i.d. perturbations from particular parametric families (typically logit). Those assumptions do imply testable restrictions. For example, McKelvey and Palfrey (1995, proof of Theorem 3) have shown that generally the set of probabilities that can form a logit QRE is a one-dimensional manifold, i.e., a set of curves, each of which implicitly defines all probabilities in terms of just one.¹¹ This can be a very strong restriction, although this is something worth checking in each application. For example, in a symmetric 2×2

¹¹For example, the behavior of a player with 3 available actions is characterized by 2 probabilities. In a logit QRE these probabilities must lie on a set of curves (often one curve) in $[0, 1]^2$. If one requires a single logit parameter to rationalize the behavior of both players (each with 3 pure strategies), equilibrium is characterized by a set of curves in $[0, 1]^4$.

game it may have limited bite, at least under the usual assumptions of symmetric equilibrium with identical distributions of perturbations for each player: in that case, the adding-up constraint already forces the 2 choice probabilities to lie on a line. However, in games with more than 2 strategies per player, this can starkly limit the outcomes that a logit QRE can rationalize.¹² With only the i.i.d. assumption, the set of probabilities consistent with QRE can become considerably larger, but still place useful limits on the behavior a QRE can explain.¹³

Because the strength of the i.i.d. (or i.i.d. logit) assumption varies with the game in question, in practice it would be useful to simulate the range of QRE outcomes possible given the particular game and distributional assumptions being considered. While informal, this could provide a sense of how to interpret the success or failure of the particular specification to rationalize the data. As an illustration, consider the following game studied by McKelvey and Palfrey (1995)

	B_1	B_2	B_3
A_1	(15, -15)	(0, 0)	(-2, 2)
A_2	(0, 0)	(-15, 15)	(-1, 1)
A_3	(1, -1)	(2, -2)	(0, 0)

with unique Nash equilibrium at (A_3, B_3) . A feature of this game is that action A_2 becomes unattractive for player 1 if there is a nontrivial chance that 2 plays B_2 . A symmetric argument applies to 2's action B_1 . When payoff perturbations are introduced to form a QRE, all actions are played with positive probability. Except when the payoff perturbations swamp the mean payoffs (so that a QRE puts nearly identical probabilities on all actions), this makes the actions A_2 and B_1 undesirable. As computations by McKelvey and Palfrey (1995, Table III and Figure 3) show, logit QRE probabilities for these actions are nearly zero for all values of the logit parameter above a certain threshold (given the symmetry of the game, they assume the same logit parameter for all players). Beyond this threshold, there is really only one probability to match in a symmetric equilibrium (since the third probability must add to 1) with the choice of the one logit parameter. Indeed, McKelvey and Palfrey show that there is a continuum of quantal response equilibria of the following form (with symmetric properties for player 2): probability of nearly zero on A_2 and essentially any division of the remaining probability between A_1 and A_3 satisfying $\Pr(A_3) \geq \Pr(A_1)$ —a condition implied by a much weaker assumption of exchangeable perturbations (discussed below).

¹²The same is true for asymmetric 2×2 games if one requires the same logit parameter to rationalize the behavior of each player. See, e.g., McKelvey, Palfrey and Weber (2000). They find that with this restriction the logit QRE fails to fit the data and thus propose allowing different logit parameters for each player.

¹³For example, the Monotonicity property described the following section must hold. See also Goeree, Holt and Palfrey (2003), who describe an additional restriction of the (implicit) assumption of i.i.d. perturbations in an asymmetric matching pennies game. They reject the assumption of QRE with i.i.d. perturbations but are able to fit the data by introducing a risk aversion parameter.

Hence, if actual play puts probability close to zero on A_2 and B_1 (something we might expect in a QRE or under other notions of how games are played), there is no significant restriction of the i.i.d. logit assumption beyond what is implied by much weaker restrictions. This game is somewhat special but, especially since it is a game that has been used to demonstrate the capabilities of the logit QRE, it points to the importance of careful attention to what types of outcomes would lead one to reject the QRE hypothesis given the game and maintained assumptions one wishes to consider.

Of course, some restrictions implied by i.i.d (or i.i.d. logit) perturbations are undesirable. In the discrete choice literature, concerns about these assumptions have long been voiced based on *a priori* considerations and on their implications for comparative statics and counterfactuals. Both the independence and identical distributions components of the i.i.d. assumption have been challenged. For example, one might expect larger payoffs to have perturbations with larger variances, or strategies that are similar to have similar perturbations. As is well known, the independence assumption has unnatural implications, including the IIA property of Luce (1958) in the case of the logit (cf. Debreu, 1960). Considerable effort has been directed at developing tractable models of random utility discrete choice that relax the i.i.d. assumption (e.g., Hausman and Wise, 1978; McFadden, 1978; Berry, Levinsohn, and Pakes, 1995; McFadden and Train, 2000, Akerberg and Rysman, 2005). In the strategic context of QRE, motivations for relaxing the i.i.d. assumption are the same: these distributions will often be too restrictive to fit a rich data set or to lead to reasonable out-of-sample predictions. Testable restrictions that do not rely on these or other arbitrary distributional assumptions may enable more meaningful evaluation of the QRE hypothesis itself.

A second question is the relationship of our result on the *falsifiability* of the QRE to *identifiability* of empirical models based on the QRE. Falsifiability and identifiability are related but distinct.¹⁴ One implication of Theorem 1 is immediate, however. If knowing the payoffs of the underlying game places no restriction on outcomes, observed outcomes cannot place any restriction on (much less identify) payoffs when they are unknown. Hence, Theorem 1 implies that observing the distribution of behavior in a single game could reveal nothing about latent expected payoffs, even when the perturbations are restricted to be draws from distributions in the set I_{J_i} or S_{J_i} . This is a more negative result than the failure of (point) identification: the data contain no information whatsoever about underlying payoffs.¹⁵ However, this implication is of little relevance in the

¹⁴In general neither identifiability nor falsifiability implies the other.

¹⁵This result, like Theorem 1, extends immediately to the “one player” discrete choice environment. It should be pointed out, however, that the empirical literature on discrete choice has rarely considered estimation using data only on the characteristics and choice probabilities for a single choice set. The econometrics and empirical literatures on discrete choice generally rely on variation in covariates and/or the choice set itself, maintaining assumptions about how the distribution of perturbations varies with these changes. We explore similar ideas to develop testable

experimental literature on QRE, since expected payoffs are easily calculated from the payoffs in the underlying game and the observed behavior of opponents.¹⁶

A third important question raised by Theorem 1 is whether there are other ways to evaluate the QRE hypothesis.¹⁷ We devote the following section to discussion of a few promising approaches.

5 Testing the QRE Hypothesis

The most promising testing approaches, in our view, come from observation of behavior in different games. This alone is not sufficient to deliver testable restrictions, since Theorem 1 implies that any behavior can be matched by appropriately selecting distributions of perturbations separately for each game. However, combining variation in the game with limits on how distribution functions can vary can enable several testing approaches, some of which build on well-known results from the discrete choice literature. The discussion here cannot be exhaustive; rather, our aim is to focus on a few approaches that may be particularly promising in practice. Two definitions will be useful for what follows:

Definition 2 (Exchangeability) *The random variables $(\epsilon_{i1}, \dots, \epsilon_{iJ_i})$ are **exchangeable** if $f_i(\epsilon_{i1}, \dots, \epsilon_{iJ_i}) = f_i(\epsilon_{i\rho(1)}, \dots, \epsilon_{i\rho(J_i)})$ for every permutation operator ρ on the set $\{1, \dots, J_i\}$.*

Definition 3 (Invariance) *The joint distribution of $(\epsilon_{i1}, \dots, \epsilon_{iJ_i})$ is **invariant** if $F_i(\epsilon_{i1}, \dots, \epsilon_{iJ_i} | u_i(\cdot)) = F_i(\epsilon_{i1}, \dots, \epsilon_{iJ_i})$ for all $(\epsilon_{i1}, \dots, \epsilon_{iJ_i})$ and all payoff functions $u_i(\cdot)$.*

Exchangeability (a.k.a. “interchangeability”) is a strong form of symmetry, requiring not only identical marginal distributions for each ϵ_{ij} , but also identical covariances, conditional moments, etc. In the more familiar and closely related discrete choice literature, exchangeability holds for the conditional logit model and also for the multinomial probit model under the restrictions $E[\epsilon_{ij}^2] = \sigma^2 \forall i, j$ and $E[\epsilon_{ij}\epsilon_{ij'}] = \rho \forall j, j' \neq j$. Nested logit and mixed logit (or probit) models, on the other hand, violate exchangeability by design.¹⁸ *Invariance* is a property requiring a similar

restrictions of QRE below.

¹⁶An open question is whether there are useful conditions under which each f_i could be identified using data from experiments. In principle, ideas from the discrete choice literature (e.g., Manski, 1988; Matzkin, 1992) might be extended. However, see the discussion in footnote 19 below. Interest in identification of f_i may be more limited in experimental settings than in applications to field data. However, if f_i were known and believed to be invariant across a set of related games, this would provide a means of making point predictions that could be tested. We discuss tests based on this “invariance” assumption below without requiring identification of f_i .

¹⁷If one takes the incomplete information interpretation of QRE, which requires an assumption of independence of payoff perturbations across players (see footnote 4), this additional assumption could be tested by testing independence of players’ actions. This requires looking at players separately, contrary to the frequent practice of examining symmetric games and pooling observations over players to maximize the number of observations.

¹⁸It is also easy to see that the Monotonicity axiom, defined below, does not hold in general for these models.

lack of sensitivity to variations in payoffs, but allows the possibility of asymmetry—for example, the possibility that perturbations are larger for some actions than others. While strong, this assumption has often been used in the econometrics literature on discrete choice models. Invariance is typically maintained in applications of the conditional logit, nested logit, and multinomial probit, for example, but partially relaxed in mixed logit/probit (random coefficients) models.

5.1 Approach 1: “Regular” QRE

Responding to an earlier draft of this paper, Goeree, Holt and Palfrey (2005) have proposed a refinement of QRE, “regular” QRE. Rather than proposing assumptions on the underlying model, however, they define the refinement with axioms on behavior. Suppose $p \in \Delta$ characterizes behavior in a QRE. As before, let $\sigma_{ij}(\bar{u}_i(p))$ represent the element of p corresponding to $\Pr(i \text{ plays } j)$. Consider the following axioms:

1. **Interiority.** $\sigma_{ij}(\bar{u}_i(p)) > 0$ for all $i, j = 1, \dots, J_i$
2. **Continuity:** $\sigma_{ij}(\bar{u}_i(p))$ is a continuous and differentiable function of $\bar{u}_i(p)$ for all $\bar{u}_i(p) \in R^{J_i}$
3. **Responsiveness:** $\frac{\partial \sigma_{ij}(\bar{u}_i(p))}{\partial \bar{u}_{ij}(p)} > 0$ for all $j = 1, \dots, J_i$
4. **Monotonicity:** $\bar{u}_{ij}(p) > \bar{u}_{ik}(p) \Rightarrow \sigma_{ij}(\bar{u}_i(p)) > \sigma_{ik}(\bar{u}_i(p))$ for all $j = 1, \dots, J_i$

Goeree et al. argue that these axioms are economically and intuitively compelling. Interiority and Continuity are natural technical properties. Responsiveness and Monotonicity are stronger properties with significant economic content. Responsiveness restricts the ways that a player’s behavior can change in response to a *ceteris paribus* change in the expected payoff from one action. Monotonicity restricts probabilistic behavior *within* a game, requiring that actions with higher expected payoffs to be played more often.

Clearly these axioms imply testable restrictions. Monotonicity can be checked directly in any game. Responsiveness concerns changes in behavior across games and is more subtle. Raising i ’s payoffs $u_i(s_{ij}, s_{-i})$ from action j in the baseline game will make j more attractive, *ceteris paribus*, but may ultimately lead to a change in the play of i ’s opponents. This will typically change $\bar{u}_{ik}(p)$ for each k , and could even cause $\bar{u}_{ij}(p)$ to be lower than in the original equilibrium.¹⁹ However,

¹⁹This feature of quantal choice in a strategic setting also suggests challenges for obtaining identification results that build on those for additive random utility models of discrete choice (e.g., Manski (1988), Matzkin (1992)) since these rely the ability to “trace out” the distribution of perturbations through sufficient variation in mean payoffs (utilities) while the distribution of perturbations is held fixed. In some games it may be difficult to generate sufficient variation in expected payoffs through manipulation of the payoffs of the underlying normal form game. Hence, even in an experimental setting, where expected payoffs can be treated as known, identification of f_i may be a challenge.

Proposition 4 of Goeree et al. demonstrates that Responsiveness is nonetheless sufficient in some games to obtain testable restrictions based on changes in the payoffs $u_i(s_i, s_{-i})$ of the baseline game. Although they discuss only one example, testable comparative statics predictions can be derived in other environments in which a change in payoffs that, *ceteris paribus*, makes action j more attractive to player i induces play by i 's opponents that also make action j (weakly) more attractive. We discuss classes of such games below.

A natural question is what assumptions on the underlying model imply the axioms, in particular the substantive Monotonicity and Responsiveness conditions. Goeree et al. show that a sufficient condition for Monotonicity is Exchangeability. They point out that Exchangeability and the implied Monotonicity restriction fail under several additive error structures that seem reasonable *a priori*. They suggest that in practice, however, violations of Monotonicity are rarely observed. A sufficient condition for Responsiveness is Invariance.²⁰

Of course, an attractive feature of the “reduced form” approach used to characterize the regular QRE is precisely the possibility of imposing restrictions on behavior without translating them into equivalent assumptions on the underlying structure.²¹ For example, it is clear that Invariance and Exchangeability are not necessary conditions for, respectively, Responsiveness and Monotonicity, and it is not obvious what weaker assumptions on perturbations would generate these restrictions.

5.2 Approach 2: Rank-Cumulative Probabilities

Start with any game Γ with payoffs given by the functions $u_i \forall i$. Now modify the game by changing the payoffs, leaving the strategy space for each player fixed. For each player i let $u'_i : \Delta \rightarrow \mathbb{R}$ give the new payoffs. We will say that these are two games that *differ only in payoffs*. Having observed repeated play of these two games, let p and p' characterize the observed behavior in each. Define the mean expected payoffs $\mu_i = \frac{1}{J_i} \sum_{j=1}^{J_i} u_i(s_{ij}, p_{-i})$ and $\mu'_i = \frac{1}{J_i} \sum_{j=1}^{J_i} u'_i(s_{ij}, p'_{-i})$. Then for all i and j let $\tilde{u}_{ij} = u_i(s_{ij}, p_{-i}) - \mu_i$ and $\tilde{u}'_{ij} = u'_i(s_{ij}, p'_{-i}) - \mu'_i$, normalizing each player's expected

²⁰We show this below. Goeree, Holt and Palfrey (2005, Propositions 5) argue that, under the usual “admissibility” conditions for the QRE, Exchangeability is sufficient for all their axioms. Their analysis maintains an implicit assumption of Invariance, and their claim is correct under this assumption.

²¹It is worth clarifying that the terms “structural” and “reduced form” here, as in Goeree et al (2005), refer to the way equilibrium is defined or restricted (not to an econometric approach), with the reduced form approach starting directly from statistical best response functions rather than generating these from payoff perturbations. Goeree et al. (2005) suggest that the structural QRE implies restrictions on behavior across games (Proposition 2), and that there are outcomes consistent with the (reduced form) regular QRE that cannot be rationalized by a structural QRE (Proposition 6). However, an assumption of Invariance is implicit in their analysis of the structural QRE, while not imposed on the regular QRE. It is immediate from Theorem 1 that without some restriction on how the joint densities f_i change when a game changes, none of the restrictions described in their Proposition 2 hold, negating the conclusion of their Proposition 6 as well.

payoff from each action by his mean. Let

$$\begin{aligned}\dot{u}_{ij} &= \tilde{u}'_{ij} - \tilde{u}_{ij} \\ d_{i(jk)} &= \tilde{u}_{ij} - \tilde{u}_{ik} \\ d'_{i(jk)} &= \tilde{u}'_{ij} - \tilde{u}'_{ik}.\end{aligned}$$

Without loss of generality, re-index i 's actions so that $\dot{u}_{i1} \geq \dot{u}_{i2} \geq \dot{u}_{i3} \geq \dots$, where some inequality must be strict except in the trivial case with $\dot{u}_{ij} = 0 \forall j$, i.e., when the games do differ, except possibly in the scaling of all payoffs. By our choice of indexing, $d'_{i(jk)} \geq d_{i(jk)} \forall j, k > j$, with at least one inequality strict except in the trivial case. Define the *rank-cumulative probabilities*, $\rho_{ik} = \sum_{l=1}^k p_{il}$ and $\rho'_{ik} = \sum_{l=1}^k p'_{il}$. For example, in the original game, ρ_{ik} gives the probability that i uses a strategy in the set $\{s_{i1}, \dots, s_{ik}\}$ (under the indexing defined above).

Theorem 2 *Consider two games that differ only in payoffs. Under the Invariance assumption, behavior consistent with QRE must produce increasing rank-cumulative probabilities, i.e., $\rho'_{ik} \geq \rho_{ik} \forall k = 1, \dots, J - 1$.*

Proof. Under the Invariance assumption, one may treat the normalized expected payoffs \tilde{u}_{ij} and \tilde{u}'_{ij} as the true expected payoffs without loss of generality. Further, we can write probabilities over realizations of ϵ_i without conditioning on which game is being played. We then have

$$\begin{aligned}\rho_{ik} &= \Pr \left(\max_{l \in \{1, \dots, k\}} \{ \tilde{u}_{il} + \epsilon_{il} \} \geq \tilde{u}_{ij} + \epsilon_{ij} \quad \forall j > k \right) \\ &= \Pr \left(\max_{l \in \{1, \dots, k\}} \{ d_{i(lj)} + \epsilon_{il} \} \geq \epsilon_{ij} \quad \forall j > k \right).\end{aligned}$$

Similarly,

$$\rho'_{ik} = \Pr \left(\max_{l \in \{1, \dots, k\}} \{ d'_{i(lj)} + \epsilon_{il} \} \geq \epsilon_{ij} \quad \forall j > k \right).$$

Since $d'_{i(lj)} \geq d_{i(lj)}$ for all $j > l$, the result follows. \square

A special case of increasing rank-cumulative probabilities arises when $u'_i(s_{ij}, p'_{-i}) > u_i(s_{ij}, p_{-i})$ for exactly one $j \in J_i$, with $u'_i(s_{ik}, p'_{-i}) = u_i(s_{ik}, p_{-i})$ for all $k \neq j$. Theorem 2 then requires the probability of play of this action to increase, giving Goeree, Holt and Palfrey's (2005) Responsiveness property. While Invariance is sufficient for Responsiveness, Theorem 2 shows that Invariance also implies a richer set of testable restrictions. These restrictions may also be more useful in practice, since they are not limited to cases in which only one of player 1's actions experiences a change in expected payoff.

The Invariance assumption may be strong, especially if one believes that the significance (e.g., variance) of the disturbance term is dependent on the "stakes" faced by the agent. It is not obvious

whether this should be the case, and indeed one interpretation of the QRE is that it provides a way for “mistakes” to be made more often when they are not very costly (when mean payoff differences are small)—a property that could be undone with perturbations scaled by the associated payoff differences. Here one can obtain a testable restriction with the somewhat weaker assumption that each f_i is invariant to variations in the *baseline* payoffs u_{-i} of *other* players. For example, consider a two-player game, where one examines behavior under variations in player 2’s payoff matrix under the assumption that f_1 does not change in response. After recalculating \bar{u}_1 based on player 2’s new behavior, it is immediate that p_{1j} must be monotonic in $(\bar{u}_{1j} - \bar{u}_{11})$ across j .

5.3 Approach 3: Block-Marschak Polynomials

The final approach we discuss is based on an extension of results obtained by Block and Marschak (1960) and Falmagne (1978) for random utility models of discrete choice.²² We first review this result and then describe an extension to strategic contexts.

Begin with a decision problem in which a utility-maximizing agent must choose one alternative from a finite choice set A . The utility from alternative $j \in A$ is given by the sum of a mean utility \bar{u}_j and a zero-mean random component ε_j . The mean utilities may or may not be known to the researcher, and no restriction is made on the joint distribution of the perturbations $\varepsilon_j, j \in A$. Let $p(A) = \{p_1(A), \dots, p_{|A|}(A)\}$ denote the resulting choice probabilities. Now consider a series of related choice experiments in which the agent chooses from restricted choice sets $B \subset A$. In these experiments, the mean utilities $\bar{u}_j, j \in B$, are identical to those in the original choice problem. We refer to this as a *sequence of choice experiments based on a master choice set A* . For each choice set B , the joint distribution of the latent utility perturbations (i.e., $\varepsilon_j, j \in B$) is also assumed to be identical to the marginal (with respect to the relevant set of choices) distribution of these perturbations in the original problem. We will refer to this as an assumption of a *fixed stochastic structure*. Under this assumption, the underlying random utility structure is held fixed across the sequence of experiments, but the choice sets are restricted to subsets of the master set A .

Let $p(B) = \{p_1(B), \dots, p_{|A|}(B)\}$ denote the corresponding choice probabilities for each choice set B , where of course $p_j(B) = 0$ whenever $j \notin B$. Let $F(B, m)$ denote the set of all subsets of B containing exactly m elements. Then for all $B \subset A$ with $B \neq A$ and any $j \in A - B$,²³ define the *Block-Marschak polynomial*

$$K_{j,B} = \sum_{k=0}^{|B|} (-1)^k \sum_{C \in F(B, |B|-k)} p_j(A - C).$$

²²See also Barbera and Pattanaik (1986) and McFadden (2005). We thank a referee for making us aware of Falmagne (1978).

²³The notation $A - B$ denotes the difference between sets A and B .




Figure 3: Each set E_B represents the region of realizations of perturbations making alternative j the utility-maximizing choice when B is the choice set, so that $p_j(B) = \Pr(E_B)$. Note that both $E_{A-\{k\}} \subset E_{A-\{k,\ell\}}$ and $E_{A-\{\ell\}} \subset E_{A-\{k,\ell\}}$ must hold. $K_{j,\{k,\ell\}}$ gives the probability of the unshaded area.

Lemma 1 (Falmagne (1978)) *Consider a sequence of choice experiments based on a master choice set A . Under the assumption of a fixed stochastic structure, choice probabilities $p(B)$, $B \subset A$, are consistent with utility maximization in a random utility model if and only if all Block-Marschak polynomials are nonnegative.*

While the compact notation for the Block-Marschak polynomials is opaque, the underlying idea is simple: the probability of choosing a given alternative j is higher when the set of other available choices is smaller. For $B = \emptyset$, $K_{j,B} = p_j(A)$, which obviously cannot be negative. If $|B| = 1$,

$$K_{j,B} = p_j(A - B) - p_j(A)$$

and the requirement is that dropping B from the choice set weakly raises the choice probability on each remaining alternative. When B contains 2 elements, k and ℓ ,

$$K_{j,B} = p_j(A - \{k, \ell\}) - [p_j(A - \{k\}) + p_j(A - \{\ell\})] + p_j(A)$$

which again must be positive (see Figure 3). For larger $|B|$, the polynomials have a higher order but a similar interpretation: the probability of a given choice goes up whenever other alternatives are removed from the choice set.

To extend Falmagne’s result to strategic settings, we will consider variation in the set of pure strategies available to a player in a game that is otherwise held fixed. This extension would be immediate were it not for the fact that changing player i ’s strategy space may lead to changes in other players’ equilibrium behavior. Since $\bar{u}_{ij}(p)$ depends on p_{-i} , this will typically lead to a violation of the condition that the expected payoff from each given action (i.e., that prior to the realization of ϵ_i) be the same across all games in which that action is available.

This can be overcome in some types of games, and the laboratory provides an ideal environment for considering them. A trivial approach is to examine the behavior of a player whose payoffs from each action are independent of the actions of his opponents. In a QRE, such a player acts as if he faces a random utility discrete choice problem, and the extension of Lemma 1 is immediate. This is not very satisfying as a test of QRE, since the essence of QRE is strategic behavior. However, games of this sort can provide more meaningful restrictions if we consider an *opponent* of players who face non-strategic decisions.

For example, consider a game Γ_0 that is *non-strategic for player 1*, i.e., letting S_{i0} denote i ’s strategy space

$$u_1(s_{1j}, s_{-i}) = u_1(s_{1j}) \text{ for all } s_{1j} \in S_{10}, s_{-i} \in S_{-i0}.$$

Suppose for simplicity that Γ_0 is a 2-player game. We focus on player 2, whose payoffs *do* depend on the actions of player 1. Consider a sequence of games $\Gamma_1, \dots, \Gamma_G$ which differ from Γ_0 only in that player 2’s strategy spaces S_{21}, \dots, S_{2G} are subsets of the original S_{20} . For each player i , the payoff function $u_i(s_i, s_{-i})$ is the same in each game.²⁴ No restriction is placed on the joint distribution of perturbations in the master game. However, analogous to the discrete choice case, we assume a *fixed stochastic structure*, i.e., in each game Γ_g the payoff perturbations ϵ_{ij} , $j \in S_{ig}$ have joint distribution equal to the marginal distribution of these same perturbations in Γ_0 . Because all of these games remain nonstrategic for player 1, player 1’s QRE behavior is the same in every game. Thus, player 2’s mean payoff \bar{u}_{2j} from any available pure strategy j is the same in each game. In a QRE player 2 anticipates the equilibrium distribution of behavior by his opponent, and his behavior is then as if he were maximizing his utility in a sequence of choice experiments based on a fixed stochastic structure, with the expected payoffs $\bar{u}_{2j}(p)$ replacing the mean utilities \bar{u}_j . Lemma 1 then can be applied, giving the following result.

Theorem 3 *Consider a game Γ_0 that is nonstrategic for all players except i and sequence of games $\Gamma_1, \dots, \Gamma_G$ that differ from Γ_0 only in that player i ’s strategy spaces S_{i1}, \dots, S_{iG} are subsets of the original S_{i0} . Let $p_{ij}(S_{ig})$ denote the probability that i plays pure strategy j in game g . Under the*

²⁴The domain of u_2 does change across games, but its value at any given profile of feasible actions does not.

assumption of a fixed stochastic structure, behavior is consistent with QRE if and only if, for all $B \subset S_{i0}$ with $B \neq S_{i0}$ and all $j \in S_{i0} - B$, $K_{j,B} = \sum_{k=0}^{|B|} (-1)^k \sum_{C \in F(B, |B|-k)} p_{ij}(S_{i0} - C) \geq 0$.

As an example, consider an n -player game Γ_0 that is nonstrategic for all players except player 1, who has pure strategies $S_{10} = \{1, 2, 3\}$. Suppose we observe probabilities $\{p_{11}(\Gamma_0), p_{12}(\Gamma_0), p_{13}(\Gamma_0)\}$ in the game Γ_0 , and $\{p_{11}(\Gamma_1), p_{12}(\Gamma_1)\}$ in game Γ_1 where player 1's strategy set is reduced to $S_{11} = \{2, 3\}$. Then Theorem 3 gives the testable restrictions

$$p_{1j}(\Gamma_1) - p_{1j}(\Gamma_0) \geq 0 \quad j \in \{2, 3\} \quad (5)$$

i.e., player 1 must be more likely to choose a given action when the set of alternative strategies available is smaller. Suppose that, in addition, we observe play probabilities $\{p_{11}(\Gamma_2), p_{13}(\Gamma_2)\}$ from Γ_2 , where player 2's strategy set is reduced to $\{1, 3\}$. Along with the two inequalities analogous to (5), this gives us the additional testable restriction

$$1 + p_{13}(\Gamma_0) \geq p_{13}(\Gamma_1) + p_{13}(\Gamma_2)$$

Note that observing player 1's behavior in three games gives 5 testable restrictions. Observing behavior in the game Γ_3 with $S_{13} = \{1, 2\}$ would give 4 more

$$\begin{aligned} p_{1j}(\Gamma_3) - p_{1j}(\Gamma_0) &\geq 0 \quad j \in \{1, 2\} \\ 1 + p_{11}(\Gamma_0) &\geq p_{11}(\Gamma_2) + p_{11}(\Gamma_3) \\ 1 + p_{12}(\Gamma_0) &\geq p_{12}(\Gamma_1) + p_{12}(\Gamma_3) \end{aligned}$$

yielding 9 restrictions from 4 treatments. Considering games with larger numbers of strategies, and running experiments with larger numbers of subsets of the original strategy set allow us to use higher-order Block-Marschak polynomials, leading to a large number of testable restrictions.

The limitation to games that are strategic for only one player can be overcome if one considers some simple sequential games.²⁵ For example, consider the class of finite "Stackelberg games," i.e., 2-period 2-player sequential move games. Let $\Gamma_0, \Gamma_1, \dots$ be a sequence of Stackelberg games with the first mover labeled player 1. Let the strategy space for player 2 be identical in all games, while the strategy space for player 1 in each game is a subset of that in the master game Γ_0 . Let the payoff functions $u_i(s_i, s_{-i})$ be the same across all games. As above, we refer to this as a *sequence of*

²⁵McKelvey and Palfrey (1998) discuss the extension of QRE to general extensive form games, using the agent-normal form representation of the game. For the Stackelberg games we consider here, the distinction between the agent-normal form and standard normal form specification is not important, since each player "moves" only once. Hence for these games the agent-QRE notion developed by McKelvey and Palfrey is an obvious extension of the QRE for normal form games discussed above.

Stackelberg games based on a master game Γ_0 . Under the assumption of a fixed stochastic structure, in each game Γ_g the payoff perturbations ϵ_{ij} , $j \in S_{ig}$ have joint distribution equal to the marginal distribution of these same perturbations in the original game Γ_0 .

It is easy to see that the sequential nature of these games leaves the play of the Stackelberg follower constant across all of these games *conditional on the action taken by player 1*. This implies that player 1’s expected payoff from taking an action s_{ij} is the same in any game g for which the action s_{ij} is available. Lemma 1 is then again immediately applicable, giving the following result.

Theorem 4 *Consider a sequence of Stackelberg games based on a master game Γ_0 . Let S_{10} denote player 1’s strategy space in the original game Γ_0 and let $p_{1j}(S_{1g})$ denote the probability that 1 plays pure strategy j in the game Γ_g in which $S_{1g} \subset S_{10}$ is his strategy space. Under the assumption of a fixed stochastic structure, behavior is consistent with QRE if and only if, for all $B \subset S_{10}$ with $B \neq S_{10}$ and all $j \in S_{10} - B$, $K_{j,B} = \sum_{k=0}^{|B|} (-1)^k \sum_{C \in F(B, |B|-k)} p_{1j}(S_{10} - C) \geq 0$.*

The maintained hypothesis for these results (the “fixed stochastic structure” assumption) is that the joint distribution of perturbations $\{\epsilon_{ij}, j \in B\}$ is the same in any game in which i ’s strategy space includes B . This assumption is implied by Invariance, but is clearly weaker. Indeed, it places no restriction on perturbations across a pair of games with different payoffs from the same profile of actions. Neither Responsiveness nor the “scale invariance” property discussed by Goeree et al. (2005) is implied by the fixed stochastic structure assumption. Further, in moving from one game to another with an enlarged set of pure strategies, the perturbations associated with the smaller set of strategies can have unrestricted covariance with those for the added strategies.²⁶ Because the Block-Marschak polynomials provide a rich set of restrictions with quite weak maintained hypotheses, we view this as a particularly promising approach for testing.²⁷

6 Conclusion

The QRE provides an appealing equilibrium notion with several compelling interpretations. While much attention has been given to the ability of the QRE to rationalize behavior observed in experiments, we have pointed out that such evidence is uninformative without significant *a priori* restrictions on the distributions of payoff perturbations. This should not be mistaken for a critique of the QRE notion itself.²⁸ Rather, our aim has been to clarify the limitations of data from

²⁶Thus, for example, the (Luce, 1959) IIA property need not hold.

²⁷For the discrete choice setting, Joe (2000) has described additional intuitive implications that can be tested if one assumes that perturbations are mutually independent.

²⁸See Ledyard (1986) for discussion in a similar spirit of the empirical restrictions imposed by Bayesian Nash equilibrium.

single games and move forward to develop approaches for more informative evaluation of QRE. In general this will require maintaining assumptions beyond those in the definition of a QRE itself, and the most appropriate set of maintained hypotheses may depend on the application. We have pointed readers to an axiomatic approach proposed recently by Goeree, Holt and Palfrey (2005) and suggested two new approaches. Each approach examines comparative statics predictions, relying explicitly or implicitly on maintained assumptions about how the distributions of payoff perturbations are related across different games. Each approach can produce a large number of testable restrictions from a relatively small number of different experimental treatments.²⁹ Currently, evidence regarding such comparative statics predictions of QRE is limited,³⁰ and we hope that attention to this topic here and in Goeree, Holt and Palfrey (2005) leads researchers to explore richer sets of testable restrictions in order to better understand the applicability of the QRE notion.

²⁹We have focused on the theoretical question of falsifiability, leaving open interesting questions of formal testing procedures. Obviously the multinomial inequality form of the implied hypotheses will make some naive tests inappropriate.

³⁰Although many papers have examined the fit of QRE in different treatments (e.g., varying payoffs), a new value of the distributional parameter(s) is usually estimated freely for each treatment. Capra, Goeree, Gomez and Holt (1999) demonstrate the ability of QRE with a single distribution of perturbations to rationalize observed comparative statics in the “traveller’s dilemma” game. Goeree, Holt and Palfrey (2002) formally test the assumption of a fixed distribution and reject. Camerer, Ho and Chong (2004) report that a fixed distribution does poorly in predicting outcomes across the different games they analyze. Like Cason and Reynolds (2005), they suggest that the distribution of perturbations required to rationalize the data varies with the scale of the payoffs. Other studies that re-estimate the distribution for each treatment (e.g., McKelvey and Palfrey (1995), Fey, McKelvey and Palfrey (1996)) report that the distribution that best explains behavior varies as players gain experience.

Appendix

Proof of Theorem 1 with $J_i > 2$ pure strategies:

(Part i) Suppose again that all ϵ_{ij} are independent draws from two-point distributions such that

$$\epsilon_{ij} = \begin{cases} \alpha_j & \text{withprob. } q_j \\ -\frac{q_j}{1-q_j}\alpha_j & \text{withprob. } 1 - q_j \end{cases}$$

for some $\alpha_j > 0$ and $q_j \in (0, 1)$ to be determined below. By construction, each ϵ_{ij} has expectation zero. Let $A_{jk} = \Pr\{\epsilon_{ij} \geq \epsilon_{ik} + H_i^{jk}(p_{-i})\}$. Now begin by fixing $\alpha_{J_i} > 0$ and $q_{J_i} \in (0, 1)$ at arbitrary values. Then for any $q_{J_i-1} \in (0, 1)$ and all sufficiently large α_{J_i-1}

$$A_{(J_i-1)J_i} = q_{J_i-1}q_{J_i} + q_{J_i-1}(1 - q_{J_i}) = q_{J_i-1}.$$

Given q_{J_i-1} , fix α_{J_i-1} at one such value, $\alpha_{J_i-1}^*$. We then also have $A_{J_i(J_i-1)} = 1 - q_{J_i-1}$. Now consider selection of α_{J_i-2} . As before, for any $q_{J_i-2} \in (0, 1)$, there exists sufficiently large α_{J_i-2} such that

$$\begin{aligned} A_{(J_i-2)(J_i-1)} &= q_{J_i-2} \\ A_{(J_i-2)J_i} &= q_{J_i-2} \\ A_{J_i(J_i-2)} &= 1 - q_{J_i-2} \\ A_{(J_i-1)(J_i-2)} &= 1 - q_{J_i-2}. \end{aligned}$$

Proceeding in this fashion, given any $q_j \forall j$, we can choose each α_j so that

$$A_{jk} = \begin{cases} q_j & \text{if } j < k \\ 1 - q_k & \text{if } j > k. \end{cases} \quad (6)$$

This construction introduces a particular second-order stochastic dominance ordering of the random variables ϵ_{ij} . With this ordering, the event

$$\left\{ \epsilon_{ij} \geq \epsilon_{ik} + H_i^{jk}(p_{-i}) \quad \forall k = 1, \dots, J_i \right\}$$

is equivalent to the event $\{\epsilon_{ij} > 0, \epsilon_{ik} < 0 \forall k < j\}$ when $j < J_i$, and to the event $\{\epsilon_{ik} < 0 \forall k < j\}$ when $j = J_i$ (realizations of ϵ_{ik} for $k > j$ do not matter). Because all ϵ_{ij} are independent, these events have probability $q_j \prod_{k < j} (1 - q_k)$ for $j < J_i$ and probability $\prod_{k < J_i} (1 - q_k)$ for $j = J_i$. So to satisfy (4), for each $j < J_i$ we set

$$q_j = \frac{p_{ij}}{1 - \sum_{k < j} p_{ik}}$$

(recall that the values of each q_j above were arbitrary). Note that $q_j \in (0, 1) \forall j$ because $p_{ij} \in (0, 1) \forall j$ and $\sum_{j=1}^{J_i} p_{ij} = 1$. Repeating this argument for each player i then shows that we can construct

distributions for each ϵ_{ij} that yield any desired probabilities as a QRE if we ignore the fact that the definition of a QRE assumed continuously distributed perturbations.³¹ However, the mixtures of Dirac-delta functions used as densities here can be replaced with mixtures of univariate normal densities (with small variances) to obtain the same result. We show this in Appendix B.³²

(Part ii) Let ξ be uniformly distributed on $[-\kappa, \kappa]$, for some $\kappa > 0$, to be chosen below. For $j = 1, \dots, J_i$ define

$$\epsilon_{ij} = \begin{cases} \xi + \delta_j & \xi + \delta_j < \kappa \\ \xi + \delta_j - 2\kappa & \xi + \delta_j > \kappa \end{cases} \quad (7)$$

where each δ_j will be a distinct value in the interval $[0, 2\kappa]$ to be determined below. Each ϵ_{ij} is then uniformly distributed on $[-\kappa, \kappa]$. Fix δ_{J_i} at zero and, without loss of generality, impose $\delta_1 > \delta_2 > \dots > \delta_{J_i}$. Suppose for the moment that $H_i^{jk} = 0$ for all j and k . One can then confirm that for each j

$$\Pr\{\epsilon_{ij} > \epsilon_{ik}, \quad k = 1, \dots, J_i\} = \frac{\delta_{j-1} - \delta_j}{2\kappa}$$

where we define $\delta_0 = 2\kappa$. Setting these probabilities equal to the given values p_{i1}, \dots, p_{iJ_i} , we obtain the solution

$$\delta_j = \left(1 - \sum_{\ell=1}^j p_{i\ell}\right) 2\kappa \quad j = 1, \dots, J_i - 1. \quad (8)$$

We now drop the assumption that each $H_i^{jk} = 0$. We do this by ensuring the equivalence

$$\{\epsilon_{ij} > \epsilon_{ik} + H_i^{jk}\} \iff \{\epsilon_{ij} > \epsilon_{ik}\} \quad (9)$$

which holds if ϵ_{ij} and ϵ_{ik} are always sufficiently different when $j \neq k$. When (8) holds we know $|\epsilon_{ij} - \epsilon_{ik}| \geq 2\kappa (\min_{j=1, \dots, J_i} p_{ij})$ for all $j \neq k$.³³ So for any $\kappa > \frac{\max_{k,j} |H_i^{jk}|}{2 \min_{j=1, \dots, J_i} p_{ij}}$, (9) holds and we have

$$\Pr\{\epsilon_{ij} > \epsilon_{ik} + H_i^{jk} : k = 1, \dots, J\} = p_{ij} \quad \forall j \neq k.$$

Repeating this construction for every player completes the proof. \square

³¹There are infinitely many other constructions since there are infinitely many ways to choose the parameters α_j (e.g., varying the starting value α_{J_i} in the proof, selecting different values of each α_j^* , or introducing the second-order stochastic dominance for any other ordering of the pure strategies).

³²It is intuitive that mixtures of normals could approximate the two-point distributions above arbitrarily well. Appendix B shows, however, that we can match the probabilities p_{ij} exactly.

³³When $\xi + \delta_j$ and $\xi + \delta_k$ both exceed κ or are both smaller than κ , this is immediate from (8). When $\xi + \delta_j > \kappa > \xi + \delta_k$, $|\epsilon_j - \epsilon_k| = |\delta_k - \delta_j + 2\kappa|$, and the claim then follows from (8).

References

- Akerberg, D. and Rysman, M. “Unobservable Product Differentiation in Discrete Choice Models: Estimating Price Elasticities and Welfare Effects,” *RAND Journal of Economics*, 36, 771–788, 2005.
- Anderson, S., DePalma, A. and Thisse, J. *Discrete Choice Theory of Product Differentiation*, Cambridge: MIT Press, 1992.
- Anderson, S., Goeree, J. and Holt, C. “Rent Seeking with Bounded Rationality: An Analysis of the All-Pay Auction,” *Journal of Political Economy*, 106(4), 828-853, August 1998.
- Anderson, S., Goeree, J. and Holt, C. “Minimum Effort Coordination Games: Stochastic Potential and the logit Equilibrium” *Games and Economic Behavior*, 34(2), 177-199, February 2001.
- Anderson S., Goeree J. and Holt C. “The Logit Equilibrium: A Perspective on Intuitive Behavioral Anomalies,” *Southern Economic Journal*, 69(1), 21-47, July 2002.
- Augereau, A., Greenstein S., and Rysman, M. “Coordination vs. Differentiation in a Standards War: 56K Modems,” *RAND Journal of Economics*, forthcoming.
- Bajari, P. “Econometrics of Sealed-bid Auctions,” *Proceedings of the Business and Economic Statistics Section of the American Statistical Association*, 1998.
- Bajari, P. and Hortacısu, A. “Are Structural Estimates of Auction Models Reasonable? Evidence from Experimental Data,” *Journal of Political Economy*, 113, 703–741, 2005.
- Barbera, S. and Pattanaik, P.K. “Falmagne and the Rationalizability of Stochastic Choices in Terms of Random Orderings,” *Econometrica*, 54:707–715.
- Baye, M. and J. Morgan. “Price Dispersion in the Lab and on the Internet,” *RAND Journal of Economics*, 35(3), 448–466, Autumn 2004.
- Berry, S. “Estimating Discrete-Choice Models of Product Differentiation,” *RAND Journal of Economics*, 25, 242-262, 1994
- Berry, S., Levinsohn, J. and Pakes, A. “Automobile Prices in Market Equilibrium,” *Econometrica*, 63(4), 841–890, July 1995.
- Block, H.D. and Marschak J. “Random Orderings and Stochastic Theories of Responses,” in Olkin et al., eds, *Contributions to Probability and Statistics: Essays in Honor of Harold Hotelling*, Stanford, 1960.
- Camerer, C. F., T. Ho, and J. K. Chong, “Behavioral Game Theory: Thinking, Learning, and Teaching,” in Steffan Huck, ed., *Advances in Understanding Strategic Behavior, Essays in Honor of Werner Guth*, Palgrave MacMillan, 2004.
- Capra, M., Goeree, J., Gomez, R., and Holt, C. “Anomalous Behavior in a Travelers Dilemma?” *American Economic Review*, 89(3), 678-690, September 1999.
- Cason, T. and Reynolds S., “Bounded rationality in laboratory bargaining with asymmetric information,” *Economic Theory*, 25, 553–574 2005.
- Chen, H., Friedman, J. and Thisse, J. “Boundedly Rational Nash Equilibrium: A Probabilistic Choice Approach,” *Games and Economic Behavior*, 18, 32–54, 1997.

- Dagsvik, J.K. "How Large is the Class of Generalized Extreme Value Random Utility Models," *Journal of Mathematical Psychology*, 39, 90–98.
- Debreu, G. "Individual Choice Behavior: A Theoretical Analysis" (Review Article) *American Economic Review*, 50, 1, 186–188 (1960).
- Dufwenberg, M., Gneezy, U., Goeree, J. and Nagel, R. "Price Floors and Competition," working paper, 2002.
- Falmagne, J.C. "A Representation Theorem for Finite Random Scale Systems," *Journal of Mathematical Psychology*, 18, 52–72 (1978).
- Fey, M., McKelvey, R.D. and Palfrey, T.R. "An Experimental Study of Constant-Sum Centipede Games," *International Journal of Game Theory*, 25(3), 269–287, 1996.
- Goeree, J. and Holt, C. "Stochastic Game Theory: For Playing Games, Not Just for Doing Theory," *Proceedings of the National Academy of Sciences*, 96, 10564–10567, September 1999.
- Goeree, J. and Holt, C. "Asymmetric Inequality Aversion and Noisy Behavior in Alternating-Offer Bargaining Games," *European Economic Review*, 44, 1079–1089, 2000.
- Goeree, J. and Holt, C., "Ten Little Treasures of Game Theory and Ten Intuitive Contradictions," *American Economic Review*, 91(5), 1402–1422, December 2001.
- Goeree, J., Holt, C. and Palfrey, T., "Quantal Response Equilibrium and Overbidding in Private-Value Auctions," *Journal of Economic Theory*, 104(1), 247–272, May 2002.
- Goeree, J., Holt, C. and Palfrey, T., "Risk Averse Behavior in Generalized Matching Pennies Games," *Games and Economic Behavior*, 45, 97–113, 2003.
- Goeree, J., Holt, C. and Palfrey, T., "Regular Quantal Response Equilibrium," *Experimental Economics*, 8, 347–367, December 2005.
- Guarnaschelli, S.; McKelvey, R.D., Palfrey, T., "An Experimental Study of Jury Decision Rules," *American Political Science Review*, 94(2), 407–423, June 2000.
- Haile, P., Hortaçsu, A., and Kosenok, G. "On the Empirical Content of Quantal Response Equilibrium," Cowles Foundation Discussion Paper No. 1432, 2003.
- Hausman, J.A. and Wise, D. A. "A Conditional Probit Model for Qualitative Choice: Discrete Decisions Recognizing Interdependence and Heterogeneous Preferences," *Econometrica*, 46(2), 403–426, March 1978.
- Joe, H. "Inequalities for Random Utility Models, with Applications to Ranking and Subset Choice Data," *Methodology and Computing in Applied Probability*, 2, 359–372, 2000.
- Joe, H. "Multivariate Extreme Value Distributions and Coverage of Ranking Probabilities," *Journal of Mathematical Psychology*, 45, 180–188, 2001.
- Ledyard, J.O. "The Scope of the Hypothesis of Bayesian Equilibrium," *Journal of Economic Theory*, 39(1), June 1986.
- Luce, R.D. *Individual Choice Behavior*, New York: Wiley, 1958.
- Manski, C.F. "Identification of Binary Response Models," *Journal of the American Statistical Association*, 83, 729–738, 1988.

- Matzkin, R. "Nonparametric and Distribution-Free Estimation of the Binary Threshold Crossing and the Binary Choice Models," *Econometrica*, 60(2), 239–270, March 1992.
- McFadden, D. "Modelling the Choice of Residential Location," in A. Karlqvist et al., eds., *Spatial Interaction Theory and Planning Models*, Amsterdam: North Holland, 1978.
- McFadden, D. "Econometric Models of Probabilistic Choice," in D. McFadden and C.F. Manski, eds, *Structural Analysis of Discrete Data with Econometric Applications*, MIT Press, 1981.
- McFadden, D. and K. Train. "Mixed MNL Models for Discrete Response," *Journal of Applied Econometrics*, 15(5), 447–470, 2000.
- McFadden, D. "Revealed Stochastic Preference: A Synthesis," *Economic Theory*, 26, 245-264, 2005.
- McKelvey, R. D. and Palfrey, T. R. "Quantal Response Equilibria for Normal Form Games," *Games and Economic Behavior*, 10(1), 6-38, 1995.
- McKelvey, R.D. and Palfrey, T.R. "Quantal Response Equilibrium in Extensive Form Games," *Experimental Economics*, 1, 9–41, 1998.
- McKelvey, R. D. and Palfrey, T. R. and Weber, R. A. "The Effect of Payoff Magnitude and Heterogeneity on Behavior in 2×2 Games with Unique Mixed Strategy Equilibria," *Journal of Economic Behavior and Organization*, 42, 523-548, 2000.
- Rosenthal, R. "A Bounded-Rationality Approach to the Study of Noncooperative Games," *International Journal of Game Theory*, 18, 273–291, 1989.
- Seim, K. "An Empirical Model of Firm Entry with Endogenous Product-Type Choices," working paper, Stanford, 2002.
- Signorino, C. "Strategic Interaction and the Statistical Analysis of International Conflict," *American Political Science Review*, 93(2), 279–297, June 1999.
- Sweeting, A. "Coordination Games, Multiple Equilibria, and the Timing of Radio Commercials," working paper, Northwestern University, 2004.