

Yale University

EliScholar – A Digital Platform for Scholarly Publishing at Yale

Cowles Foundation Discussion Papers

Cowles Foundation

1-1-2007

GMM Estimation for Dynamic Panels with Fixed Effects and Strong Instruments at Unity

Chirok Han

Peter C.B. Phillips

Follow this and additional works at: <https://elischolar.library.yale.edu/cowles-discussion-paper-series>



Part of the [Economics Commons](#)

Recommended Citation

Han, Chirok and Phillips, Peter C.B., "GMM Estimation for Dynamic Panels with Fixed Effects and Strong Instruments at Unity" (2007). *Cowles Foundation Discussion Papers*. 1892.
<https://elischolar.library.yale.edu/cowles-discussion-paper-series/1892>

This Discussion Paper is brought to you for free and open access by the Cowles Foundation at EliScholar – A Digital Platform for Scholarly Publishing at Yale. It has been accepted for inclusion in Cowles Foundation Discussion Papers by an authorized administrator of EliScholar – A Digital Platform for Scholarly Publishing at Yale. For more information, please contact elischolar@yale.edu.

**GMM ESTIMATION FOR DYNAMIC PANELS WITH FIXED EFFECTS
AND STRONG INSTRUMENTS AT UNITY**

By

Chirok Han and Peter C. B. Phillips

January 2007

COWLES FOUNDATION DISCUSSION PAPER NO. 1599



**COWLES FOUNDATION FOR RESEARCH IN ECONOMICS
YALE UNIVERSITY
Box 208281
New Haven, Connecticut 06520-8281**

<http://cowles.econ.yale.edu/>

GMM Estimation for Dynamic Panels with Fixed Effects and Strong Instruments at Unity*

Chirok Han

Victoria University of Wellington

Peter C. B. Phillips

Cowles Foundation, Yale University

University of York & University of Auckland

August, 2006

Abstract

This paper develops new estimation and inference procedures for dynamic panel data models with fixed effects and incidental trends. A simple consistent GMM estimation method is proposed that avoids the weak moment condition problem that is known to affect conventional GMM estimation when the autoregressive coefficient (ρ) is near unity. In both panel and time series cases, the estimator has standard Gaussian asymptotics for all values of $\rho \in (-1, 1]$ irrespective of how the composite cross section and time series sample sizes pass to infinity. Simulations reveal that the estimator has little bias even in very small samples. The approach is applied to panel unit root testing.

JEL Classification: C22 & C23

Key words and phrases: Asymptotic normality, Asymptotic power envelope, Moment conditions, Panel unit roots, Point optimal test, Unit root tests, Weak instruments.

*Phillips acknowledges partial support from a Kelly Fellowship and the NSF under Grant No. SES 04-142254.

1 Introduction

In simple dynamic panel models it is well-known that the usual fixed effects estimator is inconsistent when the time span is small (Nickell, 1981), as is the ordinary least squares (OLS) estimator based on first differences. In such cases, the instrumental variable (IV) estimator (Anderson and Hsiao, 1981) and generalized method of moments (GMM) estimator (Arellano and Bond, 1991) are both widely used. However, as noted by Blundell and Bond (1998), these estimators both suffer from a weak instrument problem when the dynamic panel autoregressive coefficient (ρ) approaches unity. When $\rho = 1$, the moment conditions are completely irrelevant for the true parameter ρ , and the nature of the behavior of the estimator depends on T . When T is small, the estimators are asymptotically random and when T is large the unweighted GMM estimator may be inconsistent and the efficient two step estimator (including the two stage least squares estimator) may behave in a nonstandard manner. Some special cases of such situations are studied in Staiger and Stock (1997) and Stock and Wright (2000), among others, and Han and Phillips (2006), the latter in a general context that includes some panel cases.

Methods to avoid these problems were developed in Blundell and Bond (1998) and more recently in Hsiao, Pesaran and Tahmiscioglu (2002). Blundell and Bond propose a system GMM procedure which uses moment conditions based on the level equations together with the usual Arellano and Bond type orthogonality conditions. Hsiao et al., on the other hand, consider direct maximum likelihood estimation based on the differenced data under assumed normality for the idiosyncratic errors. Both approaches yield consistent estimators for all ρ values, but there are remaining issues that have yet to be determined in regard to the limit distribution when ρ is unity and T is large.

In a recent paper dealing with the time series case, Phillips and Han (2005) introduced a differencing-based estimator in an AR(1) model for which asymptotic Gaussian-based inference is valid for all values of $\rho \in (-1, 1]$. The present paper applies those ideas to dynamic panel data models, where we show that significant advantages occur. In panels, the estimator again has a standard Gaussian limit for all ρ values including unity, it has virtually no bias except when T is very small ($T \leq 4$), and it completely avoids the usual weak instrument problem for ρ in the vicinity of unity.

As discussed later, this panel estimator makes use of moment conditions that are strong for all values of $\rho \in (-1, 1]$ under the assumption that the errors are white noise over time. (The white noise condition is stronger than that on which the usual IV/GMM approaches by Anderson and Hsiao (1981) or Arellano and Bond (1991) are based.) Under this condition, the proposed estimator is consistent, supports asymptotically valid Gaussian inference even with highly persistent panel data, and is free of initial conditions on levels. These

advantages stem from the following properties: (i) the limit distribution is continuous as the autoregressive coefficient passes through unity; (ii) the rate of convergence is the same for stationary and non-stationary panels; and (iii) differencing transformations essentially eliminate dependence on level initial conditions.

Furthermore, there are no restrictions on the number of the cross-sectional units (n) and the time span (T) other than the simple requirement that $nT \rightarrow \infty$ (and $T > 3$ or $T > 4$ depending on the presence of incidental trends). Thus, neither large T , nor large n is required for the limit theory to hold. Gaussian asymptotics apply irrespective of how the composite sample size $nT \rightarrow \infty$, including both fixed T and fixed n cases, as well as any diagonal path and relative rate of divergence for these sample dimensions. This robust feature of the asymptotics is unique to our approach and differs substantially from the existing literature, including recent contributions by Hahn and Kuersteiner (2002), Alvarez and Arellano (2003), and Moon, Perron and Phillips (2005), who analyze various cases with large n and large T . Apart from the fact that the asymptotic variance of our proposed estimator can be better estimated by different methods when n is large and T is small (because the variance evolves with T), no other modification or consideration is required in the implementation of our approach, so it is well suited to practical implementation. This wide applicability does come at a cost in efficiency for the fixed effects model and a loss of power for the incidental trends model compared with existing methods.

In what follows, section 2 considers the model and estimator for a simple dynamic model with fixed effects, where the basic idea of our transformation is explained. Section 3 deals with a dynamic panel model where exogenous variables are present, and Section 4 studies the case with incidental trends. Section 5 applies the new approach to panel unit root testing. The last section contains some concluding remarks. Proofs are in the Appendix. Throughout the paper we define $0^0 = 1$ and use T_j to denote $\max(T - j, 0)$. We assume that data are observed for $t = 0, 1, \dots, T$.

2 Simple Dynamic Panels

2.1 A New Estimator and Limit Theory

We consider the simple dynamic panel model

$$y_{it} = \alpha_i + u_{it}, \quad u_{it} = \rho u_{it-1} + \varepsilon_{it}, \quad \rho \in (-1, 1],$$

implying

$$(1) \quad y_{it} = (1 - \rho)\alpha_i + \rho y_{it-1} + \varepsilon_{it},$$

where α_i are unobservable individual effects and $\varepsilon_{it} \sim iid(0, \sigma^2)$ with finite fourth moments.

This model differs slightly in its components form from the usual dynamic panel model $y_{it} = \alpha_i + \rho y_{it-1} + \varepsilon_{it}$ in that the individual effects disappear when $\rho = 1$. This formulation is made only to guarantee continuity in the asymptotics at $\rho = 1$. When $|\rho| < 1$ the two models are not distinguishable.

As is well known, the OLS estimator based on the ‘within’ transformation yields an inconsistent estimator because the transformed regressor and the corresponding error are correlated—see Nickell (1981), among others. This bias is also not corrected by first differencing

$$(2) \quad \Delta y_{it} = \rho \Delta y_{it-1} + \Delta \varepsilon_{it},$$

because the transformation induces a correlation between Δy_{it-1} and $\Delta \varepsilon_{it}$. Instead, following Phillips and Han (2005), we transform (2) further into the form

$$(3) \quad 2\Delta y_{it} + \Delta y_{it-1} = \rho \Delta y_{it-1} + \eta_{it}, \quad \eta_{it} = 2\Delta y_{it} + (1 - \rho)\Delta y_{it-1}.$$

Then, this formulation produces the following key moment conditions.

Lemma 1 *If $E\varepsilon_{it}^2 = \sigma^2$ for all t and $E\varepsilon_{is}\varepsilon_{it} = 0$ for all $s \neq t$, then*

$$(4) \quad E g_{it}(\rho) = E \Delta y_{it-1} [2\Delta y_{it} + (1 - \rho)\Delta y_{it-1}] = 0, \quad t = 3, \dots, T$$

for every $\rho \in (-1, 1]$.

It is worth noting that the white-noise condition is required for (4). When $|\rho| < 1$, this white-noise condition is stronger than just serial uncorrelatedness (over time) which is required for the consistency of the Arellano-Bond type IV/GMM estimators for $|\rho| < 1$.

The T_1 (i.e, $T - 1$) moment conditions in (4) are strong for all $\rho \in (-1, 1]$ in the sense that the expected derivatives of the moment functions $g_{it}(\rho)$ differ from zero for all ρ as long as Δy_{it-1} has enough variation across i . This is easily verified by the calculation $E \partial g_{it}(\rho) / \partial \rho = -E(\Delta y_{it-1})^2$.

There are many ways to make use of these T_1 moment conditions. The simplest is to use pooled least squares estimation of (3), which leads to

$$\hat{\rho}_{ols} = \frac{\sum_{i=1}^n \sum_{t=2}^T \Delta y_{it-1} (2\Delta y_{it} + \Delta y_{it-1})}{\sum_{i=1}^n \sum_{t=2}^T (\Delta y_{it-1})^2},$$

which we call the *first difference least squares* (FDLS) estimator. This estimator has the following limit distribution.

Theorem 2 For each T , $\sqrt{nT_1}(\hat{\rho}_{ols} - \rho) \Rightarrow N(0, V_{ols,T})$ as $n \rightarrow \infty$ for all $\rho \in (-1, 1]$, where

$$V_{ols,T} = \frac{ET_1^{-1} \left(\sum_{t=2}^T \Delta y_{it-1} \eta_{it} \right)^2}{\left[ET_1^{-1} \sum_{t=2}^T (\Delta y_{it-1})^2 \right]^2}.$$

As $T \rightarrow \infty$, $V_T \rightarrow 2(1 + \rho)$, and furthermore, $\sqrt{nT_1}(\hat{\rho}_{ols} - \rho) \Rightarrow N(0, 2(1 + \rho))$.

The Gaussian limit theory is valid for any n/T ratio as long as $nT_1 \rightarrow \infty$, including finite T values for which the variance $V_{ols,T}$ evolves with T . Most remarkably, the joint limit as both n and T pass to infinity is identical to the limit where $T \rightarrow \infty$ individually, or the sequential limit as $T \rightarrow \infty$ and then $n \rightarrow \infty$ or the sequential limit as $n \rightarrow \infty$ and then $T \rightarrow \infty$. As a result, the limit theory is remarkably robust to different sample size constellations of (n, T) and simulations support the resulting intuition that testing based on Theorem 2 should show little size distortion.

We remark that the fourth moment condition $E\varepsilon_{it}^4 < \infty$ is required for the limit theory to hold. For small T , the variance $V_{ols,T}$ can be expressed directly in terms of the parameters, using the general formula given in (56) in the Appendix. For example, if $\varepsilon_{it} \sim N(0, \sigma^2)$ or more weakly $E\varepsilon_{it}^4 = 3\sigma^4$ and if $T = 2$, then we have $V_{ols,2} = (1 + \rho)(3 - \rho)$. For $T > 2$ (and fixed) the variances $V_{ols,T}$ are plotted in Figure 1. The expression is quite complicated for general ρ , and is unlikely to be practically useful because $V_{ols,T}$ depends on the nuisance fourth moment of ε_{it} , except for $\rho = 1$ (see below Corollary 3), and because $V_{ols,T}$ can be readily estimated by just replacing the expectation operators with averaging over i and the error process η_{it} with the residuals $\hat{\eta}_{it}$ from the regression of (3). More specifically, when n is large, $V_{ols,T}$ is estimated by


$$(5) \quad \hat{V}_{ols,T} = \frac{(nT_1)^{-1} \sum_{i=1}^n \left(\sum_{t=2}^T \Delta y_{it-1} \hat{\eta}_{it} \right)^2}{\left[(nT_1)^{-1} \sum_{i=1}^n \sum_{t=2}^T (\Delta y_{it-1})^2 \right]^2},$$

where $\hat{\eta}_{it} = 2\Delta y_{it} + \Delta y_{it-1} - \hat{\rho}_{ols} \Delta y_{it-1}$. The corresponding standard error for $\hat{\rho}_{ols}$ is

$$(6) \quad se(\hat{\rho}_{ols}) = \left[\sum_{i=1}^n \left(\sum_{t=2}^T \Delta y_{it-1} \hat{\eta}_{it} \right)^2 \right]^{1/2} / \sum_{i=1}^n \sum_{t=2}^T (\Delta y_{it-1})^2.$$

As $T \rightarrow \infty$, both $V_{ols,T}$ and $\hat{V}_{ols,T}$ converge to $2(1 + \rho)$, and the $N(0, 2(1 + \rho))$ limit holds if $T \rightarrow \infty$ whether or not $n \rightarrow \infty$. Technically speaking, the joint limit as both $n \rightarrow \infty$ and $T \rightarrow \infty$ coincides with the sequential limit as $n \rightarrow \infty$ followed by $T \rightarrow \infty$ or the sequential limit as $T \rightarrow \infty$ followed by $n \rightarrow \infty$. In this case, both (5) and $2(1 + \hat{\rho}_{ols})$ are consistent for

Figure 1: $V_{ols,T}$ for various T with normal errors. As $T \rightarrow \infty$, $V_{ols,T}$ approaches to $2(1 + \rho)$. The convergence is fast when $\rho > 0$.



$V_{ols,T}$, so either formula can be used. This indiscriminating feature is a characteristic of the new approach.

If n is small and T is large, then performance of (5) may be poor as it relies on the law of large numbers across cross sectional units. But in this case the $2(1 + \hat{\rho}_{ols})$ formula approximates the actual variances quite well. This good performance of the asymptotic theory has been confirmed in earlier simulations reported in Phillips and Han (2005) for the time series case where $n = 1$.

When $\rho = 1$, the differenced data (Δy_{it}) are *iid* over both cross-sectional and time-series dimensions, resulting in the same Gaussian limit holding as $nT \rightarrow \infty$ (more precisely, $nT_1 \rightarrow \infty$) irrespective of the n/T ratio, as given in the following result.

Corollary 3 *If $\rho = 1$, then $(nT_1)^{-1/2}(\hat{\rho}_{ols} - 1) \Rightarrow N(0, 4)$ as $nT_1 \rightarrow \infty$.*

Simulation results are given in Table 1 for $T = 2$ and $T = 24$. The exact form of $V_{ols,T}$ is provided in (56) in the Appendix. The simulated test size based on t -ratios with the standard errors obtained by (6) are given in the ‘size’ columns. The results generally reflect the asymptotic theory well, including the consistency of both the estimator and the standard error. When n is relatively small, the standard errors according to (6) slightly underestimate the true variance. The asymptotic variance formula $2(1 + \rho)$ for $\sqrt{nT_1}(\hat{\rho}_{ols} - \rho)$ from Theorem 2, which is appropriate when $T \rightarrow \infty$, obviously does not perform so well when T is as small as it is in this experiment, although the formula is surprisingly good when ρ close to unity.

Also, when $T \geq 3$, the formula $2(1+\rho)$ is rather close to the true variance for reasonably large ρ values (e.g., $\rho \geq 0.5$), at least if the errors are normally distributed. For error distributions with thicker tails one might expect that larger values of T might be needed to get a good correspondence with the asymptotic formula based on $T \rightarrow \infty$.

2.2 Increasing Efficiency

As the moment functions are correlated over t , optimal GMM, which we call *first difference GMM* (FDGMM), is possibly a more efficient alternative to pooled OLS. In order to formulate FDGMM, let D be the T_1 -vector of $E(\Delta y_{it-1})^2$ for $t = 2, \dots, T$ and Ω the $T_1 \times T_1$ matrix for $E\Delta y_{it-1}\eta_{it}\eta_{is}\Delta y_{is-1}$ for $t, s = 2, \dots, T$. Let $h = (h_2, \dots, h_T)' = \Omega^{-1}D$. Then the FDGMM estimator is the method of moments estimator using $\sum_{t=2}^T h_t \Delta y_{it-1} \eta_{it}$ as the moment function. (The FDLS estimator above corresponds to uniform weighting, i.e., $h_2 = \dots = h_T$.) In other words, the FDGMM estimation is the instrumental variable estimator of (3) using $h_t \Delta y_{it-1}$ as the instrument for Δy_{it-1} . Naturally, FDLS and FDGMM are equal if $T = 2$, where the weighting is irrelevant.

It is straightforward to make optimal GMM operational through a two-step procedure. We estimate D by $\tilde{D} = n^{-1} \sum_{i=1}^n [(\Delta y_{i1})^2, \dots, (\Delta y_{iT-1})^2]'$ and Ω by $\tilde{\Omega} = n^{-1} \sum_{i=1}^n \tilde{w}_i \tilde{w}_i'$, where $\tilde{w}_i = (\Delta y_{i1} \tilde{\eta}_{i2}, \dots, \Delta y_{iT-1} \tilde{\eta}_{iT})'$, with $\tilde{\eta}_{it}$ denoting the residuals from (3) using an initial consistent estimate $\tilde{\rho}$ (e.g., $\hat{\rho}_{ols}$). Let $\tilde{h} = (\tilde{h}_2, \dots, \tilde{h}_T)' = \tilde{\Omega}^{-1} \tilde{D}$. Then the second step efficient GMM estimator is obtained by pooled IV regression of (3) using $\tilde{h}_t \Delta y_{it-1}$ as instrument for Δy_{it-1} .

Let us still assume that T is small. Let $\hat{\rho}_{gmm}$ be the FDGMM estimator. Then obviously $(nT_1)^{1/2}(\hat{\rho}_{gmm} - \rho) \Rightarrow N(0, V_{gmm})$ where $V_{gmm} = (T_1^{-1} D' \Omega^{-1} D)^{-1}$. In the case of the two-step efficient GMM, this variance can be estimated by replacing D with \tilde{D} and Ω by $n^{-1} \sum_{i=1}^n \hat{w}_i \hat{w}_i'$, where $\hat{w}_i = \Delta y_{it-1} \hat{\eta}_{it}$ with $\hat{\eta}_{it} = 2\Delta y_{it} + (1 - \hat{\rho}_{gmm})\Delta y_{it-1}$.

Because $D = d\boldsymbol{\iota}$ for some constant d where $\boldsymbol{\iota}$ is the T_1 vector of ones, we have

$$(7) \quad \frac{V_{gmm}}{V_{ols,T}} = \frac{(D' \Omega^{-1} D)^{-1}}{[D' \boldsymbol{\iota} (\boldsymbol{\iota}' \Omega \boldsymbol{\iota})^{-1} \boldsymbol{\iota}' D]^{-1}} = \frac{T_1^2}{\boldsymbol{\iota}' \Omega \boldsymbol{\iota} \boldsymbol{\iota}' \Omega^{-1} \boldsymbol{\iota}} \leq 1,$$

where the last inequality boundary is obtained by the usual algebra in proving the asymptotic efficiency of optimal GMM.

Even though the optimal GMM estimator may have a smaller asymptotic variance than the OLS estimator, the efficiency gain looks marginal in our case. When $\rho = 1$, it can be shown that OLS equals the optimal GMM (because $h_2 = \dots = h_T$). For other ρ values the variance ratio (7) is evaluated in Figure 2 in the case of standard normal errors with $\rho = -0.5, 0, 0.5, 0.9, 1$ and $T = 2, 3, \dots, 100$. The lowest variance ratio is approximately 0.99, which is obtained at $\rho = 0.5$ and $T = 5$, indicating that the efficiency gain of optimal

Table 1: FDLS for $\varepsilon_{it} \sim N(0, 1)$. Simulations conducted using `Gauss` with 10,000 iterations. The limit variance $V_{ols,T}$ (denoted by V in this table) is calculated by (56). The sizes of test based on the t -ratios using (6) are listed in the ‘size’ columns.


| $T = 2$ | | | | | | | | | |
|--------------------|-------|------------|--------------------------|--------|------------|--------------------------|--------|------------|-------|
| $\rho = 0 (V = 3)$ | | | $\rho = -0.5 (V = 1.75)$ | | | $\rho = -0.9 (V = 0.39)$ | | | |
| n | mean | nT_1 var | size | mean | nT_1 var | size | mean | nT_1 var | size |
| 50 | 0.000 | 3.189 | 0.077 | -0.499 | 1.857 | 0.076 | -0.900 | 0.401 | 0.071 |
| 100 | 0.002 | 3.081 | 0.063 | -0.500 | 1.822 | 0.063 | -0.900 | 0.387 | 0.058 |
| 200 | 0.000 | 3.031 | 0.054 | -0.500 | 1.734 | 0.054 | -0.900 | 0.392 | 0.051 |
| 400 | 0.000 | 3.032 | 0.053 | -0.499 | 1.779 | 0.054 | -0.900 | 0.387 | 0.055 |

| $\rho = 0.5 (V = 3.75)$ | | | $\rho = 0.9 (V = 3.99)$ | | | $\rho = 1 (V = 4)$ | | | |
|-------------------------|-------|------------|-------------------------|-------|------------|--------------------|-------|------------|-------|
| n | mean | nT_1 var | size | mean | nT_1 var | size | mean | nT_1 var | size |
| 50 | 0.498 | 3.968 | 0.075 | 0.901 | 4.049 | 0.067 | 1.001 | 4.066 | 0.068 |
| 100 | 0.500 | 3.808 | 0.064 | 0.900 | 4.041 | 0.061 | 1.001 | 3.980 | 0.058 |
| 200 | 0.501 | 3.823 | 0.057 | 0.899 | 3.962 | 0.055 | 1.000 | 4.047 | 0.057 |
| 400 | 0.500 | 3.846 | 0.054 | 0.900 | 3.930 | 0.052 | 1.001 | 4.098 | 0.058 |

| $T = 24$ | | | | | | | | | |
|------------------------|-------|------------|---------------------------|--------|------------|---------------------------|--------|------------|-------|
| $\rho = 0 (V = 2.043)$ | | | $\rho = -0.5 (V = 1.074)$ | | | $\rho = -0.9 (V = 0.313)$ | | | |
| n | mean | nT_1 var | size | mean | nT_1 var | size | mean | nT_1 var | size |
| 50 | 0.001 | 2.057 | 0.061 | -0.498 | 1.091 | 0.062 | -0.899 | 0.329 | 0.069 |
| 100 | 0.000 | 2.077 | 0.059 | -0.499 | 1.095 | 0.056 | -0.899 | 0.321 | 0.060 |
| 200 | 0.000 | 2.039 | 0.052 | -0.500 | 1.053 | 0.049 | -0.900 | 0.325 | 0.057 |
| 400 | 0.000 | 2.035 | 0.050 | -0.500 | 1.079 | 0.051 | -0.900 | 0.324 | 0.050 |

| $\rho = 0.5 (V = 2.987)$ | | | $\rho = 0.9 (V = 3.758)$ | | | $\rho = 1 (V = 4)$ | | | |
|--------------------------|-------|------------|--------------------------|-------|------------|--------------------|-------|------------|-------|
| n | mean | nT_1 var | size | mean | nT_1 var | size | mean | nT_1 var | size |
| 50 | 0.500 | 2.986 | 0.059 | 0.900 | 3.735 | 0.059 | 1.000 | 3.988 | 0.059 |
| 100 | 0.500 | 3.036 | 0.054 | 0.900 | 3.723 | 0.053 | 1.000 | 4.039 | 0.057 |
| 200 | 0.500 | 3.036 | 0.056 | 0.900 | 3.685 | 0.050 | 1.000 | 3.841 | 0.045 |
| 400 | 0.500 | 3.073 | 0.054 | 0.900 | 3.817 | 0.053 | 1.000 | 4.071 | 0.055 |

Figure 2: Variance ratio V_{gmm}/V_{ols} for normal errors for $T = 2, 3, \dots, 100$. The minimum efficiency of FDLs relative to FDGMM is approximately 0.99 with the low point being attained at $\rho = 0.5$ and $T = 4$. The efficiency gain of FDGMM over FDLs is marginal.



GMM over OLS is marginal. But note that this simulation result applies only to normally distributed errors. From additional experiments (not reported here) it was found that the efficiency of GMM over OLS is responsive to kurtosis, but for reasonable degrees of kurtosis, the efficiency gain of FDGMM remains marginal. For example, when $var((\varepsilon_i/\sigma)^2) - 2 = 5$, the minimal V_{gmm}/V_{ols} ratio is approximately 0.98.

Because the performance of the feasible two-step GMM estimator may deteriorate due to inaccurate estimation of the covariance matrix, the two-step efficient GMM may yield a poorer estimator than OLS when the efficiency gain of the infeasible optimal GMM is marginal. When ε_{it} is normally distributed, this is likely to be the case. According to simulations not reported here, the two-step efficient GMM (using OLS as the first step estimator) looks less efficient than OLS for a wide range of ρ and T values up to quite a large n . So we generally recommend FDLs over FDGMM for practical use.

It is interesting to view FDLs in the context of method of moments and compare it with other consistent estimators. For the model $y_{it} = \alpha_i + u_{it}$, $u_{it} = \rho u_{it-1} + \varepsilon_{it}$, under the further assumption that α_i is uncorrelated with ε_{it} (and u_{i0}), we get the moments

$$E y_{it} y_{is} = E \alpha_i^2 + \sigma^2 \rho^{|t-s|} / (1 - \rho^2), \quad |\rho| < 1,$$

$$E y_{it} y_{is} = E \alpha_i^2 + E u_{i0}^2 + \sigma^2 (s \wedge t), \quad \rho = 1,$$

for all $t, s = 0, 1, \dots, T$, which in turn provide $(T + 1)(T + 2)/2$ distinct moments. Next,

rewrite the moments in terms of $(y_{i0}, \Delta y'_i)' = (y_{i0}, \Delta y_{i1}, \dots, \Delta y_{iT})'$ as

$$\begin{aligned} Ey_{i0}^2 &= E\alpha_i^2 + \rho^2 Eu_{i0}^2 + \sigma^2, \\ Ey_{i0}\Delta y_{it} &= -\sigma^2 \rho^{t-1}/(1+\rho), \quad t \geq 1, \\ E(\Delta y_{it})^2 &= 2\sigma^2/(1+\rho), \\ E\Delta y_{it}\Delta y_{is} &= -\sigma^2 \rho^{|t-s|-1}(1-\rho)/(1+\rho), \quad t \neq s, \end{aligned}$$

or in matrix form as

$$(8) \quad E \begin{bmatrix} y_{i0} \\ \Delta y_i \end{bmatrix} \begin{bmatrix} y_{i0} \\ \Delta y_i \end{bmatrix}' = \frac{\sigma^2}{1+\rho} \begin{bmatrix} \xi_\rho & -1 & -\rho & \cdots & -\rho^{T_1} \\ -1 & 2 & -(1-\rho) & \cdots & -\rho^{T_2}(1-\rho) \\ -\rho & -(1-\rho) & 2 & \cdots & -\rho^{T_3}(1-\rho) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\rho^{T_1} & -\rho^{T_2}(1-\rho) & -\rho^{T_3}(1-\rho) & \cdots & 2 \end{bmatrix},$$

where $\xi_\rho = (E\alpha_i^2 + \rho^2 Eu_{i0}^2 + \sigma^2)(1+\rho)/\sigma^2$. (Note that the moments Ey_{i0}^2 and $E\Delta y_i y_{i0}$ depend on the condition that the α_i are uncorrelated with $u_{i,-1}$ and ε_{it} but the moments $E\Delta y_i \Delta y_i$ do not.) Rewriting the moments $Ey_i y'_i$ as (8) does not waste any information because we can recover the original moments $Ey_i y'_i$ by a linear transformation. Now, among these $(T+1)(T+2)/2$ distinct moments, Ey_{i0}^2 contains the nuisance parameters $E\alpha_i^2$ and $Eu_{i,-1}^2$ and, in fact, only Ey_{i0}^2 does so. Thus this element does not contribute to the estimation of ρ and can be safely ignored. Now, in what follows, we will show that each conventional (consistent) estimator can be derived as a (generalized) method of moments estimator using a subset of the above moment conditions.

Let us first consider the conventional IV/GMM estimators such as those of Anderson and Hsiao (1981) and Arellano and Bond (1991). The moment conditions

$$(9) \quad Ef_{i1t}(\rho) = Ey_{i0}(\Delta y_{it} - \rho \Delta y_{it-1}) = 0, \quad t \geq 2,$$

are obtained by combining any two consecutive elements of the first column (except for Ey_{i0}^2). Furthermore, any lower off-diagonal element of $E\Delta y_i \Delta y'_i$ and the element below it provides the moment condition

$$(10) \quad Eh_{ist}(\rho) = E\Delta y_{is}(\Delta y_{it} - \rho \Delta y_{it-1}) = 0, \quad s < t - 1,$$

for some s and t . These moment conditions are strong when $\rho < 1$ so the IV/GMM estimators are consistent. But if $\rho \simeq 1$, then the moment conditions are weakly identifying, and in case $\rho = 1$, each moment condition in (9) and (10) fails to identify the true parameter because then $Ef_{i1t}(\rho) \equiv 0$ and $Eh_{ist}(\rho) \equiv 0$. When $\rho = 1$, if T is small and fixed, then the limit distribution of the GMM estimator is nondegenerate due to the lack of identification

(see, e.g., Phillips, 1989, Staiger and Stock, 1997). But if T is large, then the unweighted GMM using these moment conditions converges to zero (under some regularity conditions for $u_{i0} = y_{i0} - \alpha_i$), because the unweighted criterion function satisfies the convergence

$$q_n^{-1} \left\{ \sum_{t=2}^T \left[n^{-1/2} \sum_{i=1}^n f_{i1t}(\rho) \right]^2 + \sum_{s < t-1} \left[n^{-1/2} \sum_{i=1}^n h_{ist}(\rho) \right]^2 \right\} \rightarrow_p (1 + \rho^2)\sigma^4,$$

(if the initial condition that $(nT)^{-1/2} \sum_{i=1}^n u_{i,-1} \rightarrow_p 0$ holds under other regularity conditions) due to the accumulating signal variability and despite the fact that each moment condition fails to identify any ρ value, and this limit is minimized at zero (see Han and Phillips (2006) for a detailed study of such situations), where $q_n = (T - 1) + T(T + 1)/2$ is the total number of moment conditions. So IV/GMM estimation based on (9) and (10) can hardly be used successfully when ρ may take values near unity. The behavior of the two step efficient GMM estimator in this case has not been determined.

It is remarkable that Arellano and Bond (1991)'s estimator does not make full use of all the moment conditions implied by the off-diagonal elements of (8). Taking the example of $T = 2$, besides $Ey_{i0}(\Delta y_{i2} - \rho \Delta y_{i1}) = 0$, the moment condition $E\Delta y_{i2}(y_{i0} - \rho y_{i1}) = 0$ also holds. (But it should also be noted that $Ey_{i0}(\Delta y_{i2} - \rho \Delta y_{i1}) = 0$ if the ε_{it} are not autocorrelated (over t) while the moment condition that $E\Delta y_{i2}(y_{i0} - \rho y_{i1}) = 0$ requires homoskedasticity over time also. See Ahn and Schmidt (1995) for the complete list of moment conditions implied by various set of assumptions.) These additional moment conditions certainly improve the quality of Arellano and Bond's estimator when $|\rho| < 1$, but do not help solve the problem at $\rho = 1$.

Unlike this IV/GMM estimator which uses the off-diagonal elements of (8), our approach works from the moment conditions on the diagonal elements of $E\Delta y_i \Delta y_i'$. Each diagonal element of $E\Delta y_i \Delta y_i'$ in (8) and the element right below it construct a moment condition in (4), i.e., $(1 - \rho)E(\Delta y_{it-1})^2 + 2E\Delta y_{it-1} \Delta y_{it} = 0$ or equivalently $E\Delta y_{it-1}[2\Delta y_{it} + (1 - \rho)\Delta y_{it-1}] = 0$. It is also interesting that more moment conditions can be obtained by combining the diagonal elements and their left elements, viz.,

$$(11) \quad E g_{it}^*(\rho) = E\Delta y_{it}[2\Delta y_{it-1} + (1 - \rho)\Delta y_{it}] = 0, \quad t = 2, \dots, T.$$

These moment conditions are symmetric to (4) and are obtained by swapping the roles of Δy_{it} and Δy_{it-1} . We may consider GMM estimation or OLS estimation using these additional moment conditions together with those in (4). Simulations illustrate that the efficiency gain over $\hat{\rho}_{ols}$ by considering (11) is considerable when $T = 2$ especially for negative ρ , but the contribution of these additional moment conditions diminishes with T . Furthermore, when $\rho = 1$, $g_{it}(\rho)$ of (4) and $g_{it}^*(\rho)$ of (11) are asymptotically identical when evaluated

Table 2: Efficiency gain by adding $g_{it}^*(\rho)$

| $\rho \setminus T$ | 2 | 3 | 4 | 5 | 7 | 10 | 20 |
|--------------------|------|------|------|------|------|------|------|
| -0.5 | 0.44 | 0.47 | 0.56 | 0.62 | 0.73 | 0.79 | 0.90 |
| 0.0 | 0.75 | 0.78 | 0.86 | 0.88 | 0.92 | 0.95 | 0.98 |
| 0.5 | 0.93 | 0.95 | 0.97 | 0.98 | 0.99 | 0.99 | 1.00 |
| 0.9 | 0.99 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| 1.0 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |

Variance ratios $\text{var}(\hat{\rho}_{ols}^{**})/\text{var}(\hat{\rho}_{ols})$ are reported.

at the true parameter, so the moment conditions are singular and the traditional feasible optimal GMM procedure does not provide standard asymptotics should we use both (4) and (11). In that case, procedures based on analytically calculated optimal weights (so the weights are a function of ρ) could avoid the singularity, but this sort of general analytic weighting scheme cannot be implemented because the optimal weighting matrix depends on the nuisance parameter $E\varepsilon_{it}^4$. When $\varepsilon_{it} \sim N(0, \sigma^2)$, however, the variances of $g_{it}(\rho)$ and $g_{it}^*(\rho)$ are almost equal except for $\rho \simeq -1$ according to simulations, implying that the unweighted sum $g_{it}(\rho) + g_{it}^*(\rho)$ is an (almost) optimal transformation of $g_{it}(\rho)$ and $g_{it}^*(\rho)$. Especially, when $\sum_{t=2}^T g_{it}(\rho)$ is used (as in the derivation of $\hat{\rho}_{ols}$) instead of each $g_{it}(\rho)$ separately, the unweighted sum $\sum_{t=2}^T [g_{it}(\rho) + g_{it}^*(\rho)]$ is an almost optimal exactly identifying moment condition with normal errors. This leads to a natural OLS regression of the pooled dependent variable $(2\Delta y_{it} + \Delta y_{it-1}, 2\Delta y_{it-1} + \Delta y_{it})'$ on the correspondingly pooled independent variable $(\Delta y_{it-1}, \Delta y_{it})'$. Let us denote this estimator by $\hat{\rho}_{ols}^{**}$. According to simulations, when $\rho = 0$ and $T = 2$, the variance ratio $\text{var}(\hat{\rho}_{ols}^{**})/\text{var}(\hat{\rho}_{ols})$ is about 0.75, meaning that by additionally using the moment condition that $Eg_{i3}^*(\rho) = 0$, we can reduce 25% of the variance. For other values of ρ and T , Table 2 reports the ratio of the variance of the OLS estimator based on $g_{it}(\rho)$ to the variance of the OLS estimator based on both $g_{it}(\rho)$ and $g_{it}^*(\rho)$. Note again that this result applies only to normal errors, and when the tails of the error distribution are thicker (e.g., the t_5 distribution), $g_{it}^*(\rho)$ will have larger variance than $g_{it}(\rho)$, and as a result, the estimator $\hat{\rho}_{ols}^{**}$ may be even less efficient than $\hat{\rho}_{ols}$ because of the failure of optimal weighting.

In summary, the loss of efficiency from using OLS rather than GMM is marginal, and the gain from adding $g_{it}^*(\rho)$ is not big enough compared with the possible risk of singularity (in case of GMM) and efficiency loss (in case of pooled OLS). In view of these many considerations, the original FDLS method (which yields $\hat{\rho}_{ols}$) is again recommended for practical use.

2.3 More Moment Conditions

We may want to use all the moment conditions f_{i1t} , h_{ist} , g_{it} and their mirror images. This is done in Ahn and Schmidt (1995). The GMM estimator based on all these moment conditions is asymptotically efficient when $|\rho| < 1$ and T^2 is small relative to n . However, this approach faces the many instrument problem when T is large and the weak instrument problem when $\rho \simeq 1$. Noting that some moment conditions are strong for all ρ , we may calculate the weighted sum of all the moment conditions using a nonrandom weighting function of ρ (and only ρ with no other nuisance parameters). Nickell's (1981) analysis allows for this kind of method, which is a method of moments estimator (let us call it the *fixed effects method of moments*, or FEMM for short, estimator) due to the fact that

$$\frac{\sum_{t=1}^T E\dot{y}_{it-1}(\ddot{y}_{it} - \rho\dot{y}_{it-1})}{\sum_{t=1}^T E\dot{y}_{it-1}^2} = d_T(\rho),$$

where $\ddot{y}_{it} = y_{it} - T^{-1} \sum_{t=1}^T y_{it}$, $\dot{y}_{it-1} = y_{it-1} - T^{-1} \sum_{t=1}^T y_{it-1}$, and

$$d_T(\rho) = - \left(\frac{1+\rho}{T_1} \right) \tilde{d}_T(\rho) / \left[1 - \frac{2\rho\tilde{d}_T(\rho)}{(1-\rho)T_1} \right],$$

with $\tilde{d}_T(\rho) = 1 - T^{-1}(1-\rho)^{-1}(1-\rho^T)$. (Here $T_1 = T - 1$ as before.) The above moment condition can be written as

$$(12) \quad E \sum_{t=1}^T \dot{y}_{it-1} \{ \ddot{y}_{it} - [\rho + d_T(\rho)] \dot{y}_{it-1} \} = 0.$$

In order to estimate ρ , we can set the sample moment to zero and use a nonlinear algorithm to locate the root. It is also possible to estimate $\rho + d_T(\rho)$ by linear fitting (which is the usual within-group estimator) and then correct the bias by applying the inverse mapping of $\rho \mapsto \rho + d_T(\rho)$, but we do not consider this method because estimating the variance of the resulting estimator is not straightforward in that case.

Now, because \ddot{y}_{it} and \dot{y}_{it-1} can be expressed in terms of Δy_{is} as

$$\ddot{y}_{it} = \frac{1}{T} \sum_{s=2}^T (s-1) \Delta y_{is} - \sum_{s=t+1}^T \Delta y_{is} \quad \text{and} \quad \dot{y}_{it} = \frac{1}{T} \sum_{s=1}^{T-1} s \Delta y_{is} - \sum_{s=t+1}^{T-1} \Delta y_{is},$$

the moment condition (12) can be expressed as a linear combination of the elements of $E\Delta y_i \Delta y_i'$ of (8).

When $T = 2$, the moment condition for FEMM is equal to the moment condition for $\hat{\rho}_{ols}$, and the FEMM estimator is algebraically equal to $\hat{\rho}_{ols}$. If $|\rho| < 1$ in this case, the GMM estimator based on (4) and (11) is more efficient than FEMM when the errors are normally

distributed as we have seen before. This example is interesting because it shows that the bias corrected within estimator (FEMM) may not be efficient for small T .

When $T = 3$, the moment condition for FEMM is ‘factorized’ as

$$[(1 - \rho)(3 + \rho)]^{-1} E\{(2 - 3\rho - \rho^2)g_{i2}(\rho) + 2g_{i2}^*(\rho) + (3 + \rho)(1 - \rho)[g_{i3}(\rho) + h_{i13}(\rho)]\} = 0,$$

where $g_{it}(\rho)$, $g_{it}^*(\rho)$ and $h_{ist}(\rho)$ are defined in (4), (11) and (10), respectively. The leading factor $[(1 - \rho)(3 + \rho)]^{-1}$ is ignored (we can do so because the probability of $\hat{\rho} = 1$ is zero) to handle the $\rho = 1$ case, and we have

$$(13) \quad E\{(2 - 3\rho - \rho^2)g_{i2}(\rho) + 2g_{i2}^*(\rho) + (3 + \rho)(1 - \rho)[g_{i3}(\rho) + h_{i13}(\rho)]\} = 0.$$

In (13) it is remarkable and important that $h_{i13}(\rho)$ is multiplied by the factor $1 - \rho$, which makes $h_{i13}(\rho)$ relevant when $\rho = 1$. That is, when the true parameter is unity, $Eh_{i13}(\rho) = 0$ for all ρ so $h_{i13}(\rho)$ is irrelevant, but because of the $1 - \rho$ factor, the sample moment function $n^{-1} \sum_{i=1}^n (1 - \rho)h_{i13}(\rho)$ involving the irrelevant $h_{i13}(\rho)$ is always set to zero at $\rho = 1$, which is the true parameter. This happens for all T , ensuring consistency of the FEMM estimator for all ρ . Interesting as it is, we do not further pursue this issue here.

2.4 The Explosive Case

When $\rho > 1$, the differences Δy_{it} continue to manifest explosive behavior and all the good properties of FDLS do not hold. The estimator is inconsistent and the limit depends on the n/T ratio and the initial status $u_{i,-1}$. The following lemma is indicative of what happens in this case.

Lemma 4 *If $\rho > 1$, then*

$$E \frac{1}{T_1} \sum_{t=2}^T (\Delta y_{it-1})^2 = (\rho + 1)^{-1} [2 + \nu(\rho, T)] \sigma^2,$$

$$E \frac{1}{T_1} \sum_{t=2}^T \Delta y_{it-1} \eta_{it} = \nu(\rho, T) \sigma^2,$$

where

$$\nu(\rho, T) = \left[\frac{\rho^2(\rho^{2T_1} - 1)\delta_\rho}{T_1(\rho + 1)} \right], \quad \delta_\rho = (\rho^2 - 1)Eu_{i,-1}^2/\sigma^2 + 1.$$

When T is fixed, this lemma implies, by the law of large numbers, that

$$\text{plim}_{n \rightarrow \infty} \hat{\rho}_{ols} = \rho + \frac{(\rho + 1)\nu(\rho, T)}{2 + \nu(\rho, T)}.$$

Thus, in the case where ρ is close to unity and $Eu_{i,-1}^2$ is small so that $\nu(\rho, T)$ is negligible, then the inconsistency of $\hat{\rho}_{ols}$ is also small. For a given ρ , the bias of $\hat{\rho}_{ols}$ is bigger when T is large than when T is small. If ρ is even closer to unity and such that $\sqrt{n}\nu(\rho, T) \rightarrow 0$, then the bias of $\sqrt{nT_1}(\hat{\rho}_{ols} - \rho)$ is negligible, leading to a Gaussian limit whose variance changes continuously as ρ deviates slightly from unity into the explosive area. This observation implies that the limit distribution of Theorem 2 is continuous as ρ passes through unity to very mildly explosive cases.

The case with large T and fixed n can be analyzed by Theorems 2 and 3 of Phillips and Han (2005) who consider the time series case. Again the asymptotics are continuous as ρ passes through unity under the initial condition that $(\rho_T - 1)u_{i,-1}^2 \rightarrow_p 0$ where $\rho = \rho_T \downarrow 1$.

If both n and T are large, the $N(0, 2(1 + \rho))$ asymptotics are still continuous as ρ passes through the boundary of unity into the explosive area, though the boundary of ρ for the continuous asymptotics is then correspondingly narrower on the explosive side of unity.

2.5 Heterogeneity and Cross Section Dependence

The remainder of this section considers issues of cross-sectional heteroskedasticity, heterogeneity, and cross section dependence.

If the error variance $E\varepsilon_{it}^2$ is different across i , then Theorem 2 still applies as long as the Lindeberg condition holds, with the only modification to $V_{ols,T}$ being the more general expression

$$\lim_{n \rightarrow \infty} \frac{(nT_1)^{-1} \sum_{i=1}^n E(\sum_{t=2}^T \Delta y_{it-1} \eta_{it})^2}{(nT_1)^{-1} \sum_{i=1}^n \sum_{t=2}^T (\Delta y_{it-1})^2}.$$

Computation of standard errors by (6) remains valid.

If the AR coefficient changes across i so that the model involves $u_{it} = \rho_i u_{it-1} + \varepsilon_{it}$, then we have the limit

$$\hat{\rho}_{ols} \rightarrow_p \text{plim}_{n \rightarrow \infty} \sum_{i=1}^n w_i \rho_i,$$

if the limit on the right hand side exists, where

$$w_i = \frac{\sum_{t=2}^T (\Delta y_{it-1})^2}{\sum_{i=1}^n \sum_{t=2}^T (\Delta y_{it-1})^2}.$$

Noting that $E(\Delta y_{it-1})^2 = 2\sigma_i^2/(1 + \rho_i)$, we see that an individual unit with smaller ρ_i is given a bigger weight if there is no heteroskedasticity.

To allow for cross section dependence, let $\varepsilon_{it} = \sum_{k=1}^K \lambda_{ik} f_{kt} + v_{it}$, where the f_{kt} are common shocks *iid* over t and independent of other shocks, λ_{ik} are the factor loadings, and

the v_{it} are *iid*. Since the compounded error ε_{it} is still white noise, the moment conditions (4) still hold. However, consistency does not hold unless $T \rightarrow \infty$ or $K \rightarrow \infty$ (where K is the number of common factors). The case $K = 1$ is exemplary. Here $(\Delta y_{it-1})^2$ involves $(\lambda_{i1} f_{1t})^2$ and the randomness in the double sum

$$(nT_1)^{-1} \sum_{i=1}^n \sum_{t=2}^T (\lambda_{i1} f_{1t})^2 = n^{-1} \sum_{i=1}^n \lambda_{i1}^2 \cdot T_1^{-1} \sum_{t=2}^T f_{1t}^2$$

persists unless T is large (c.f. Phillips and Sul, 2004). When T is small, as is the case in typical microeconomic work, K may be considered large and each factor loading is assumed to be sufficiently small to ensure convergence. (Then the variation of the cross section dependence term $\sum_{k=1}^K \lambda_{ik} f_{it}$ is correspondingly controlled.) As long as the common factors f_{kt} are white noise over time, the moment conditions (4) hold, and under further regularity to ensure convergence, the FDLS estimator would be consistent. See Phillips and Sul (2004) for details on panel models with these characteristics.

3 Dynamic Panels with Exogenous Variables

This section considers the model with exogenous variables $y_{it} = \alpha_i + \beta' x_{it} + u_{it}$ where $u_{it} = \rho u_{it-1} + \varepsilon_{it}$ for $\rho \in (-1, 1]$, which may be transformed to

$$(14) \quad y_{it} = (1 - \rho)\alpha_i + \beta'(x_{it} - \rho x_{it-1}) + \rho y_{it-1} + \varepsilon_{it}.$$

Let $z_{it} = y_{it} - \beta' x_{it}$. Then the model (14) is written as $z_{it} = (1 - \rho)\alpha_i + \rho z_{it-1} + \varepsilon_{it}$, which is the same as (1) with y_{it} replaced with z_{it} . By applying the same transformation, we get

$$2\Delta z_{it} + \Delta z_{it-1} = \rho \Delta z_{it-1} + \eta_{it}, \quad \eta_{it} = 2\Delta \varepsilon_{it} + (1 + \rho)\Delta z_{it-1}.$$

For all ρ , Δz_{it-1} and η_{it} are uncorrelated and these moment conditions are strong for all ρ , as before. Next, if we allow α_i to be arbitrarily correlated with x_{it} , then we can apply the within transformation to (14) giving

$$\ddot{y}_{it} - \rho \dot{y}_{it-1} = (\ddot{x}_{it} - \rho \dot{x}_{it-1})' \beta + \ddot{\varepsilon}_{it},$$

where $\ddot{y}_{it} = y_{it} - T^{-1} \sum_{s=1}^T y_{is}$, $\dot{y}_{it-1} = y_{it-1} - T^{-1} \sum_{s=1}^T y_{is-1}$, and so on. From the fact that the within-group estimator is efficient when α_i is allowed to be correlated with x_{it} in the usual linear panel data model (see Im, Ahn, Schmidt and Wooldridge, 1999), we propose the following exactly identifying moment conditions:

$$(15) \quad E \sum_{t=2}^T \Delta z_{it-1} [(2\Delta z_{it} + \Delta z_{it-1}) - \rho \Delta z_{it-1}] = 0,$$

$$(16) \quad E \sum_{t=1}^T (\ddot{x}_{it} - \rho \dot{x}_{it-1}) [(\ddot{y}_{it} - \rho \dot{y}_{it-1}) - (\ddot{x}_{it} - \rho \dot{x}_{it-1})' \beta] = 0,$$

where $\Delta z_{it} = \Delta y_{it} - \Delta x'_{it}\beta$. Note that when there is no exogenous variable, the first moment condition (15) leads to the OLS estimator of the previous section. We can estimate ρ and β by the method of moments. In practice, after (15) and (16) are rewritten in terms of the sample moment conditions, the parameters can be estimated by estimating ρ and β iteratively between (15) and (16) if the procedure converges.

For the asymptotic variance we can use the usual “ $(D'\Omega^{-1}D)^{-1}$ ” formula where D contains the expected scores of (15) and (16) and Ω is the variance matrix of those moment conditions, both evaluated at the true parameter. Conveniently, D is block diagonal and if ε_{it} has zero third moment, then so is the Ω matrix, separating the estimation of ρ from the estimation of β . (See the Appendix for further details.) As a result, we can treat the final estimator $\hat{\beta}$ as the true β parameter in computing the Δz_{it} 's and then estimate ρ and compute its standard error; similarly, we can treat the final $\hat{\rho}$ as the true ρ parameter and compute the standard errors of $\hat{\beta}$ using usual within-group estimation technique after transforming x_{it} and y_{it} to $x_{it} - \hat{\rho}x_{it-1}$ and $y_{it} - \hat{\rho}y_{it-1}$, respectively.

4 Incidental Trends

When the model includes incidental trends so that $y_{it} = \alpha_i + \gamma_i t + u_{it}$, $u_{it} = \rho u_{it-1} + \varepsilon_{it}$, it may be written in the form

$$(17) \quad y_{it} = (1 - \rho)\alpha_i + \rho\gamma_i + (1 - \rho)\gamma_i t + \rho y_{it-1} + \varepsilon_{it}.$$

Double differencing eliminates the combined fixed effects, giving

$$(18) \quad \Delta^2 y_{it} = \rho \Delta^2 y_{it-1} + \Delta^2 \varepsilon_{it},$$

which implies that

$$\Delta^2 y_{it} = \sum_{j=0}^{\infty} \rho^j \Delta^2 \varepsilon_{it-j} = \varepsilon_{it} - (2 - \rho)\varepsilon_{it-1} + (1 - \rho)^2 \sum_{j=0}^{\infty} \rho^j \varepsilon_{it-j-2}.$$

When $\rho = 1$, we have $\Delta^2 y_{it} = \Delta \varepsilon_{it}$, so $E(\Delta^2 y_{it-1})^2 = 2\sigma^2$ and $E\Delta^2 y_{it-1} \Delta^2 \varepsilon_{it} = -3\sigma^2$. When $|\rho| < 1$, we have

$$E(\Delta^2 y_{it-1})^2 = \left[1 + (2 - \rho)^2 + \frac{(1 - \rho)^4}{1 - \rho^2} \right] \sigma^2 = \frac{2(3 - \rho)\sigma^2}{1 + \rho},$$

and

$$E\Delta^2 y_{it-1} \Delta^2 \varepsilon_{it} = -(4 - \rho)\sigma^2$$

so these formulae cover the case of $\rho = 1$. Thus

$$E\Delta^2 y_{it-1} \tilde{\eta}_{it} = 0, \quad \tilde{\eta}_{it} = 2\Delta^2 \varepsilon_{it} + \frac{(1+\rho)(4-\rho)}{3-\rho} \Delta^2 y_{it-1}.$$

This corresponds to transforming (18) to the model in differences

$$(19) \quad 2\Delta^2 y_{it} + \Delta^2 y_{it-1} = \theta \Delta^2 y_{it-1} + \tilde{\eta}_{it}, \quad \theta = -\frac{(1-\rho)^2}{3-\rho}.$$

Correspondingly, the *double difference least squares* (DDLs) estimator $\hat{\theta}$ is

$$\hat{\theta} = \frac{\sum_{i=1}^n \sum_{t=3}^T \Delta^2 y_{it-1} (2\Delta^2 y_{it} + \Delta^2 y_{it-1})}{\sum_{i=1}^n \sum_{t=3}^T (\Delta^2 y_{it-1})^2}.$$

Note that $\theta \in (-1, 0]$ for all $\rho \in (-1, 1]$ and $\theta = 0$ if $\rho = 1$. When point estimation of ρ is of interest, we can simply run pooled OLS on (19) to get $\hat{\theta}$ and censor at 0 and -1 , and then recover $\hat{\rho}$ from $\hat{\theta}$ by $\hat{\rho} = \frac{1}{2}[2 + \hat{\theta} - (\hat{\theta}^2 - 8\hat{\theta})^{1/2}]$. Alternatively, we may also consider method of moments estimation based on (19) using the moment condition

$$E \sum_{t=3}^T \Delta^2 y_{it-1} \left(2\Delta^2 y_{it} + \left[1 + \frac{(1-\rho)^2}{3-\rho} \right] \Delta y_{it-1} \right) = 0.$$

Some caution is needed here because the parameter $\theta = -(1-\rho)^2/(3-\rho)$ can never exceed zero for all $\rho \in (-1, 1]$ and therefore the sample moment function may not attain zero at any parameter value.

For testing, we can rely on the asymptotic distribution

$$\sqrt{nT_2}(\hat{\theta} - \theta) \Rightarrow N(0, W_{ols,T})$$

for some positive $W_{ols,T}$. (Asymptotic normality is established in Theorems 5 and 6 in the Appendix.) Just as for the case without incidental trends, the limit theory is continuous and joint as $n \rightarrow \infty$ or $T \rightarrow \infty$ or both pass to infinity with $W_{ols,T}$ evolving with T . However, as is apparent from the relation $\theta = -\frac{(1-\rho)^2}{3-\rho}$, an $O(n^{-1/4}T^{-1/4})$ neighborhood of $\rho = 1$ corresponds to an $O(n^{-1/2}T^{-1/2})$ neighborhood of $\theta = 0$, and for $\rho = 1$, it is easily seen that the rate of convergence of $\hat{\rho}$ is at the slower $n^{1/4}T_2^{1/4}$ rate, while that of $\hat{\theta}$ is $n^{1/2}T_2^{1/2}$. For $\rho < 1$, the rates of convergence of $\hat{\rho}$ and $\hat{\theta}$ are both $n^{1/2}T_2^{1/2}$. Hence, there is a deficiency in the convergence rate for $\hat{\rho}$ around unity.

When T is small, $W_{ols,T}$ depends on $E\varepsilon_{it}^4$ and the algebraic form of $W_{ols,T}$ for small T is too complicated to be of interest. But when n is large, as in typical microeconomic projects, the standard error of $\hat{\theta}$ is easily calculated by

$$(20) \quad se(\hat{\theta}) = \frac{\sqrt{\sum_{i=1}^n (\sum_{t=3}^T \Delta^2 y_{it-1} \hat{\eta}_{it})^2}}{\sum_{i=1}^n \sum_{t=3}^T (\Delta^2 y_{it-1})^2},$$

with $\hat{\eta}_{it}$ denoting the residuals from the regression of (19). On the other hand, in macroeconometrics where T is often moderately large, then $\sqrt{nT_2}(\hat{\theta} - \theta) \Rightarrow N(0, \lim_{T \rightarrow \infty} W_{ols,T})$, where

$$\lim_{T \rightarrow \infty} W_{ols,T} = (1 + \rho)^2 (3 - \rho)^{-2} \sum_1^k b_k^2,$$

with

$$\begin{aligned} b_1 &= 2(3 - \rho) + (1 - \rho)^2 - [(2 - \rho) + 2(1 - \rho)^2/(1 + \rho)]\phi, \\ b_2 &= -(2 - \rho)[1 + (1 - \rho)^2] + (1 - \rho)^3\phi/(1 + \rho), \\ b_k &= \rho^{k-3}(1 - \rho)^3[(1 - \rho) + \rho\phi/(1 + \rho)], \quad k \geq 3, \end{aligned}$$

and with $\phi = (4 - \rho)(1 + \rho)/(3 - \rho)$. (Phillips and Han, 2005, derive this limit in the time series case.) As presented in Theorems 5 and 6 in the Appendix, the convergence is uniform in the sense that the limit of the variance is the variance of the limit distribution as $T \rightarrow \infty$.

As a special case, if $\rho = 1$, then

$$(21) \quad \sqrt{nT_2}\hat{\theta} \Rightarrow N(0, 2 + \kappa_4/T_2)$$

for all T as $n \rightarrow \infty$, where $\kappa_4 = \text{var}(\varepsilon_{it}^2)/2\sigma^4$, and the limit distribution as $T \rightarrow \infty$ is simply $N(0, 2)$ whether n is large or small.

5 Panel Unit Root Testing

5.1 Fixed Effects Model

The inferential apparatus and its limit theory may be applied directly to panel unit root testing. Consider the fixed effects panel $y_{it} = (1 - \rho)\alpha_i + \rho_i y_{it-1} + \varepsilon_{it}$ where the ε_{it} are *iid*. The unit root null hypothesis is that $\rho_i = 1$ for all i . The OLS estimator $\hat{\rho}_{ols}$ and its limit theory in Theorem 2 and Corollary 3 can form the basis of a statistical test. More precisely, the test statistic is $\hat{\tau}_0 := (nT_1)^{1/2}(\hat{\rho}_{ols} - 1)/2$. This test statistic is derived under the assumption of cross-sectional homoskedasticity. If the variances $\sigma_i^2 := E\varepsilon_{it}^2$ differ across i and $\rho_i \equiv 1$, then the standard deviation of the limit distribution of $\hat{\rho}_{ols}$ is approximated by

$$(22) \quad \frac{2(\sum_{i=1}^n \sigma_i^4)^{1/2}}{T_1^{1/2} \sum_{i=1}^n \sigma_i^2} = \frac{2}{(nT_1)^{1/2}} \times \frac{(n^{-1} \sum_{i=1}^n \sigma_i^4)^{1/2}}{n^{-1} \sum_{i=1}^n \sigma_i^2},$$

which is larger than the simple standard error form $2/(nT_1)^{1/2}$ obtained for homoskedastic data. Under heteroskedasticity, the $\hat{\tau}_0$ statistic can be computed by using formula (6) for the

standard error, or by replacing σ_i^2 in (22) with the estimate $\hat{\sigma}_i^2 = T^{-1} \sum_{t=1}^T (\Delta y_{it})^2$. Under the null hypothesis, $\hat{\tau}_0 \Rightarrow N(0, 1)$, and under the alternative hypothesis that $\rho_i < 1$ for some i , $\text{plim } \hat{\rho}_{ols} < 1$, and as a result, $\hat{\tau}_0 \rightarrow_p -\infty$ as $nT \rightarrow \infty$. So, the test is consistent for any passage of $nT \rightarrow \infty$.

Unlike Levin and Lin (1992) or Im, Pesaran and Shin (1997), this test does not require any restriction on the path to infinity such as $T \rightarrow \infty$ and $n/T \rightarrow 0$. Only the composite divergence $nT \rightarrow \infty$ is required, and there is virtually no size distortion when T is small. On the other hand, while the point optimal test for a unit root has local power in a neighborhood of unity that shrinks at the rate $n^{-1/2}T^{-1}$ (see Moon and Perron, 2004, and Moon, Perron and Phillips, 2006b), the $\hat{\tau}_0$ test has only trivial asymptotic power in $n^{-1/2}T^{-1}$ neighborhoods and non-trivial local asymptotic power in $n^{-1/2}T^{-1/2}$ neighborhoods of unity. So in this model at least, there is an infinite power deficiency in neighborhoods of order $n^{-1/2}T^{-1}$.

This deficiency is partly due to the fact that $\hat{\rho}_{ols}$ depends only on differenced data, which thereby reduces the signal relative to that of point optimal and other tests which make direct use of the nonstationary regressor. But that is not the only reason, and interestingly a common point optimal test based on the differenced data may attain the optimal rate of $n^{1/2}T$. To illustrate this possibility, consider the common point likelihood ratio test for the case where the ε_{it} are *iid* $N(0, \sigma^2)$ and σ^2 is known. The test for $H_0 : n^{1/2}T(1 - \rho) = 0$ against the alternative $H_1 : n^{1/2}T(1 - \rho) = c$ using the differenced data Δy_{it} is based on the likelihood ratio

$$(23) \quad U_{nT}(c) = 2 \left[\log L(1 - n^{-1/2}T^{-1}c) - \log L(1) \right],$$

where $\log L(\cdot)$ is calculated based on the joint distribution of $\Delta y_i = (\Delta y_{i1}, \dots, \Delta y_{iT})'$, i.e.,

$$\log L(\rho) = -\frac{nT}{2} \log(2\pi) - \frac{nT}{2} \log \sigma^2 - \frac{n}{2} \log |\Omega_T(\rho)| - \frac{1}{2\sigma^2} \sum_{i=1}^n \Delta y_i' \Omega_T(\rho)^{-1} \Delta y_i,$$

with

$$(24) \quad \Omega_T(\rho) = (1 + \rho)^{-1} \text{Toeplitz} \{2, -1(1 - \rho), -\rho(1 - \rho), \dots, -\rho^{T-2}(1 - \rho)\}.$$

In the above, the notation $\text{Toeplitz}\{a_0, \dots, a_{m-1}\}$ denotes the $m \times m$ symmetric Toeplitz matrix whose (i, j) element is $a_{|i-j|}$. As shown in the appendix, when $n^{1/2}T(1 - \rho) = c$ (i.e., under the alternative hypothesis), we have

$$(25) \quad U_{nT}(c) \rightarrow_d N(c^2/8, c^2/2).$$

So if the null hypothesis of $n^{1/2}T(1 - \rho) = 0$ is tested against the alternative that $n^{1/2}T(1 - \rho) = \tilde{c}$ such that the null hypothesis is rejected for $U_{nT}(\tilde{c}) \geq \bar{z}_\alpha c / \sqrt{2}$ where $\Phi(-\bar{z}_\alpha) = \alpha$,

and if the \tilde{c} happens to equal the true c , then the local power of this test is

$$\Phi\left(\frac{c}{4\sqrt{2}} - \bar{z}_\alpha\right).$$

For all $c > 0$, this local power function resides below the power envelope $\Phi(c/\sqrt{2} - \bar{z}_\alpha)$ based on the level data for large T obtained by Moon, Perron and Phillips (2006b). Nonetheless, it is remarkable that the optimal rate can be attained by using differenced data.

As discussed so far, the deficiency of testing using $\hat{\rho}_{ols}$ comes partly from differencing and partly from inefficient use of the moment conditions. This fact may at first seem to contradict our earlier observation that $\hat{\rho}_{ols}$ is almost as good as the (infeasible) optimal GMM estimator based on the differenced data (e.g., Figure 2). Nonetheless, it seems that maximum likelihood estimation on the differenced model combines the moment conditions so cleverly that, at $\rho = 1$ (at which point the levels MLE is superconsistent), the otherwise useless moment conditions contribute to estimating $\rho = 1$. (See the last part of section 2.3.) A full analysis and comparison of panel MLE in levels and differences that explores this issue will be useful and interesting, and deserves a separate research paper.

To return to testing based on the FDLS estimator, the deficiency in power based on this procedure may be interpreted as a cost arising from the simplicity of the $\hat{\tau}_0$ test, the uniform convergence rate of the estimator and its robustness to the asymptotic expansion path for (n, T) . Table 3 reports the simulated size and power of the $\hat{\tau}_0$ test, in comparison to Im et al.'s (2003) test, for the data generating process $y_{it} = (1 - \rho_i)\alpha_i + \rho_i y_{it-1} + \sigma_i \varepsilon_{it}$ with alternative parameter settings, where α_i and ε_i are standard normal and $\sigma_i \sim U(0.5, 1.5)$. To simulate power, the cases $\rho_i \equiv 0.9$ and $\rho_i \sim U(0.9, 1)$ are considered. Panel length is chosen to be $T = 6$ and $T = 25$, choices that roughly illustrate size and power for small and moderate T . ($T = 6$ is the smallest value covered by Table 1 of Im et al., 2003.) Note that the $\hat{\tau}_0$ test does not require bias adjustment. It is also remarkable that the $\hat{\tau}_0$ test seems to have better power than the IPS test when T is small. But with larger T ($T = 25$ in the simulation), the IPS test has better power, which is related to the $O(\sqrt{nT})$ convergence rate of the FDLS estimator.

5.2 Incidental Trends Model

Next consider the case where incidental trends are present, as laid out in Section 4. Let $\hat{\theta}$ be the pooled OLS estimator from the regression of (19). Noting that $\rho = 1$ corresponds to $\theta = 0$, we can base the panel unit root test on the statistic $\hat{\tau}_1 := \hat{\theta}/se(\hat{\theta}) \Rightarrow N(0, 1)$, where $se(\hat{\theta})$ is given in (20) when n is large or $se(\hat{\theta}) = \sqrt{2/nT_2}$ when T is large. (Again note that (20) is robust to the presence of cross-sectional heteroskedasticity.) The null hypothesis $H_0 : \rho_i = 1$ for all i is rejected if $\hat{\tau}_1$ is less than the left-tailed critical value from

Table 3: Simulated Size and Power of Unit Root Tests with Incidental Intercepts case.

DGP: $y_{it} = \alpha_i + u_{it}$, $u_{it} = \rho_i u_{it-1} + \sigma_i \varepsilon_{it}$
 $\alpha_i, \varepsilon_{it} \sim iid N(0, 1)$, $\sigma_i \sim iid U(0.5, 1.5)$
 Significance level: 5%

$T = 6$

| n | $\rho_i \equiv 1$ | | $\rho_i \equiv 0.9$ | | $\rho_i \sim U(0.9, 1)$ | |
|-----|-------------------|------|---------------------|-------|-------------------------|-------|
| | HP | IPS | HP | IPS | HP | IPS |
| 50 | 6.09 | 5.83 | 20.13 | 10.76 | 12.13 | 7.95 |
| 100 | 6.00 | 5.47 | 28.37 | 14.77 | 13.37 | 9.20 |
| 200 | 5.30 | 5.56 | 42.88 | 22.05 | 18.49 | 11.52 |
| 400 | 5.54 | 5.97 | 66.30 | 35.02 | 26.77 | 15.63 |

$T = 25$

| n | $\rho_i \equiv 1$ | | $\rho_i \equiv 0.9$ | | $\rho_i \sim U(0.9, 1)$ | |
|-----|-------------------|------|---------------------|-------|-------------------------|-------|
| | HP | IPS | HP | IPS | HP | IPS |
| 50 | 6.17 | 5.14 | 49.56 | 84.26 | 22.66 | 33.61 |
| 100 | 5.89 | 5.34 | 72.64 | 98.63 | 30.01 | 54.62 |
| 200 | 5.05 | 5.31 | 93.00 | 99.99 | 47.09 | 81.03 |
| 400 | 4.91 | 5.70 | 99.73 | 100.0 | 71.05 | 97.83 |

the standard normal distribution. When some ρ_i are smaller than unity, or equivalently when some θ_i are smaller than 0, $\hat{\theta}$ converges in probability to the limit of $\sum_{i=1}^n \tilde{w}_i \theta_i$ where $\tilde{w}_i = \sum_{t=3}^T (\Delta^2 y_{it-1})^2 / \sum_{i=1}^n \sum_{t=3}^T (\Delta^2 y_{it-1})^2$. As long as a non-negligible portion of individual units i have $\rho_i < 1$, this limit is strictly negative, and hence the test statistic $\hat{\tau}_1$ diverges to $-\infty$ in probability. Thus, the test $\hat{\tau}_1$ is consistent regardless of the existence of incidental trends.

Local power of the test is relatively weak, which can be explained by the definition of θ in (19). Since $\hat{\theta}$ has an $(nT_2)^{1/2}$ rate of convergence the test should have some local power when θ is in an $O(n^{-1/2}T_2^{-1/2})$ neighborhood of zero. But that is the case when ρ is in an $O(n^{-1/4}T_2^{-1/4})$ neighborhood of unity, which shrinks at a far slower rate than the optimal $n^{-1/4}T^{-1}$ rate attained by a point optimal test when the incidental trends are extracted from the panel data as described in Moon, Perron and Phillips (2006b).

It is interesting that there may exist a unit root test based on the double differenced data which has local power in a neighborhood of unity shrinking at the $n^{-1/2}T_1^{-1/2}$ rate when $T_1/\sqrt{n} \rightarrow 0$ and at a correspondingly faster rate when T grows faster. This possibility can again be illustrated by considering a likelihood ratio test with double differenced data under the Gaussianity assumption such that

$$(26) \quad \Delta^2 y_i = (\Delta^2 y_{i2}, \dots, \Delta^2 y_{iT})' \sim N(0, \sigma^2 \tilde{\Omega}_T(\rho)), \quad \tilde{\Omega}_T(\rho) = \text{Toeplitz}(\omega_0(\rho), \dots, \omega_{T_2}(\rho)),$$

where

$$\begin{aligned}\omega_0(\rho) &= 2(3 - \rho)/(1 + \rho), \\ \omega_1(\rho) &= -(4 - 3\rho + \rho^2)/(1 + \rho), \\ \omega_j(\rho) &= \rho^{j-2}(1 - \rho)^3/(1 + \rho), \quad j = 2, 3, \dots, T_2.\end{aligned}$$

(The ω_j 's are straightforwardly calculated from (58) in the appendix.) The associated log-likelihood function is

$$(27) \quad \log \tilde{L}(\rho, \sigma^2) = -\frac{nT_1}{2} \log(2\pi) - \frac{nT_1}{2} \log(\sigma^2) - \frac{n}{2} \log |\tilde{\Omega}_T(\rho)| \\ - \frac{1}{2\sigma^2} \sum_{i=1}^n (\Delta^2 y_i)' \tilde{\Omega}_T(\rho)^{-1} (\Delta^2 y_i).$$

If σ^2 is known, then the common point optimal test for $H_0 : n^{-1/2}T_1^{-1/2}(1 - \rho) = 0$ against $H_1 : n^{-1/2}T_1^{-1/2}(1 - \rho) = c$ is based on

$$(28) \quad \tilde{U}_{nT}(c) = 2 \left[\log \tilde{L}(1 - n^{-1/2}T_1^{-1/2}c, \sigma^2) - \log \tilde{L}(1, \sigma^2) \right].$$

Let the true ρ be $\rho = \rho_{nT} = 1 - (nT_1)^{-1/2}c$ for some $c \in (0, 1]$, so the alternative hypothesis of the likelihood ratio test coincides with the data generating process. Then, as derived in the appendix,

$$(29) \quad \tilde{U}_{nT}(c) \rightarrow_d N(c^2/2, 2c^2), \text{ if } n^{-1/2}T_1 \rightarrow 0.$$

Note that in the above asymptotics $\rho_{nT} = 1 - (nT_1)^{-1/2}c$ is the true parameter used in generating $\Delta^2 y_{it}$. So if the null hypothesis that $(nT_1)^{1/2}(1 - \rho) = 0$ is tested against the alternative that $(nT_1)^{1/2}(1 - \rho) = c$, in such a way that the null hypothesis is rejected when $U_{nT}(\rho_{nT}) \geq \sqrt{2}c\bar{z}_\alpha$ (with \bar{z}_α again denoting the $100\alpha\%$ critical value for the standard normal distribution), then the size α asymptotic local power is

$$\Phi \left(\frac{c}{2\sqrt{2}} - \bar{z}_\alpha \right).$$

This finding is potentially important because it reveals the possibility that an optimal test based on double differenced data (which would have non-trivial local power in an $n^{-1/2}T_1^{-1/2}$ neighborhood of unity) would outperform the point optimal test (which is known to have local power in a neighborhood shrinking at the $n^{-1/4}T^{-1}$ rate) when the panel is wide and short. In effect, if this conjecture is true, then the asymptotic power envelope in panel unit root tests will depend on the manner in which the sample size parameters pass to infinity.

Unfortunately this possibility is not realized in the case of the $\hat{\tau}_1$ test, and considering its local power properties it may be natural to conclude that this test is less useful. Nonetheless,

its straightforward and general Gaussian asymptotics and accurate size properties make it at least an *ex tempore* method for simple diagnostic purposes, especially for the case where n is large compared to T .

Table 4 reports simulation results for $\hat{\tau}_1$ applied to the data generating process with incidental trends for small T . Tests that require large T for valid size (e.g., the Ploberger and Phillips, 2002, test in the simulation) look biased and certainly perform poorly with small T , but are considerably more powerful with large T . (Simulation results with large T are not reported here.) On the other hand, Breitung's (2000) unbiased (UB) test, which is based on

$$(30) \quad (nT_2)^{-1/2} \sum_{i=1}^n \sum_{t=3}^T x_{it}^* y_{it}^*, \quad x_{it}^* = \lambda'_{1t} \Delta^2 y_i, \quad y_{it}^* = \lambda'_{2t} \Delta^2 y_i$$

for specially chosen λ_{1t} and λ_{2t} such that $E x_{it}^* y_{it}^* = 0$ (see Breitung, 2000; the proof of the validity of this expression is available upon request), performs well with small T with better local power than $\hat{\tau}_1$. The greater power of the UB test can be ascribed to several causes. First, the UB test is based on the special choice of λ_{1t} and λ_{2t} , which is more efficient at unity than our approach. Naturally, the power gain from this source comes at the cost that the test is available only for the null hypothesis of unity. For other null hypotheses (e.g., $H_0 : \rho = 0$), the UB test cannot be used, though some modification of the test might be possible, of course, in this case. Another relevant explanation is that the UB test statistic estimates the variance of (30) in a more efficient, but somewhat unintuitive way, which is valid only when the errors are cross-sectionally homoskedastic and no skewness and extra kurtosis are present.

In the simulations, the UB test and the PP (Ploberger and Phillips, 2002) are computed with σ_i^2 known. To correspondingly tweak the performance of the $\hat{\tau}_1$ test and effect a fairer comparison with the UB test, a variant of the $\hat{\tau}_1$ test (denoted as HP* in Table 4) is introduced, which is

$$\hat{\tau}_1^* = \frac{\sum_{i=1}^n \sum_{t=3}^T \sigma_i^{-2} \Delta^2 y_{it-1} (2\Delta^2 y_{it} + \Delta^2 y_{it-1})}{\sqrt{[(8 + 4/T_2)/6] \sum_{i=1}^n \sum_{t=3}^T \sigma_i^{-2} (2\Delta^2 y_{it} + \Delta^2 y_{it-1})^2}}.$$

This is asymptotically standard normal under the null of unity if $E \varepsilon_{it}^3 = 0$ and $E \varepsilon_{it}^4 = 3\sigma_i^4$.

The case $T = 3$ is particularly useful in illustrating the comparison of the UB and $\hat{\tau}_1^*$ tests. In this case, $x_{i3}^* = (2\Delta^2 y_{i2} + \Delta^2 y_{i3})$ and $y_{i3}^* = \Delta^2 y_{i3}$ for the UB test, so the moment condition the UB test is based on is

$$E \Delta^2 y_{i3} (2\Delta^2 y_{i2} + \Delta^2 y_{i3}) = 0 \quad \text{if } \rho = 1,$$

which is the ‘mirror image’ (obtained by swapping the roles of $\Delta^2 y_{i2}$ and $\Delta^2 y_{i3}$) of our moment condition

$$E\Delta^2 y_{i2}(2\Delta^2 y_{i3} + \Delta^2 y_{i2}) = 0 \text{ if } \rho = 1.$$

So the UB test and the $\hat{\tau}_1^*$ test (HP*) should manifest similar power performance, which indeed proves to be so in simulations. But when $T > 3$ (e.g., $T = 5$ in the simulation) the power of the UB test exceeds that of $\hat{\tau}_1$ and $\hat{\tau}_1^*$, which can be attributed to the first cause mentioned above.

It is also worth noting that both the UB test and the $\hat{\tau}_1$ test have trivial local power in the neighborhood of unity shrinking at the \sqrt{n} rate for fixed T . This is related with the fact that the mean function of the ‘numerators’ of the tests have zero slope at unity. Interestingly, the LM test statistic based on the normal distribution $\Delta^2 y_i \sim N(0, \tilde{\Omega}(\rho))$ is identically zero under $H_0 : \rho = 1$ (a proof is available on request), an outcome that seems to be related to the trivial local power in the $O(n^{-1/2})$ neighborhood for fixed T .

Notwithstanding the above discussion and simulation findings, the power envelope analysis given earlier seems to imply the existence of a most powerful test based on double differenced data that may have nontrivial power in an $O(n^{-1/2})$ neighborhood of unity with T fixed. Increasing power to approach this power envelope would involve using other moment conditions (e.g., by the use of MLE with double differenced data). This interesting issue presents a major challenge for future research.

6 Conclusion

This paper develops a simple GMM estimator for dynamic panel data models, which is largely free from bias as the AR coefficient approaches unity, and which yields standard Gaussian asymptotics for all values of ρ and without any discontinuity at unity. The limit theory is also robust in the sense that it performs well under all possible passages to infinity, including $n \rightarrow \infty$, $T \rightarrow \infty$ and all diagonal paths. The method also extends in a straightforward manner to cases with exogenous variables, cross section dependence, and incidental trends.

The approach leads to standard Gaussian panel unit root tests. These tests do not suffer from size distortion regardless of the n/T ratio. Illustration of power properties of some infeasible likelihood ratio tests indicates that the optimal convergence rate can perhaps be achieved using (double) differenced data while at the same time preserving standard Gaussian limits. Such tests can be expected to be particularly useful when n is large and T is small or moderate, and to outperform existing point optimal tests and exceed the usual power envelope for such sample size configurations. Extension of the present line of research in this direction is a major challenge and is left for future work.

Table 4: Simulated Size and Power of Unit Root Tests with Incidental Trends case.

$$\text{DGP: } y_{it} = \alpha_i + \gamma_i t + u_{it}, \quad u_{it} = \rho_i u_{it-1} + \sigma_i \varepsilon_{it}$$

$$\alpha_i, \gamma_i, \varepsilon_{it} \sim iid N(0, 1), \quad \sigma_i \sim iid U(0.5, 1.5),$$

Significance level: 5%

$T = 3$

| n | $\rho_i \equiv 1$ | | | | $\rho_i \equiv 0.5$ | | | | $\rho_i \sim U(0.5, 1)$ | | | |
|-----|-------------------|------|------|-------|---------------------|-------|-------|------|-------------------------|-------|-------|------|
| | HP | HP* | UB | PP | HP | HP* | UB | PP | HP | HP* | UB | PP |
| 50 | 6.93 | 4.72 | 4.84 | 12.59 | 13.19 | 19.62 | 19.51 | 0.00 | 8.41 | 9.93 | 10.35 | 0.53 |
| 100 | 6.32 | 4.89 | 4.64 | 15.57 | 15.11 | 25.15 | 25.19 | 0.00 | 8.89 | 11.93 | 11.22 | 0.09 |
| 200 | 5.27 | 4.75 | 4.98 | 22.32 | 20.02 | 33.70 | 33.35 | 0.00 | 9.50 | 13.35 | 12.84 | 0.01 |
| 400 | 5.25 | 5.09 | 4.79 | 31.35 | 28.75 | 46.25 | 45.58 | 0.00 | 11.29 | 16.59 | 16.21 | 0.00 |

$T = 5$

| n | $\rho_i \equiv 1$ | | | | $\rho_i \equiv 0.5$ | | | | $\rho_i \sim U(0.5, 1)$ | | | |
|-----|-------------------|------|------|------|---------------------|-------|-------|------|-------------------------|-------|-------|------|
| | HP | HP* | UB | PP | HP | HP* | UB | PP | HP | HP* | UB | PP |
| 50 | 6.03 | 5.55 | 5.40 | 5.94 | 20.51 | 31.44 | 47.67 | 0.00 | 9.97 | 13.40 | 17.60 | 0.17 |
| 100 | 5.25 | 5.16 | 5.78 | 6.84 | 28.59 | 46.25 | 69.10 | 0.00 | 12.03 | 16.87 | 23.65 | 0.06 |
| 200 | 5.11 | 5.35 | 5.37 | 7.35 | 43.84 | 64.14 | 89.20 | 0.00 | 14.59 | 21.74 | 33.27 | 0.01 |
| 400 | 4.84 | 5.15 | 5.16 | 9.12 | 67.10 | 85.08 | 98.98 | 0.00 | 19.94 | 30.36 | 49.43 | 0.00 |

Note: PP: Ploberger and Phillips (2002);

$$\text{HP}^* = \frac{\sum_i \sum_t \sigma_i^{-2} \Delta^2 y_{it-1} (2\Delta^2 y_{it} + \Delta^2 y_{it-1})}{\sqrt{[(8 + 4/T_2)/6] \sum_i \sum_t \sigma_i^{-2} (2\Delta^2 y_{it} + \Delta^2 y_{it-1})^2}};$$

HP*, UB and PP: Calculated with σ_i known.

A Proofs

The assumed model in the fixed effects case is $y_{it} = \alpha_i + u_{it}$, $u_{it} = \rho u_{it-1} + \varepsilon_{it}$, where $\varepsilon_{it} \sim iid(0, \sigma^2)$. Obviously we can express, for all $\rho \in (-1, 1]$,

$$(31) \quad \Delta y_{it-1} = \sum_{j=0}^{\infty} \rho^j \Delta \varepsilon_{it-1-j} = \varepsilon_{it-1} - (1 - \rho) \sum_{j=1}^{\infty} \rho^{j-1} \varepsilon_{it-j-1},$$

and

$$(32) \quad \eta_{it} = 2\Delta \varepsilon_{it} + (1 + \rho)\Delta y_{it-1} = 2\varepsilon_{it} - (1 - \rho)\varepsilon_{it-1} - (1 - \rho^2) \sum_{j=1}^{\infty} \rho^{j-1} \varepsilon_{it-j-1},$$

where $0 \cdot \pm\infty = 0$.

Proof of Lemma 1. If $\rho = 1$, then $\Delta y_{it} = \varepsilon_{it}$, and $\eta_{it} = 2\varepsilon_{it}$. So we have $E\Delta y_{it-1}\eta_{it} = 2E\varepsilon_{it-1}\varepsilon_{it} = 0$. If $|\rho| < 1$, then $E\Delta y_{it-1}\eta_{it} = -(1 - \rho)\sigma^2 + (1 - \rho)(1 - \rho^2)(1 - \rho^2)^{-1}\sigma^2 = 0$. ■

Next, we present some lemmas that are useful in proving the theorems. Let $T_m = \max(T - m, 0)$. Let $X_{it} = \sum_0^\infty c_j \varepsilon_{it-j}$ and $Y_{it} = \sum_0^\infty d_j \varepsilon_{it-j}$ where $\varepsilon_{it} \sim iid(0, \sigma^2)$. We will frequently assume that

$$(33) \quad \sum_0^\infty (|c_s| + |d_s|) < \infty,$$

$$(34) \quad \sum_0^\infty s(c_s^2 + d_s^2) < \infty.$$

Theorem 5 *If $E\varepsilon_{it}^2 < \infty$, then under (34),*

$$\frac{1}{nT_m} \sum_{i=1}^n \sum_{t=m+1}^T X_{it}^2 \rightarrow_p \sigma^2 \sum_0^\infty c_j^2$$

for any small m (e.g., 2 or 3), as $nT_m \rightarrow \infty$.

Of course, the part of condition (34) related to d_s is not relevant in this case.

Proof. If T is fixed and $n \rightarrow \infty$, then by Khinchine's law of large numbers,

$$\frac{1}{T_m} \sum_{t=m+1}^T \left[\frac{1}{n} \sum_{i=1}^n X_{it}^2 \right] \rightarrow_p \frac{1}{T_m} \sum_{t=m+1}^T EX_{it}^2 = EX_{it}^2 = \sigma^2 \sum_0^\infty c_j^2,$$

where the first equality on the right hand side holds because of stationarity. If n is fixed and $T \rightarrow \infty$, then the result follows from the convergence $T_m^{-1} \sum_{t=m+1}^T X_{it}^2 \rightarrow_p \sigma^2 \sum_0^\infty c_j^2$ for each i under the stated conditions (Theorem 3.7 of Phillips and Solo (1992)) and the

cross-sectional *iid* assumption. If both n and T increase to infinity, by Theorem 1 of Phillips and Moon (1999), it suffices to show that (a) $\limsup_{n,T} n^{-1} \sum_{i=1}^n EZ_{iT}\{Z_{iT} > n\delta\} = 0$ for all $\delta > 0$ where $Z_{iT} = T_m^{-1} \sum_{t=m+1}^T X_{it}^2$ (a notation that is used only in this proof), because all other conditions in Phillips and Moon (1999)'s theorem are obviously satisfied. Because the Z_{iT} are *iid* across i , condition (a) is equivalent to $EZ_{1T}\{Z_{1T} > n\delta\} = 0$ for all $\delta > 0$, which is implied by the uniform integrability of Z_{1T} (over T). Since $Z_{1T} \rightarrow_p \sigma^2 \sum_0^\infty c_j^2$, the uniform integrability of Z_{1T} (over T) is equivalent to the convergence $EZ_{1T} \rightarrow \sigma^2 \sum_0^\infty c_j^2$, which is obviously true because $EZ_{1T} = \sigma^2 \sum_0^\infty c_j^2$ for all T . ■

Next, we establish a panel CLT for the sample covariance of X_{it} and Y_{it} . We assume that X_{it} and Y_{it} are uncorrelated by imposing the condition

$$(35) \quad \sum_0^\infty c_j d_j = 0.$$

Theorem 6 *Let $\Pi_{j,r} = c_j d_{j+r} + c_{j+r} d_j$. If $E\varepsilon_{it}^4 < \infty$, then under (33), (34) and (35), for any fixed T and small m (e.g., 2 or 3),*

$$(36) \quad U_{nT} := (nT_m)^{-1/2} \sum_{i=1}^n \sum_{t=m+1}^T X_{it} Y_{it} \Rightarrow N(0, V_T)$$

as $n \rightarrow \infty$, where

$$V_T = A_T \text{var}(\varepsilon_{it}^2) + B_T \sigma^4,$$

with

$$(37) \quad A_T = \sum_0^\infty c_j^2 d_j^2 + \frac{2}{T_m} \sum_{t=m+2}^T \sum_{r=1}^{t-m-1} \sum_{j=0}^\infty c_j d_j c_{j+r} d_{j+r},$$

$$(38) \quad B_T = \sum_{j=0}^\infty \sum_{r=1}^\infty \Pi_{j,r}^2 + \frac{2}{T_m} \sum_{t=m+2}^T \sum_{k=1}^{t-m-1} \sum_{j=0}^\infty \sum_{r=1}^\infty \Pi_{j,r} \Pi_{j+k,r}.$$

Furthermore,

$$(39) \quad V_T \rightarrow \sigma^4 B = \sigma^4 \sum_{r=1}^\infty \left(\sum_{j=0}^\infty \Pi_{j,r} \right)^2,$$

as $T \rightarrow \infty$. Whether or not $n \rightarrow \infty$, $U_{nT} \Rightarrow N(0, \sigma^4 B)$ as $T \rightarrow \infty$.

Proof. When T is small, (36) follows from the central limit law for *iid* variates because fourth moments are finite. The variance V_T is computed as follows. Since

$$(40) \quad X_{it} Y_{it} = \sum_{j=0}^\infty c_j d_j \varepsilon_{it-j}^2 + \sum_{j=0}^\infty \sum_{r=1}^\infty \Pi_{j,r} \varepsilon_{it-j} \varepsilon_{it-j-r},$$

we have

$$(41) \quad C_0 = \text{var}(X_{it}Y_{it}) = \left(\sum_{j=0}^{\infty} c_j^2 d_j^2 \right) \text{var}(\varepsilon_{it}^2) + \left(\sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r}^2 \right) \sigma^4.$$

Also

$$(42) \quad X_{it-k}Y_{it-k} = \sum_{j=0}^{\infty} c_j d_j \varepsilon_{it-k-j}^2 + \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r} \varepsilon_{it-k-j} \varepsilon_{it-k-j-r},$$

and from (40) and (42), $C_k := \text{cov}(X_{it}Y_{it}, X_{it-k}Y_{it-k})$ is

$$C_k = \left(\sum_{j=0}^{\infty} c_j d_j c_{j+k} d_{j+k} \right) \text{var}(\varepsilon_{it}^2) + \left(\sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r} \Pi_{j+k,r} \right) \sigma^4.$$

Now

$$(43) \quad T_m^{-1} \text{var} \left(\sum_{t=m+1}^T X_{it}Y_{it} \right) = C_0 + \frac{2}{T_m} \sum_{t=m+2}^T \sum_{k=1}^{t-m-1} C_k = A_T \text{var}(\varepsilon_{it}^2) + B_T \sigma^4,$$

as stated. Lemmas 7 and 8 below respectively show that $A_T \rightarrow 0$ and $B_T \rightarrow B$ under (33) and (34), and thus the convergence (39) is obvious from (43).

Now we prove the central limit theory as $T \rightarrow \infty$. If n is fixed, then the result follows from Theorem 6 of Phillips and Han (2005), implied by (34) and the finiteness of the second moments. For the case both n and T increase, we will apply the Lindeberg central limit theorem to the row-wise independent array $\{n^{-1/2}W_{Ti}\}$, where $W_{Ti} = T_m^{-1/2} \sum_{t=m+1}^T X_{it}Y_{it}$ (notation that is used only in this proof). The Lindeberg condition is

$$(44) \quad \frac{1}{n} \sum_{i=1}^n EW_{Ti}^2 \{W_{Ti}^2 > n\epsilon\} \rightarrow 0, \text{ for all } \epsilon > 0$$

(e.g., Kallenberg (2002, Theorem 5.12)). Because the random variables are *iid* across i , (44) reduces to $EW_{T1}^2 \{W_{T1}^2 > n\epsilon\} \rightarrow 0$ for all $\epsilon > 0$ (note that T depends on n), which is again implied by

$$\sup_T EW_{T1}^2 \{W_{T1}^2 > n\epsilon\} \rightarrow 0 \text{ for all } \epsilon > 0.$$

This last condition holds if W_{T1}^2 (as a sequence indexed by T) is uniformly integrable over T . When a positive random variable converges in distribution, uniform convergence is equivalent to convergence of the means (Kallenberg (2002, Lemma 4.11)). In the case of W_{T1}^2 , we have $W_{T1} \rightarrow_d W_1 \sim N(0, \sigma^4 B)$ by Theorem 6 of Phillips and Han (2005), and by the continuous mapping theorem, $W_{T1}^2 \rightarrow_d W_1^2$. But by the first part of the theorem,

$$EW_{T1}^2 = A_T \text{var}(\varepsilon_{it}^2) + \sigma^4 B_T \rightarrow \sigma^4 B_T = EW_1^2,$$

and the joint limit theory follows straightforwardly. \blacksquare

The next two lemmas respectively show that $A_T \rightarrow 0$ and $B_T \rightarrow B$ as $T \rightarrow \infty$, as indicated above.

Lemma 7 *Under (34) and (35), $\lim_{T \rightarrow \infty} A_T = 0$.*

Proof. Let $f_j = c_j d_j$ (notation used only in this proof). For any sequence $\{a_r\}$,

$$(45) \quad \sum_{t=m+2}^T \sum_{r=1}^{t-m-1} a_r = \sum_{s=1}^{T_m-1} (T_m - s) a_s,$$

and thus we have

$$\begin{aligned} A_T &= \sum_0^\infty f_j^2 + \frac{2}{T_m} \sum_{j=0}^\infty \sum_{s=1}^{T_m-1} (T_m - s) f_j f_{j+s} \\ &= \left[\sum_0^\infty f_j^2 + 2 \sum_{j=0}^\infty \sum_{s=1}^{T_m-1} f_j f_{j+s} \right] - 2 \left[\frac{1}{T_m} \sum_{j=0}^\infty \sum_{s=1}^{T_m-1} s f_j f_{j+s} \right] \\ &= \left[\sum_0^\infty f_j^2 + 2 \sum_{j=0}^\infty \sum_{s=1}^\infty f_j f_{j+s} \right] - 2 \sum_{j=0}^\infty \sum_{s=T_m}^\infty f_j f_{j+s} - 2 \left[\frac{1}{T_m} \sum_{j=0}^\infty \sum_{s=1}^{T_m-1} s f_j f_{j+s} \right] \\ &= A_{1T} - 2A_{2T} - 2A_{3T}, \text{ say.} \end{aligned}$$

Here $A_{1T} = (\sum_0^\infty f_j)^2 = (\sum_0^\infty c_j d_j)^2 = 0$ because of (35), and therefore it suffices to show that $A_{2T} \rightarrow 0$ and $A_{3T} \rightarrow 0$ as $T \rightarrow \infty$. First,

$$|A_{2T}| \leq \sum_{j=0}^\infty \sum_{s=T_m}^\infty |f_j f_{j+s}| \leq \sum_{j=0}^\infty |f_j| \sum_{s=T_m}^\infty |f_s| \rightarrow 0,$$

because

$$(46) \quad \sum_0^\infty |f_j| = \sum_0^\infty |c_j d_j| \leq \left(\sum_0^\infty c_j^2 \right)^{1/2} \left(\sum_0^\infty d_j^2 \right)^{1/2} < \infty,$$

by (34). Next,

$$|A_{3T}| \leq T_m^{-1} \sum_{j=0}^\infty \sum_{s=1}^{T_m} s |f_j f_{j+s}| = T_m^{-1} \sum_{j=0}^\infty |f_j| \sum_{s=1}^{T_m} s |f_{j+s}| \leq T_m^{-1} \left(\sum_0^\infty |f_j| \right) \sum_0^\infty s |f_s|.$$

But $\sum_0^\infty |f_j| < \infty$ by (46), and $\sum_0^\infty s |f_s| \leq (\sum_0^\infty s c_s^2)^{1/2} (\sum_0^\infty s d_s^2)^{1/2} < \infty$ by (34). So $A_{3T} = O(T_m^{-1}) \rightarrow 0$. \blacksquare

Lemma 8 Under (33) and (34), $\lim_{T \rightarrow \infty} B_T = B = \sum_{r=1}^{\infty} \left(\sum_{j=0}^{\infty} \Pi_{j,r} \right)^2 < \infty$.

Proof. The finiteness of B is proved in Theorem 6 of Phillips and Han (2005). To establish convergence, note that

$$B = \sum_{r=1}^{\infty} \left(\sum_{j=0}^{\infty} \Pi_{j,r} \right)^2 = \sum_{r=1}^{\infty} \sum_{j=0}^{\infty} \Pi_{j,r}^2 + 2 \sum_{r=1}^{\infty} \sum_{j=0}^{\infty} \sum_{k=1}^{\infty} \Pi_{j,r} \Pi_{j+k,r},$$

so we have

$$B_T - B = -2 \left(\sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r} \Pi_{j+k,r} - T_m^{-1} \sum_{t=m+2}^T \sum_{k=1}^{t-m-1} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r} \Pi_{j+k,r} \right) = -2D_T,$$

say. Using (45) again, we have

$$\begin{aligned} D_T &= \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r} \Pi_{j+k,r} - \sum_{k=1}^{T_m-1} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r} \Pi_{j+k,r} + \frac{1}{T_m} \sum_{k=1}^{T_m-1} k \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r} \Pi_{j+k,r} \\ &= \sum_{k=T_m}^{\infty} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r} \Pi_{j+k,r} + \frac{1}{T_m} \sum_{k=1}^{T_m-1} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} k \Pi_{j,r} \Pi_{j+k,r} = D_{1T} + D_{2T}, \text{ say.} \end{aligned}$$

As shown below, we have

$$(47) \quad \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} |\Pi_{j,r} \Pi_{j+k,r}| < \infty \text{ and } \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} k |\Pi_{j,r} \Pi_{j+k,r}| < \infty,$$

which imply that $D_{1T} \rightarrow 0$ and $D_{2T} = O(T_m^{-1}) \rightarrow 0$.

Now we prove (47). For the first part of (47), note that $\sum_{k=1}^{\infty} |\Pi_{j+k,r}| \leq \sum_{k=1}^{\infty} |\Pi_{k,r}| \leq \sum_{j=0}^{\infty} |\Pi_{j,r}|$, so

$$(48) \quad \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} |\Pi_{j,r} \Pi_{j+k,r}| \leq \sum_{r=1}^{\infty} \left(\sum_{j=0}^{\infty} |\Pi_{j,r}| \right)^2.$$

Now let $a_j = |c_j| + |d_j|$, implying that $|\Pi_{j,r}| \leq |c_j d_{j+r}| + |c_{j+r} d_j| \leq a_j a_{j+r}$. Then (33) and (34) imply that

$$(49) \quad \sum_0^{\infty} a_s < \infty, \quad \sum_0^{\infty} a_s^2 < \infty, \quad \sum_0^{\infty} s a_s^2 < \infty,$$

where the second and third results hold because $a_j^2 \leq 2(c_j^2 + d_j^2)$. Now, the right hand side of (48) is bounded by

$$\sum_{r=1}^{\infty} \left(\sum_{j=0}^{\infty} a_j a_{j+r} \right)^2 \leq \sum_{r=1}^{\infty} \left(\sum_{j=0}^{\infty} a_j^2 \right) \left(\sum_{j=0}^{\infty} a_{j+r}^2 \right) = \left(\sum_0^{\infty} a_j^2 \right) \left(\sum_1^{\infty} s a_s^2 \right) < \infty,$$

by (49). So the first part of (47) is proved.

For the second part of (47),

$$\begin{aligned} \sum_{k=1}^{\infty} k |\Pi_{j+k,r}| &\leq \sum_{k=1}^{\infty} k a_{j+k} a_{j+k+r} \leq \left(\sum_{k=1}^{\infty} k a_{j+k}^2 \right)^{1/2} \left(\sum_{k=1}^{\infty} k a_{j+k+r}^2 \right)^{1/2} \\ &\leq \sum_{k=1}^{\infty} k a_{j+k}^2 \leq \sum_{k=1}^{\infty} (k+j) a_{k+j}^2 \leq \sum_{k=1}^{\infty} k a_k^2, \end{aligned}$$

and therefore the second part of (47) is

$$\sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} k |\Pi_{j,r} \Pi_{j+k,r}| \leq \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} |\Pi_{j,r}| \left(\sum_{k=1}^{\infty} k a_k^2 \right) \leq \left(\sum_{j=0}^{\infty} a_j \right)^2 \left(\sum_{k=1}^{\infty} k a_k^2 \right) < \infty$$

by (49). ■

We apply Theorems 5 and 6 to the components of the FDLS estimator $\hat{\rho}_{ols}$.

Lemma 9 $(nT_1)^{-1} \sum_{i=1}^n \sum_{t=2}^T (\Delta y_{it-1})^2 \rightarrow_p 2\sigma^2/(1+\rho)$ as $nT_1 \rightarrow \infty$.

Proof. Because of (31), we invoke Theorem 5 with $c_0 = 0$, $c_1 = 1$, and $c_j = -(1-\rho)\rho^{j-2}$ for $j \geq 2$. The calculation of $\sum_0^\infty c_j^2$ for the limit is then obvious. ■

For the central limit theorem, we have the following lemma.

Lemma 10 *If $E\varepsilon_{it}^4 < \infty$, then as $n \rightarrow \infty$, $(nT_1)^{-1/2} \sum_{i=1}^n \sum_{t=2}^T \Delta y_{it-1} \eta_{it}$ converges to a normal distribution with variance $8\sigma^4/(1+\rho)$ and furthermore $(nT_1)^{-1/2} \sum_{i=1}^n \sum_{t=2}^T \Delta y_{it-1} \eta_{it} \Rightarrow N(0, 8\sigma^4/(1+\rho))$ whether $n \rightarrow \infty$ or not.*

Proof. Because the coefficients of the lag polynomials for Δy_{it-1} and η_{it} decay exponentially, conditions (33) and (34) are satisfied. Condition (35) holds by Lemma 1. The results follow from Theorem 6. The variances for finite T and infinite T are calculated below after Theorem 2 is proved. ■

Proof of Theorem 2. This follows from Lemmas 9 and 10. ■

The Variance of FDLS

We calculate the variance of the FDLS estimator $\hat{\rho}_{ols}$ in terms of the parameters using the expressions in Theorem 6. Let $\rho_1 = 1 - \rho$ and $\rho_2 = 1 - \rho^2$. The lag polynomial coefficients

Table 5: Coefficients of lag polynomials for Δy_{it-1} and η_{it} .

| | 0 | 1 | 2 | 3 | 4 | j |
|------------------|---|-----------|-----------|---------------|-----------------|---------------------|
| Δy_{t-1} | 0 | 1 | $-\rho_1$ | $-\rho\rho_1$ | $-\rho^2\rho_1$ | $-\rho^{j-2}\rho_1$ |
| η_t | 2 | $-\rho_1$ | $-\rho_2$ | $-\rho\rho_2$ | $-\rho^2\rho_2$ | $-\rho^{j-2}\rho_2$ |

$$\rho_1 = 1 - \rho, \rho_2 = 1 - \rho^2$$

of Δy_{t-1} and η_t are tabulated in Table 5, and the corresponding $\Pi_{j,r}$ terms are tabulated in Table 6. So

$$(50) \quad \sum_0^{\infty} c_j^2 d_j^2 = \rho_1^2 + \rho_1^2 \rho_2^2 (1 + \rho^4 + \dots) = 2(1 - \rho)^2 / (1 + \rho^2),$$

$$(51) \quad \begin{aligned} \sum_0^{\infty} c_j d_j c_{j+k} d_{j+k} &= -\rho^{2(k-1)} [\rho_1^2 \rho^2 + \rho^2 \rho_1^2 \rho_2^2 (1 + \rho^4 + \dots)] \\ &= -\rho^{2(k-1)} (1 - \rho)^2 (1 - \rho^2) / (1 + \rho^2), \quad k \geq 1, \end{aligned}$$

where $0^0 = 1$. As well

$$(52) \quad \begin{aligned} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r}^2 &= [4 + 4\rho_1^2 (1 + \rho^2 + \dots)] + 4\rho^2 \rho_1^2 (1 + \rho^2 + \dots) \\ &\quad + 4\rho^2 \rho_1^2 \rho_2^2 (1 + \rho^2 + \dots) (1 + \rho^4 + \dots) \\ &= 4 \left[1 + \frac{(1 + \rho^2)(1 - \rho)}{1 + \rho} + \frac{\rho^2 (1 - \rho)^2}{1 + \rho^2} \right], \end{aligned}$$

and (after some algebra)

$$(53) \quad \begin{aligned} \sum_{j=0}^{\infty} \sum_{r=1}^{\infty} \Pi_{j,r} \Pi_{j+k,r} &= -4\rho(1 - \rho) / (1 + \rho) \{k = 1\} + 4\rho^{2k-3} (1 - \rho)^2 \{k > 1\} \\ &\quad - 4\rho^{2k} (1 - \rho)^2 / (1 + \rho^2), \quad k \geq 1. \end{aligned}$$

Table 6: The $\Pi_{j,r}$ terms.

| $r \setminus j$ | 0 | 1 | 2 | 3 | 4 | 5 |
|-----------------|----------------|------------------|-----------------------|-----------------------|-----------------------|---------|
| 1 | 2 | $-2\rho\rho_1$ | $2\rho\rho_1\rho_2$ | $2\rho^3\rho_1\rho_2$ | $2\rho^5\rho_1\rho_2$ | \dots |
| 2 | $-2\rho_1$ | $-2\rho^2\rho_1$ | $2\rho^2\rho_1\rho_2$ | $2\rho^4\rho_1\rho_2$ | $2\rho^6\rho_1\rho_2$ | \dots |
| 3 | $-2\rho\rho_1$ | $-2\rho^3\rho_1$ | $2\rho^3\rho_1\rho_2$ | $2\rho^5\rho_1\rho_2$ | $2\rho^7\rho_1\rho_2$ | \dots |
| \vdots | \vdots | \vdots | \vdots | \vdots | \vdots | |

The calculation of A_T and B_T of Theorem 6 (with $m = 1$) is tedious. We will use

$$\sum_{t=3}^T \sum_{k=1}^{t-2} \rho^{2(k-1)} = \frac{T_2}{1-\rho^2} - \left(\frac{\rho^2}{1-\rho^2} \right) \frac{1-\rho^{2T_2}}{1-\rho^2},$$

and

$$\sum_{t=3}^T \sum_{k=2}^{t-2} \rho^{2(k-2)} = \sum_{t=4}^T \sum_{k=2}^{t-2} \rho^{2(k-2)} = \frac{T_3}{1-\rho^2} - \left(\frac{\rho^2}{1-\rho^2} \right) \frac{1-\rho^{2T_3}}{1-\rho^2}.$$

From the definition of A_T , (50) and (51), we have

$$\begin{aligned} A_T &= \frac{2(1-\rho)^2}{1+\rho^2} - \frac{2}{T_1} \left[\frac{T_2}{1-\rho^2} - \left(\frac{\rho^2}{1-\rho^2} \right) \frac{1-\rho^{2T_2}}{1-\rho^2} \right] \frac{(1-\rho)^2(1-\rho^2)}{1+\rho^2} \\ (54) \quad &= \frac{2}{T_1} \left[\frac{(1-\rho)^2}{1+\rho^2} + \frac{\rho^2(1-\rho)(1-\rho^{2T_2})}{(1+\rho)(1+\rho^2)} \right], \end{aligned}$$

and

$$\begin{aligned} B_T &= 4 \left[1 + \frac{(1+\rho^2)(1-\rho)}{1+\rho} + \frac{\rho^2(1-\rho)^2}{1+\rho^2} \right] \\ &\quad + \frac{8}{T_1} \left\{ -\frac{T_2\rho(1-\rho)}{1+\rho} \{T > 3\} + \left[\frac{T_3(1-\rho)}{1+\rho} - \frac{\rho^2(1-\rho^{2T_3})}{(1+\rho)^2} \right] \rho \right. \\ &\quad \left. - \left[\frac{T_2(1-\rho)}{1+\rho} - \frac{\rho^2(1-\rho^{2T_2})}{(1+\rho)^2} \right] \frac{\rho^2}{1+\rho^2} \right\}, \end{aligned}$$

which equals

$$\begin{aligned} (55) \quad B_T &= 4 \left[1 + \frac{(1+\rho^2)(1-\rho)}{1+\rho} + \frac{\rho^2(1-\rho)^2}{1+\rho^2} - \left(\frac{2}{T_1} \right) \frac{\rho(1-\rho)}{1+\rho} \{T \geq 3\} \right. \\ &\quad \left. - 2 \left(\frac{T_2}{T_1} \right) \frac{\rho^2(1-\rho)}{(1+\rho)(1+\rho^2)} + \left(\frac{2}{T_1} \right) \frac{\rho^4(1-\rho^{2T_2})}{(1+\rho)^2(1+\rho^2)} - \left(\frac{2}{T_1} \right) \frac{\rho^3(1-\rho^{2T_3})}{(1+\rho)^2} \right], \end{aligned}$$

where A_T and B_T are defined in Theorem 6 and $m = 1$ is used.

The limit variance (as $n \rightarrow \infty$) of $(nT_1)^{1/2}(\hat{\rho}_{ols} - \rho)$ is now obtained by multiplying $A_T \text{var}(\varepsilon_{it}^2) + B_T \sigma^4$ by $(1+\rho)^2/4\sigma^4$, viz.,

$$\begin{aligned} (56) \quad V_{ols,T} &= \frac{1}{T_1} \left[\frac{(1-\rho^2)^2}{1+\rho^2} - \frac{\rho^2(1-\rho^2)(1-2\rho^{2T_2})}{1+\rho^2} \right] (1/2) \text{var}(\varepsilon_{it}^2/\sigma^2) \\ &\quad + (1+\rho)^2 + (1+\rho^2)(1-\rho^2) + \frac{\rho^2(1-\rho^2)^2}{1+\rho^2} - \left(\frac{2}{T_1} \right) \rho(1-\rho^2) \{T \geq 3\} \\ &\quad - 2 \left(\frac{T_2}{T_1} \right) \frac{\rho^2(1-\rho^2)}{1+\rho^2} + \left(\frac{2}{T_1} \right) \frac{\rho^4(1-\rho^{2T_2})}{1+\rho^2} - \left(\frac{2}{T_1} \right) \rho^3(1-\rho^{2T_3}). \end{aligned}$$

As a special case, if $T = 2$ and $\varepsilon_{it} \sim N(0, \sigma^2)$, then $\text{var}(\varepsilon_{it}^2/\sigma^2) = 2$ and $V_{ols,T}$ is simplified to

$$(57) \quad V_{ols,T} = (3-\rho)(1+\rho), \quad T = 2, \quad \varepsilon_{it} \sim N(0, \sigma^2).$$

It is also easily verified that $V_{ols,T} \rightarrow 2(1+\rho)$ as $T \rightarrow \infty$.

The Variance of DDLS When $\rho = 1$

Next, we calculate the variance of the DDLS (double-differencing least squares) estimator $\hat{\theta}$ for $\rho = 1$. According to Phillips and Han (2005), $\Delta^2 y_{it-1} = \sum_0^\infty c_j \varepsilon_{it-j}$ and $\tilde{\eta}_{it} = \sum_0^\infty d_j \varepsilon_{it-j}$, where

$$(58) \quad c_0 = 0, \quad c_1 = 1, \quad c_2 = -(2 - \rho), \quad c_k = \rho^{k-3}(1 - \rho)^2, \quad k \geq 3,$$

$$(59) \quad d_0 = 2, \quad d_1 = -4 + \phi c_1, \quad d_2 = 2 + \phi c_2, \quad d_k = \phi c_k, \quad k \geq 3,$$

with $\phi = (4 - \rho)(1 + \rho)/(3 - \rho)$ and $0^0 = 1$. When $\rho = 1$, we have

$$c_0 = 0, \quad c_1 = 1, \quad c_2 = -1, \quad c_k = 0, \quad k \geq 3,$$

$$d_0 = 2, \quad d_1 = -1, \quad d_2 = -1, \quad d_k = 0, \quad k \geq 3.$$

So

$$\sum_0^\infty c_j^2 d_j^2 = 2 \quad \text{and} \quad \sum_{t=m+2}^T \sum_{r=1}^{t-m-1} \sum_{j=0}^\infty c_j d_j c_{j+r} d_{j+r} = -T_{m+1},$$

and because

$$\Pi_{0,1} = 2, \quad \Pi_{0,2} = -2, \quad \text{and} \quad \Pi_{j,r} = 0 \quad \text{for all other } j \text{ and } r,$$

we have

$$\sum_{j=0}^\infty \sum_{r=1}^\infty \Pi_{j,r}^2 = 8, \quad \sum_{t=m+2}^T \sum_{k=1}^{t-m-1} \sum_{j=0}^\infty \sum_{r=1}^\infty \Pi_{j,r} \Pi_{j+k,r} = 0.$$

For the DDLS estimator $\hat{\theta}$, we use $m = 2$, and thus, by Theorem 6, when $\rho = 1$ and $T > 2$,

$$(nT_2)^{-1/2} \sum_{i=1}^n \sum_{t=3}^T \Delta^2 y_{it-1} \tilde{\eta}_{it} \rightarrow_d N\left(0, 2 \text{var}(\varepsilon_{it}^2)/T_2 + 8\sigma^4\right),$$

and because $\text{plim}_{nT \rightarrow \infty} (nT_2)^{-1} \sum_{i=1}^n \sum_{t=3}^T (\Delta^2 y_{it-1})^2 = 2\sigma^4$ by Theorem 5, we have $\theta = 1$ and as $nT \rightarrow \infty$,

$$(nT_2)^{1/2} \hat{\theta} \rightarrow_d N\left(0, \text{var}(\varepsilon_{it}^2/\sigma^2)/2T_2 + 2\right).$$

Proof of Lemma 4. Recall that the model is $y_{it} = \alpha_i + u_{it}$ with $u_{it} = \rho u_{it-1} + \varepsilon_{it}$. Let the sequence be initialized at u_{i0} . Then

$$u_{it} = \rho^t u_{i0} + \sum_{j=0}^{t-1} \rho^j \varepsilon_{it-j},$$

and therefore

$$\Delta y_{it-1} = \Delta u_{it-1} = \rho^{t-2}(\rho - 1)u_{i0} + \varepsilon_{it-1} + (\rho - 1) \sum_{j=1}^{t-2} \rho^{j-1} \varepsilon_{it-j-1},$$

and

$$\begin{aligned} E(\Delta y_{it-1})^2 &= (\rho - 1)^2 \rho^{2(t-2)} E u_{i0}^2 + [1 + (\rho - 1)^2 (1 + \rho^2 + \dots + \rho^{2(t-3)})] \sigma^2 \\ (60) \quad &= (\rho - 1)^2 \rho^{2(t-2)} E u_{i0}^2 + (\rho + 1)^{-1} [2 + (\rho - 1) \rho^{2(t-2)}] \sigma^2 \\ &= 2\sigma^2 / (\rho + 1) + \left(\frac{\rho - 1}{\rho + 1} \right) \rho^{2(t-2)} \sigma^2 \xi_\rho, \end{aligned}$$

where $(\rho^2 - 1)E u_{i0}^2 / \sigma^2 + 1$. (We can verify the calculation by checking with the case of $\rho < 1$, where $\xi_\rho = 0$ and therefore $E(\Delta y_{it-1})^2 = 2\sigma^2 / (1 + \rho)$ as in Lemma 9.) Next, because $\eta_{it} = 2\Delta \varepsilon_{it} + (1 + \rho)\Delta y_{it-1}$, we have

$$(61) \quad E \Delta y_{it-1} \eta_{it} = (\rho - 1) \rho^{2(t-2)} \sigma^2 \xi_\rho.$$

The results follow from (60) and (61). ■

Next we show that the moment conditions (15) and (16) make the expected first derivative matrix (D) block diagonal and if $E \varepsilon_{it}^3 = 0$ then the variance-covariance matrix (Ω) is block diagonal too when evaluated at the true parameter. For convenience, we write (15) and (16) again as

$$\begin{aligned} E A_{it}(\rho, \beta) &= E \sum_{t=2}^T \Delta z_{it-1} [2\Delta z_{it} + (1 - \rho)\Delta z_{it}] = 0, \\ E B_{it}(\rho, \beta) &= E \sum_{t=1}^T (\ddot{x}_{it} - \rho \dot{x}_{it-1}) [(\ddot{y}_{it} - \rho \dot{y}_{it-1}) - (\ddot{x}_{it} - \rho \dot{x}_{it-1})' \beta] = 0, \end{aligned}$$

where $\Delta z_{it} = \Delta y_{it} - \Delta x'_{it} \beta$.

For the D matrix, we need to show that $E \partial A_{it} / \partial \beta = 0$ and $E \partial B_{it} / \partial \rho = 0$ when evaluated at the true parameter. Because $\Delta z_{it} = \Delta u_{it}$ and $\ddot{y}_{it} = (\ddot{x}_{it} - \rho \dot{x}_{it-1})' \beta + \rho \dot{y}_{it-1} + \ddot{\varepsilon}_{it}$ when evaluated at the true parameter, we have

$$\begin{aligned} E \frac{\partial A_{it}}{\partial \beta} &= -E \sum_{t=2}^T \left[\Delta x_{it-1} \eta_{it} + \Delta u_{it} \{ 2\Delta x_{it} + (1 - \rho)\Delta x_{it-1} \} \right] = 0, \\ E \frac{\partial B_{it}}{\partial \rho} &= -E \sum_{t=1}^T \left[\dot{x}_{it-1} \ddot{\varepsilon}_{it} + (\ddot{x}_{it} - \rho \dot{x}_{it-1}) \dot{u}_{it-1} \right] = 0, \end{aligned}$$

where $\eta_{it} = 2\Delta u_{it} + (1 - \rho)\Delta u_{it-1}$ and $\dot{u}_{it-1} = u_{it-1} - T^{-1} \sum_{s=1}^T u_{is-1} = \dot{y}_{it-1} - \dot{x}'_{it-1}\beta$. So the D matrix is block diagonal. Next, for the Ω matrix, at the true parameter,

$$EB_{it}A_{it} = E \left[\sum_{t=1}^T (\ddot{x}_{it} - \rho \ddot{x}_{it-1}) \ddot{\varepsilon}_{it} \right] \left[\sum_{t=2}^T \Delta u_{it-1} \eta_{it} \right] = 0,$$

when $E\varepsilon_{it} = 0$, $E\varepsilon_{it}^3 = 0$ and ε_{it} are *iid*. So the Ω matrix is also block diagonal under the zero third moment assumption.

Now let us obtain the limit behavior of $U_{nT}(c)$ of (23) under the local to unity setting $\rho = \rho_{nT} = 1 - n^{-1/2}T^{-1}c$. Clearly,

$$U_{nT}(c) = -n \log (|\Omega_T(\rho_{nT})|/|\Omega_T(1)|) - \frac{1}{\sigma^2} \sum_{i=1}^n (\Delta y_i)' [\Omega_T(\rho_{nT})^{-1} - \Omega_T(1)^{-1}] (\Delta y_i),$$

where $\Omega_T(\cdot)$ is defined in (24). Under the alternative hypothesis that $n^{1/2}T(1 - \rho) = c$,

$$\begin{aligned} EU_{nT}(c) &= n \left(\text{tr} \Omega_T(\rho_{nT}) - T - \log |\Omega_T(\rho_{nT})| \right) \\ &= n \left(\frac{(1 - \rho_{nT})T}{1 + \rho_{nT}} - \log \left[1 + \frac{(1 - \rho_{nT})T}{1 + \rho_{nT}} \right] \right), \end{aligned}$$

where the determinant is calculated in Han (2006, Theorem 2). Because

$$x - \log(1 + x) = x^2/2 - o(x^2)$$

when x is close to zero, we have

$$EU_{nT}(c) = \frac{nT^2(1 - \rho_{nT})^2}{2(1 + \rho_{nT})^2} + o \left(\frac{nT^2(1 - \rho_{nT})^2}{(1 + \rho_{nT})^2} \right) = \frac{c^2}{2(1 + \rho_{nT})^2} + o(1) \rightarrow \frac{c^2}{8}.$$

The variance is calculated from the fact that $\text{var}(z'Az) = 2\text{tr}(A'A)$ if $z \sim N(0, I)$ and A is symmetric. In our case,

$$\sigma^{-2} \Delta y_i' [\Omega_T(\rho)^{-1} - \Omega_T(1)^{-1}] \Delta y_i = z_i' [I_T - \Omega_T(\rho)] z_i = z_i' Q_T z_i, \quad \text{say,}$$

where $z_i = \sigma^{-1} \Omega_T(\rho)^{-1/2} \Delta y_i \sim N(0, I)$ and

$$Q_T = \text{Toeplitz}\{1, 1, \rho, \dots, \rho^{T_2}\}(1 - \rho)/(1 + \rho),$$

so that

$$\begin{aligned} \text{var}(U_{nT}(c)) &= 2n \text{tr}(Q_T' Q_T) = 2 \left[T + 2(T_1 + T_2 \rho_{nT}^2 + \dots + \rho_{nT}^{2T_2}) \right] \frac{(1 - \rho_{nT})^2}{(1 + \rho_{nT})^2} \\ (62) \quad &= \frac{2nT(1 - \rho_{nT})^2}{(1 + \rho_{nT})^2} + \frac{4n(1 - \rho_{nT})^2}{(1 + \rho_{nT})^2} \left[\frac{T_1}{1 - \rho_{nT}^2} - \frac{\rho_{nT}^2(1 - \rho_{nT}^{2T_1})}{(1 - \rho_{nT}^2)^2} \right]. \end{aligned}$$

The first term of (62) is $[1 + o(1)]c^2/2T$ when $n^{1/2}T(1 - \rho_{nT}) = c$. (The notation $o(1)$ means here that the relevant term disappears as $n \rightarrow \infty$.) The second term can be handled by the following lemma.

Lemma 11 For every c ,

$$n(1 - \rho_{nT}^{2T_1}) = 2\sqrt{n}(T_1/T)c - (T_1/T)^2(2 - T_1^{-1})c^2 + O(n^{-1/2}).$$

Proof. From the binomial expansion of $\rho_{nT}^{2T_1} = (1 - n^{-1/2}T^{-1}c)^{2T_1}$, we have

$$n(1 - \rho_{nT}^{2T_1}) = -n \sum_{k=1}^{2T_1} \binom{2T_1}{k} (-n^{-1/2}T^{-1}c)^k.$$

The first two terms of the lemma correspond to $k = 1$ and $k = 2$ respectively. For the remainder term, note that

$$n \sum_{k=3}^{2T_1} \binom{2T_1}{k} (n^{-1/2}T^{-1}c)^k \leq \sum_{k=3}^{2T_1} \frac{n^{1-k/2}(2c)^k}{k!} \leq n^{-1/2} \sum_{k=0}^{\infty} \frac{(2c)^k}{k!} = n^{-1/2}e^{2c},$$

which is $O(n^{-1/2})$ as stated. ■

Whether T is fixed or $T \rightarrow \infty$ as $n \rightarrow \infty$, we have, after some algebra,

$$\text{var}(U_{nT}(c)) \rightarrow c^2/2.$$

In the derivation of this variance, we can also verify that it is uniformly bounded, which is sufficient to invoke the Lindeberg CLT for $U_{nT}(c)$. Thus,

$$U_{nT}(c) \rightarrow_d N(c^2/8, c^2/2).$$

Now establish a similar CLT for $\tilde{U}_{nT}(c)$ of (28) under the local to unity setting $\rho = \rho_{nT} = 1 - c/\sqrt{nT_1}$ for some $c > 0$. The algebra is tedious, and is therefore separated into several pieces in the following lemmas. First, to evaluate $E\tilde{U}_{nT}(c)$ when $1 - (nT_1)^{-1/2}c$ is the true ρ parameter, we obtain an algebraic expression of $\tilde{\Omega}_T(1)^{-1}$.

Lemma 12 Let $A_T = \text{Toeplitz}\{T_1, T_1 - 1, \dots, 1\}$, $\bar{A}_{T,0} = A_T$ and

$$\bar{A}_{T,j} = \begin{bmatrix} O_{j \times j} & O_{j \times (T_1 - 2j)} & O_{j \times j} \\ O_{(T_1 - 2j) \times j} & A_{T-2j} & O_{(T_1 - 2j) \times j} \\ O_{j \times j} & O_{j \times (T_1 - 2j)} & O_{j \times j} \end{bmatrix}, \quad j = 1, 2, \dots, [(T_1 - 1)/2],$$

where $[x]$ denotes the greatest integer not exceeding x and $O_{p \times q}$ denotes the $p \times q$ matrix of zeros. Then

$$\tilde{\Omega}_T(1)^{-1} = T^{-1} \sum_{j=0}^{[T_2/2]} \bar{A}_{T,j}.$$

Proof. Use mathematical induction. ■

Lemma 13 *Let $S_k = k^2 + (k - 2)^2 + \dots + (1 \text{ or } 2)^2$, where the last term is 1 if k is odd and 2 otherwise. Then $S_k = k(k + 1)(k + 2)/6$.*

Proof. We use mathematical induction as follows. First, $S_1 = 1 = 1 \cdot 2 \cdot 3/6$ and $S_2 = 4 = 2 \cdot 3 \cdot 4/6$. Next, suppose that $S_k = k(k + 1)(k + 2)/6$ is true. Then

$$S_{k+2} = (k + 2)^2 + S_k = (k + 2)^2 + k(k + 1)(k + 2)/6 = (k + 2)(k + 3)(k + 4)/6,$$

which gives the result. ■

Now we consider $E\tilde{U}_{nT}(c)$ where $n^{1/2}T_1^{1/2}(\rho - 1) = c > 0$. Let

$$(63) \quad \begin{aligned} \mu_T(\rho) &= \text{tr} \left\{ [\tilde{\Omega}_T(\rho) - \tilde{\Omega}_T(1)]\tilde{\Omega}_T(1)^{-1} \right\} - \log |\tilde{\Omega}_T(\rho)\tilde{\Omega}_T(1)^{-1}| \\ &= \text{tr} \left[\tilde{\Omega}_T(\rho)\tilde{\Omega}_T(1)^{-1} \right] - T_1 - \log |\tilde{\Omega}_T(\rho)| + \log T, \end{aligned}$$

so $E\tilde{U}_{nT}(c) = n\mu_T(\rho_{nT})$. Let $\mu_{1T}(\rho) = \text{tr} \tilde{\Omega}_T(\rho)\tilde{\Omega}_T(1)^{-1} - T_1$ and $\mu_{2T}(\rho) = \log |\tilde{\Omega}_T(\rho)| - \log T$ so that $\mu_T(\rho) = \mu_{1T}(\rho) - \mu_{2T}(\rho)$. We shall first investigate $\mu_{1T}(\rho)$ and $\mu_{2T}(\rho)$ separately.

Lemma 14 *We have*

$$\mu_{1T}(\rho) = \frac{T_1(1 - \rho)}{1 + \rho} + \frac{2}{1 + \rho} \left(1 - \frac{1}{T} \sum_{k=0}^{T_1} \rho^k \right) = f_T(\rho) + g_T(\rho), \text{ say.}$$

Proof. By Lemma 12, we have

$$\mu_{1T}(\rho) = \frac{1}{T} \sum_{j=0}^{[T_2/2]} \text{tr} \tilde{\Omega}_T(\rho)\bar{A}_{T,j} - T_1 = \frac{1}{T} \sum_{j=0}^{[T_2/2]} \text{tr} \tilde{\Omega}_{T-2j}(\rho)A_{T-2j} - T_1.$$

The result can be obtained through some tedious workings using $\text{tr} \tilde{\Omega}_{k+2}A_{k+2} = \sum_{j=0}^{k-1} \omega_j(k - j)^2$ and Lemma 13. ■

The determinant of $\tilde{\Omega}_T(\rho)$ is calculated in Han (2006, Theorem 2) as follows.

Lemma 15 *We have*

$$\begin{aligned} \frac{|\tilde{\Omega}_T(\rho)|}{|\tilde{\Omega}_T(1)|} &= 1 + \frac{1}{12(1 + \rho)} \left[8(2 + \rho)(1 - \rho)T_1 + (7 + \rho)(1 - \rho)^2T_1^2 + (1 - \rho)^3T_1^3 \right] \\ &= 1 + h_T(\rho), \text{ say.} \end{aligned}$$

Remember that $\mu_{2T}(\rho) = \log |\tilde{\Omega}_T(\rho)|/|\tilde{\Omega}_T(1)|$ and $\mu_T(\rho) = \mu_{1T}(\rho) - \mu_{2T}(\rho)$, i.e.,

$$\mu_T(\rho) = f_T(\rho) + g_T(\rho) - \log[1 + h_T(\rho)],$$

where $f_T(\cdot)$ and $g_T(\cdot)$ are defined in Lemma 14 and $h_T(\cdot)$ in Lemma 15. Thus

$$(64) \quad E\tilde{U}_{nT}(\rho_{nT}) = n\mu_T(\rho_{nT}) = nf_T(\rho_{nT}) + ng_T(\rho_{nT}) - n\log[1 + h_T(\rho_{nT})].$$

For the first term of (64) we only note that

$$(65) \quad nf_T(\rho_{nT}) = (nT_1)^{1/2}c/(1 + \rho_{nT})$$

when $\rho_{nT} = 1 - (nT_1)^{-1/2}c$. To handle $ng_T(\rho_{nT})$, let $\Pi_{T,k} = \prod_{j=1}^{k-1}(1 - j/T_1) = T_1!/(T_1 - k)!T_1^k$ for $2 \leq k \leq T_1$ and $\Pi_{T,1} = 1$. Let $M_{nT} = T_1/\sqrt{n}$. We consider the case $M_{nT} \rightarrow 0$.

Lemma 16 *If $T_1/\sqrt{n} \rightarrow 0$ and $c \in (0, 1]$, then*

$$ng_T(\rho_{nT}) = nf_T(\rho_{nT}) + \frac{c^2}{3(1 + \rho_{nT})} - \frac{T_1c^2}{3(1 + \rho_{nT})} + \frac{\Pi_{T,3}M_{nT}T_1^{1/2}c^3}{12(1 + \rho_{nT})} + o(1).$$

Proof. Let $a_{nT} = 1 - \rho_{nT} = (nT_1)^{-1/2}c$. Then

$$\begin{aligned} n \left(1 - \frac{1}{T} \sum_{j=0}^{T_1} \rho_{nT}^j \right) &= \frac{n}{a_{nT}T} [(1 - a_{nT})^T - (1 - Ta_{nT})] \\ &= \frac{n}{a_{nT}T} \sum_{k=2}^T \binom{T}{k} (-a_{nT})^k = \frac{n}{a_{nT}T} \sum_{k=1}^{T_1} \binom{T_1 + 1}{k + 1} (-a_{nT})^{k+1} \\ &= -n \sum_{k=1}^{T_1} \frac{1}{k + 1} \binom{T_1}{k} (-a_{nT})^k, \end{aligned}$$

where the second line comes from the binomial expansion of $(1 - a_{nT})^T$. In the last summation, the first three terms (corresponding to $k = 1$ through $k = 3$) construct the important terms in the lemma. (It is only a matter of calculation.) Truncating the terms for k exceeding 3 is valid because the term for $k = 4$ is $O(M_{nT}^2) = o(1)$ and the sum of the rest is at most $O(n^{1-5/4})$. \blacksquare

Now the last term $n\log[1 + h_T(\rho_{nT})]$ is to be expanded. When $T_1/n \rightarrow 0$, we have $h_T(\rho_{nT}) \rightarrow 0$ as $n \rightarrow \infty$ (see (69) below), and therefore we may expand the logarithm by

$$(66) \quad \log(1 + x) = \sum_{k=1}^K \frac{(-1)^{k-1}x^k}{k} + o(x^K),$$

which is valid if x is close to zero. The choice of K can be made as follows. Since $\rho_{nT} = 1 - c/\sqrt{nT_1}$, we have

$$(67) \quad n(1 - \rho_{nT})^k T_1^k = n^{1-k/2} T_1^{k/2} c^k.$$

Especially, $n(1 - \rho_{nT})^4 T_1^4 = M_{nT}^2 c^4$, and $n(1 - \rho_{nT})^5 T_1^5 = M_{nT}^3 T_1^{-1/2} c^5 = M_{nT}^{5/2} n^{-1/4} c^5$. This fact, together with (66), gives us

$$(68) \quad n \log[1 + h_T(\rho_{nT})] = n \sum_{k=1}^3 \frac{(-1)^{k-1} h_T(\rho_{nT})^k}{k} + o(1),$$

when $c \in (0, 1]$. (Expand up to the fourth order. Then the remainder is $o(M_{nT}^2)$ and the fourth term itself is $O(M_{nT}^2)$.) So we shall expand it up to the third order. Due to Lemma 15 and (67), we have

$$(69) \quad h_T(\rho_{nT}) = \frac{1}{12(1 + \rho_{nT})} \left[8(2 + \rho_{nT})n^{-1/2} T_1^{1/2} c + (7 + \rho_{nT})n^{-1} T_1 c^2 + n^{-3/2} T_1^{3/2} c^3 \right].$$

Since any term involved with $n^{-k/2} T_1^{k/2} c^k$ for $k > 3$ is negligible when multiplied by n , we have the following results.

Lemma 17 *If $T_1/\sqrt{n} \rightarrow 0$ and $c \in (0, 1]$, then*

$$\begin{aligned} nh_T(\rho_{nT}) &= 2f_T(\rho_{nT}) + \frac{2T_1 c^2}{3(1 + \rho_{nT})} - \frac{2c^2}{3(1 + \rho_{nT})} + \frac{M_{nT} T_1^{1/2} c^3}{12(1 + \rho_{nT})} - o(1), \\ nh_T(\rho_{nT})^2 &= T_1 c^2 + \frac{2M_{nT} T_1^{1/2} c^3}{3} + o(1), \\ nh_T(\rho_{nT})^3 &= M_{nT} T_1^{1/2} c^3 + o(1). \end{aligned}$$

Proof. By direct calculation. ■

Now we can obtain the limiting mean of $\tilde{U}_{nT}(c)$.

Lemma 18 *If $T_1/\sqrt{n} \rightarrow 0$, then $E\tilde{U}_{nT}(c) \rightarrow c^2/2$.*

Proof. Combine (64), Lemmas 16 and 17, and (68). Simplify using the fact that

$$\frac{T_1}{1 + \rho_{nT}} = \frac{T_1}{2} + \frac{(T_1/n)^{1/2} c}{2(1 + \rho_{nT})}, \quad \text{and} \quad \frac{T_1^{1/2}}{1 + \rho_{nT}} = \frac{T_1^{1/2}}{2} + \frac{n^{-1/2} c}{2(1 + \rho_{nT})},$$

and the result follows immediately. ■

For the variance of $\tilde{U}_{nT}(c)$, recall as above that the variance of $z'Az$ where $z \sim N(0, I_m)$ is $2\text{tr}(A'A)$. So we have

$$\text{var}(\tilde{U}_{nT}(c)) = 2n\text{tr}(\tilde{Q}'\tilde{Q}^\circ), \quad \tilde{Q}^\circ = \tilde{\Omega}_T(\rho_{nT})^{1/2}[\tilde{\Omega}_T(\rho_{nT})^{-1} - \tilde{\Omega}_T(1)^{-1}]\tilde{\Omega}_T(\rho_{nT})^{1/2}.$$

Because $\text{tr}(AB) = \text{tr}(BA)$ when AB and BA are both defined, we have

$$(70) \quad \text{var}(\tilde{U}_{nT}(c)) = 2n\text{tr}(\tilde{Q}\tilde{Q}), \quad \tilde{Q} = [\tilde{\Omega}_T(\rho_{nT}) - \tilde{\Omega}_T(1)]\tilde{\Omega}_T(1)^{-1}.$$

Obviously,

$$\tilde{\Omega}_T(\rho) - \tilde{\Omega}_T(1) = \text{Toeplitz}\{\omega_0^*(\rho), \omega_1^*(\rho), \omega_2(\rho), \dots, \omega_{T_2}(\rho)\},$$

where $\omega_0^*(\rho) = 4(1 - \rho)/(1 + \rho)$ and $\omega_1^*(\rho) = -(3 - \rho)(1 - \rho)/(1 + \rho)$. Split

$$(71) \quad \begin{aligned} \tilde{\Omega}_T(\rho) - \tilde{\Omega}_T(1) &= \text{Toeplitz}\{4, -2, 0, \dots, 0\} \cdot (1 - \rho)/(1 + \rho) \\ &\quad - \text{Toeplitz}\{0, 1, 0, \dots, 0\} \cdot (1 - \rho)^2/(1 + \rho) \\ &\quad + \text{Toeplitz}\{0, 0, 1, \rho, \dots, \rho^{T_4}\} \cdot (1 - \rho)^3/(1 + \rho) \\ &= (1 + \rho)^{-1}[(1 - \rho)D_{1T} - (1 - \rho)^2D_{2T} + (1 - \rho)^3D_{3T}], \text{ say.} \end{aligned}$$

We observe that $D_{1T} = 2\tilde{\Omega}_T(1)$. Other terms are involved but can be shown to be negligible when (70) is calculated if $T_1/\sqrt{n} \rightarrow 0$.

Lemma 19 *The following are true:*

- (i) $n(1 - \rho_{nT})^2 \text{tr} D_{1T} \tilde{\Omega}_T(1)^{-1} D_{1T} \tilde{\Omega}_T(1)^{-1} = 4c^2$;
- (ii) $0 < n(1 - \rho_{nT})^4 \text{tr} D_{2T} \tilde{\Omega}_T(1)^{-1} D_{2T} \tilde{\Omega}_T(1)^{-1} \leq n^{-1} T^2 c^4 / 4$;
- (iii) $0 < n(1 - \rho_{nT})^6 \text{tr} D_{3T} \tilde{\Omega}_T(1)^{-1} D_{3T} \tilde{\Omega}_T(1)^{-1} \leq n^{-1} T^2 c^4 / 4$;
- (iv) $0 < n(1 - \rho_{nT})^3 \text{tr} D_{1T} \tilde{\Omega}_T(1)^{-1} D_{2T} \tilde{\Omega}_T(1)^{-1} \leq n^{-1/2} T^{1/2} c^3$;
- (v) $0 < n(1 - \rho_{nT})^4 \text{tr} D_{1T} \tilde{\Omega}_T(1)^{-1} D_{3T} \tilde{\Omega}_T(1)^{-1} \leq n^{-1/2} T^{1/2} c^4$;
- (vi) $0 < n(1 - \rho_{nT})^5 \text{tr} D_{2T} \tilde{\Omega}_T(1)^{-1} D_{3T} \tilde{\Omega}_T(1)^{-1} \leq n^{-1} T^2 c^4 / 4$.

Proof. (i) Obvious because $1 - \rho_{nT} = (nT_1)^{-1/2}c$ and $D_{1T} = 2\tilde{\Omega}_T(1)$. (ii) Because all the elements of D_{2T} and $\tilde{\Omega}_T(1)^{-1}$ are positive, the trace is no bigger than $\text{tr}(D_{2T}B D_{2T}B)$ for any B whose elements are no smaller than the elements of $\tilde{\Omega}_T(1)^{-1}$. By Lemma 12, the elements of $\tilde{\Omega}_T(1)^{-1}$ are bounded by $T/4$, so

$$\text{tr} D_{2T} \tilde{\Omega}_T(1)^{-1} D_{2T} \tilde{\Omega}_T(1)^{-1} \leq (T^2/16) \text{tr} D_{2T} \iota' D_{2T} \iota = (T^2/16) (\iota' D_{2T} \iota)^2 = T^2 T_2^2 / 4.$$

The result follows immediately because $n(1 - \rho_{nT})^4 = n^{-1}T_1^{-2}c^4$. (iii) Similarly, we have

$$(72) \quad \text{tr} D_{3T} \tilde{\Omega}_T(1)^{-1} D_{3T} \tilde{\Omega}_T(1)^{-1} \leq (T^2/16)(\iota' D_{3T} \iota)^2.$$

But

$$\iota' D_{3T} \iota = 2(T_3 + T_4 \rho_{nT} + T_5 \rho_{nT}^2 + \cdots + \rho_{nT}^{T_4}) = 2 \left[\frac{T_2}{1 - \rho_{nT}} - \frac{1 - \rho_{nT}^{T_2}}{(1 - \rho_{nT})^2} \right] \leq \frac{2T_2}{1 - \rho_{nT}},$$

thus (72) is bounded by $T^2 T_2^2 / 4(1 - \rho_{nT})^2$, implying the result. The remainder are similarly proved. \blacksquare

Now the limiting variance is straightforwardly obtained.

Lemma 20 *If $T_1/\sqrt{n} \rightarrow 0$, then $\text{var} \tilde{U}_{nT}(c) \rightarrow 2c^2$.*

Proof. By (70) and (71),

$$(73) \quad \begin{aligned} \text{var} \tilde{U}_{nT}(c) = 2(1 + \rho_{nT})^{-2} & \left[n(1 - \rho_{nT})^2 \text{tr} D_{1T} \tilde{\Omega}_T(1)^{-1} D_{1T} \tilde{\Omega}_T(1)^{-1} \right. \\ & + n(1 - \rho_{nT})^4 \text{tr} D_{2T} \tilde{\Omega}_T(1)^{-1} D_{2T} \tilde{\Omega}_T(1)^{-1} \\ & + n(1 - \rho_{nT})^6 \text{tr} D_{3T} \tilde{\Omega}_T(1)^{-1} D_{3T} \tilde{\Omega}_T(1)^{-1} \\ & - 2n(1 - \rho_{nT})^3 \text{tr} D_{1T} \tilde{\Omega}_T(1)^{-1} D_{2T} \tilde{\Omega}_T(1)^{-1} \\ & + 2n(1 - \rho_{nT})^4 \text{tr} D_{1T} \tilde{\Omega}_T(1)^{-1} D_{3T} \tilde{\Omega}_T(1)^{-1} \\ & \left. - 2n(1 - \rho_{nT})^5 \text{tr} D_{2T} \tilde{\Omega}_T(1)^{-1} D_{3T} \tilde{\Omega}_T(1)^{-1} \right]. \end{aligned}$$

By Lemma 19 above, only the first term is important when $T_1/\sqrt{n} \rightarrow 0$. More specifically,

$$\text{var} \tilde{U}_{nT}(c) = 2(1 + \rho_{nT})^{-2} [4c^2 + o(1)] \rightarrow 2c^2$$

when $n^{-1/2}T_1 \rightarrow 0$. \blacksquare

With the limiting mean and variance in hand, we can now prove (29) by establishing asymptotic normality for $\tilde{U}_{nT}(\rho_{nT})$ as follows.

Proof of (29). Thanks to Lemmas 18 and 20, it remains to show the asymptotic normality of $\tilde{U}_{nT}(c)$. Rewrite

$$\begin{aligned} \tilde{U}_{nT}(c) &= - \sum_{i=1}^n \left\{ (\sigma^{-1} \Delta^2 y_i)' [\tilde{\Omega}_T(\rho_{nT})^{-1} - \tilde{\Omega}_T(1)^{-1}] (\sigma^{-1} \Delta^2 y_i)' \right. \\ &\quad \left. + [\log |\tilde{\Omega}_T(\rho_{nT})| - \log |\tilde{\Omega}_T(1)|] \right\} \\ &= - \sum_{i=1}^n \psi_{nT,i} \quad \text{say.} \end{aligned}$$

For asymptotic normality, we need to prove that the Lindeberg condition holds for the array $(\psi_{nT,i})$, viz.,

$$\sum_{i=1}^n E\psi_{nT,i}^2 \{\psi_{nT,i}^2 > \epsilon\} = En\psi_{nT,i}^2 \{n\psi_{nT,i}^2 > n\epsilon\} \rightarrow 0 \text{ for all } \epsilon > 0.$$

This condition is satisfied because $\limsup_{n,T} nE\psi_{nT,i}^2 < \infty$ due to (73) and Lemma 19 when $n^{-1/2}T_1 \rightarrow 0$. ■

References

- Ahn, S. C. and P. Schmidt (1995). Efficient Estimation of Models for Dynamic Panel Data. *Journal of Econometrics*, 68, 5–27.
- Alvarez, J. and M. Arellano (2003). The Time Series and Cross-Section Asymptotics of Dynamic Panel Data Estimators. *Econometrica*, 71(4), 1121–1159.
- Anderson, T. W. and C. Hsiao (1981). Estimation of Dynamic Models with Error Components. *Journal of American Statistical Association*, 76, 598–606.
- Arellano, M. and S. Bond (1991). Some Tests of Specification for Panel Data: Monte Carlo Evidence and an Application to Employment Equations. *Review of Economic Studies*, 58, 277–297.
- Blundell, R. and S. Bond (1998). Initial Conditions and Moment Restrictions in Dynamic Panel Data Models. *Journal of Econometrics*, 87, 115–143.
- Breitung, J. (2000). The Local Power of Some Unit Root Tests for Panel Data, in: B. Baltagi (ed.), *Nonstationary Panels, Panel Cointegration, and Dynamic Panels, Advances in Econometrics, Vol. 15*, JAI: Amsterdam, 161–178.
- Hahn, J. and G. Kuersteiner (2002). Asymptotically Unbiased Inference for a Dynamic Panel Model with Fixed Effects When Both N and T are Large. *Econometrica*, 70(4), 1639–1657.
- Han, C. (2006). Determinant of Covariance Matrix of Differenced AR(1) Processes. Unpublished manuscript.
- Han, C. and P. C. B. Phillips (2006). GMM with Many Moment Conditions. *Econometrica*, 74, 147–192.
- Hsiao, C., M. H. Pesaran, and A. K. Tahmiscioglu (2002). Maximum Likelihood Estimation of Fixed Effects Dynamic Panel Data Models Covering Short Time Periods. *Journal of Econometrics*, 109, 107–150.

- Im, K. S., S. C. Ahn, P. Schmidt, and J. Wooldridge (1999). Efficient Estimation of Panel Data Models with Strictly Exogenous Explanatory Variables—A Monte Carlo Study. *Journal of Econometrics*, 93(1), 177–201.
- Im, K. S., M. H. Pesaran, and Y. Shin (1997). Testing for Unit Roots in Heterogeneous Panels. Working paper, University of Cambridge.
- Kallenberg, O. (2002). *Foundations of Modern Probability*, second edition. Springer.
- Levin, A. and C. F. Lin (1992). Unit Root Test in Panel Data: Asymptotic and Finite Sample Properties. University of California at San Diego, Discussion paper #92-93.
- Moon, H.R. and B. Perron (2004): Asymptotic Local Power of Pooled t-ratio Tests for Unit Roots in Panels with Fixed Effects, *Mimeo*.
- Moon, H. R., B. Perron, and P. C. B. Phillips (2006a). On the Breitung Test for Panel Unit Roots and Local Asymptotic Power. *Econometric Theory*, forthcoming.
- Moon, H. R., B. Perron, and P. C. B. Phillips (2006b). Incidental Trends and the Power of Panel Unit Root Tests. Working paper.
- Nickell, S. (1981). Biases in Dynamic Models with Fixed Effects. *Econometrica*, 49, 1417–1426.
- Phillips, P. C. B. (1989). Partially identified econometric models, *Econometric Theory* 5, 181–240.
- Phillips, P. C. B. and C. Han (2005). Gaussian Inference in AR(1) Times Series with or without Unit Root. Unpublished manuscript.
- Phillips, P. C. B. and H. R. Moon (1999). Linear Regression Limit Theory for Nonstationary Panel Data. *Econometrica*, 67(5), 1057–1111.
- Phillips, P. C. B. and V. Solo (1992). Asymptotics for Linear Processes. *The Annals of Statistics*, 20(2), 971–1001.
- Phillips, P. C. B. and D. Sul (2004). Bias in Dynamic Panel Estimation with Fixed Effects, Incidental Trends and Cross Section Dependence. Working paper.
- Ploberger, W. and P. C. B. Phillips (2002). Optimal Testing for Unit Roots in Panel Data, *Mimeo*.
- Staiger, D. and J. H. Stock (1997). Instrumental Variables Regression with Weak Instruments. *Econometrica*, 65(3), 557–586.
- Stock, J. H. and J. H. Wright (2000). GMM with Weak Identification. *Econometrica*, 68(5), 1055–1096.