

Old Dominion University
ODU Digital Commons

Rehabilitation Sciences Faculty Publications

Rehabilitation Sciences

2020

Corticospinal Activity During A Single-Leg Stance In People With Chronic Ankle Instability

Masafumi Terada

Kyle B. Kosik

Ryan S. McCann
Old Dominion University, rmccann@odu.edu

Colin Drinkard

Phillip A. Gribble

Follow this and additional works at: https://digitalcommons.odu.edu/pt_pubs



Part of the [Musculoskeletal System Commons](#), [Rehabilitation and Therapy Commons](#), and the [Sports Sciences Commons](#)

Original Publication Citation

Terada, M., Kosik, K. B., McCann, R. S., Drinkard, C., & Gribble, P. A. (2020). Corticospinal activity during a single-leg stance in people with chronic ankle instability. *Journal of Sport and Health Science*.
<https://doi.org/http://dx.doi.org/10.1016/j.jshs.2020.08.008>

This Article is brought to you for free and open access by the Rehabilitation Sciences at ODU Digital Commons. It has been accepted for inclusion in Rehabilitation Sciences Faculty Publications by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.



Original article

Corticospinal activity during a single-leg stance in people with chronic ankle instability

Masafumi Terada^{a,*}, Kyle B. Kosik^b, Ryan S. McCann^c, Colin Drinkard^d, Phillip A. Gribble^b

^a College of Sport and Health Science, Ritsumeikan University, Kusatsu, Shiga 525-8577, Japan

^b Department of Athletic Training & Clinical Nutrition, University of Kentucky, Lexington, KY 40536-0200, USA

^c School of Rehabilitation Sciences, Old Dominion University, Norfolk, VA 23529, USA

^d Westlake High School Athletics, Austin, TX 78746, USA

Received 22 January 2020; revised 22 May 2020; accepted 22 July 2020

2095-2546/© 2020 Published by Elsevier B.V. on behalf of Shanghai University of Sport. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Abstract

Purpose: The aim of the study was to determine whether corticospinal excitability and inhibition of the tibialis anterior during single-leg standing differs among individuals with chronic ankle instability (CAI), lateral ankle sprain copers, and healthy controls.

Methods: Twenty-three participants with CAI, 23 lateral ankle sprain copers, and 24 healthy control participants volunteered. Active motor threshold (AMT), normalized motor-evoked potential (MEP), and cortical silent period (CSP) were evaluated by transcranial magnetic stimulation while participants performed a single-leg standing task.

Results: Participants with CAI had significantly longer CSP at 100% of AMT and lower normalized MEP at 120% of AMT compared to lateral ankle sprain copers (CSP_{100%}: $p = 0.003$; MEP_{120%}: $p = 0.044$) and controls (CSP_{100%}: $p = 0.041$; MEP_{120%}: $p = 0.006$).

Conclusion: This investigation demonstrated altered corticospinal excitability and inhibition of the tibialis anterior during single-leg standing in participants with CAI. Further research is needed to examine the effects of corticospinal maladaptations to motor control of the tibial anterior on postural control performance in those with CAI.

Keywords: Ankle sprain; Joint instability; Motor cortex; Postural control

1. Introduction

The most prevalent musculoskeletal injury is the lateral ankle sprain.^{1–3} Up to 75% of lateral ankle sprainers will suffer from recurrent injury and lifelong residual symptoms (i.e., perceived instability),^{4–6} commonly referred to as chronic ankle instability (CAI).⁷ The long-term negative consequences of CAI are evident because patients with CAI continue to experience persistent disability for more than 7 years after an initial lateral ankle sprain,⁴ limiting their physical activity⁸ and leading to decreased health-related quality of life.^{4,9,10} Furthermore, recent evidence has shown that CAI subsequently accelerates ankle joint cartilage degeneration.^{11–13} Therefore, CAI is a pathological condition that could negatively impact long-term joint and general health.

Sensorimotor maladaptations associated with CAI can manifest into self-reported functional disability and threaten quality of life.^{14–16} Optimal neuromuscular control is achieved with proper integration of the sensory inputs from somatosensory structures and motor outputs from the central nervous system.^{17–19} Although sensory dysfunction following an initial lateral ankle sprain may contribute to functional limitations associated with CAI,²⁰ it has been proposed that alterations in supraspinal motor control may also compromise physical and self-reported function,^{21–25} as evidenced by decreased corticospinal excitability of the fibularis longus^{14,26} and tibialis anterior,²⁶ as well as compensatory muscle activation patterns of proximal musculature in patients with CAI.²⁷ Furthermore, previous investigations have shown the inefficiency of the central nervous system in the controlling reflexive response of the ankle stabilizers in those with CAI.²⁸

Proper efferent signaling from the corticospinal system is important for proper muscle recruitment and movement

Peer review under responsibility of Shanghai University of Sport.

*Corresponding author: M. Terada

E-mail address: mterada@fc.ritsumei.ac.jp (M. Terada).

<https://doi.org/10.1016/j.jshs.2020.08.008>

corrections,^{29,30} and the primary motor cortex acts in an executive role for neuromuscular control.³¹ Although motor outputs from corticospinal pathways are most often measured during a seated position using transcranial magnetic stimulation (TMS),^{14,32} it lacks a sense of functionality and limits our understanding as to how altered corticospinal excitability and inhibition are associated with CAI during a postural task. TMS measures may depend on change in body position and the type of task;³³ therefore, the findings of altered corticospinal excitability in previous studies¹⁴ may not adequately reflect neuromuscular function during a balance task. Previous studies have attempted to examine the descending corticospinal pathways during a double-leg stance task,^{29,30} but only in healthy control participants. Therefore, there is a lack of information on how the corticospinal excitability and inhibition of the ankle stabilizers are affected in individuals with CAI during a challenging postural task. Better understanding of the link between CAI and corticospinal excitability during a balance task will aid in the understanding of how corticospinal deficits are associated with CAI, leading to improved clinical interventions for CAI.

There is a known cohort of individuals with histories of lateral ankle sprain, but there is an absence of CAI-related characteristics, such as ankle dysfunction, complaints of perceived instability, and episodes of giving-way. The populations who do not develop CAI after a lateral ankle sprain have been categorized as lateral ankle sprain copers.^{34–40} Lateral ankle sprain copers restore sensorimotor control and functional levels similar to a population with no history of lateral ankle sprains.⁴¹ Therefore, a recent study⁴¹ has focused on identifying differences in characteristics between lateral ankle sprain copers and patients with CAI in order to understand the underlying mechanism that elucidates why some individuals develop CAI while others do not. Understanding the coping mechanism may guide future research focused on interventions to convert patients with CAI into lateral ankle sprain copers, with the long-term goal of reducing disability and maintaining joint health.

Therefore, the purpose of this study was to determine whether corticospinal excitability and inhibition of the tibialis anterior during single-leg standing differs among individuals with CAI, lateral ankle sprain copers, and healthy controls. We hypothesized that individuals with CAI would have decreased corticospinal excitability and increased corticospinal inhibition of the tibialis anterior during single-leg standing compared to lateral ankle sprain copers and healthy controls; however, there would be no difference in our selected variables between the lateral ankle sprain copers and healthy control groups. The tibialis anterior muscle was chosen and evaluated for this study because it has a significant role in maintaining stability during a single-leg stance,^{42,43} and altered motor control patterns of the tibialis anterior during standing tasks have been documented in individuals with CAI.^{27,44–47} It is believed that deficits in tibialis anterior function can impair one's ability to accelerate the center of mass in the direction of the support limb, which may increase a risk for the contralateral ankle sprain.^{48–50} Additionally, the tibialis anterior muscle is 1 of the dynamic stabilizers in the ankle and

protects the ankle joint against injury because it eccentrically controls ankle plantar flexion that is a common mechanism of ankle injury.

2. Methods

2.1. Study design

This study was conducted with a single-blinded case-control design. A single investigator screened participants for inclusion criteria, while 2 investigators responsible for measuring corticospinal excitability and inhibition of the tibialis anterior were blinded to group membership. Participants reported to the research laboratory for a single testing session. Specifically, the second investigator (KBK) was responsible for recording and analyzing all of the primary outcome measures, whereas the primary author (MT) was responsible for coil placement during the TMS testing. We estimated a sample size of 20 participants in each group (60 total) from corticospinal excitability and inhibition data from previous studies,^{14,25,32,51} a predetermined α level of 0.05, and an estimated power of 0.80.

2.2. Participants

Seventy physically active participants from the university community volunteered for the current study (Table 1). Participants' physical activity levels were determined by using the Godin Leisure-Time Exercise Questionnaire, and being physically active was defined as having a Godin Leisure-Time score of 24 or above.⁵² All participants read and signed an informed written consent approved by the University of Kentucky Institutional Review Board prior to the study. All participants had (1) no diagnosed balance or vestibular disorders, (2) no history of self-reported low back pain, (3) no history of surgery in the lower extremity, (4) no history of a concussion in the past 6 months, and (5) no history of any self-reported musculoskeletal or neurovascular injuries or disorders in the lower extremities in the previous 2 years other than lateral ankle sprains. All participants met additional inclusion criteria for TMS in accordance with the TMS safety guidelines outlined by the National Institutes of Neurological Disorders and Stroke.⁵³

After enrolling the participants and screening self-reported questionnaires, the participants were initially separated by previous history of lateral ankle sprain. Participants without histories of lateral ankle sprain were placed in the healthy control group. Participants reporting a previous history of lateral ankle sprain were screened and included in the CAI group based on the guidelines endorsed by the International Ankle Consortium.⁵⁴ The specific inclusion criteria for participants with CAI included: (1) a previous history of at least 1 significant lateral ankle sprain resulting in swelling, pain, and temporary loss of function; (2) a history of feelings of giving-way at least twice in the past 6 months; (3) ongoing perceived ankle instability and dysfunction during daily activities; and (4) a score of ≥ 5 on the Ankle Instability Instrument (AI), ≥ 11 on the Identification of Functional Ankle

Table 1
Demographic and ankle injury characteristics for chronic ankle instability (CAI), Coper, and control groups (mean \pm SD).

	CAI	Coper	Control
<i>n</i>	23 (3 male, 20 female)	23 (6 male, 17 female)	24 (9 male, 15 female)
Age (year)	23.17 \pm 3.54	24.74 \pm 5.25	21.08 \pm 1.86
Height (cm)	168.63 \pm 7.26	166.79 \pm 8.29	169.17 \pm 10.18
Body mass (kg)	73.01 \pm 15.32	68.34 \pm 16.08	66.99 \pm 13.04
Body mass index (kg/m ²)	25.64 \pm 4.94	24.35 \pm 4.02	23.27 \pm 2.91
Godin Leisure-time Exercise Questionnaire	58.05 \pm 24.18	62.55 \pm 46.97	65.32 \pm 27.80
AII*	6.35 \pm 1.72	2.39 \pm 0.94	0.00
IdFAI*	19.87 \pm 4.44	4.70 \pm 3.71	0.00
CAIT*	14.43 \pm 5.34	27.09 \pm 4.07	30.00 \pm 0.00
Number of lateral ankle sprains	4.09 \pm 3.71	2.04 \pm 1.87	–
Time since last ankle sprain (month)	42.00 \pm 37.84	99.91 \pm 74.34	–
Number of giving-way episodes in past 6 months	9.13 \pm 14.57 (range: 2–72)	0.13 \pm 0.34 (range:0–1)	–
Modified physical activity because of CAI	Yes: 12; No:11	Yes: 0; No: 23	–
Using protective ankle devices	Yes: 6; No: 17	Yes: 2; No: 21	Yes: 2; No: 22
Feel a risk for injury when playing sports	Yes: 15; No:8	Yes: 1; No:22	–
Concerned with surrounding environment (i.e., walking on icy surfaces)	Yes: 14; No:9	Yes: 1; No: 22	–
Have received rehabilitation from an allied health care professional	Yes: 12; No: 11	Yes: 7; No: 16	–

* $p < 0.05$, significant differences in AII, IdFAI, CAIT between groups.

Abbreviations: AII = Ankle Instability Instrument; IdFAI = Identification of Functional Ankle Instability; CAIT = Cumberland Ankle Instability Tool.

Instability, and ≤ 24 on the Cumberland Ankle Instability Tool.⁵⁴ The CAI group included 23 participants. No participant with CAI had acutely sprained an ankle in the previous 3 months before testing. If a participant had a history of bilateral ankle injury, we measured the limb with the greatest amount of self-reported instability. A test limb for the control group was randomly selected.

Participants who did not meet the CAI inclusion criteria were then screened according to the lateral ankle sprain coper inclusion criteria. Twenty-six participants were included in the lateral ankle sprain coper group, defined as participants with a history of lateral ankle sprains but no reported episodes of giving-way, perceived instability, loss of function, or modification of physical activity. Lateral ankle sprain copers had to (1) have no history of feelings of giving-way fewer than twice in the past 12 months; (2) have no ongoing perceived ankle instability and dysfunction during daily and physical activities; and (3) score < 5 on the AII and < 11 on the Identification of Functional Ankle Instability. We did not use the Cumberland Ankle Instability Tool scores to classify participants as lateral ankle sprain copers because it has been reported that the cutoff score of ≥ 25 resulted in a high rate of false-positives in the lateral ankle sprain coper group.⁵⁵ No participant in the lateral ankle sprain coper group had acutely sprained an ankle in the previous 12 months before testing.

The control group included 24 participants. The specific inclusion criteria for participants in the control group were (1) no history of lateral ankle sprain; (2) a score of 0 on both the AII and Identification of Functional Ankle Instability; and (3) a score of 30 on CAIT. Means and standard deviations for participant demographics and ankle injury information are found in Table 1.

2.3. Experimental procedures

Participants reported to the research laboratory for the single TMS testing session. Two 10 mm pre-gelled Ag/AgCl electromyography (EMG) electrodes (EL503; BIOPAC Systems, Goleta, CA, USA) were placed 1.75 mm apart over the midpoint of the tibialis anterior muscle belly, and a ground electrode was placed over the contralateral medial malleolus.⁵⁶ The areas were shaved, abraded with fine sandpaper, and cleaned with isopropyl alcohol wipes prior to electrode placement. EMG signals were recorded at a sampling rate of 2000 Hz (gain set at 1000, common mode rejection ratio = 110 dB, input impedance = 1 MOhms) (EMG100C; BIOPAC Systems). A 16-bit converter (MP150; BIOPAC Systems) was used to process analog-to-digital signal conversion. EMG signals were filtered with a high pass of 10 Hz and low pass of 500 Hz. EMG, and stimulation signals were visualized through AcqKnowledge 4.1 Software (BIOPAC Systems).

Participants wore a Lycra swim cap (Sprint Aquatics, Rothhammer International, San Luis Obispo, CA, USA) that helps to mark the approximate primary motor cortex location so that the location of a double-cone coil did not change during the experiment. On the cap, a standard 1 cm² dot grid was marked, and 2 straight lines were drawn vertically in the sagittal and frontal planes. These lines intersected over the vertex of the skull: one line separating the hemispheres sagittally and the other connecting the apexes of the ears bisecting the other line. The double-cone coil was positioned over the intersected lines and dot grids.

A Magstim 200 (Magstim, Whitland, Wales, UK) was used to produce a magnetic stimulus (max 2.0 Tesla) over the primary motor cortex contralateral to the test limb. During the TMS testing, participants stood in a barefoot single-leg stance on the middle of a force platform (AccuSway Plus, AMTI,




Fig. 1. Experiment set-up.

Watertown, MA, USA) integrated with Balance Clinic software (AMTI), kept their hands on the waist, and stared at an X in front of them while keeping their foot flat on the force platform (Fig. 1). An investigator held the TMS coil to secure it on the participant's head while providing the magnetic stimuli. We used the force platform to visually monitor center-of-pressure displacement to ensure that participants remained still while in position and that the investigator holding the coil did not influence participant's performance during the TMS testing. The magnetic stimulation was provided approximately 3 s after participants assumed the single-leg balance position. To minimize the potential effect of fatigue, a minimum of 30 s rest was provided between each trial. The trial was discarded and repeated if (1) the nontested limb made contact with the force platform; (2) the nontested limb made contact with the stance limb; (3) the participant took a step with the stance limb; (4) the participant removed the hands from the chest; or (5) the participant abducted the hip or laterally flexed the trunk into > 30 degrees. To determine the optimal stimulating point location, a series of magnetic stimuli at 50% of the maximal stimulator output was delivered at a number of locations on the grid until the largest and most consistent motor-evoked potentials (MEPs) were observed.⁵⁷ Once the optimal stimulating location was detected, an investigator stabilized the TMS coil at this location for all subsequent tests.

The active motor threshold (AMT) was assessed by using the method previously described.⁵⁷ Briefly, the MEP threshold was calculated by determining the average peak amplitude of the background EMG signal collected while participants performed the single-leg balance without magnetic stimulus.⁵⁷ The cut-off threshold was set 2 SDs above this EMG amplitude.⁵⁷ The AMT was determined as the lowest stimulator intensity required to elicit at least four of 8 MEPs whose peak-to-peak amplitudes exceeded this MEP threshold.^{57,58} A higher AMT indicates decreased corticospinal excitability because a greater stimulator intensity is required to excite the neurons within the corticospinal pathway.⁵⁹ Once AMT was established, 8 stimuli were delivered at intensities of 100% of AMT and 120% of AMT, and peak-to-peak MEP amplitudes were recorded for each trial. The amplitude of the 8 MEPs at the

intensities of 100% of AMT and 120% of AMT were averaged and normalized to mean amplitudes of background EMG activity that was recorded 50 ms prior to a stimulus in order to minimize the effect of background EMG on the MEP amplitude.⁶⁰ The MEP amplitude provides an estimate of the overall excitability of the corticospinal tract.⁵⁹

Cortical silent period (CSP) can be used to evaluate inhibition within the corticospinal pathway, which may be mediated through γ -aminobutyric acid-B receptors.⁶¹ The CSP period of the tibialis anterior was obtained at intensities of 100% and 120% of AMT and was measured as the time from the end of the MEP to a return of the mean EMG signal plus 2 times the standard deviation of the baseline (pre-stimulus) EMG signal.⁶² The CSP was reported in ms. A longer CSP indicates greater corticospinal inhibition to the tibialis anterior.

2.4. Statistical analysis

All statistical analyses were performed using SPSS software (Version 23.0; IBM, Armonk, NY, USA). Significance was set at $p < 0.05$ for all analyses. We could not collect the CSP data from a few of participants (9 participants at 100% intensities of AMT and 10 participants at 120% intensities of AMT) because the EMG signal during the expected CSP time window was obscured, resulting in the landmark to define CSP being absent. Therefore, 13% of CSP at 100% intensities of AMT (9/70) and 14% of CSP at 120% intensities of AMT (10/70) in the data set were imputed using the multiple imputation method with 5 repetitions.⁶³

Based on an analysis of the data using a Kolmogorov-Smirnov Z test for normality,⁶⁴ we found that MEP measures were not normally distributed ($p < 0.05$). Therefore, a separate independent-samples Kruskal-Wallis test was performed to compare MEP measures among the CAI, lateral ankle sprain copers, and control groups. A Mann-Whitney U test was conducted as *post hoc* analysis in the case of statistical significance. Last, r effect sizes (Z/\sqrt{n}) were calculated to determine the magnitude of significant group differences. Effect sizes were interpreted as small ($r = 0.10$ – 0.29), moderate ($r = 0.30$ – 0.49), large ($r = 0.50$ – 0.69), or very large ($r > 0.70$).⁶⁵

Other outcome measures (AMT, CSP at 100% and 120% intensities of AMT, and background EMG) were compared among the CAI, lateral ankle sprain copers, and control groups using separate one-way analysis of variance models. Bonferroni *post hoc* testing was conducted as needed. Last, Cohen d effect sizes using the pooled standard deviations were calculated for outcome measures that were normally distributed,⁶⁶ along with 95% confidence intervals, to determine the magnitude of difference in outcome variables between groups. The strength of effect sizes was interpreted as weak ($d < 0.40$), moderate ($d = 0.40$ – 0.79), or large ($d \geq 0.80$).⁶⁶

3. Results

Between-group differences were observed for CSP at 100% intensities of AMT ($p = 0.003$) (Fig. 2A) and MEP at 120% intensities of AMT ($p = 0.021$) (Fig. 2B) of the tibialis anterior.




Fig. 2. Corticospinal excitability and inhibition results (mean ± SD). (A) Quantified at 100% of active motor threshold. (B) Quantified at 120% of active motor threshold. AMT = active motor threshold; CAI = chronic ankle instability; CSP = cortical silent period; MEP = motor evoked potential.

Post hoc analysis revealed that participants with CAI had significantly longer CSP at 100% intensities of AMT and lower normalized MEP at 120% intensities of AMT compared to lateral ankle sprain copers (CSP at 100% intensities of AMT: $p=0.003$; MEP at 120% intensities of AMT: $p=0.044$) and controls (CSP at 100% intensities of AMT: $p=0.041$; MEP at 120% intensities of AMT: $p=0.006$), which were supported by moderate to strong effect sizes (Table 2) (Figs. 2 and 3). There were no differences between the lateral ankle sprain copers and control groups in CSP at 100% intensities of AMT ($p=1.000$) and MEP at 120% intensities of AMT ($p=0.774$).

AMT was not significantly different among the groups (CAI = $35.57\% \pm 9.52\%$, lateral ankle sprain copers = $36.70\% \pm 11.59\%$, control = $35.88\% \pm 11.54\%$; $p=0.936$). There were no differences in CSP at 120% intensities of AMT ($p=0.975$) and MEP at 100% intensities of AMT ($p=0.425$) between the CAI, lateral ankle sprain copers, and control groups (Fig. 2A and 2B). For background EMG, no differences were observed between groups (CAI = 0.28 ± 0.20 mV, lateral ankle sprain copers = 0.27 ± 0.20 mV, control = 0.17 ± 0.13 mV; $p=0.07$). All effect size comparisons for AMT, CSP at

Table 2

Pairwise comparisons and effect sizes for outcome measures that were not normally distributed.

	Mann-Whitney U	Significance	Effect size
Normalized MEP at 100% of AMT			
CAI vs. control	$Z = -0.80$	$p = 0.440$	$r = 0.11$
CAI vs. LAS copers	$Z = -0.44$	$p = 0.660$	$r = 0.07$
LAS copers vs. control	$Z = -0.29$	$p = 0.774$	$r = 0.05$
Normalized MEP at 120% of AMT			
CAI vs. control	$Z = -2.74$	$p = 0.006$	$r = 0.40$
CAI vs. LAS copers	$Z = -2.01$	$p = 0.044$	$r = 0.32$
LAS copers vs. control	$Z = -0.27$	$p = 0.790$	$r = 0.04$

Abbreviations: AMT = active motor threshold; CAI = chronic ankle instability; LAS = lateral ankle sprain; MEP, motor-evoked potential.




Fig. 3. Cohen d effect sizes and associated 95% confidence intervals. CAI = chronic ankle instability; CSP = cortical silent period.

120% intensities of AMT, and MEP at 100% intensities of AMT were weak, with 95% confidence intervals that crossed 0 (Table 2) (Fig. 3).

4. Discussion

The association between CAI and altered neuromuscular control of the tibialis anterior during sophisticated tasks incorporating postural control and gait has been well documented;^{27,44,45} however, our understanding of the neurophysiological mechanism underlying this alteration is undiscovered. Researchers have hypothesized that altered supraspinal organization of the central nervous system may influence motor control in individuals with CAI.²¹ Our study examined differences in corticospinal excitability and inhibition during a single-leg stance between individuals with and without CAI. We observed that participants with CAI demonstrated longer CSP at 100% of AMT and lower MEP at 120% of AMT in the tibialis anterior during the single-leg standing compared to lateral ankle sprain copers and controls, supporting the previous speculation of centrally mediated changes within corticospinal pathways.¹⁴ These results indicate that participants with CAI may have an increase in the sensitivity

of inhibitory intermediate neurons and a decrease in the sensitivity of excitatory neurons in the corticospinal pathway. To our knowledge, we are the first to investigate corticospinal excitability and inhibition during a balance task in a CAI population.

4.1. Potential corticospinal maladaptations

Declines in sensorimotor control during a single-leg stance have been consistently observed in individuals with CAI.^{41,67} Van Deun et al.⁴⁵ observed altered muscle activation patterns of the tibialis anterior during the transition from double-leg to single-leg standing in individuals with CAI. Furthermore, individuals with CAI have demonstrated the inability to control spinal reflex excitability of the ankle stabilizers during postural transitions,²⁸ potentially suggesting an altered supraspinal mechanism within the central nervous system. Impairments in the somatosensory system at the damaged ankle joint due to the presence of CAI may decrease sensory inputs to the central nervous system, which could create a chronic central-mediated alteration in neuromuscular function.²¹ Altered corticospinal adaptations in the motor control of the tibialis anterior observed in our study indicate that individuals with CAI may encounter more difficulty in controlling the tibialis anterior muscle during single-leg stance, indicating that movement patterns are altered during a balance task in those with CAI.

4.2. Corticospinal inhibition

Increased CSP in the tibialis anterior muscle of participants with CAI at the intensity of 100% of AMT indicates greater corticospinal inhibition to the corresponding muscle.⁶¹ However, there were no differences in CSP at 120% of AMT among the CAI, lateral ankle sprain copers, and control groups. This may be attributable to the effect of TMS intensity on CSP duration. CSP duration is prolonged with higher TMS intensity.⁶⁸ At lower TMS intensity, inhibitory intermediate neurons in the corticospinal pathway may already be activated in participants with CAI. As the intensity increases, inhibitory neurons in the pathway are more activated in lateral ankle sprain copers and control participants; thus, they become more similar to those with CAI. The higher intensities likely have less effect on CSP in those with CAI because there may not be many additional inhibitory neurons left to activate.

4.3. Corticospinal excitability

Similar to our results, Needle et al.³² reported that AMT of the tibialis anterior was not different between the CAI and control groups. AMT is considered to be an outcome measure that estimates the membrane excitability of pyramidal neurons.⁵⁹ Based on findings from Needle et al.³² and our current study, individuals with CAI do not require higher magnetic stimulation in order to excite the pyramidal cells in the primary motor cortex, possibly indicating no effects of CAI on corticomotor excitability. However, we observed less normalized MEP amplitude at 120% AMT in the tibialis anterior for participants with CAI than for lateral ankle sprain copers and controls.

Decreased MEPs at 120% of AMT may indicate that the degree of motor responses in the tibialis anterior with high TMS intensity may be small during a single-leg stance in participants with CAI.⁶⁹ MEPs represent the magnitude of motor outputs induced by magnetic stimulus that is relayed to the muscles via the entire corticospinal system.³² This decrease in the percentage of outgoing motor information being delivered to the tibialis anterior through the corticospinal system is, therefore, interpreted as a decrease in the corticospinal excitability of this muscle in the CAI group.

Although normalized MEP amplitude at 120% AMT was decreased in the CAI group relative to the lateral ankle sprain copers and control groups, the amplitude of responses at 100% AMT was not different among the groups. These results indicate that the excitation of the number of a central core of tibialis anterior motor neurons that arise from the primary motor cortex may not be different at 100% of AMT among the groups. However, the higher normalized MEPs observed at 120% of AMT in the lateral ankle sprain copers and control groups may be attributed to additional recruitment of inherently less excitable neurons in the primary motor cortex. Recording MEP amplitudes at multiple intensities is considered a stimulus response, which takes into account both the response of neurons with lower and higher thresholds and the potentials increase in the MEP amplitude of the signal as the intensity of TMS increases.^{70,71} The higher TMS intensity can recruit more motor neurons that are less excitable or located farther from the center of activation by the magnetic stimulus, thereby increasing the MEP amplitude.⁷⁰ Therefore, decreased MEPs at 120% of AMT in the CAI group may indicate that a smaller portion of motor neurons in the corticospinal system could be excited by a high level of excitation to the primary cortex.

4.4. Potential coping mechanism and clinical implications

Corticospinal excitability and inhibition of the tibialis anterior in lateral ankle sprain copers more closely resembled that of control participants compared to participants with CAI. This provides a potentially significant explanation for the differences between lateral ankle sprain copers and CAI patients. Corticospinal outcomes observed during the single-leg stance in lateral ankle sprain copers is suggestive of adequate motor outputs delivered to the tibialis anterior through the corticospinal pathway following lateral ankle sprain. This may help lateral ankle sprain copers to make postural control adjustments and maintain ankle stability. Optimal postural control requires proper transfer of motor information to the postural muscles from the corticospinal system in order to achieve proper muscle recruitment during postural control.¹⁷ Researchers have shown better postural control performance in lateral ankle sprain copers than in participants with CAI.⁴¹ It is possible that lateral ankle sprain copers could successfully reorganize corticospinal activity following the ankle injury, leading to appropriate postural-control strategies. Our findings suggest that treatment and rehabilitation aimed at improving postural control in patients with CAI might include manipulation of

corticospinal excitability and inhibition. However, because of our cross-sectional, retrospective study design, it remains unknown exactly how lateral ankle sprain copers have adopted neuromuscular strategies without perceived instability following lateral ankle sprains. Thus, future prospective studies should explore the time-course of the corticospinal changes after an initial lateral ankle sprain and the response to interventions during the rehabilitation process.

Understanding the effect of ankle joint injury on the corticospinal pathway may be critical in the advancement of our knowledge of neural mechanisms causing function limitations. However, we did not include spatiotemporal measures of a single-leg stance performance in the current investigation. We do not know whether the corticospinal alterations observed in the tibialis anterior actually influence postural control and movement patterns during the single-leg stance. Therefore, future studies should incorporate these measures to further determine the level of CAI's influence on corticospinal pathway alterations and how they influence functional measures.

4.5. Limitations

We acknowledge that more specific neurophysiological mechanisms responsible for the observed corticospinal outcomes cannot be determined using only the single-pulse TMS paradigm. The MEP and CSP can be modulated at both the cortical and spinal levels,^{59,70} and we did not quantify the Hoffmann reflex (spinal reflex excitability) of the tibialis anterior during the task. Thus, it is difficult to identify more specific local modulation in the cortical or spinal regions that contribute to CAI. A paired-pulse TMS paradigm provides additional information related to intracortical excitability and inhibition during a postural control task. Therefore, in order to parse out specifically where a change has occurred, further research is needed to investigate the association between the CAI and intracortical measures using paired-pulse TMS paradigms.

We carefully and visually monitored center-of-pressure displacement using the force platform to ensure that participants remained still, but we did not truly control postural sway direction in the single-leg stance when delivering magnetic stimulation. Previous studies have shown that postural sway direction influenced the size of MEP amplitude due to a change in the length of muscle fibers and/or spinal or subcortical modulation.^{30,72} Specifically, the tibialis anterior fascicles become shortened when swaying in the forward direction,⁴² and this shortening of the muscle alters proprioceptive feedback which, in turn, influences corticospinal excitability.^{30,72} It is possible that lateral ankle sprain copers and control participants were swaying backward at the time of stimulation, whereas those with CAI were swaying in the forward direction. However, the background EMG that was recorded 50 ms prior was not different among groups ($p = 0.07$). Therefore, the between-group differences in our selected corticospinal measures were likely to be attributed to the presence of CAI.

Although we did not acquire corticospinal variables during sitting, it is important to note that results from our study may not be representation of corticospinal characteristics assessed during sitting. Because posture and standing tasks influence TMS measures of corticospinal excitability and inhibition,³³ TMS measures obtained in the single-leg stance may be more relevant and sensitive indices of neuromuscular characteristics during functionally relevant posture compared to measurements during sitting.⁷³ If background activation levels are matched between standing and sitting positions, corticospinal characteristics may not be different between standing and sitting.^{33,73} However, it is not clear how the presence of CAI alters posture-related effects on corticospinal excitability and inhibition of the tibialis anterior muscle. Therefore, future studies are needed to compare corticospinal excitability and inhibition between sitting and standing in a population with CAI.

Evaluation of corticospinal excitability and inhibition of the tibialis anterior muscle using TMS has been demonstrated to be reliable (intraclass correlation coefficient = 0.89–0.98).⁷⁴ However, we did not assess intra- and inter-rater reliability of the selected TMS variables during the standing condition. Furthermore, our study included only the tibialis anterior muscle because it is an important ankle and postural stabilizer. The fibularis longus muscles and soleus also have vital roles for ankle stabilization and maintenance of upright stance.⁷⁵ Further investigation is needed to include quantification of corticospinal excitability and inhibition of other ankle joint stabilizers during a balance task. Last, in our study we did not assess the presence of mechanical instability in the CAI group, and mechanical instability following an initial ankle sprain remains an important factor in understanding the CAI paradigm.

5. Conclusion

Our study examined differences in corticospinal excitability and inhibition during single-leg stance between individuals with and without CAI. Participants with CAI demonstrated longer CSP at 100% of AMT and lower MEP at 120% of AMT in the tibialis anterior during the single-leg balance compared to lateral ankle sprain copers and controls. Our investigation found that corticospinal excitability and inhibition in the tibialis anterior muscle was altered in CAI participants during the single-leg stance, illustrating centrally mediated changes within corticospinal pathways. Altered corticospinal adaptations to motor control of the tibialis anterior indicate that those with CAI may encounter more difficulty in controlling the tibialis anterior muscle during a single-leg stance, potentially influencing postural control performance in those with CAI. Future studies should incorporate alternative analyses to further determine the level of influence of the motor control via the corticospinal pathway on postural control performance in CAI populations.

Acknowledgment

The authors thank Dr. Abbey Thomas, PhD, ATC, for assistance with aspects of the study design.

Authors' contributions

MT, KBK, and PAG participated in the design of the study and contributed to data collection and data reduction/analysis and to the interpretation of results; RSM participated in the design of the study and interpretation of results; and CD participated in the design of the study and contributed to data collection. All authors contributed to the manuscript writing. All authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Competing interests

The authors declare that they have no competing interests.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.jshs.2020.08.008.

References

- Doherty C, Delahunt E, Caulfield B, Hertel J, Ryan J, Bleakley C. The incidence and prevalence of ankle sprain injury: a systematic review and meta-analysis of prospective epidemiological studies. *Sports Med* 2014;**44**:123–40.
- Fong DT, Man CY, Yung PS, Cheung SY, Chan KM. Sport-related ankle injuries attending an accident and emergency department. *Injury* 2008;**39**:1222–7.
- Hootman J, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: summary and recommendations for injury prevention initiatives. *J Athl Train* 2007;**42**:311–9.
- Gribble PA, Bleakley CM, Caulfield BM, Docherty CL, Fourchet F, Fong DT, et al. Evidence review for the 2016 international ankle consortium consensus statement on the prevalence, impact and long-term consequences of lateral ankle sprains. *Br J Sports Med* 2016;**50**:1496–505.
- Konradsen L, Bech L, Ehrenbjerg M, Nickelsen T. Seven years follow-up after ankle inversion trauma. *Scand J Med Sci Sports* 2002;**12**:129–35.
- Anandacoomarasamy A, Barnsley L. Long term outcomes of inversion ankle injuries. *Br J Sports Med* 2005;**39**:e14. doi:10.1136/bjism.2004.011676.
- Delahunt E, Coughlan GF, Caulfield B, Nightingale EJ, Lin CW, Hiller CE. Inclusion criteria when investigating insufficiencies in chronic ankle instability. *Med Sci Sports Exerc* 2010;**42**:2106–21.
- Hubbard-Turner T, Turner MJ. Physical activity levels in college students with chronic ankle instability. *J Athl Train* 2015;**50**:742–7.
- Arnold BL, Wright CJ, Ross SE. Functional ankle instability and health-related quality of life. *J Athl Train* 2011;**46**:634–41.
- Houston MN, Van Lunen BL, Hoch MC. Health-related quality of life in individuals with chronic ankle instability. *J Athl Train* 2014;**49**:758–63.
- Golditz T, Steib S, Pfeifer K, Uder M, Gelse K, Janka R, et al. Functional ankle instability as a risk factor for osteoarthritis: using t2-mapping to analyze early cartilage degeneration in the ankle joint of young athletes. *Osteoarthr Cartilage* 2014;**22**:1377–85.
- Hirose K, Murakami G, Minowa T, Kura H, Yamashita T. Lateral ligament injury of the ankle and associated articular cartilage degeneration in the talocrural joint: anatomic study using elderly cadavers. *J Orthop Sci* 2004;**9**:37–43.
- Valderrabano V, Hintermann B, Horisberger M, Fung TS. Ligamentous posttraumatic ankle osteoarthritis. *Am J Sports Med* 2006;**34**:612–20.
- Pietrosimone BG, Gribble PA. Chronic ankle instability and corticomotor excitability of the fibularis longus muscle. *J Athl Train* 2012;**47**:621–6.
- Harkey M, McLeod MM, Terada M, Gribble PA, Pietrosimone BG. Quadratic association between corticomotor and spinal-reflexive excitability and self-reported disability in participants with chronic ankle instability. *J Sport Rehabil* 2016;**25**:137–45.
- Houston MN, Hoch JM, Gabriner ML, Kirby JL, Hoch MC. Clinical and laboratory measures associated with health-related quality of life in individuals with chronic ankle instability. *Phys Ther Sport* 2015;**16**:169–75.
- Riemann BL, Lephart SM. The sensorimotor system, part i: the physiologic basis of functional joint stability. *J Athl Train* 2002;**37**:71–9.
- McKeon PO, Wikstrom EA. Sensory-targeted ankle rehabilitation strategies for chronic ankle instability. *Med Sci Sports Exerc* 2016;**48**:776–84.
- Wikstrom EA, McKeon PO. Predicting balance improvements following stars treatments in chronic ankle instability participants. *J Sci Med Sport* 2017;**20**:356–61.
- Powell MR, Powden CJ, Houston MN, Hoch MC. Plantar cutaneous sensitivity and balance in individuals with and without chronic ankle instability. *Clin J Sport Med* 2014;**24**:490–6.
- Hertel J. Sensorimotor deficits with ankle sprains and chronic ankle instability. *Clin Sports Med* 2008;**27**:353–70.
- Kim KM, Kim JS, Cruz-Diaz D, Ryu S, Kang M, Taube W. Changes in spinal and corticospinal excitability in patients with chronic ankle instability: a systematic review with meta-analysis. *J Clin Med* 2019;**8**:1037. doi:10.3390/jcm8071037.
- Harkey M, McLeod MM, Terada M, Gribble PA, Pietrosimone BG. Quadratic association between corticomotor and spinal-reflexive excitability and self-reported disability in participants with chronic ankle instability. *J Sport Rehabil* 2016;**25**:137–45.
- Kosik KB, Terada M, Drinkard CP, McCann RS, Gribble PA. Potential corticomotor plasticity in those with and without chronic ankle instability. *Med Sci Sports Exerc* 2016;**49**:141–9.
- Terada M, Bowker S, Thomas AC, Pietrosimone B, Hiller CE, Gribble PA. Corticospinal excitability and inhibition of the soleus in individuals with chronic ankle instability. *PM R* 2016;**8**:1090–6.
- Nanbancha A, Tretriluxana J, Limroongreungrat W, Sinsurin K. Decreased supraspinal control and neuromuscular function controlling the ankle joint in athletes with chronic ankle instability. *Eur J Appl Physiol* 2019;**119**:2041–52.
- Rios JL, Gorges AL, dos Santos MJ. Individuals with chronic ankle instability compensate for their ankle deficits using proximal musculature to maintain reduced postural sway while kicking a ball. *Hum Mov Sci* 2015;**43**:33–44.
- Kim KM, Ingersoll CD, Hertel J. Altered postural modulation of Hoffmann reflex in the soleus and fibularis longus associated with chronic ankle instability. *J Electromyogr Kinesiol* 2012;**22**:997–1002.
- Solopova IA, Kazennikov OV, Deniskina NB, Levik YS, Ivanenko YP. Postural instability enhances motor responses to transcranial magnetic stimulation in humans. *Neurosci Lett* 2003;**337**:25–8.
- Tokuno CD, Taube W, Cresswell AG. An enhanced level of motor cortical excitability during the control of human standing. *Acta Physiol (Oxf)* 2009;**195**:385–95.
- Jacobs JV, Horak FB. Cortical control of postural responses. *J Neural Transm (Vienna)* 2007;**114**:1339–48.
- Needle AR, Palmer JA, Kesar TM, Binder-MacLeod SA, Swanik CB. Brain regulation of muscle tone in healthy and functionally unstable ankles. *J Sport Rehabil* 2013;**22**:202–11.
- Soto O, Valls-Sole J, Shanahan P, Rothwell J. Reduction of intracortical inhibition in soleus muscle during postural activity. *J Neurophysiol* 2006;**96**:1711–7.
- Wikstrom EA, Brown CN. Minimum reporting standards for copers in chronic ankle instability research. *Sports Med* 2014;**44**:251–68.
- Bowker S, Terada M, Thomas AC, Pietrosimone BG, Hiller CE, Gribble PA. Neural excitability and joint laxity in chronic ankle instability, copers, and control groups. *J Athl Train* 2016;**51**:336–43.
- Croy T, Saliba SA, Saliba E, Anderson MW, Hertel J. Differences in lateral ankle laxity measured via stress ultrasonography in individuals with chronic ankle instability, ankle sprain copers, and healthy individuals. *J Orthop Sports Phys Ther* 2012;**42**:593–600.
- Doherty C, Bleakley C, Hertel J, Caulfield B, Ryan J, Delahunt E. Locomotive biomechanics in persons with chronic ankle instability and lateral ankle sprain copers. *J Sci Med Sport* 2016;**19**:524–30.

38. Liu W, Jain TK, Santos M, Heller D, Hiller C. Important issues concerning use of term “copers” in chronic ankle instability research. *Sports Med* 2014;**44**:1775–6.
39. Wikstrom EA, Tillman MD, Chmielewski TL, Cauraugh JH, Naugle KE, Borsa P. Discriminating between copers and people with chronic ankle instability. *J Athl Train* 2012;**47**:136–42.
40. McCann RS, Crossett ID, Terada M, Kosik KB, Bolding BA, Gribble PA. Hip strength and star excursion balance test deficits of patients with chronic ankle instability. *J Sci Med Sport* 2017;**20**:992–6.
41. Wikstrom EA, Fournier KA, McKeon PO. Postural control differs between those with and without chronic ankle instability. *Gait Posture* 2010;**32**:82–6.
42. Day JT, Lichtwark GA, Cresswell AG. Tibialis anterior muscle fascicle dynamics adequately represent postural sway during standing balance. *J Appl Physiol (1985)* 2013;**115**:1742–50.
43. Di Giulio I, Maganaris CN, Baltzopoulos V, Loram ID. The proprioceptive and agonist roles of gastrocnemius, soleus and tibialis anterior muscles in maintaining human upright posture. *J Physiol* 2009;**587**:2399–416.
44. Feger MA, Donovan L, Hart JM, Hertel J. Lower extremity muscle activation during functional exercises in patients with and without chronic ankle instability. *PM R* 2014;**6**:602–11.
45. Van Deun S, Staes FF, Stappaerts KH, Janssens L, Levin O, Peers KKH. Relationship of chronic ankle instability to muscle activation patterns during the transition from double-leg to single-leg stance. *Am J Sports Med* 2007;**35**:274–81.
46. Lofvenberg R, Karrholm J, Sundelin G, Ahlgren O. Prolonged reaction time in patients with chronic lateral instability of the ankle. *Am J Sports Med* 1995;**23**:414–7.
47. Wikstrom EA, Bishop MD, Inamdar AD, Hass CJ. Gait termination control strategies are altered in chronic ankle instability subjects. *Med Sci Sports Exerc* 2010;**42**:197–205.
48. Sousa ASP, Silva M, Gonzalez S, Santos R. Bilateral compensatory postural adjustments to a unilateral perturbation in subjects with chronic ankle instability. *Clin Biomech (Bristol, Avon)* 2018;**57**:99–106.
49. Sousa ASP, Valente I, Pinto A, Soutelo T, Silva M. Short and medium latency responses in participants with chronic ankle instability. *J Athl Train* 2018;**53**:679–86.
50. Mitchell A, Dyson R, Hale T, Abraham C. Biomechanics of ankle instability. Part 1: reaction time to simulated ankle sprain. *Med Sci Sports Exerc* 2008;**40**:1515–21.
51. McLeod MM, Gribble PA, Pietrosimone BG. Chronic ankle instability and neural excitability of the lower extremity. *J Athl Train* 2015;**50**:847–53.
52. Godin G. The Godin–Shephard leisure–time physical activity questionnaire. *Health Fit J Can* 2011;**4**:18–22.
53. Rossi S, Hallett M, Rossini PM, Pascual-Leone A. Safety of TMS Consensus Group. Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clin Neurophysiol* 2009;**120**:2008–39.
54. Gribble PA, Delahunt E, Bleakley CM, Caulfield B, Docherty CL, Fong DT, et al. Selection criteria for patients with chronic ankle instability in controlled research: a position statement of the international ankle consortium. *J Athl Train* 2014;**49**:121–7.
55. Wright CJ, Arnold BL, Ross SE, Linens SW. Recalibration and validation of the Cumberland ankle instability tool cutoff score for individuals with chronic ankle instability. *Arch Phys Med Rehabil* 2014;**95**:1853–9.
56. Palmieri RM, Ingersoll CD, Hoffman MA. The Hoffmann reflex: methodologic considerations and applications for use in sports medicine and athletic training research. *J Athl Train* 2004;**39**:268–77.
57. Stevens-Lapsley JE, Thomas AC, Hedgecock JB, Kluger BM. Corticospinal and intracortical excitability of the quadriceps in active older and younger healthy adults. *Arch Gerontol Geriatr* 2013;**56**:279–84.
58. Temesi J, Gruet M, Rupp T, Verges S, Millet GY. Resting and active motor thresholds versus stimulus-response curves to determine transcranial magnetic stimulation intensity in quadriceps femoris. *J Neuroeng Rehabil* 2014;**11**:40. doi:10.1186/1743-0003-11-40.
59. Groppa S, Oliviero A, Eisen A, Quartarone A, Cohen LG, Mall V, et al. A practical guide to diagnostic transcranial magnetic stimulation: report of an ifcn committee. *Clin Neurophysiol* 2012;**123**:858–82.
60. Knikou M, Hajela N, Mummidisetty CK. Corticospinal excitability during walking in humans with absent and partial body weight support. *Clin Neurophysiol* 2013;**124**:2431–8.
61. Werhahn KJ, Kunesch E, Noachtar S, Benecke R, Classen J. Differential effects on motorcortical inhibition induced by blockade of gaba uptake in humans. *J Physiol* 1999;**517**:591–7.
62. Garvey MA, Ziemann U, Becker DA, Barker CA, Bartko JJ. New graphical method to measure silent periods evoked by transcranial magnetic stimulation. *Clin Neurophysiol* 2001;**112**:1451–60.
63. Schlomer GL, Bauman S, Card NA. Best practices for missing data management in counseling psychology. *J Couns Psychol* 2010;**57**:1–10.
64. Mishra P, Pandey CM, Singh U, Gupta A, Sahu C, Keshri A. Descriptive statistics and normality tests for statistical data. *Ann Card Anaesth* 2019;**22**:67–72.
65. Rosenthal JA. Qualitative descriptors of strength of association and effect size. *J Soc Serv Res* 1996;**21**:37–59.
66. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. Hillsdale, NJ: L. Erlbaum Associates; 1988.
67. McKeon PO, Hertel J. Spatiotemporal postural control deficits are present in those with chronic ankle instability. *BMC Musculoskelet Disord* 2008;**9**:76. doi:10.1186/1471-2474-9-76.
68. Kojima S, Onishi H, Sugawara K, Kirimoto H, Suzuki M, Tamaki H. Modulation of the cortical silent period elicited by single- and paired-pulse transcranial magnetic stimulation. *BMC Neurosci* 2013;**14**:43. doi:10.1186/1471-2202-14-43.
69. del Olmo MF, Reimunde P, Viana O, Acero RM, Cudeiro J. Chronic neural adaptation induced by long-term resistance training in humans. *Eur J Appl Physiol* 2006;**96**:722–8.
70. Hallett M. Transcranial magnetic stimulation: a primer. *Neuron* 2007;**55**:187–99.
71. Ridding MC, Rothwell JC. Stimulus/response curves as a method of measuring motor cortical excitability in man. *Electroencephalogr Clin Neurophysiol* 1997;**105**:340–4.
72. Sekiguchi H, Nakazawa K, Suzuki S. Differences in recruitment properties of the corticospinal pathway between lengthening and shortening contractions in human soleus muscle. *Brain Res* 2003;**977**:169–79.
73. Kesar TM, Eicholtz S, Lin BJ, Wolf SL, Borich MR. Effects of posture and coactivation on corticomotor excitability of ankle muscles. *Restor Neurol Neurosci* 2018;**36**:131–46.
74. Cacchio A, Cimmini N, Alosi P, Santilli V, Marrelli A. Reliability of transcranial magnetic stimulation-related measurements of tibialis anterior muscle in healthy subjects. *Clin Neurophysiol* 2009;**120**:414–9.
75. Lakie M, Caplan N, Loram ID. Human balancing of an inverted pendulum with a compliant linkage: neural control by anticipatory intermittent bias. *J Physiol* 2003;**551**:357–70.