









# Mapping sites of gibberellin biosynthesis in the *Arabidopsis* root tip

Richard Barker<sup>1,2</sup> , Maria Nieves Fernandez Garcia<sup>3</sup> , Stephen J. Powers<sup>1</sup> , Simon Vaughan<sup>1</sup> , Malcolm J. Bennett<sup>2</sup> , Andrew L. Phillips<sup>1</sup> , Stephen G. Thomas<sup>1</sup>  and Peter Hedden<sup>1,4,5</sup> 

<sup>1</sup>Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ, UK; <sup>2</sup>Plant & Crop Sciences, School of Biosciences, University of Nottingham, Sutton Bonington, LE12 5RD, UK; <sup>3</sup>Department of Abiotic Stress and Plant Pathology, Centro de Edafología y Biología Aplicada del Segura (CSIC), Murcia, Spain; <sup>4</sup>Laboratory of Growth Regulators, Centre of the Region Haná for Biotechnological and Agricultural Research, Institute of Experimental Botany, Czech Academy of Sciences, Olomouc CZ-783 71, Czech Republic; <sup>5</sup>Faculty of Science, Palacký University, Olomouc CZ-783 71, Czech Republic

## Summary

Author for correspondence:

Peter Hedden

Email: [peter.hedden@rothamsted.ac.uk](mailto:peter.hedden@rothamsted.ac.uk)

Received: 6 July 2020

Accepted: 5 September 2020

New Phytologist (2020)

doi: 10.1111/nph.16967

**Key words:** gibberellin action, gibberellin metabolism, root apical meristem, root elongation zone, tissue-specific gibberellin depletion, tissue-specific mutant rescue.

- Root elongation depends on the action of the gibberellin (GA) growth hormones, which promote cell production in the root meristem and cell expansion in the elongation zone. Sites of GA biosynthesis in the roots of 7-d-old *Arabidopsis thaliana* seedlings were investigated using tissue-specific GA inactivation in wild-type (Col-0) or rescue of GA-deficient dwarf mutants.
- Tissue-specific GA depletion was achieved by ectopic expression of the GA-inactivating enzyme AtGA2ox2, which is specific for C<sub>19</sub>-GAs, and AtGA2ox7, which acts on C<sub>20</sub>-GA precursors. In addition, tissue-specific rescue of *ga20ox* triple and *ga3ox* double mutants was shown. Furthermore, GUS reporter lines for major GA20ox, GA3ox and GA2ox genes were used to observe their expression domains in the root.
- The effects of expressing these constructs on the lengths of the root apical meristem and cortical cells in the elongation zone confirmed that roots are autonomous for GA biosynthesis, which occurs in multiple tissues, with the endodermis a major site of synthesis.
- The results are consistent with the early stages of GA biosynthesis within the root occurring in the meristematic region and indicate that the penultimate step of GA biosynthesis, GA 20-oxidation, is required in both the meristem and elongation zone.

## Introduction

The action of the plant hormone gibberellin (GA) is necessary for normal root growth, although lower GA concentrations are required to achieve maximal rates of root elongation than for the shoot (Tanimoto, 1994, 2012). Indeed, supraoptimal GA concentrations can be inhibitory for root growth (Inada & Shimmen, 2000; Coelho *et al.*, 2013). Root elongation proceeds by two mechanisms: cell replication within the root apical meristem (RAM) and cell elongation in the elongation zone (EZ). The RAM is the source of new cells within the root and plays a critical role in defining the lineages of the cell files that result from meristematic divisions. It is located at the distal tip of the root, next to a nondividing group of cells known as the quiescent centre (QC) (Nawy *et al.*, 2005; Dinneny & Benfey, 2008). As their neighbouring initial cells divide, one of the daughter cells is disconnected from the QC and then differentiates into distinct tissues. The cell files are attached and cannot move relative to one another, consequentially causing a spatial relationship that is indicative of the cell's age, with the youngest cells near the root tip getting progressively older as their distance from the QC increases (Benfey & Scheres, 2000). Tissue-specific suppression of GA signal transduction has demonstrated that this hormone

determines both the size of the RAM, that is the number of divisions before the cells exit into the transition zone (Achard *et al.*, 2009; Ubeda-Tomas *et al.*, 2009) and the final cell length achieved within the EZ (Ubeda-Tomas *et al.*, 2008). In both cases, GA acts in the endodermis to allow the coordinated growth of the root cell files (Ubeda-Tomas *et al.*, 2008, 2009).


Comparison of the growth of cultured roots of wild-type and GA-deficient tomato indicated that roots are autonomous for GA production, which is close to saturating for wild-type root growth (Butcher *et al.*, 1990). It was shown from grafting experiments with *Arabidopsis thaliana* (*Arabidopsis*) that the root stocks could restore the growth of GA-deficient scions by supplying the GA precursor GA<sub>12</sub>, while later metabolites are not mobile (Regnault *et al.*, 2015). GA<sub>12</sub> is biosynthesised from the common diterpene precursor *trans*-geranylgeranyl diphosphate by the sequential action of the terpene cyclases *ent*-copalyl diphosphate synthase (CPS) and *ent*-kaurene synthase and the cytochrome P450 monooxygenases *ent*-kaurene oxidase and *ent*-kaurenoic acid oxidase (reviewed by Hedden & Thomas, 2012). GA<sub>12</sub> is converted to the biologically active GA<sub>4</sub> by the 2-oxoglutarate-dependent dioxygenases (2ODDs) GA 20-oxidase (GA20ox) and GA 3-oxidase (GA3ox) (Fig. 1). These last two enzymes are encoded by small gene families, comprising five and four members,

respectively, in *Arabidopsis* (Hedden & Phillips, 2000). Two further clades of 2ODDs deactivate GAs by 2 $\beta$ -hydroxylation, members of one acting primarily on C<sub>19</sub>-GAs, including the biologically active forms, and of the other acting on C<sub>20</sub>-GA precursors, including GA<sub>12</sub>. In *Arabidopsis* these two GA 2-oxidase (GA2ox) clades contain five and four functional members, respectively (Thomas *et al.*, 1999; Hedden & Phillips, 2000; Schomburg *et al.*, 2003; Lange & Pimenta-Lange, 2020). The 2ODDs are major sites of regulation of GA content, with GA2ox activity limiting the concentration of biologically active GAs in many species, including *Arabidopsis* (Huang *et al.*, 1998; Coles *et al.*, 1999). Members of the 2ODD families show specificity in terms of regulation and tissue expression, but there is partial redundancy, with three GA2ox enzymes, GA2ox1, -2, -3 (Rieu *et al.*, 2008; Plackett *et al.*, 2012), and the two GA3ox paralogues GA3ox1 and -2 (Mitchum *et al.*, 2006), largely responsible for bioactive GA production required for the proper development of vegetative tissues, including root growth, in *Arabidopsis*.

Based on transcript abundance, the root tip was indicated to be the main site of GA biosynthesis in the *Arabidopsis* root, while several GA2ox genes were expressed in the mature region of the root (Dugardeyn *et al.*, 2008). Transcripts for a majority of GA-biosynthetic genes, including *CPS*, are more abundant in the endodermis than other tissues, while GA2ox1 mRNA

accumulates in the cortex. The indication that the endodermis may be a major site of GA biosynthesis is of particular interest given that functional studies revealed that this tissue is also the site of GA action (Ubeda-Tomas *et al.*, 2008, 2009). Furthermore, the accumulation of fluorescence in the endodermis when GA-fluorescein probes were applied to *Arabidopsis* roots suggested the existence of mechanisms to concentrate GAs within this tissue (Shani *et al.*, 2013). Indeed, the transcript for the NPF3 transporter, which was shown to transport the GA<sub>3</sub> fluorescent probe as well as GA<sub>4</sub> and other C<sub>19</sub>-GAs into *Xenopus* oocytes, accumulates to higher levels in the endodermis than in other tissues (Tal *et al.*, 2015). Rizza *et al.* (2017) used a fluorescence resonance energy transfer (FRET)-based, nuclear-targeted biosensor (nlsGPS1) to visualise and provide relative quantification of nuclear GA<sub>4</sub> levels within the *Arabidopsis* root. The output from the sensor indicated a gradient of bioactive GA concentration along the primary root axis increasing with distance from the QC and peaking in the EZ. In contrast with the results reported by Shani *et al.* (2013), this peak of GA concentration was observed not only in the endodermis, but also in the cortical and epidermal cells.

Here we describe further approaches to define the sites of GA biosynthesis and action within the *Arabidopsis* root. We use tissue-specific promoters to drive expression of a C<sub>20</sub>-GA2ox,



**Fig. 1** Biosynthetic pathway from GA<sub>12</sub> to GA<sub>4</sub> and 2 $\beta$ -hydroxy catabolites, indicating the reactions catalysed by AtGA20ox1, AtGA3ox1, AtGA2ox2 and AtGA2ox7.

*AtGA2ox7* and a C<sub>19</sub>-GA2ox, *AtGA2ox2* in the Col-0 ecotype, the first expected to act at the site of synthesis, whereas the second could act at the site of synthesis and the site of action. The same promoters are used to express functional *AtGA20ox1* in the *atga20ox1*, *atga20ox2*, *atga20ox3* triple mutant and to express *AtGA3ox1* in the *atga3ox1*, *atga3ox2* double mutant, both stacked mutants being extreme dwarfs with short roots (Mitchum *et al.*, 2006; Plackett *et al.*, 2012). The degree of rescue should indicate the location of the reactions lacking in the mutant lines. The extent to which shoot-derived GAs contribute to promotion of root growth is determined by driving expression of the same effector genes from the shoot-specific CHLOROPHYLL A/B-BINDING PROTEIN (CAB) promoter.

## Materials and Methods

### Plant material

*Arabidopsis thaliana* ecotype Col-0 was used as wild-type in all experiments. The *ga20ox1 ga20ox2 ga20ox3* triple mutant (Plackett *et al.*, 2012) and the *AtGA20ox1::GUS* (Hay *et al.*, 2002) and *AtGA20ox2::GUS* (Frigerio *et al.*, 2006) reporter lines are as previously published. The *ga3ox1 ga3ox2* double mutant and *AtGA3ox1::GUS* and *AtGA3ox2::GUS* lines are as described in Mitchum *et al.* (2006) and were provided by Dr Tai-ping Sun, who also provided the *gal-3* (Col-0) mutant line containing the mutation in CPS that was originally found in *Ler* and was backcrossed to Col-0 six times (Tyler *et al.*, 2004). The reporter lines are translational fusions incorporating promoter and exon and intron sequences. The *AtGA2ox6::GUS* transcriptional reporter line was provided by Dr Sharyn Perry (Wang *et al.*, 2004).

### Growth conditions and replication

Seeds to be grown for seed production, crossing or characterisation were surface sterilised with 10% bleach, rinsed and then imbibed in water or 50  $\mu\text{M}$  GA<sub>4</sub> for 3 d at 4°C in the dark before being sown on Levington's compost. Seeds imbibed in GA<sub>4</sub> were rinsed five times in H<sub>2</sub>O before planting. Plants were grown in a controlled environment in 16-h d (150  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) at 24°C : 18°C, day : night temperatures. For measurement of root growth, after imbibition on 50  $\mu\text{M}$  GA<sub>4</sub> and washing, seeds were grown on medium containing 0.5% Gelrite, 1% sucrose, half-strength Murashige–Skoog medium at pH 5.8 on plates placed at an angle of 10° to the vertical in continuous light (150  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ). For measurement of root length, two (pseudoreplicate) or three plants of each line were grown on each plate so that the number of plates used was also the biological replication. Hence, there were five or six biological replicates per line for measurement of root length with two or three roots (plants) per biological replicate. For meristem and cell size measurements, a single plant per plate was used and there were five biological replicates. Also for cell size, five (pseudoreplicate) cells were measured for each biological replicate plant. Randomised complete block designs were used with the plates being statistical blocks. For comparison of the effects of each promoter on root length,

meristem size and final cortical cell length, the median lines in terms of root length for each promoter were grown together on plates, one line for each promoter per plate with 15 plates as replicates.

### Activity of recombinant AtGA2ox2 fusions and AtGA2ox7

To produce plasmid constructs allowing expression of GA2ox2 fused to EYFP, the *EYFP* coding region was amplified by PCR and cloned into pET-32a (Novagen). Primer sequences are listed in Supporting Information Table S1. The *AtGA2ox2* coding region was amplified from a cDNA clone by PCR (Thomas *et al.*, 1999) and cloned into the pET-32a constructs containing *EYFP* sequences to produce the N- and C-terminal expression constructs N-RB-YFP-*AtGA2ox2*:pET-32a and C-RB-*AtGA2ox2*:YFP:pET-32a. *AtGA2ox7* was amplified from 35S::GA2ox7 in pRMA1, provided by Dr R. Amasino (Schomburg *et al.*, 2003), to introduce *Bam*HI and *Sal*I restriction sites and then ligated into pET-32a. The constructs were expressed in *E. coli* Rosetta 2 (DE3 pLysS) cells (Merck), lysates prepared and GA2ox activity determined by incubating with <sup>14</sup>C-labelled GA substrates (Prof. L. N. Mander, Australian National University, Canberra, ACT, Australia) as previously described (Ward *et al.*, 2010). When necessary, product identity was confirmed by combined gas chromatography–mass spectrometry as described previously (MacMillan *et al.*, 1997), except that samples were run on a MAT95XP mass spectrometer coupled to Trace GC (ThermoElectron) (Rieu *et al.*, 2008) operated in full-scan mode.

### Production of transgenic Arabidopsis with tissue-specific expression

The constructs for Arabidopsis transformation were produced using the Gateway cloning system. The *YFP:AtGA2ox2* fusion was inserted downstream of the tissue-specific promoters within the Gateway pENTR11 entry plasmid (Marques-Bueno *et al.*, 2016) acquired from the Nottingham stock centre (NASC). *AtGA2ox2* was then excised using *Bam*HI and *Sal*I and replaced with *AtGA2ox7*, *AtGA20ox1* and *AtGA3ox1*. The constructs were transferred to the pGBW7 binary vector following a standard Invitrogen LR reaction. The constructs in pGBW7 were transformed into *Agrobacterium tumefaciens* strain GV3101, and then into Arabidopsis by the floral dip method (Clough & Bent, 1998). Plants were selected on hygromycin to obtain at least 10 independent T1 lines, segregation analysis was then performed to create at least four stable single insertion homozygous lines for physiological analysis. Stable homozygous, single insertion lines were taken through to the T4 generation before subsequent genotyping and phenotyping.

### Plant phenotype analysis

To examine root growth, seedlings were grown on solid MS Gelrite medium for 7 d so that the growth rates of GA-deficient mutants reached maximum (Achard *et al.*, 2009). Seedlings were then mounted on slides in 10  $\mu\text{M}$  propidium iodide solution to

achieve counterstaining of cell walls around the cortical cells. Confocal microscopy images were acquired using a Leica TCS-SP confocal microscope (Leica, Milton Keynes, UK). EYFP was excited at 488 nm from an argon laser and propidium iodide was visualised at 514 nm. Fluorescence emission for the EYFP was collected between 505 and 530 nm. Fluorescence images were processed using Leica LAS AF LIGHT and Adobe PHOTOSHOP. Subsequently, images were analysed using LAS AF lite, allowing meristem and cell size to be measured (Fig. S1). Measurements of cell length were taken after root hairs became visible.

Plants for assessment of shoot phenotype were grown on compost in 5.5-cm pots as described above and photographed 3 and 6 wk after germination.

### Histochemical GUS assays and microscopy

To visualise GUS expression, whole plants were placed in a substrate buffer (100 mM sodium phosphate, pH 7, 0.5 mM potassium ferrocyanide, 0.5 mM potassium ferricyanide, and 1 mM 5-bromo-4-chloro-3-indolyl glucuronide) and were incubated overnight in the dark at 37°C. Chlorophyll was cleared from stained tissue by incubating in 70% (v/v) ethanol. Whole plants stained with GUS were observed and photographed after decolouration using an Olympus SZ-PT stereomicroscope.

For sectioning, GUS samples were fixed with 2.5% glutaraldehyde and 4% paraformaldehyde in 0.1 M sodium phosphate buffer (pH 7.2) for 2.5 h at 4°C. After this, three washes with phosphate buffer were performed. All fixed tissues were dehydrated in a graded series of ethanol, then infiltrated, first with a propylene oxide and then with propylene oxide and Spurr's resin mixture. The samples were then immersed in Spurr's resin overnight at 4°C. Finally, the samples were embedded in Spurr resin. Blocks were sectioned in semithin sections on a Leica EM UC6 ultramicrotome. Semithin sections (1 µm thick) of material prepared for GUS expression were mounted in dibutylphthalate polystyrene xylene and observed with a Leica DMR light microscope. The micrographs were captured using Leica QM500 image analysis software.

### Statistical analysis

Analysis of variance was used to test (using the F-test) the overall statistical significance of difference between lines for the root, meristem and cell length data. The analysis took account of the design structure with plates being blocks and sections on plates being plots, with any variation due to pseudoreplicates (roots within plants, or cells within plants) being separated from the biological variation. The standard error of the difference (SED) on the residual degrees of freedom (*df*) from analysis of variance (ANOVA) was used to calculate the least significant difference (LSD) at the 5% ( $P=0.05$ ) and 1% ( $P=0.01$ ) levels of significance. The LSDs were used to assign statistical significance to differences between the means of most relevant pairs of lines. Homogeneity of variance over the lines was checked by inspecting plots of residuals and, when required, a natural log transformation was applied with comparisons of means being made on

the transformed scale (see figure legends). The GENSTAT (2009, 12<sup>th</sup> Edn; VSN International Ltd, Hemel Hempstead, UK) statistical package was used for this analysis.

## Results


### Expression of GA-metabolism reporter genes in roots

First the expression domains of GA 2ODD genes in the root were investigated using GUS reporters. For *AtGA20ox1* and *AtGA20ox2*, the reporters comprised 1.5 kbp of promoter sequence and the first two exons and introns upstream of GUS (Plackett *et al.*, 2012). GUS staining for the *AtGA20ox1* reporter indicated expression in both the meristem and EZ with less expression in the transition zone (Fig. 2a). Within the EZ, expression was confined to the cortical cells (Fig. 2b,c). GUS staining for the *AtGA20ox2* reporter was much weaker than for *AtGA20ox1*, but some staining was detected in the endodermis of the EZ (Fig. 2d). Although we were not able to determine its expression profile, data reported by Dugaredeyn *et al.* (2008) showed the expression of *AtGA20ox3* in roots with a similar distribution to *AtGA20ox2*. Mitchum *et al.* (2005) reported the expression from a *AtGA3ox1::GUS* reporter within the stele throughout the root, except in the tip (meristem and EZ). A root cross-section in the translational fusion reporter line indicates that expression is in the pericycle, particularly at the metaxylem poles, with some staining in the adjacent endodermal cells (Fig. 2e). Mitchum *et al.* (2005) found a similar expression profile in roots for *AtGA3ox2*, with additional GUS expression in the EZ, QC and columella, but not in the meristem. In cross-section through the EZ, expression is indicated throughout the stele and also in the endodermis (Fig. 2f).

Expression analysis by real-time quantitative RT-PCR indicates that *GA2ox6* is the most highly expressed *C<sub>19</sub>-GA2ox* gene in roots (Rieu *et al.*, 2008). We examined the expression domain of this GA-inactivation gene within the root using a *GA2ox6::GUS* transcriptional reporter line (Wang *et al.*, 2004). Expression was present throughout the differentiated root and absent from the root tip (Fig. 2g) but, in contrast with *GA3ox1*, was present in all cell types (Fig. 1h).

### Defining promoters for targeted tissue- and stage-specific root expression

To determine the relevance of the observed expression domains in the root, the sites of GA production necessary for promotion of root growth were defined by tissue-specific GA inactivation or mutant rescue. The promoters selected to drive tissue-specific expression are listed in Table 1. A 2-kb promoter sequence from the *GLABRA2* (*GL2*) gene was used to confine expression to epidermal cells of the root. Expression occurs in cells outside a periclinal cortical cell wall and extends from the meristem to the root hair zone, with most intense expression in the differentiation zone (Masucci *et al.*, 1996). The *GL2* expression domain is defined by a 500-bp fragment located between positions -840 and -1340 in the 5' region of the gene (Hung *et al.*, 1998).



**Fig. 2** Reporter gene (*GUS*) expression in roots of 7-d-old seedlings of *Arabidopsis thaliana*. (a–c) pAtGA20ox1, (d) pAtGA20ox2, (e) pAtGA3ox1, (f) pAtGA3ox2, (g, h) pAtGA20x6. With the exception of pAtGA20x6:*GUS*, the reporter constructs consisted of translational fusions including the final intron.

*Co2* expression within the root is confined to the meristematic region of the cortex, its root expression domain conferred by a 500-bp promoter sequence immediately upstream of the 5' end of the open reading frame (Heidstra *et al.*, 2004). Publicly available data <https://www.arabidopsis.org/servlets/TairObject?type=locus&name=At1g62500> indicate the expression of *Co2* also in numerous aerial organs, including juvenile rosette leaves and inflorescence, with peaks of expression within the carpel and ovary. *Co2* is highly expressed at the shoot apex during the vegetative, inflorescence and transition stages, more specifically within the rib meristem and peripheral zone of the shoot apical meristem (SAM). The *Co2* promoter was first used by Mustroph *et al.* (2009) as part of a cell sorting experiment investigating plants' response to hypoxia.

The *CoR* promoter was shown by fluorescence-activated cell sorting of root cells to be expressed specifically within the cortex

**Table 1** Genes for which expression is targeted to specific tissues within the *Arabidopsis* root.

Tissue targeted	Gene with tissue-specific expression	ATG code	References
Epidermis	GL2 – GLABRA2	At1g79840	Masucci <i>et al.</i> (1996)
Cortex – Meristematic region	Co2 – Unknown protein	At1g62500	Heidstra <i>et al.</i> (2004)
Cortex – Elongation zone	CoR – Unknown protein	At1g09750	Dinneny <i>et al.</i> (2008)
Endodermis	SCR – SCARECROW	At3g54220	Di Laurenzio <i>et al.</i> (1996)
Stele	SHR – SHORTROOT	At4g37650	Helariutta <i>et al.</i> (2000)
Green tissue/shoots	CAB – Chlorophyll <i>a/b</i> -binding protein	At1g29920	Puente <i>et al.</i> (1996)

of the EZ (Dinneny *et al.*, 2008). Outside of the root it is expressed in germinating seeds and seedlings, leaves, reproductive organs and in the rib meristem and peripheral region of the SAM. A 500-bp sequence upstream of the start codon was used as this has been shown by green fluorescent protein (GFP)-targeted expression to confirm its expression pattern in the root (Vasava *et al.*, 2018).

*SCARECROW* (*SCR*) encodes a GRAS transcription factor that is expressed in cortex/endodermal initials of the developing embryo and then solely in the endodermis (Di Laurenzio *et al.*, 1996). A 2-kb 5' upstream sequence of the *SCR* gene was used to drive expression, which as well as in the root, occurs in vegetative and reproductive aerial organs in the cell layer surrounding the vascular tissues (Wysocka-Diller *et al.*, 2000).

We used a 2-kb promoter sequence upstream of the start codon of *SHORTROOT* (*SHR*), which is expressed in the stele (Helariutta *et al.*, 2000). Its product moves from the stele into the adjacent endodermis to regulate *SCR* expression (Nakajima *et al.*, 2001). It is expressed also in aerial tissues, being strongly expressed in the peripheral zone of the SAM (Yadav *et al.*, 2009) and in the leaf vasculature (Dhondt *et al.*, 2010).

In order to assess the contribution of shoot-derived GAs to root growth, a 1892-bp 5' upstream sequence of the *CAB* gene was used to drive expression in green tissues in the absence of significant root expression. The *CAB* gene responds to many of the developmental signals required for chloroplast development and green tissue specification (Puente *et al.*, 1996). Experiments using *CAB1::GUS* fusions have shown that in wild-type *Arabidopsis* *CAB* is first expressed in the cotyledons in response to light, and after further growth the expression is confined to the photosynthetic tissue (Yadav *et al.*, 2002).


### Tissue-specific GA inactivation

To inactivate C<sub>19</sub>-GAs and C<sub>20</sub>-GAs by 2β-hydroxylation, *Arabidopsis* Col-0 was transformed with *AtGA2ox2* and *AtGA2ox7* constructs, respectively (see Fig. 1). Their cDNAs were

transformed as 5' translational fusions with *YFP* in constructs containing the promoters listed in the previous section (Table S2). *AtGA2ox2* has been previously shown by *in vitro* enzyme assays with bacterial expressed protein to be specific for  $C_{19}$ -GAs (Thomas *et al.*, 1999). To confirm that *YFP* fusions retained enzyme activity, the activity of N- and C-fusions produced in *E. coli* was compared with that of *AtGA2ox2* against  $[17-^{14}C]GA_9$  and  $[17-^{14}C]GA_4$  as substrates. HPLC analysis of the products demonstrated that both *YFP* fusion proteins were active with both substrates (Fig. S2). Therefore, it was decided to use N-terminal *YFP* fusions for each of the effector proteins. The assumption that the fusion proteins retained enzymatic function is supported by their expression from the *CAB* promoter, which produced the expected shoot phenotype (to be described later). The specificity of *AtGA2ox7* was confirmed by incubating recombinant enzyme with  $^{14}C$ -labelled substrates: the  $C_{20}$ -GAs,  $GA_{12}$  and  $GA_{24}$  and the  $C_{19}$ -GAs,  $GA_9$  and  $GA_4$  (Fig. S3). While both  $C_{20}$ -GAs were  $2\beta$ -hydroxylated by the enzyme, we could detect no activity against  $GA_9$  and  $GA_4$ .  $GA_{12}$  was metabolised by *AtGA2ox7* to  $GA_{110}$  ( $2\beta$ -hydroxy $GA_{12}$ ), and  $GA_{24}$  was converted to a product consistent with the previously undescribed  $2\beta$ -hydroxy $GA_{24}$ .

Three to six homozygous T3 lines with single inserts were selected for each transformed construct and the presence of the transgene was confirmed by PCR. The expression domains within the root were confirmed by imaging *YFP* fluorescence using confocal microscopy (shown for representative lines for each promoter in Fig. S4). Root lengths, as well as meristem and final cortical cell lengths (see Fig. S1) were determined in seedlings 7 d after germination and compared with those for *Col-0* and the highly GA-deficient *ga1-3* mutant. The data for three representative *YFP-AtGA2ox2*-expressing lines for each construct are shown in Fig. S5, while the median lines in terms of root length for each *AtGA2ox2*-containing construct are compared in Fig. 3. With the exception of *CAB*, each promoter driving *YFP-AtGA2ox2* resulted in a reduction in root length, with *SCR* having the most severe effect and *SHR* producing a relatively small reduction. *SCR*, *Co2* and *GL2*, but not the other promoters, reduced meristem size, while *SCR* and particularly *CoR* reduced final cell length. The equivalent data for the *promoter:YFP-AtGA2ox7* lines are presented in Fig. S6 and Fig. 4. The effect of these constructs on root growth was less severe than for the *AtGA2ox2* constructs, but some reduction in root length was obtained with *SCR*, *CoR*, *Co2* and *GL2*. Significant ( $P < 0.01$ , LSD) reductions in meristem size compared with *Col-0* were found with the *SCR*, *Co2* and *GL2* promoters, while *SCR*, *CoR* and *GL2* also produced shorter cortical cells.

The effect of the ectopic gene expression on growth of the shoot was assessed at 3 and 6 wk after sowing on compost. Results for the *AtGA2ox2* effector are shown in Fig. S7. Most severe dwarfism was obtained with the *CAB* promoter, while some lines transformed with *CoR:YFP-GA2ox2* also exhibited substantial dwarfism. Mild dwarfism was noted for plants harbouring genes driven by the other promoters, except for *SHR:YFP-GA2ox2* plants, which were identical to *Col-0*. In contrast with *AtGA2ox2*, expression of *AtGA2ox7* produced substantial shoot




**Fig. 3** Comparison of the primary root parameters for median *Col-0* lines expressing *YFP-AtGA2ox2* with those of *Col-0* and *ga1-3*(*Col-0*) in 7-d-old seedlings of *Arabidopsis thaliana*. (a) Primary root length. (b) Proximal meristem length. (c) Final cortical cell length. Seedlings were grown on vertical plates and lengths calculated using *IMAGEJ* software. † Indicates that the transgenic lines are statistically significantly different from *ga1-3* and \* indicates that they are statistically significantly different from *Col-0*. 12 plants for each line were measured for root length with two plants (roots) per biological replicate ( $n = 6$ ), five plants for each line were measured for meristem and cell size ( $n = 5$ ), with five cells per biological replicate being measured. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; †,  $P < 0.05$ ; ††,  $P < 0.01$  according to least significant difference (LSD) values (see Supporting Information Notes S1 for means, LSDs and df). Error bars indicate  $\pm$  SE.

dwarfism regardless of the promoter (Fig. S8). In this case, *CAB* and *Co2* were the most effective promoters, with *SCR* and *GL2* intermediate, and *CoR* and *SHR* producing milder dwarfism.

#### Rescue of the *ga20ox1*, *ga20ox2*, *ga20ox3* triple mutant

Loss of *AtGA20ox1*, *AtGA20ox2* and *AtGA20ox3*, which are three of the five *GA20ox* genes in *Arabidopsis*, produces a severely dwarfed, sterile plant with reduced root length and seeds which do not germinate without application of GA (Plackett *et al.*, 2012). The triple mutant was transformed with genes in which *YFP-AtGA20ox1* was downstream of the same tissue-specific



**Fig. 4** Comparison of the primary root parameters for median lines expressing each *YFP-AtGA2ox7* construct with Col-0 and *ga1-3* in 7-d-old seedlings of *Arabidopsis thaliana*. (a) Primary root length. (b) Proximal meristem length. (c) Final cortical cell length. Seedlings were grown on vertical plates and lengths calculated using IMAGEJ software. † Indicates that the transgenic lines are statistically significantly different from *ga1-3* and \* indicates that they are statistically significantly different from Col-0. Twelve plants for each line were measured for root length with two plants (roots) per biological replicate ( $n = 6$ ), five plants for each line were measured for meristem and cell size ( $n = 5$ ), with five cells per biological replicate being measured. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; †,  $P < 0.05$ ; ††,  $P < 0.01$  according to least significant difference (LSD) values (see Supporting Information Notes S1 for means, LSDs and df). Error bars indicate  $\pm$  SE.

promoters described above. The root expression domains were confirmed in representative plants for each construct by confocal microscopy (Fig. S9). Total root lengths, meristem and final cortical cell lengths were determined 7 d after germination and compared with Col-0 and the triple *ga20ox* mutant. The data for three lines are presented in Fig. S10, and for the median lines for each construct, Col-0 and triple *ga20ox* mutant in Fig. 5. Complete rescue of root growth, including meristem size and cortical cell length, was obtained with the *SCR* promoter, while *Co2*, *GL2* and *CAB* provided partial rescue of root length. In the case of *Co2* and *GL2*, both meristem and cell lengths were promoted, while the *CAB* promoter increased only cell length. No rescue was obtained with the *SHR* or *CoR* promoter. The failure of *CoR*:*YFP-AtGA20ox1* to rescue cell length, whereas there was slight

rescue by the *Co2* promoter, suggested that cell elongation required GA 20-oxidation in the meristem. Indeed, full rescue of cell elongation and root growth was obtained by combining *CoR*:*YFP-AtGA20ox1* with *Co2*:*YFP-AtGA20ox1* in the *ga20ox* triple mutant (Fig. 6).

The effect of the transgenes on growth of the shoots is illustrated in Fig. S11, in which the lines are compared with Col-0 and GA-deficient triple *ga20ox* and *ga1-3* mutants. Growth of plants expressing *CAB*:*YFP-GA20ox1* was slightly greater than for the wild-type, and the *SCR* and *Co2* promoters gave almost full mutant rescue. Expression from the *GL2* promoter also gave substantial rescue, but the *CoR* and *SHR* promoters driving *YFP-GA20ox1* produced no growth stimulation in the triple *ga20ox* mutant.


### Rescue of the *ga3ox1 ga3ox2* double mutant

The same tissue-specific promoters were used to drive expression of *YFP:AtGA3ox1* in the *ga3ox1 ga3ox2* double mutant; the spatial distribution of YFP in roots of the transgenic lines was determined by confocal microscopy (Fig. S12). Apart from in their expected root expression domains, the *GL2* and *CAB* promoters also produced fluorescence in the columella. Total root lengths, meristem and final cortical cell lengths for three lines are shown in Fig. S13, and the median plants for each construct are compared with Col-0 and the double *ga3ox* mutant in Fig. 7. All promoters driving *YFP:AtGA3ox1* rescued root length to some extent, with roots in the *SHR*, *SCR*, *CoR* and *GL2* promoter lines not being significantly ( $P > 0.05$ , LSD) shorter than the Col-0 wild-type. In the *Co2* and *CAB* lines there was partial rescue of root length. Meristem length in the *SHR*, *SCR* and *Co2* lines was not significantly different ( $P > 0.05$ , LSD) from that in Col-0, while there was partial rescue of meristem size in the *CoR* and *GL2* lines, but none in the *CAB* lines. There was substantial rescue of final cortical cell length with all promoters, with only the *Co2* line producing cells significantly ( $P < 0.05$ , LSD) shorter than those in Col-0.

Similar to what was observed for root growth, the constructs fully restored growth of the shoots, regardless of which promoter was used (Fig. S14).

### Discussion

Rizza *et al.* (2017), using a nuclear-targeted biosensor nlsGPS1, reported a gradient of bioactive GA from the root tip to the EZ that was exaggerated when GA<sub>4</sub> was applied to the root. This indicates the existence of a transport system that concentrates bioactive GAs in this zone. The results of Shani *et al.* (2013) from application of fluorescein-conjugated GA<sub>4</sub> and GA<sub>3</sub> probes to the *Arabidopsis* root support this conclusion, although, while Shani *et al.* (2013) found accumulation of the probe within the endodermis, the results of Rizza *et al.* (2017) suggest wider distribution of GA within the EZ. However, these findings are not in agreement with the model proposed by Band *et al.* (2012), which predicts that GA concentration is diluted as cells expand, as a potential mechanism to explain growth cessation. Furthermore,




**Fig. 5** Comparison of the primary root parameters in 7-d-old seedlings of *Arabidopsis thaliana* for median lines expressing each YFP-AtGA20ox1 construct in the *ga20ox1,2,3* triple mutant compared with those of Col-0 and the triple *ga20ox* mutant. (a) Primary root length. (b) Proximal meristem length. (c) Final cortical cell length. Seedlings were grown on vertical plates and lengths calculated using IMAGEJ software. † Indicates that the transgenic lines are significantly different from and *atga20ox1,2,3* and \* indicates that they are significantly different from Col-0. 12 plants for each line were measured for root length with two plants (roots) per biological replicate ( $n = 6$ ), five plants for each line were measured for meristem and cell size ( $n = 5$ ), with five cells per biological replicate being measured. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; †,  $P < 0.05$ ; ††,  $P < 0.01$  according to least significant difference (LSD) values (see Supporting Information Notes S1 for means, LSDs and df). Error bars indicate  $\pm$  SE.

it has remained unclear as to whether bioactive GAs are synthesised *in situ* or are translocated to the sites of action from other tissues as final products or as precursors. The existence of a concentrating mechanism within the root, as suggested by Rizza *et al.* (2017) and Shani *et al.* (2013), is compatible with movement to the EZ from another site. In the present study to investigate the sites of GA biosynthesis in the root, the promoters chosen to provide cell-specific GA inactivation or mutant rescue were based on their established expression domains within the root, which were confirmed by expressing YFP fusions. This multipronged approach should enhance the robustness of the results, which could be influenced by differences in the expression levels of the ectopic genes compared with those of the native GA-biosynthesis genes.

The selected promoters are also active in aerial tissues and several of the constructs modified shoot as well as root growth. To

determine the extent to which root growth is influenced by the shoot, the same effector open reading frames were expressed from the shoot-specific *CAB* promoter. While expression of the GA inactivation *AtGA20ox2* and *AtGA20ox7* cDNAs from *CAB* resulted in severe dwarfing of the shoot, these constructs had no effect on root growth, confirming that roots can be autonomous for GA production. It has been reported that application of the GA-biosynthesis inhibitor uniconazole to shoots of *Arabidopsis* seedlings stimulates root elongation and upregulates the feedback-regulated *GA20ox* and *GA3ox* genes in roots (Bidadi *et al.*, 2009). Bidadi *et al.* (2009) argued that, under standard conditions, root growth is promoted partially by GA or a precursor transported from the shoot, but that, in the absence of this source of promotion, the plant can enhance *in situ* GA production to (more than) compensate. Grafting experiments have shown that






**Fig. 6** Combining cortical *GA20ox* expression in the meristem and elongation zone of 7-d-old seedlings of *Arabidopsis thaliana*. (a) YFP fluorescence in roots of a cross between *Co2:YFP:AtGA20ox1* and *CoR:YFP:AtGA20ox2* and in the parent lines. (b) Comparison of the lengths of the primary root, proximal meristem and final cortical cells in the cross and parent lines compared with those in *Col-0* and the triple *atga20ox1,2,3* mutant. Seedlings were grown on vertical plates and lengths calculated using IMAGEJ. † Indicates that the transgenic lines are significantly different from *ga1-3* and \* indicates that they are significantly different from *Col-0*. 12 plants for each line were measured for root length with two plants (roots) per biological replicate ( $n = 6$ ), five plants for each line were measured for meristem and cell size ( $n = 5$ ), with five cells per biological replicate being measured. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; ††,  $P < 0.01$  according to least significant difference (LSD) values (see Supporting Information Notes S1 for means, LSDs and df). Error bars indicate  $\pm$  SE.

the shoot can provide the GA precursor  $GA_{12}$  to rescue root growth in GA mutants with lesions early in the GA-biosynthetic pathway, while later intermediates and the active hormone are

apparently not transported (Regnault *et al.*, 2015). Transformation of the *ga20ox* triple mutant with *CAB::YFP:AtGA20ox1* and of the *ga3ox* double mutant with *CAB::YFP:AtGA3ox1* gave



**Fig. 7** Comparison of the primary root parameters in 7-d-old seedlings of *Arabidopsis thaliana* for median lines expressing each *YFP-AtGA3ox1* construct in the *ga3ox1,2* double mutant compared with those of Col-0 and the double *ga3ox* mutant. (a) Primary root length. (b) Proximal meristem length. (c) Final cortical cell length. Seedlings were grown on vertical plates and lengths calculated using IMAGEJ. † Indicates that the transgenic lines are statistically significantly different from the double *ga3ox* mutant and \* indicates that they are statistically significantly different from Col-0. Twelve plants for each line were measured for root length with two plants (roots) per biological replicate ( $n = 6$ ), five plants for each line were measured for meristem and cell size ( $n = 5$ ), with five cells per biological replicate being measured. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; ††,  $P < 0.01$  according to least significant difference (LSD) values (see Supporting Information Notes S1 for means, LSDs and df). Error bars indicate  $\pm$  SE.

partial recovery of root length and cell length, but not meristem size, indicating that shoot-derived  $C_{19}$ -GAs can make a small contribution to root growth, even if  $GA_{12}$  is the main form transported from the shoots.


Gibberellin signalling is required for normal cell production in the meristem (Achard *et al.*, 2009; Ubeda-Tomas *et al.*, 2009) and for cell elongation (Ubeda-Tomas *et al.*, 2008), with the endodermis identified as the major site of GA action. Our results based on rescue of the triple *ga20ox* mutant indicate that GA 20-oxidation in the endodermis (SCR promoter), cortex (Co2) or

epidermis (GL2) is sufficient to stimulate root growth, with the endodermal activity having the largest influence (Fig. 5). On the basis of meta-analysis of gene expression data, Dugardeyn *et al.* (2008) concluded that many of the genes that influence GA production in the *Arabidopsis* root are most highly expressed in the endodermis, but also have expression in the cortex. Expression of *AtGA2ox2* and *AtGA2ox7* from the *Co2* promoter, which is active in cortical cells of the meristem, reduced meristem size, but not cell length, while expression of these effectors from *CoR*, which promotes expression in the cortex of the EZ, reduced cell length,

but not meristem size. This is consistent with GA being required in both the meristem and EZ for normal root growth. It also suggests that the cortex is an important site of GA biosynthesis. However, while *AtGA20ox1* expression from *Co2* gave partial rescue of both meristem and cell size in the triple *ga20ox* mutant, expression of *AtGA20ox1* from *CoR* gave no rescue, even of cortical cell length. Based on GUS staining in lines expressing the *AtGA20ox1* reporter, the endogenous gene is expressed specifically in cortical cells of the EZ as well as in the meristem (Fig. 2a–c). The unexpected lack of rescue by *CoR::YFP::AtGA20ox1* may indicate insufficient availability of its C<sub>20</sub>-GA substrates in the EZ, which may be provided by the meristem. This was tested by crossing *CoR::YFP::AtGA20ox1* with *Co2::YFP::AtGA20ox1* to restore GA20ox activity in the cortex of the meristem and EZ. Full rescue of the root phenotype in the cross confirmed that GA biosynthesis, and potentially activity within the meristem is necessary for cell elongation. If the meristem provides C<sub>20</sub>-GA substrate for GA 20-oxidation in the EZ cortex, this would not be GA<sub>12</sub>, which would be depleted by GA20ox activity, when in fact *Co2::YFP::AtGA20ox1* restored some cell elongation, although this could also be explained by provision of active GA to the EZ. Full rescue of cell elongation was obtained with *CoR::YFP::AtGA3ox1* in the *ga3ox1/ga3ox2* background, indicating the presence of its substrate GA<sub>9</sub> in the cortex and that its product GA<sub>4</sub> is able to move to the site of action in the endodermis. It is also possible that GA action in the cortex enables some cell expansion. This rescue of cell elongation by *CoR::YFP::AtGA3ox1* indicates that GA biosynthesis and/or action in the meristem is not required for cell elongation, although *CoR::YFP::AtGA3ox1* also provided partial rescue of meristem size suggesting movement of GA<sub>4</sub> from the EZ to the meristem. It is necessary to consider the upregulation of *GA20ox* genes in the GA-deficient mutants due to feedback regulation (Phillips *et al.*,

1995) that could increase GA<sub>9</sub> production and availability in cells where it is normally limiting.

On the basis of reporter gene expression, Mitchum *et al.* (2006) showed that *AtGA3ox1* is expressed in the mature, non-elongating region of the root and our results with the same reporter lines indicate that this expression is in the pericycle cells of the stele. Using an equivalent reporter gene, Bidadi *et al.* (2009) obtained GUS staining in the EZ and root tip after shoot-applied uniconazole. This indicates that the gene is expressed in these regions, albeit normally at very low levels. *AtGA3ox2* is also expressed in the mature root, but in addition is expressed in the EZ, QC and columella, but apparently not in the meristem (Mitchum *et al.*, 2006). Taken with the nlsGPS1 reporter data, this could indicate that small quantities of GA are activated around the QC providing a low concentration of GA required for cell division. Then, as the cells reach the transition zone, the higher expression of *GA3ox1* increases the amount of bioactive GA to promote cell elongation. We show that within the EZ, *AtGA3ox2* is expressed in the stele and endodermis, with some expression in cortical cells. This expression domain is consistent with the complete rescue of root growth in *ga3ox1 ga3ox2* by expression of *AtGA3ox1*, which encodes a functionally identical enzyme to *AtGA3ox2*, from the *SHR* (stele), *SCR* (endodermis) and *CoR* (cortex) promoters. Expression in the epidermis from the *GL2* promoter also gave strong rescue of root growth, although there is no indication that *GA3ox* genes are normally expressed in this cell file. This rescue demonstrates that these tissues have access to the *GA3ox* substrate GA<sub>9</sub> and that the product GA<sub>4</sub> is accessible to the site of action in the endodermis. The absence of *AtGA3ox1* causes a small reduction in root growth, whereas loss of *AtGA3ox2* has no effect (Mitchum *et al.*, 2006). This would suggest that GA<sub>4</sub> produced from GA<sub>9</sub> in the stele makes a contribution to promotion of root growth. While there



**Fig. 8** Summary of the results showing the root domains in which expression of *AtGA20ox2* or *AtGA20ox7* produced strong (red), partial (yellow) or no inhibition of root growth (blue), or in which expression of *AtGA20ox1* or *AtGA3ox1* produced complete (red), partial (yellow) or no rescue of root growth in the *ga20ox* triple or *ga3ox* double mutants, respectively.

is evidence for the long-distance movement of GA<sub>12</sub>, but not of GA<sub>9</sub>, from the shoot to roots (Regnault *et al.*, 2015), there may be some limited transport of GA<sub>9</sub> from the shoot or it may be produced in the phloem. However, *SHR::YFP:AtGA20ox1* failed completely to rescue root growth in the triple *ga20ox* mutant, indicating that the *SHR* expression domain does not overlap with those of *AtGA20ox1*, *AtGA20ox2* or *AtGA20ox3*.

Full rescue of the growth of aerial organs in the *ga3ox1 ga3ox2* mutant was obtained with all promoters driving *AtGA3ox1* expression, while no rescue of the triple *ga20ox* mutant occurred when *AtGA20ox1* was expressed from *SHR* or *CoR*. The other promoters gave different degrees of triple *ga20ox* rescue, with *CAB* and *SCR* proving the most effective. It would appear that the cells expressing *SHR* and *CoR* have limited access to GA<sub>12</sub> or other C<sub>20</sub>-GA intermediates, whereas GA<sub>9</sub> is freely available in all the expression domains tested. It is also possible that GA<sub>9</sub> is not released from these cells or, less likely, that there is cell-specific instability of GA20ox1. Expression of the C<sub>20</sub>-GA-inactivation cDNA *AtGA2ox7* was generally more effective than that of the *C<sub>19</sub>-GA2ox*, *AtGA2ox2* at reducing shoot growth with almost all promoters, except for *CAB*, which gave strong dwarfism with both effector genes. This may reflect the strength and broad expression domain of the *CAB* promoter in shoots. For *AtGA2ox7* expression, the *SHR* and *CoR* promoters were the least effective, consistent with the *GA20ox* mutant rescue and reinforcing the conclusion that the stele and cortex of the EZ have least access to C<sub>20</sub>-GAs. By contrast, with the exception of *CAB*, the *CoR* promoter driving *YFP:AtGA2ox2* was the most dwarfing, with *SCR* having some effect. If there is high mobility between tissues of GA<sub>9</sub> and GA<sub>4</sub>, as suggested by the *ga3ox* mutant rescue, then, providing their formation occurs in more than one tissue type, inactivation restricted to specific tissues will be ineffective. The dwarfing obtained with *CoR* and *SCR* promoters could reflect the fact that they are active at, or close to, the site of GA action.

In conclusion, the results, which are summarised in Fig. 8, support previous indications that roots and shoots are autonomous for GA production and that GA is required for cell division in the root meristem and elongation in the EZ. The cortex and endodermis are important sites of GA biosynthesis, but GA biosynthesis (*GA20ox* activity) and/or cell elongation in the cortex of the EZ are dependent on GA production and/or action in the meristem. The meristem may also supply GA and precursors to the EZ. While transport of C<sub>20</sub>-GAs or earlier precursors of GA biosynthesis between root tissues is restricted, the rescue of root growth by *AtGA3ox1* in the *ga3ox* double mutant indicates efficient movement of C<sub>19</sub>-GAs between tissues. Partial restoration of cell elongation, but not cell division, by shoot expression of *GA20ox1* or *GA3ox1* in the respective mutants indicate some movement of C<sub>19</sub>-GAs from shoot to root, despite GA<sub>12</sub> being the major transport form (Regnault *et al.*, 2015).

## Acknowledgements









We are grateful to Dr Tai-ping Sun, Duke University, for the *ga3ox1 ga3ox2* and *ga1-3*(Col-0) mutants, to Dr Sharyn Perry,

University of Kentucky, for the *AtGA2ox6::GUS* reporter line, to Dr Susana Ubeda-Tomas, University of Nottingham, for the *pENTR11:SCR:GIDI:YFP* plasmid, to Dr Richard Amasino, University of Wisconsin, for the *35S:AtGA2ox7* plasmid and to I. Pearman, A. Griffin and other glasshouse staff at Rothamsted Research for plant husbandry. This work was supported by a Lawes Agricultural Trust Studentship to RB and by grant P16508, funded by the Biotechnology and Biological Sciences Research Council of the UK, which also provides support to Rothamsted Research through funding of the 20:20 Wheat® (BBS/E/C/00005202) and Designing Future Wheat Integrated Strategic Programmes (BBS/E/C/00010220; ALP and ST). PH acknowledges funding from the Czech Science Foundation (grant nos. 18-10349S and 20-17984S) and the European Regional Developmental Fund Project 'Centre for Experimental Plant Biology' no. CZ.02.1.01/0.0/0.0/16\_019/0000738.

## Author contributions

RB, MJB, ALP, SGT and PH planned and designed the research, RB and MNFG performed the experiments, SV, MJB, ALP, SGT and PH supervised the research, SJP carried out the statistical analysis. RB and PH wrote the manuscript, which was edited by SJP, MJB, ALP and SGT.

## ORCID

Richard Barker  <https://orcid.org/0000-0002-9951-1382>  
 Malcolm J. Bennett  <https://orcid.org/0000-0003-0475-390X>  
 Maria Nieves Fernandez Garcia  <https://orcid.org/0000-0002-5860-2173>  
 Peter Hedden  <https://orcid.org/0000-0002-4216-7488>  
 Andrew L. Phillips  <https://orcid.org/0000-0002-6406-7015>  
 Stephen J. Powers  <https://orcid.org/0000-0003-3349-4425>  
 Stephen G. Thomas  <https://orcid.org/0000-0001-6731-2699>  
 Simon Vaughan  <https://orcid.org/0000-0002-1550-6915>

## References

- Achard P, Gusti A, Cheminant S, Alioua M, Dhondt S, Coppens F, Beemster GTS, Genschik P. 2009. Gibberellin signaling controls cell proliferation rate in *Arabidopsis*. *Current Biology* 19: 1188–1193.
- Benfey PN, Scheres B. 2000. Primer – root development. *Current Biology* 10: R813–R815.
- Bidadi H, Yamaguchi S, Asahina M, Satoh S. 2009. Effects of shoot-applied gibberellin/gibberellin-biosynthesis inhibitors on root growth and expression of gibberellin biosynthesis genes in *Arabidopsis thaliana*. *Plant Root* 4: 4–11.
- Butcher DN, Clark JA, Lenton JR. 1990. Gibberellins and the growth of excised tomato roots – comparison of gib-1 mutant and wild-type and responses to applied GA<sub>3</sub> and 2S,3S paclobutrazol. *Journal of Experimental Botany* 41: 715–722.
- Clough SJ, Bent AF. 1998. Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *The Plant Journal* 16: 735–743.
- Coelho M, Colebrook EH, Lloyd DPA, Webster CP, Mooney SJ, Phillips AL, Hedden P, Whalley WR. 2013. The involvement of gibberellin signalling in the effect of soil resistance to root penetration on leaf elongation and tiller number in wheat. *Plant & Soil* 371: 81–94.

- Coles JP, Phillips AL, Croker SJ, Garcia Lepe R, Lewis MJ, Hedden P. 1999. Modification of gibberellin production and plant development in *Arabidopsis* by sense and antisense expression of gibberellin 20-oxidase genes. *The Plant Journal* 17: 547–556.
- Dhondt S, Coppens F, De Winter F, Swarup K, Merks RMH, Inze D, Bennett MJ, Beechster GTS. 2010. SHORT-ROOT and SCARECROW regulate leaf growth in *Arabidopsis* by stimulating S-phase progression of the cell cycle. *Plant Physiology* 154: 1183–1195.
- Di Lorenzo L, Wysocka-Diller J, Malamy JE, Pysh L, Helariutta Y, Freshour G, Hahn MG, Feldmann KA, Benfey PN. 1996. The SCARECROW gene regulates an asymmetric cell division that is essential for generating the radial organization of the *Arabidopsis* root. *Cell* 86: 423–433.
- Dinneny JR, Benfey PN. 2008. Plant stem cell niches: standing the test of time. *Cell* 132: 553–557.
- Dinneny JR, Long TA, Wang JY, Jung JW, Mace D, Pointer S, Barron C, Brady SM, Schiefelbein J, Benfey PN. 2008. Cell identity mediates the response of *Arabidopsis* roots to abiotic stress. *Science* 320: 942–945.
- Dugardeyn J, Vandebussche F, Van Der Straeten D. 2008. To grow or not to grow: what can we learn on ethylene-gibberellin cross-talk by *in silico* gene expression analysis? *Journal of Experimental Botany* 59: 1–16.
- Frigerio M, Alabadi D, Perez-Gomez J, Garcia-Carcel L, Phillips AL, Hedden P, Blazquez MA. 2006. Transcriptional regulation of gibberellin metabolism genes by auxin signaling in *Arabidopsis*. *Plant Physiology* 142: 553–563.
- Hay A, Kaur H, Phillips A, Hedden P, Hake S, Tsiantis M. 2002. The gibberellin pathway mediates KNOTTED1-type homeobox function in plants with different body plans. *Current Biology* 12: 1557–1565.
- Hedden P, Phillips AL. 2000. Gibberellin metabolism: new insights revealed by the genes. *Trends in Plant Science* 5: 523–530.
- Hedden P, Thomas SG. 2012. Gibberellin biosynthesis and its regulation. *Biochemical Journal* 444: 11–25.
- Heidstra R, Welch D, Scheres B. 2004. Mosaic analyses using marked activation and deletion clones dissect *Arabidopsis* SCARECROW action in asymmetric cell division. *Genes & Development* 18: 1964–1969.
- Helariutta Y, Fukaki H, Wysocka-Diller J, Nakajima K, Jung J, Sena G, Hauser MT, Benfey PN. 2000. The SHORT-ROOT gene controls radial patterning of the *Arabidopsis* root through radial signaling. *Cell* 101: 555–567.
- Huang SS, Raman AS, Ream JE, Fujiwara H, Cerny RE, Brown SM. 1998. Overexpression of 20-oxidase confers a gibberellin-overproduction phenotype in *Arabidopsis*. *Plant Physiology* 118: 773–781.
- Hung CY, Lin Y, Zhang M, Pollock S, Marks MD, Schiefelbein J. 1998. A common position-dependent mechanism controls cell-type patterning and GLABRA2 regulation in the root and hypocotyl epidermis of *Arabidopsis*. *Plant Physiology* 117: 73–84.
- Inada S, Shimmen T. 2000. Regulation of elongation growth by gibberellin in root segments of *Lemna minor*. *Plant & Cell Physiology* 41: 932–939.
- Lange T, Pimenta Lange MJ. 2020. The multifunctional dioxygenases of gibberellin synthesis. *Plant & Cell Physiology*. doi: 10.1093/pcp/pcaa051.
- MacMillan J, Ward DA, Phillips AL, Sanchez Beltran MJ, Gaskin P, Lange T, Hedden P. 1997. Gibberellin biosynthesis from gibberellin A<sub>12</sub>-aldehyde in endosperm and embryos of *Marah macrocarpus*. *Plant Physiology* 113: 1369–1377.
- Marques-Bueno MM, Morao AK, Cayrel A, Platre MP, Barberon M, Caillieux E, Colot V, Jaillais Y, Roudier F, Vert G. 2016. A versatile Multisite Gateway-compatible promoter and transgenic line collection for cell type-specific functional genomics in *Arabidopsis*. *The Plant Journal* 85: 320–333.
- Masucci JD, Rerie WG, Foreman DR, Zhang M, Galway ME, Marks MD, Schiefelbein JW. 1996. The homeobox gene *GLABRA2* is required for position-dependent cell differentiation in the root epidermis of *Arabidopsis thaliana*. *Development* 122: 1253–1260.
- Mitchum MG, Yamaguchi S, Hanada A, Kuwahara A, Yoshioka Y, Kato T, Tabata S, Kamiya Y, Sun TP. 2006. Distinct and overlapping roles of two gibberellin 3-oxidases in *Arabidopsis* development. *The Plant Journal* 45: 804–818.
- Mustroph A, Zanetti MW, Jang CJH, Holtan HE, Repetti PP, Galbraith DW, Girke T, Bailey-Serres J. 2009. Profiling transcriptomes of discrete cell populations resolves altered cellular priorities during hypoxia in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* 106: 18843–18848.
- Nakajima K, Sena G, Nawy T, Benfey PN. 2001. Intercellular movement of the putative transcription factor SHR in root patterning. *Nature* 413: 307–311.
- Nawy T, Lee JY, Colinas J, Wang JY, Thongrod SC, Malamy JE, Birnbaum K, Benfey PN. 2005. Transcriptional profile of the *Arabidopsis* root quiescent center. *Plant Cell* 17: 1908–1925.
- Phillips AL, Ward DA, Uknes S, Appleford NEJ, Lange T, Huttly A, Gaskin P, Graebe JE, Hedden P. 1995. Isolation and expression of three gibberellin 20-oxidase cDNA clones from *Arabidopsis*. *Plant Physiology* 108: 1049–1057.
- Plackett ARG, Powers SJ, Fernandez-Garcia N, Urbanova T, Takebayashi Y, Seo M, Jikumaru Y, Benlloch R, Nilsson O, Ruiz-Rivero O *et al.* 2012. Analysis of the developmental roles of the *Arabidopsis* gibberellin 20-oxidases demonstrates that GA20ox1, -2, and -3 are the dominant paralogs. *Plant Cell* 24: 941–960.
- Puente P, Wei N, Deng XW. 1996. Combinatorial interplay of promoter elements constitutes the minimal determinants for light and developmental control of gene expression in *Arabidopsis*. *EMBO Journal* 15: 3732–3743.
- Regnault T, Daviere JM, Wild M, Sakvarelidze-Achard L, Heintz D, Bergua EC, Diaz IL, Gong F, Hedden P, Achard P. 2015. The gibberellin precursor GA<sub>12</sub> acts as a long-distance growth signal in *Arabidopsis*. *Nature Plants* 1: 15073.
- Rieu I, Ruiz-Rivero O, Fernandez-Garcia N, Griffiths J, Powers SJ, Gong F, Linhartova T, Eriksson S, Nilsson O, Thomas SG *et al.* 2008. The gibberellin biosynthetic genes *AtGA20ox1* and *AtGA20ox2* act, partially redundantly, to promote growth and development throughout the *Arabidopsis* life cycle. *The Plant Journal* 53: 488–504.
- Rizza A, Walia A, Lanquar V, Frommer W, Jones A. 2017. In vivo gibberellin gradients visualized in rapidly elongating tissues. *Nature Plants* 3: 803–813.
- Schomburg FM, Bizzell CM, Lee DJ, Zeevaert JAD, Amasino RM. 2003. Overexpression of a novel class of gibberellin 2-oxidases decreases gibberellin levels and creates dwarf plants. *Plant Cell* 15: 151–163.
- Shani E, Weinstain R, Zhang Y, Castillejo C, Kaiserli E, Chory J, Tsien RY, Estelle M. 2013. Gibberellins accumulate in the elongating endodermal cells of *Arabidopsis* root. *Proceedings of the National Academy of Sciences, USA* 110: 4834–4839.
- Tal I, Zhang Y, Jørgensen ME, Pisanty O, Barbosa ICR, Zourelidou M, Regnault T, Crocoll C, Olsen CE, Weinstain R *et al.* 2015. The *Arabidopsis* NPF3 protein is a GA transporter. *Nature Communications* 7: 11486.
- Tanimoto E. 1994. Interaction of gibberellin A<sub>3</sub> and ancymidol in the growth and cell-wall extensibility of dwarf pea roots. *Plant & Cell Physiology* 35: 1019–1028.
- Tanimoto E. 2012. Tall or short? Slender or thick? A plant strategy for regulating elongation growth of roots by low concentrations of gibberellin. *Annals of Botany* 110: 373–381.
- Thomas SG, Phillips AL, Hedden P. 1999. Molecular cloning and functional expression of gibberellin 2-oxidases, multifunctional enzymes involved in gibberellin deactivation. *Proceedings of the National Academy of Sciences, USA* 96: 4698–4703.
- Tyler L, Thomas SG, Hu JH, Dill A, Alonso JM, Ecker JR, Sun TP. 2004. DELLA proteins and gibberellin-regulated seed germination and floral development in *Arabidopsis*. *Plant Physiology* 135: 1008–1019.
- Ubeda-Tomas S, Federici F, Casimiro I, Beechster GTS, Bhalerao R, Swarup R, Doerner P, Haseloff J, Bennett MJ. 2009. Gibberellin signaling in the endodermis controls *Arabidopsis* root meristem size. *Current Biology* 19: 1194–1199.
- Ubeda-Tomas S, Swarup R, Coates J, Swarup K, Laplaze L, Beechster GTS, Hedden P, Bhalerao R, Bennett MJ. 2008. Root growth in *Arabidopsis* requires gibberellin/DELLA signalling in the endodermis. *Nature Cell Biology* 10: 625–628.
- Vaseva II, Qudeimat E, Potuschak T, Du Y, Genschik P, Vandebussche F, van der Straeten D. 2018. The plant hormone ethylene restricts *Arabidopsis* growth via the epidermis. *Proceedings of the National Academy of Sciences, USA* 115: E4130–E4139.
- Wang H, Caruso LV, Downie AB, Perry SE. 2004. The embryo MADS domain protein AGAMOUS-Like 15 directly regulates expression of a gene encoding an enzyme involved in gibberellin metabolism. *Plant Cell* 16: 1206–1219.
- Ward DA, MacMillan J, Gong F, Phillips AL, Hedden P. 2010. Gibberellin 3-oxidases in developing embryos of the southern wild cucumber, *Marah macrocarpus*. *Phytochemistry* 71: 2010–2018.

Wysocka-Diller JW, Helariutta Y, Fukaki H, Malamy JE, Benfey PN. 2000.

Molecular analysis of SCARECROW function reveals a radial patterning mechanism common to root and shoot. *Development* 127: 595–603.

Yadav RK, Girke T, Pasala S, Xie MT, Reddy V. 2009. Gene expression map of the Arabidopsis shoot apical meristem stem cell niche. *Proceedings of the National Academy of Sciences, USA* 106: 4941–4946.

Yadav V, Kundu S, Chattopadhyay D, Negi P, Wei N, Deng XW, Chattopadhyay S. 2002. Light regulated modulation of Z-box containing promoters by photoreceptors and downstream regulatory components, COP1 and HY5, in Arabidopsis. *The Plant Journal* 31: 741–753.

## Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

**Fig. S1** The root tip indicating the regions measured.

**Fig. S2** HPLC-radiochromatograms from incubations of [ $^{17-14}\text{C}$ ]GA<sub>9</sub> and -GA<sub>4</sub> with cell lysates from *E. coli* transformed with *AtGA2ox2* and *YFP* fusions.

**Fig. S3** HPLC-radiochromatograms from incubations of cell lysates from *E. coli* transformed with *AtGA2ox7* and  $^{14}\text{C}$ -labelled C<sub>19</sub>-GAs, GA<sub>9</sub> and GA<sub>4</sub> and C<sub>20</sub>-GAs, GA<sub>12</sub> and GA<sub>24</sub>.

**Fig. S4** YFP fluorescence in roots of representative lines expressing *YFP-AtGA2ox2* or *YFP-AtGA2ox7* in Col-0, indicating expression domains.

**Fig. S5** Comparison of the primary root parameters for three lines expressing *YFP-AtGA2ox2* from tissue-specific promoters in Col-0 with those of Col-0 and *ga1-3* in 7-d-old seedlings.

**Fig. S6** Comparison of the primary root parameters for three lines expressing *YFP-AtGA2ox7* from tissue-specific promoters in Col-0 with those of Col-0 and *ga1-3* in 7-d-old seedlings.

**Fig. S7** The effect on shoot growth in Col-0 of expressing *AtGA2ox2* from tissue-specific promoters. Four to six independent transformants are shown for each construct with Col-0 and *ga1-3*.

**Fig. S8** The effect on shoot growth in Col-0 of expressing *AtGA2ox7* from tissue-specific promoters. Four to six independent

transformants are shown for each construct with Col-0 and *ga1-3*.

**Fig. S9** YFP fluorescence in roots of representative lines expressing *YFP-AtGA2ox1* in the *ga2ox1,2,3* triple mutant, indicating expression domains.

**Fig. S10** Comparison of the primary root parameters for three lines expressing *YFP-AtGA2ox1* from tissue-specific promoters in the *ga2ox1 ga2ox2 ga2ox3* triple mutant with those of Col-0 and the triple *ga20ox* mutant in 7-d-old seedlings.

**Fig. S11** The degree of shoot growth rescue of the *ga20ox1 ga20ox2 ga20ox3* triple mutant from expressing *AtGA20ox1* from tissue-specific promoters.

**Fig. S12** YFP fluorescence in roots of representative lines expressing *YFP-AtGA3ox1* in the *ga3ox1,2* double mutant, indicating expression domains.

**Fig. S13** Comparison of the primary root parameters for three lines expressing *YFP-AtGA3ox1* from tissue-specific promoters in the *ga3ox1 ga3ox2* double mutant with those of Col-0 and the double *ga3ox* mutant in 7-d-old seedlings.

**Fig. S14** The degree of shoot growth rescue of the *ga3ox1 ga3ox2* double mutant from expressing *AtGA3ox1* from tissue-specific promoters.

**Notes S1** Means, least significant differences (LSDs) and degrees of freedom (*df*) for the data presented in Figs 3–7.

**Table S1** Primer sequences for amplification of cDNAs and promoters.

**Table S2** Constructs for tissue-specific expression of GA-metabolism cDNA-GFP fusions indicating the promoter expression domains in roots.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.