

# האוניברסיטה העברית בירושלים

## THE HEBREW UNIVERSITY OF JERUSALEM

---

### ON OPTIMAL ALLOCATION OF A CONTINUOUS RESOURCE USING AN ITERATIVE APPROACH AND TOTAL POSITIVITY

By

JAY BARTROFF, LARRY GOLDSTEIN, YOSEF  
RINOTT and ESTER SAMUEL-CAHN

Discussion Paper # 530

January 2010

מרכז לחקר הרציונליות

CENTER FOR THE STUDY  
OF RATIONALITY

---

Feldman Building, Givat-Ram, 91904 Jerusalem, Israel  
PHONE: [972]-2-6584135      FAX: [972]-2-6513681  
E-MAIL:                      [ratio@math.huji.ac.il](mailto:ratio@math.huji.ac.il)  
URL:                         <http://www.ratio.huji.ac.il/>

# On Optimal Allocation of a Continuous Resource Using an Iterative Approach and Total Positivity

Jay Bartroff<sup>1</sup>, Larry Goldstein<sup>1</sup>, Yosef Rinott<sup>2</sup> and Ester Samuel-Cahn<sup>3</sup>

<sup>1</sup> Department of Mathematics, University of Southern California, Los Angeles, California, USA

<sup>2</sup> Department of Statistics and Center for the Study of Rationality,  
The Hebrew University of Jerusalem, Israel, and LUISS, Rome

<sup>3</sup> Department of Statistics and Center for the Study of Rationality,  
The Hebrew University of Jerusalem, Israel

December 21, 2009

## Abstract

We study a class of optimal allocation problems, including the well-known Bomber Problem, with the following common probabilistic structure. An aircraft equipped with an amount  $x$  of ammunition is intercepted by enemy airplanes arriving according to a homogenous Poisson process over a fixed time duration  $t$ . Upon encountering an enemy, the aircraft has the choice of spending any amount  $0 \leq y \leq x$  of its ammunition, resulting in the aircraft's survival with probability equal to some known increasing function of  $y$ . Two different goals have been considered in the literature concerning the optimal amount  $K(x, t)$  of ammunition spent: (i) Maximizing the probability of surviving for time  $t$ , which is the so-called Bomber Problem, and (ii) maximizing the number of enemy airplanes shot down during time  $t$ , which we call the Fighter Problem. Several authors have attempted to settle the following conjectures about the monotonicity of  $K(x, t)$ : [A]  $K(x, t)$  is decreasing in  $t$ , [B]  $K(x, t)$  is increasing in  $x$ , and [C] the amount  $x - K(x, t)$  held back is increasing in  $x$ . [A] and [C] have been shown for the Bomber Problem with discrete ammunition, while [B] is still an open question. In this paper we consider both time and ammunition continuous, and for the Bomber Problem prove [A] and [C], while for the Fighter we prove [C] in general, and that [A] holds for one special case and [B] for another. These proofs involve showing that the optimal survival probability and optimal number shot down are totally positive of order 2 (TP<sub>2</sub>) in the Bomber and Fighter Problems, respectively. The TP<sub>2</sub> property is shown by constructing convergent sequences of approximating functions through an iterative operation which preserves TP<sub>2</sub> and other properties.

---

<sup>0</sup>AMS 2000 subject classifications. Primary 60G40, Secondary 62L05, 91A60

<sup>0</sup>Key words and phrases: bomber problem, log-concave, optimal allocation, sequential optimization, total positivity

# 1 Introduction

Throughout this paper we write decreasing for non-increasing and increasing for non-decreasing. When we use strict monotonicity, we will say so explicitly. The allocation problem discussed in this paper can be described in terms of the following example: an aircraft, which is at flying time  $t$  away from its final destination, is equipped with an amount  $x$  of ammunition. It is confronted by enemy airplanes whose appearance is driven by a time-homogenous Poisson process with known intensity. By adjusting the units of time, we assume without loss of generality that the intensity is one. It is assumed that the encounters are instantaneous, that is, that their duration time is zero. The question of interest is how much of the presently available ammunition should be spent when confronted by an enemy while in “state”  $(x, t)$ . We consider two problems, the Bomber Problem and the Fighter Problem, with this common probabilistic setup. For the Bomber Problem, the goal is to maximize the probability  $P(x, t)$  of reaching its destination. For the Fighter Problem, the goal is to maximize the expected number  $N(x, t)$  of enemy airplanes shot down. In both problems there is given an increasing function  $a(y)$  which, in the Bomber Problem is the probability that the Bomber *survives* an enemy encounter in which it expends  $0 \leq y \leq x$  units of its ammunition, while in the Fighter Problem  $a(y)$  is the probability that the Fighter *destroys* an enemy when it expends  $y$  units of its ammunition. Although our results below are proved when  $a(y)$  is general, the canonical example considered in the literature for  $a(y)$  in the Bomber Problem is

$$1 - (1 - u)e^{-y} \tag{1}$$

for some fixed  $u \in [0, 1]$ , which can be interpreted as the Bomber’s  $y$  units of ammunition destroying the enemy with probability  $1 - e^{-y}$ , while otherwise allowing the enemy to launch a counterattack which succeeds with probability  $1 - u$ .

For the Fighter, where  $a(y)$  is the probability of destroying the enemy, the probability that the Fighter survives an encounter is  $a(y) + u(1 - a(y))$ , since the enemy is hit with probability  $a(y)$ , and otherwise, with probability  $1 - a(y)$  the enemy counterattacks, and misses with probability  $u$ . When taking the canonical choice  $a(y) = 1 - e^{-y}$ , the Fighter’s survival is given by (1).

While the optimal ammunition spending strategies  $K(x, t)$  may differ for the Bomber and Fighter Problems, it is perhaps intuitively obvious that the following three monotonicity properties, first posed by Klinger and Brown (1968) for the Bomber Problem, should hold for both:

[A]:  $K(x, t)$  is decreasing in  $t$ .

[B]:  $K(x, t)$  is increasing in  $x$ .

[C]:  $x - K(x, t)$  is increasing in  $x$ .

Roughly speaking, [A] states that the closer one is to the destination, the more one should spend, [B] states that the more ammunition one has, the more one should spend, and [C] states that the more ammunition one has, the more one should retain for possible future encounters; replacing “more” by “not less” makes these characterizations precise. Although we write  $K(x, t)$  for the two different goals, no confusion should arise, as it will always be clear which of the problems we are discussing, or that we are discussing them simultaneously. An optimal policy may not be unique. If it is not unique,  $K(x, t)$  will always denote the minimal optimal amount of ammunition needed.

Although the paper is written in *plane* English, the settings above serve only as illustrations of the general method which can be applied to a wide class of problems where some limited resource

needs to be optimally allocated over time or space to maximize the probability of a system's survival, which clearly also have more peaceful applications.

Of the two problems, the Bomber Problem has received most attention. For discrete ammunition, such as missiles, it is first treated in Klinger and Brown (1968), who prove that [B] implies [A]. A clearer picture emerges in Samuel (1970), who proves [A] without assuming [B]. Samuel's (1970) formula (3.9) also proves [C]. Conjecture [B] has remained elusive, and in fact remains in doubt. Simons and Yao (1990) argue that when both time and ammunition are continuous, then [B] is equivalent to the log concavity of  $P(x, t)$  in  $x$  for each fixed  $t$ . They also show that when both time and ammunition are discrete, there exist parameter values for which log concavity fails. However, they are unable to supply an example where [B] fails.

Regarding the Fighter Problem, we are unaware of this variation on the Bomber Problem being discussed in a finite time  $t$  horizon setting. For an infinite time horizon, an optimal policy can be written simply as  $K(x)$ . Weber (1985) shows that the monotonicity of  $K(x)$  fails to hold for discrete ammunition when  $a$  is given by (1) with  $u = 0$ , and Shepp et al. (1991) show that monotonicity fails for continuous ammunition. In a setting where both ammunition and time are discrete, the duration  $T$  is an unknown exponential random variable with known mean, and geometric arrivals of the enemy, Shepp et al. (1991) also show that there exist parameter values for which [B] fails. Though this variant is on a finite time interval of length  $T$ , the optimal policy does not depend on  $t$  and so resembles an infinite time horizon problem. On the other hand, Bartroff et al. (2009) show for the Bomber Problem with continuous time and ammunition that there exists a certain region of the  $(x, t)$ -space over which  $K(x, t) = x$ , a monotone function of  $x$ , and hence [B] holds in this region.

In the present paper we consider the "doubly continuous" setting where ammunition and time are continuous variables. Given an increasing function  $a(y)$  taking values in  $[0, 1]$ , the dynamic programming equation for the Bomber Problem giving the optimal survival probability  $P(x, t)$  is

$$P(x, t) = \int_0^t \sup_{0 \leq y \leq x} a(y)P(x - y, t - s)e^{-s}ds + e^{-t} \quad \text{for } (x, t) \in \mathbb{R}^+ \times \mathbb{R}^+.$$

The explanation is as follows. There are no encounters in a time interval of length  $t$  with probability  $e^{-t}$ , yielding the final term. Otherwise, an encounter occurs at time  $s \in [0, t]$  with density  $e^{-s}$ , at which time the amount  $y \in [0, x]$  will be chosen to maximize the probability  $a(y)$  of surviving the current encounter times the probability of future survival under an optimal policy. A simple change of variables yields

$$P(x, t) = \int_0^t \sup_{0 \leq y \leq x} a(y)P(x - y, s)e^{-(t-s)}ds + e^{-t} \quad \text{for } (x, t) \in \mathbb{R}^+ \times \mathbb{R}^+. \quad (2)$$

Given an increasing function  $a(y)$  taking values in  $[0, 1]$  and a fixed value  $u \in [0, 1]$ , the dynamic programming equation for the Fighter Problem giving the optimal expected number  $N(x, t)$  of enemies shot down is

$$N(x, t) = \int_0^t \sup_{0 \leq y \leq x} \{a(y) + [a(y) + u(1 - a(y))]N(x - y, s)\}e^{-(t-s)}ds \quad \text{for } (x, t) \in \mathbb{R}^+ \times \mathbb{R}^+, \quad (3)$$

which can be interpreted as follows. Encounters occur with density  $e^{-s}$ , and expending amount  $y$  the Fighter gains a single hit if he destroys the enemy, which happens with probability  $a(y)$ . The Fighter will gain an expected additional  $N(x - y, t - s)$  future hits if he destroys the enemy

or otherwise if the enemy's counterattack fails, which happens with probability  $u$ . A change of variables then yields (3).

By putting both the Bomber and the Fighter in a common general framework, we show that the solutions to (2) and (3) exist and are unique and continuous. In particular, the supremum is attained in both (2) and (3), and hence sup can be replaced by max in each. The optimal policy  $K(x, t)$  is then defined as the minimal value for which the max in the integrands in (2) and (3) is attained, respectively.

Recall that a nonnegative function  $Q(x, t)$  is *totally positive of order 2*, written  $TP_2(x, t)$  or simply  $TP_2$ , if

$$Q(x', t')Q(x, t) \geq Q(x', t)Q(x, t') \quad \text{whenever } x < x' \text{ and } t < t', \quad (4)$$

and strictly  $TP_2$  if  $\geq$  in (4) can be replaced by  $>$ . To prove [A] for the Bomber Problem, we establish the  $TP_2$  property of  $P(x, t)$  in Section 3.

**Theorem 1.1.** *Let  $a(y)$  be a uniformly continuous log-concave function. Then a unique bounded solution  $P(x, t)$  to equation (2) exists, satisfies  $P(x, t) \in [0, 1]$ , is continuous and  $TP_2$ .*

**Theorem 1.2.** *If  $a(y)$  is a uniformly continuous log-concave function, then [A] holds for the doubly continuous Bomber Problem.*

Simons and Yao (1990) claim that [C] can be shown to hold for continuous  $x$  and  $t$  by arguments similar to the ones they provide for a case where both  $x$  and  $t$  are discrete; in Section 3 we provide a rigorous proof of [C] for the doubly continuous case. The discrete  $x$  analogues to Theorems 1.1 and 1.2 were proved in Samuel (1970).

Regarding the Fighter in general, we have the following result.

**Theorem 1.3.** *If  $a(y)$  is uniformly continuous, then for any  $u \in [0, 1]$  a bounded solution  $N(x, t)$  to equation (3) exists, is unique, satisfies  $N(x, t) \in [0, t]$ , and is continuous.*

For the Fighter Problem we do not resolve the question of whether the general solution to (3) is  $TP_2$ , but instead examine two special cases in more detail:

i) *The Frail Fighter*: the case  $u = 0$ . Once the Frail Fighter fails to shoot down an enemy, he himself is shot down with probability one. Equation (3) then simplifies to

$$N(x, t) = \int_0^t \sup_{0 \leq y \leq x} a(y)[1 + N(x - y, s)]e^{-(t-s)} ds \quad \text{for } (x, t) \in \mathbb{R}^+ \times \mathbb{R}^+. \quad (5)$$

**Theorem 1.4.** *If  $a(y)$  is a uniformly continuous log-concave function, then the solution  $N(x, t)$  is  $TP_2$ , and property [A] holds for the Frail Fighter.*

The Frail Fighter is discussed in Section 4, where [C] is also established.

ii) *The Invincible Fighter*: the case  $u = 1$ . As the name suggests, the Invincible Fighter cannot be shot down. Equation (3) simplifies to

$$N(x, t) = \int_0^t \sup_{0 \leq y \leq x} [a(y) + N(x - y, s)]e^{-(t-s)} ds \quad \text{for } (x, t) \in \mathbb{R}^+ \times \mathbb{R}^+. \quad (6)$$

**Theorem 1.5.** *If  $a(y)$  is a uniformly continuous concave function then property [B] holds for the Invincible Fighter.*

The Invincible Fighter is discussed in Section 5, where we also show [C] and that the maximum  $y$  determining  $K(x, t)$  is uniquely attained. It should be noted that for none of the problems considered here have we managed to show that [A] and [B] hold simultaneously.

While in the discrete ammunition case one can demonstrate claims by using induction on the number of available units (e.g., Simons and Yao, 1990), here we must take a different approach. In Section 2 we prove the existence and uniqueness of solutions to equations which have the form of (2) or (3) by following the outline suggested by Weber (Simons and Yao, 1990, p. 431) and construct a sequence of functions through an iteration which converges exponentially fast to the solution.

We illustrate our approach using the Bomber Problem; similar remarks apply to the Fighter. First, consider a slightly transformed version of (2). For ease of notation, for functions  $a(\cdot)$  and  $Q(\cdot, \cdot)$  let

$$(a \star Q)(x, s) = \sup_{0 \leq y \leq x} a(y)Q(x - y, s), \quad (7)$$

the sup-convolution of  $a(\cdot)$  and  $Q(\cdot, s)$ . Letting

$$\bar{P}(x, t) = e^t P(x, t), \quad (8)$$

substitution into (2) results in the integral equation

$$\bar{P}(x, t) = \int_0^t (a \star \bar{P})(x, s) ds + 1 \quad \text{for } (x, t) \in \mathbb{R}^+ \times \mathbb{R}^+. \quad (9)$$

Clearly there exists a unique, continuous solution  $\bar{P}$  to (9) satisfying  $0 \leq \bar{P}(x, t) \leq e^t$  if and only if there exists a unique, continuous solution  $P$  to (2) satisfying  $0 \leq P(x, t) \leq 1$ ; Theorem 2.1 proves the former and provides an iteration method to approximate the solution. Theorem 3.1 provides further properties of the solution.

To prove the claims of Theorem 3.1, consider the sequence of functions  $\{\bar{P}_m\}_{m \geq 0}$  generated with some initial  $\bar{P}_0(x, t)$  by the recursion

$$\bar{P}_{m+1}(x, t) = \int_0^t (a \star \bar{P}_m)(x, s) ds + 1 \quad \text{for } m \geq 0. \quad (10)$$

In Section 2, by showing that the operation which gives  $\bar{P}_{m+1}$  from  $\bar{P}_m$  is a type of contraction, we are able to prove the existence and uniqueness of the solution to (10) by the method of the contraction mapping theorem. In Section 3 we prove that various operations preserve the  $\text{TP}_2$  and other properties, allowing us to prove inductively that  $\bar{P}_m$  possesses a particular property for all  $m \geq 0$ , that may then be inherited by  $\bar{P}$ . We remark in Section 6 why this same approach appears to fail to give [B]. Nevertheless, the sequence of approximate solutions, which are guaranteed to converge in the supremum norm at an exponential rate by Theorem 2.1, may yet have uses in addition to that of proving [A].

## 2 Existence, uniqueness, and approximating sequences

We will put our models into the following common framework. Suppose that for given functions  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  and  $Q : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ , an operation  $\otimes$  returns a function  $a \otimes Q : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$ .

Although our results in this section are more general, we will be interested in the specific cases where  $(a \otimes Q)(x, t)$  is one of the following:

$$\begin{aligned}
(Bo) \quad & \sup_{0 \leq y \leq x} a(y)Q(x-y, t), \\
(F_0) \quad & \sup_{0 \leq y \leq x} a(y)[e^t + Q(x-y, t)] \\
(F_1) \quad & \sup_{0 \leq y \leq x} [a(y)e^t + Q(x-y, t)], \\
(F_u) \quad & \sup_{0 \leq y \leq x} \{a(y)e^t + [a(y) + u(1-a(y))]Q(x-y, t)\}.
\end{aligned} \tag{11}$$

Case  $(Bo)$  corresponds to the Bomber Problem (9), case  $(F_u)$  to the general Fighter equation (3) after a rescaling by  $e^t$  as in (8),  $(F_0)$  the case  $u = 0$  of  $(F_u)$ , the Frail Fighter, and case  $(F_1)$  the case  $u = 1$  of  $(F_u)$ , the Invincible Fighter.

Given  $a$  and  $\otimes$  we will consider operators mapping the collection of functions  $Q : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$  to itself. For the Fighter Problem, these operators will be of the form

$$\mathcal{G}(x, t, Q) = \int_0^t (a \otimes Q)(x, s) ds, \tag{12}$$

and for the Bomber Problem,

$$\mathcal{F}(x, t, Q) = \int_0^t (a \otimes Q)(x, s) ds \quad \text{and} \quad \mathcal{G}(x, t, Q) = \mathcal{F}(x, t, Q) + 1. \tag{13}$$

For simplicity, in what follows we may write  $\mathcal{G}(Q)$  in place of  $\mathcal{G}(x, t, Q)$ . For a given function  $Q_0(x, t)$  on  $\mathbb{R}^+ \times \mathbb{R}^+$ , we study the sequence of iterates of  $\mathcal{G}$  given by

$$Q_{m+1}(x, t) = \mathcal{G}(x, t, Q_m) \quad \text{for } m \geq 0. \tag{14}$$

The main result of this section, given in Theorems 2.1 and 2.2, proves the existence and uniqueness of a solution  $\bar{Q}$  to the equation

$$\bar{Q}(x, t) = \mathcal{G}(x, t, \bar{Q}), \tag{15}$$

and the exponential rate of convergence of the sequence  $Q_m$  to  $\bar{Q}$  when  $Q_0$  is continuous and  $a$  and  $\otimes$  satisfy certain continuity-type properties. These conditions make  $\mathcal{G}$  into a type of contraction, allowing us to apply the methods of the contraction mapping theorem (see, for instance, Luenberger (1969)) to prove that sequences of functions generated by the iterates of  $\mathcal{G}$  on some initial function converge to a unique fixed point of  $\mathcal{G}$ .

For a function  $Q$  bounded over all compact domains  $D \subset \mathbb{R}^+ \times \mathbb{R}^+$ , let

$$\|Q\|_D = \sup_{(y,s) \in D} |Q(y, s)|.$$

For  $Q$  and  $R$  any two functions defined on  $D \subset \mathbb{R}^+ \times \mathbb{R}^+$  and  $\delta \in [0, \infty)$ , let

$$\mathbf{d}_{D, \delta}^1(Q, R) = \sup_{\{(x,t), (x',t) \in D : |x-x'| \leq \delta\}} |Q(x, t) - R(x', t)|,$$

and

$$\mathbf{d}_{D, \delta}(Q, R) = \sup_{\{(x,t), (x',t') \in D : |x-x'| \leq \delta, |t-t'| \leq \delta\}} |Q(x, t) - R(x', t')|.$$

**Definition 2.1.** For  $\kappa \geq 0$  we will say that  $\otimes$  is of  $\kappa$ -contraction type if for all compact domains  $D = [0, X] \times [0, T]$  there exist  $\beta \geq 0$  and  $\eta(\delta)$ , defined for  $\delta \geq 0$  and satisfying

$$\eta(0) = 0 \quad \text{and} \quad \lim_{\delta \downarrow 0} \eta(\delta) = 0,$$

such that for any functions  $Q$  and  $R$  on  $\mathbb{R}^+ \times \mathbb{R}^+$ ,

$$\mathbf{d}_{D,\delta}^1((a \otimes Q), (a \otimes R)) \leq \kappa \mathbf{d}_{D,\delta}^1(Q, R) + \eta(\delta)(\|Q\|_D \vee \|R\|_D + \beta), \quad (16)$$

for all  $\delta \in [0, \infty)$ , where  $a \vee b$  is the maximum of  $a$  and  $b$ . We say that  $\otimes$  is of contraction type if it is of  $\kappa$ -contraction type for some  $\kappa \geq 0$ .

**Lemma 2.1.** Let  $\mathcal{G}$  be given by (12) or (13) for  $\otimes$  of  $\kappa$ -contraction type. Let  $X \geq 0$ ,  $T' \geq T \geq 0$  be arbitrary, and set  $D' = [0, X] \times [0, T']$  and  $D = [0, X] \times [0, T]$ . If  $Q$  and  $R$  are any functions defined on  $D'$ , then

$$\|\mathcal{G}(Q) - \mathcal{G}(R)\|_{D'} \leq \kappa T \|Q - R\|_D + \kappa(T' - T) \|Q - R\|_{D'}, \quad (17)$$

and when  $a(y)$  is bounded, for all  $\delta \geq 0$ , the inequality

$$\mathbf{d}_{D,\delta}(\mathcal{G}(Q), \mathcal{G}(R)) \leq \kappa T \mathbf{d}_{D,\delta}^1(Q, R) + \omega(\delta)(\|Q\|_D \vee \|R\|_D + \gamma) \quad (18)$$

holds for some  $\gamma \geq 0$  and  $\omega(\delta)$  satisfying  $\omega(0) = 0$  and  $\lim_{\delta \downarrow 0} \omega(\delta) = 0$ .

*Proof.* Clearly it suffices to prove the claim for  $\mathcal{G}$  of the form (12). To obtain (17) let  $(x, t) \in D'$ . Applying (16) with  $\delta = 0$ , and noting  $\mathbf{d}_{D,0}^1(Q, R) = \|Q - R\|_D$ , we have

$$\begin{aligned} & |\mathcal{G}(x, t, Q) - \mathcal{G}(x, t, R)| \\ & \leq \int_0^t |(a \otimes Q)(x, s) - (a \otimes R)(x, s)| ds \leq \int_0^{T'} |(a \otimes Q)(x, s) - (a \otimes R)(x, s)| ds \\ & = \int_0^T |(a \otimes Q)(x, s) - (a \otimes R)(x, s)| ds + \int_T^{T'} |(a \otimes Q)(x, s) - (a \otimes R)(x, s)| ds \\ & \leq \int_0^T \kappa \mathbf{d}_{D,0}^1(Q, R) ds + \int_T^{T'} \kappa \mathbf{d}_{D',0}^1(Q, R) ds \\ & \leq \kappa T \|Q - R\|_D + \kappa(T' - T) \|Q - R\|_{D'}. \end{aligned}$$

Taking supremum on the left hand side over  $(x, t) \in D'$ , we obtain (17).

To obtain (18), let  $\delta \geq 0$  and suppose  $(x, t), (x', t') \in D$  satisfy  $|x - x'| \leq \delta$  and  $|t - t'| \leq \delta$ ; assume first that  $t' \geq t$ . Then, applying (16), for each of the four cases of (11) we have

$$\begin{aligned} & |\mathcal{G}(x', t', Q) - \mathcal{G}(x, t, R)| \\ & \leq \int_0^t |(a \otimes Q)(x', s) - (a \otimes R)(x, s)| ds + \int_t^{t'} |(a \otimes Q)(x', s)| ds \\ & \leq T (\kappa \mathbf{d}_{D,\delta}^1(Q, R) + \eta(\delta)(\|Q\|_D \vee \|R\|_D) + \beta) + \delta \|a\| (e^T + \|Q\|_D), \end{aligned}$$

where  $\|a\|$  is the supremum of  $a$  on  $[0, \infty)$ . If we assume that  $t' \leq t$  we obtain a similar bound, except that the last term has  $\|R\|_D$  in place of  $\|Q\|_D$ . Taking the above two cases into account and taking supremum on the left hand side, we obtain (18) with  $\gamma = \beta + e^T$  and  $\omega(\delta) = T\eta(\delta) + \delta\|a\|$ . ■



**Definition 2.2.** If  $\mathcal{G}$  satisfies the conclusions of Lemma 2.1 for some  $\kappa \geq 0$  and some  $\gamma \geq 0$  and  $\omega(\delta)$ , we say  $\mathcal{G}$  is of  $\kappa$ -contraction type. We say that  $\mathcal{G}$  is of contraction type if it is of  $\kappa$ -contraction type for some  $\kappa \geq 0$ .

Tautologically, under Definition 2.2, Lemma 2.1 says that if  $\mathcal{G}$  is given by (12) with  $\otimes$  of contraction type, then  $\mathcal{G}$  is of contraction type.

**Proposition 2.1.** If  $\mathcal{G}_1$  and  $\mathcal{G}_2$  are of  $\kappa_1$  and  $\kappa_2$ -contraction type, respectively, then  $\mathcal{G}_1 + \mathcal{G}_2$  is of contraction  $\kappa_1 + \kappa_2$  type, and  $\alpha\mathcal{G}_1$  is of  $|\alpha|\kappa_1$ -contraction type for any  $\alpha \in \mathbb{R}$ . Hence the collection of contraction type operators form a vector space over  $\mathbb{R}$ . The constant operator  $\mathcal{G}(\cdot) = 1$  is a contraction operator of type 0.

*Proof.* The triangle inequality shows the first claim, the remaining claims are self evident.  $\blacksquare$

We note that Proposition 2.1 shows that if  $\mathcal{G}$  is of contraction type then so is  $\mathcal{G} + 1$ . The next corollary shows that any  $\mathcal{G}$  of contraction type preserves continuity.

**Corollary 2.1.** If  $a(y)$  is bounded,  $Q$  is continuous on  $\mathbb{R}^+ \times \mathbb{R}^+$ , and  $\mathcal{G}$  is of contraction type, then  $\mathcal{G}(Q)$  is continuous.

*Proof.* Given  $\varepsilon > 0$ , by the uniform continuity of  $Q$  in any compact domain  $D$  of the form  $[0, X] \times [0, T]$ , and, since  $\omega(\delta) \downarrow 0$  as  $\delta \downarrow 0$ , there exists  $\delta > 0$  such that  $\kappa T \mathbf{d}_{D, \delta}^1(Q, Q) < \varepsilon/2$  and  $\omega(\delta)(\|Q\|_D + \gamma) < \varepsilon/2$ . Hence  $\mathbf{d}_{D, \delta}(\mathcal{G}(Q), \mathcal{G}(Q)) < \varepsilon$  by (18).  $\blacksquare$

**Theorem 2.1.** Let  $Q_0(x, t)$  be any continuous function on  $\mathbb{R}^+ \times \mathbb{R}^+$ , and let  $Q_m, m = 1, 2, \dots$ , be given by (14) for some  $\mathcal{G}$  of  $\kappa$ -contraction type. Then  $Q_m, m = 0, 1, \dots$ , are continuous and converge uniformly on compact subsets of  $\mathbb{R}^+ \times \mathbb{R}^+$  to a continuous function  $\bar{Q}$  that satisfies (15).

Additionally, for every compact domain  $D \subset \mathbb{R}^+ \times \mathbb{R}^+$  there exists a constant  $C > 0$  and  $\theta \in (0, 1)$  such that

$$\|\bar{Q} - Q_m\|_D \leq C\theta^m \quad \text{for all } m = 0, 1, \dots$$

*Proof.* Let  $\kappa > 0$  as otherwise  $Q_m = 0$  for all  $m \geq 1$  and the result is trivial. Let  $X \geq 0$  be arbitrary, and let  $\tilde{T}$  be the supremum over all  $T \geq 0$  such that there exists  $C > 0$  and  $\theta \in (0, 1)$  such that

$$c_{m, T} \leq C\theta^m \quad \text{for } m = 1, 2, \dots, \text{ where } c_{m, T} = \|Q_m - Q_{m-1}\|_{[0, X] \times [0, T]}. \quad (19)$$

We first prove that  $\tilde{T} = \infty$ . Let  $\theta \in (0, 1)$ . Using (17) of Lemma 2.1 with  $T = 0$  and  $T' = \theta/\kappa$  we obtain

$$c_{m, T'} \leq \left(\frac{c_{1, T'}}{\theta}\right) \theta^m \quad \text{for } m = 1, 2, \dots$$

by induction, first noting that it holds trivially for  $m = 1$ . Hence  $\tilde{T} > 0$ .

If (19) holds for some  $C > 0$ ,  $\vartheta \in (0, 1)$  and  $T > 0$  then, for  $T' = T + \vartheta/\kappa$ , using (17) again

$$c_{m, T'} \leq \kappa T c_{m-1, T} + \kappa(T' - T)c_{m-1, T'} \leq \kappa T C \vartheta^{m-1} + \vartheta c_{m-1, T'} \quad \text{for } m \geq 2. \quad (20)$$

The inequality

$$c_{m, T'} \leq C' m \vartheta^{m-1} \quad \text{where } C' = \max\{\kappa T C, c_{1, T'}\}$$

holds trivially for  $m = 1$ , and now induction, using (20), shows it holds for all  $m \geq 1$ . Elementary calculus now shows that  $c_{m, T'} \leq C' m \vartheta^{m-1}$  for  $m \geq 1$  implies that for any  $\theta \in (\vartheta, 1)$  there exists  $C$  such that  $c_{m, T'} \leq C\theta^m$  for all  $m \geq 1$ . Hence (19) holds when replacing  $T$  by  $T'$ , and therefore assuming  $\tilde{T} < \infty$  leads to a contradiction. Now note that (19) holds when replacing  $[0, X] \times [0, T]$

by any compact  $D$ , and  $c_{m,T}$  by  $c_{m,D} = \|Q_m - Q_{m-1}\|_D$ , as  $D \subset [0, X] \times [0, T]$  for some  $X$  and  $T$  sufficiently large.

By Corollary 2.1 and induction, the functions  $Q_m$ ,  $m = 0, 1, \dots$ , are continuous and by (19) they form a Cauchy sequence in the supremum norm on any compact domain  $D$ . Therefore  $Q_m$ ,  $m = 0, 1, \dots$ , converges uniformly on any compact domain to a continuous limit function  $\bar{Q}$ . The function  $\bar{Q}$  solves (15) as, applying the triangle inequality and then (17) of Lemma 2.1 with  $T' = T$  yields

$$\begin{aligned} \|\mathcal{G}(\bar{Q}) - \bar{Q}\|_D &\leq \|\mathcal{G}(\bar{Q}) - Q_{m+1}\|_D + \|Q_{m+1} - \bar{Q}\|_D \\ &= \|\mathcal{G}(\bar{Q}) - \mathcal{G}(Q_m)\|_D + \|Q_{m+1} - \bar{Q}\|_D \\ &\leq \kappa T \|\bar{Q} - Q_m\|_D + \|Q_{m+1} - \bar{Q}\|_D, \end{aligned}$$

whose right hand side converges to zero as  $m \rightarrow \infty$ . Hence  $\mathcal{G}(\bar{Q}) = \bar{Q}$ .

Lastly, by the triangle inequality, for all  $n \geq m$ , by (19)

$$\|Q_n - Q_m\|_D \leq C \sum_{j=m+1}^n \theta^j \leq \left( \frac{C}{1-\theta} \right) \theta^{m+1},$$

so letting  $n \rightarrow \infty$  yields the final claim of the theorem.  $\blacksquare$

**Theorem 2.2.** *If  $\mathcal{G}$  is of contraction type, equation (15) has a unique solution that is bounded over all compact domains of  $\mathbb{R}^+ \times \mathbb{R}^+$ .*

*Proof.* Let  $Q$  and  $R$  be two solutions to (15) that are bounded over all compact subsets of  $\mathbb{R}^+ \times \mathbb{R}^+$ . Let  $D = [0, X] \times [0, T]$  and  $D' = [0, X] \times [0, T']$ , for some  $X \geq 0$  and  $T' \geq T \geq 0$ . Taking  $T = 0$  and  $\kappa T' < 1$  in (17) yields  $\|Q - R\|_{D'} = 0$ .

Replacing  $T$  and  $T'$  by  $T'$  and  $T''$ , respectively, in (17), where  $0 \leq \kappa(T'' - T') < 1$ , it is easy to see that for the larger domain  $D'' = [0, X] \times [0, T'']$  we now obtain

$$\|Q - R\|_{D''} = \|\mathcal{G}(Q) - \mathcal{G}(R)\|_{D''} \leq \kappa(T'' - T') \|Q - R\|_{D''}. \quad (21)$$

This implies that  $\|Q - R\|_{D''} = 0$ , and it follows that  $\|Q - R\|_D = 0$  holds for any domain  $[0, X] \times [0, T]$ , and hence for any compact domain  $D$ .  $\blacksquare$

In the cases of interest to us here, the operator  $\mathcal{G}$  has some additional properties, in particular, monotonicity and a type of boundedness.

**Definition 2.3.** *We say that an operator  $\mathcal{G}$  is  $B$ -bounded for a continuous function  $B : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$  if*

$$Q \leq B \quad \text{implies} \quad \mathcal{G}(Q) \leq B.$$

*We say that  $\mathcal{G}$  is monotone if*

$$Q \leq R \quad \text{implies} \quad \mathcal{G}(Q) \leq \mathcal{G}(R).$$

Clearly the operator  $\mathcal{G}$  of (12) is monotone for all four choices in (11); hence so is  $\mathcal{G}$  of (13) with  $\otimes$  given by case (Bo). The following proposition is immediate from the preceding definitions.

**Proposition 2.2.** *Let  $\mathcal{G}$  be a monotone,  $B$ -bounded operator of contraction type, and let  $Q_m$ ,  $m = 0, 1, \dots$ , be the functions given by (14) when  $Q_0(x, t) \equiv 0$ , and  $Q'_m$  the functions when  $Q'_0(x, t) = B(x, t)$ . Then the unique solution  $\bar{Q}$  to (15) satisfies*

$$0 \leq Q_m(x, t) \leq \bar{Q}(x, t) \leq Q'_m(x, t) \leq B(x, t) \quad \text{for all } m = 0, 1, \dots$$

If the iteration (14) is used for numerical approximation of the solution  $\overline{Q}$  to (15), then Proposition 2.2 allows for the verification of the precision of the calculation, as it provides a lower and upper bound for  $\overline{Q}$ ; the distance between these bounds therefore gives the approximation error.

We now show that when  $a$  is a bounded function,  $\mathcal{G}$  of (12) is of contraction type in all cases of (11), and that the same is true for  $\mathcal{G}$  of (13) with  $\otimes$  given in case (Bo).

**Lemma 2.2.** *Let  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be uniformly continuous and bounded by  $\kappa$ . The operator  $\mathcal{G}$  given by (12) is of  $\kappa$ -contraction type in cases (Bo) and ( $F_0$ ) of (11), of 1-contraction type in case ( $F_1$ ), and of  $[(1-u)\kappa + u]$ -contraction type in case ( $F_u$ ). The operator  $\mathcal{G}$  given by (13) with  $\otimes$  as in case (Bo) is of  $\kappa$  contraction type.*

*Proof.* For the first claim, with  $\otimes$  given by case (Bo) of (11) it suffices, by Lemma 2.1, to prove that  $\otimes$  is of  $\kappa$ -contraction type. Let a compact  $D = [0, X] \times [0, T]$  and  $\delta \geq 0$  be given and  $(x, t), (x', t) \in D$  with  $|x - x'| \leq \delta$ . For a given  $\epsilon > 0$  let  $y^* \in [0, x']$  achieve  $\sup_{0 \leq y \leq x'} a(y)Q(x' - y, t)$  to within  $\epsilon$ , and set  $y_1^* = \min\{y^*, x\}$ . Hence

$$\begin{aligned} & (a \otimes Q)(x', t) - (a \otimes R)(x, t) \\ & \leq a(y^*)Q(x' - y^*, t) - a(y_1^*)R(x - y_1^*, t) + \epsilon \\ & = a(y^*)[Q(x' - y^*, t) - R(x - y_1^*, t)] + (a(y^*) - a(y_1^*))R(x - y_1^*, t) + \epsilon. \end{aligned}$$

Since  $0 \leq a(y) \leq \kappa$  for all  $y \geq 0$  and  $|(x' - y^*) - (x - y_1^*)| \leq \delta$ , with

$$\eta(\delta) = \sup_{|y' - y| \leq \delta} |a(y') - a(y)|,$$

we obtain

$$(a \otimes Q)(x', t) - (a \otimes R)(x, t) \leq \kappa \mathbf{d}_{D, \delta}^1(Q, R) + \eta(\delta) \|R\|_D. \quad (22)$$

Similarly, there exists another  $y^* \in [0, x]$  and corresponding  $y_1^* \in [0, x']$  such that

$$(a \otimes Q)(x', t) - (a \otimes R)(x, t) \geq a(y_1^*)Q(x' - y_1^*, t) - a(y^*)R(x - y^*, t) - \epsilon,$$

which yields a bound of the form (22) with the roles of  $Q$  and  $R$  interchanged. Since  $\epsilon$  is arbitrary

$$|(a \otimes Q)(x', t) - (a \otimes R)(x, t)| \leq \kappa \mathbf{d}_{D, \delta}^1(Q, R) + \eta(\delta) (\|Q\|_D \vee \|R\|_D).$$

Taking supremum on the left hand side and noting that  $\eta(0) = 0$  and  $\lim_{\delta \downarrow 0} \eta(\delta) = 0$  by the uniform continuity of  $a(y)$  yields the desired conclusion. The first and final claims of the lemma are now immediate.

In case ( $F_u$ ) we have in a similar way

$$\begin{aligned} & (a \otimes Q)(x', t) - (a \otimes R)(x, t) \\ & \leq a(y^*)e^t + [a(y^*) + u(1 - a(y^*))]Q(x - y^*, t) - a(y_1^*)e^t - [a(y_1^*) + u(1 - a(y_1^*))]R(x - y_1^*, t) + \epsilon. \end{aligned}$$

Adding and subtracting terms as before, we obtain

$$\begin{aligned} & (a(y^*) - a(y_1^*))e^t + [(1 - u)a(y^*) + u][Q(x - y^*, t) - R(x - y_1^*, t)] \\ & + [(a(y^*) - a(y_1^*))(1 - u)]R(x - y_1^*, t) + \epsilon \\ & \leq [(1 - u)\kappa + u] \mathbf{d}_{D, \delta}^1(Q, R) + \eta(\delta)e^T (\|R\|_D + 1) + \epsilon, \end{aligned}$$

since  $|(1 - u)a(y^*) + u| \leq (1 - u)\kappa + u$ . The result follows as for case (Bo). Now cases ( $F_0$ ) and ( $F_1$ ) follow by specializing ( $F_u$ ) to  $u = 0$  and  $u = 1$ , respectively.  $\blacksquare$

**Lemma 2.3.** *Let  $\mathcal{G}$  be given by (13) for case (Bo) of (11). If  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is bounded by  $\kappa$  then  $\mathcal{G}$  is  $e^{\kappa t}$ -bounded.*

*Proof.* If  $Q(x, s) \leq e^{\kappa s}$ , using that  $0 \leq a(y) \leq \kappa$  for all  $y \geq 0$ , we have for all  $x \geq 0$ ,

$$\mathcal{G}(x, t, Q) \leq \kappa \int_0^t \sup_{0 \leq y \leq x} Q(y, s) ds + 1 \leq \kappa \int_0^t e^{\kappa s} ds + 1 = e^{\kappa t},$$

proving the boundedness claim. ■

**Lemma 2.4.** *Consider case ( $F_u$ ) of (11) with  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  bounded by  $\kappa$ , and let  $\alpha = (1 - u)\kappa + u$ . Then  $\mathcal{G}$  of (12) is  $B$ -bounded where  $B(x, t) = B_\alpha(t)$  is given by*

$$B_\alpha(t) = \begin{cases} \kappa t e^t & \alpha = 1 \\ \frac{\kappa}{1-\alpha} (e^t - e^{\alpha t}) & \alpha \neq 1. \end{cases}$$

*In particular, in case ( $F_0$ ) the operator  $\mathcal{G}$  is  $B_\kappa(t)$ -bounded, and in case ( $F_1$ ) is  $\kappa t e^t$ -bounded.*

*Proof.* It is easily checked in both the  $\alpha = 1$  and  $\alpha \neq 1$  cases that if  $Q(x, t) \leq B_\alpha(t)$ , then

$$\mathcal{G}(x, t, Q) \leq \int_0^t (\kappa e^s + \alpha B_\alpha(s)) ds = B_\alpha(t).$$

The following corollary summarizes the results above. ■

**Corollary 2.2.** *Let  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be uniformly continuous and bounded by  $\kappa$ . Then, with  $\mathcal{G}$  given by (12) and  $\otimes$  any of the four cases of (11), or  $\mathcal{G}$  given by (13) with  $\otimes$  given in case (Bo) of (11), there exists a continuous solution to the equation  $\mathcal{G}(Q) = Q$ , and there are no other solutions bounded over compact domains. Moreover, with  $Q_0(x, t)$  any continuous function from  $\mathbb{R}^+ \times \mathbb{R}^+$  to  $\mathbb{R}^+$ , the sequence of iterates of  $\mathcal{G}$  given by  $Q_{m+1} = \mathcal{G}(Q_m)$  for  $m \geq 0$  converges uniformly to the solution of  $\mathcal{G}(Q) = Q$  at an exponential rate. For every  $m$ , the solution to  $\mathcal{G}(Q) = Q$  is bounded below by the  $m^{\text{th}}$  term of the sequence initialized with  $Q_0(x, t) = 0$ , and if  $\mathcal{G}$  is  $B$ -bounded, then it is also bounded above by the  $m^{\text{th}}$  term of the sequence initialized with  $Q_0(x, t) = B(x, t)$ .*

We conclude this section with a proof of Theorem 1.3.

*Proof of Theorem 1.3.* Let  $\mathcal{G}$  be given by (12) and  $\otimes$  by ( $F_u$ ) of (11) for  $u \in [0, 1]$ . By Lemma 2.2, when  $a$  is a probability, the operator  $\mathcal{G}$  is  $t e^t$ -bounded. By Corollary 2.2 the unique solution  $\bar{N}$  to  $\mathcal{G}(Q) = Q$  exists, is continuous and satisfies  $0 \leq \bar{N}(x, t) \leq t e^t$ . Now letting  $\bar{N}(x, t) = e^t N(x, t)$  as in (8), we have  $N(x, t) \in [0, t]$ , as claimed. ■

### 3 The Bomber

In this section we consider the dynamic programming equation (9), and specialize the results of the previous section to  $\otimes$  given by case (Bo) of (11). Let  $\mathcal{F}(Q)$  and  $\mathcal{G}(Q)$  be given by (13). In particular, equation (9) may be written more compactly as  $\mathcal{G}(\bar{P}) = \bar{P}$ . Throughout this section  $a(y)$  can be interpreted as the probability of the Bomber surviving an enemy encounter after expending an amount  $y$  of its ammunition, although our results hold in greater generality, in particular, for any bounded log-concave  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ . We require strict log-concavity for Theorem 3.2. For example,

the function  $a(y)$  in (1) is strictly log-concave in  $y > 0$  for any  $u < 1$ , being the composition of a strictly concave function and a strictly increasing and concave function, the log.

We show that  $\mathcal{F}$  and  $\mathcal{G}$  preserve  $TP_2$  and other properties, which are used to prove Theorem 1.2, that [A] holds for the Bomber problem. The Fighter Problems that we treat in subsequent sections have much in common with the present one. We begin by collecting some facts about  $TP_2$  and log concave functions in the following lemma; the claims which cannot be easily verified using definition (4) can be found in Karlin (1968). For part 1 of Lemma 3.1 see also Schoenberg (1951). Note also that part 2 is a special case of part 3.

**Lemma 3.1.** *1. A nonnegative function  $a(y)$  is (strictly) log-concave if and only if  $a(x - y)$  is (strictly)  $TP_2(x, y)$ .*

*2. The convolution of log-concave functions is log-concave.*

*3. (The composition formula) If  $J(x, y)$  is  $TP_2(x, y)$ ,  $L(y, z)$  is  $TP_2(y, z)$ , then  $M(x, z) = \int J(x, y)L(y, z)d\sigma(y)$  is  $TP_2(x, z)$ , where  $\sigma$  is a nonnegative sigma-finite measure on  $\mathbb{R}$ .*

*4. If  $J(x, y)$  and  $L(x, y)$  are  $TP_2(x, y)$ , then so is their product.*

*5. If  $J(x, y)$  is  $TP_2(x, y)$  then  $J(x, y)^p$  for any positive  $p$ , and  $J(x, y)f(x)g(y)$  for any nonnegative functions  $f$  and  $g$ , are  $TP_2$ .*

Given a  $TP_2$  function  $Q$  defined on  $\mathbb{R}^+ \times \mathbb{R}^+$  we extend it to  $\mathbb{R} \times \mathbb{R}$  by setting it equal to zero where it was not defined before; the resulting function is  $TP_2$  on  $\mathbb{R} \times \mathbb{R}$ .

The next lemma follows directly from (12) and (13).

**Lemma 3.2.** *If  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  and  $Q(x, t)$  is nonnegative and increasing in  $x$  then  $\mathcal{F}(Q)$  and  $\mathcal{G}(Q) = \mathcal{F}(Q) + 1$  are nonnegative and increasing in  $x$  and  $t$ .*

The following key lemma gives conditions under which the  $TP_2$  property is preserved by the sup-convolution. We recall that  $\otimes$  in this section is given by (Bo) of (11).

**Lemma 3.3.** *If  $Q$  is a continuous, nonnegative  $TP_2$  function on  $\mathbb{R}^+ \times \mathbb{R}^+$  and  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is log-concave, then  $a \otimes Q$  is  $TP_2$ .*

*Proof.* Let  $(x, t) \in \mathbb{R}^+ \times \mathbb{R}^+$  and  $p > 0$  be arbitrary. Making the change of variable  $v = x - y$  and writing the limits of integration using the indicator  $I(x, v) = \mathbf{1}(0 \leq v \leq x)$  we obtain

$$\int_0^x [a(y)Q(x - y, s)]^p dy = \int_{-\infty}^{\infty} a(x - v)^p I(x, v) Q(v, s)^p dv.$$

One can verify directly from (4) that  $I(x, v)$  is  $TP_2(x, v)$ . Since  $a(x)$  is log-concave on  $\mathbb{R}^+$ , defining  $a(x) = 0$  for  $x < 0$  we have  $a(x)$  log-concave on all of  $\mathbb{R}$ , and so the product  $a(x - v)I(x, v)$ , and therefore also  $a(x - v)^p I(x, v)$ , are  $TP_2(x, v)$  on  $\mathbb{R} \times \mathbb{R}$ . Hence the integrand is a product of a  $TP_2(x, v)$  function with a  $TP_2(v, s)$  function, and so the integral is  $TP_2(x, s)$  by part 3 (the composition formula) of Lemma 3.1. Raising to the power  $1/p$ , we therefore have

$$\left( \int_0^x [a(y)Q(x - y, s)]^p dy \right)^{1/p} \text{ is } TP_2(x, s) \text{ for all } p > 0. \quad (23)$$

Since  $a(y)$  is log-concave on  $\mathbb{R}^+$  it is continuous in  $(0, x]$  and bounded on  $[0, x]$ , with a possible discontinuity at 0, in which case there is a downward jump. Hence for all fixed  $x$  and  $s$  the product

$a(y)Q(x - y, s)$  has the same properties. Therefore the limit of the  $L^p$  norms (23) of the function  $a(y)Q(x - y, s)$  of  $y$  on  $[0, x]$  converges to its essential supremum which coincides with its maximum value over  $y$ . Using the nonnegativity of  $a(\cdot)$  and  $Q(\cdot, \cdot)$ , we therefore have, by the preservation of the  $TP_2$  property under limits, that

$$\lim_{p \rightarrow \infty} \left( \int_0^x [a(y)Q(x - y, s)]^p dy \right)^{1/p} = (a \otimes Q)(x, s) \quad \text{is } TP_2(x, s).$$

■

**Lemma 3.4.** *If  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is log-concave and  $Q$  is  $TP_2$  then  $\mathcal{F}(Q)$  is  $TP_2$ .*

*Proof.* By Lemma 3.3  $(a \otimes Q)(x, s)$  is  $TP_2(x, s)$ . As  $I(s, t) = \mathbf{1}(0 \leq s \leq t)$  is  $TP_2(s, t)$ , the integral of the product

$$\mathcal{F}(x, t, Q) = \int_0^t (a \otimes Q)(x, s) ds = \int_{-\infty}^{\infty} (a \otimes Q)(x, s) I(s, t) ds,$$

is  $TP_2(x, t)$ .

■

**Lemma 3.5.** *If  $Q(x, t)$  is nonnegative, increasing in  $x$  and  $t$ , and  $TP_2$ , then so is  $Q(x, t) + 1$ .*

*Proof.* It is clear that  $Q(x, t) + 1$  is nonnegative and increasing in  $x$  and  $t$  whenever  $Q(x, t)$  is. To demonstrate the  $TP_2$  property it is required to show that

$$(Q(x', t') + 1)(Q(x, t) + 1) \geq (Q(x', t) + 1)(Q(x, t') + 1),$$

where  $x' > x$  and  $t' > t$ . The monotonicity and nonnegativity of  $Q$  imply that the result is trivial if  $Q(x', t') = 0$ . Otherwise using  $Q(x', t')Q(x, t) \geq Q(x', t)Q(x, t')$  we obtain

$$\begin{aligned} & (Q(x', t') + 1)(Q(x, t) + 1) - (Q(x', t) + 1)(Q(x, t') + 1) \\ & \geq (Q(x', t') + 1) \left( \frac{Q(x', t)Q(x, t')}{Q(x', t')} + 1 \right) - (Q(x', t) + 1)(Q(x, t') + 1) \\ & = (Q(x', t') - Q(x', t))(Q(x', t') - Q(x, t')) / Q(x', t') \geq 0 \end{aligned}$$

where the last inequality follows since

$$Q(x', t') \geq \max\{Q(x', t), Q(x, t')\},$$

as  $Q$  is increasing in both  $x$  and  $t$ .

■

**Theorem 3.1.** *If  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a uniformly continuous log-concave function then the solution  $\bar{P}$  to  $\mathcal{G}(Q) = Q$  is  $TP_2$ .*

*Proof.* Let  $\bar{P}_0 = 0$  and  $\bar{P}_{m+1} = \mathcal{G}(\bar{P}_m)$  for  $m = 1, 2, \dots$ . We first claim that the functions  $\{\bar{P}_m\}_{m=0}^{\infty}$  are nonnegative, increasing in both  $x$  and  $t$ , and  $TP_2$ . Clearly the claim is true for  $\bar{P}_0$ . Assume the claim is true for  $\bar{P}_m$  for some  $m \geq 0$ . By Lemma 3.2  $\mathcal{F}(\bar{P}_m)$  and  $\bar{P}_{m+1}$  are nonnegative and increasing in  $x$  and  $t$ , and by Lemma 3.4  $\mathcal{F}(\bar{P}_m)$  is  $TP_2$ , and now Lemma 3.5 shows that  $\bar{P}_{m+1} = \mathcal{F}(\bar{P}_m) + 1$  is  $TP_2$ , thus completing the proof of the claim. That  $\bar{P}$  is  $TP_2$  follows, as  $\bar{P}_m$  converges uniformly to  $\bar{P}$  by Corollary 2.2, and the  $TP_2$  property is preserved under (even) pointwise limits.

■

We may now prove Theorem 1.1:

*Proof of Theorem 1.1.* By Lemmas 2.3 and 2.2, when  $a$  is a probability, the operator  $\mathcal{G}$  is of 1-contraction type and  $e^t$ -bounded. By Corollary 2.2 the unique solution  $\bar{P}$  to  $\mathcal{G}(Q) = Q$  exists, is continuous and satisfies  $0 \leq \bar{P}(x, t) \leq e^t$ . Now recalling (8), that is, that  $\bar{P}(x, t) = e^t P(x, t)$ , we have  $P(x, t) \in [0, 1]$ . Theorem 3.1 gives that  $\bar{P}$  is TP<sub>2</sub>, hence  $P$  is TP<sub>2</sub> by Property 5 of Lemma 3.1. ■

The next lemma modifies the result in Samuel (1970), page 158, to continuous ammunition and time, thus allowing us to show Theorem 1.2.

**Lemma 3.6.** *Let  $R(x, z, t)$  be a positive continuous function of  $x \geq z \geq 0$  and  $t \geq 0$  such that the ratio  $R(x, z, t)/R(x, z', t)$  is increasing in  $t$  whenever  $x \geq z' \geq z \geq 0$ , and let*

$$k(x, t) = \min\{y \in [0, x] : R(x, y, t) = \max_{0 \leq z \leq x} R(x, z, t)\}.$$

*Then  $k(x, t)$  is decreasing in  $t$ .*

*Proof.* For the sake of contradiction, assume that there exists  $t_1 < t_2$  and  $z_1 < z_2 \leq x$  such that  $k(x, t_1) = z_1 < z_2 = k(x, t_2)$ . Then the first equality gives  $R(x, z_1, t_1)/R(x, z_2, t_1) \geq 1$ , as the maximum of  $R(x, z, t_1)$  is attained at  $z = z_1$ , whereas the second equality gives  $R(x, z_1, t_2)/R(x, z_2, t_2) < 1$  as  $k(x, t_2)$  is the minimum over all  $y$  which attain the maximum of  $R(x, z, t_2)$ . This contradicts the assumption that  $R(x, z, t)/R(x, z', t)$  is increasing in  $t$ . ■

*Proof of Theorem 1.2.* By Theorem 1.1, the solution  $P(x, t)$  of (2) is TP<sub>2</sub>; also from (2),  $P(x, t) \geq e^{-t}$  for all  $(x, t) \in \mathbb{R}^+ \times \mathbb{R}^+$ . In view of definition (4), the ratio  $P(x', t)/P(x, t)$  is increasing in  $t$  whenever  $x < x'$ , or

$$\frac{P(x - z, t)}{P(x - z', t)}, \quad \text{and therefore} \quad \frac{a(z)P(x - z, t)}{a(z')P(x - z', t)},$$

is increasing in  $t$  when  $0 \leq z < z' \leq x$ . Applying Lemma 3.6 with the continuous function  $R(x, z, t) = a(z)P(x - z, t)$  now yields that

$$K(x, t) = \min\{y \in [0, x] : a(y)P(x - y, t) = \max_{0 \leq z \leq x} a(z)P(x - z, t)\} \quad (24)$$

is decreasing in  $t$ , and [A] holds. ■

We now consider [C] for the Bomber.

**Theorem 3.2.** *If  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a strictly log-concave function, then conjecture [C] holds for the doubly continuous Bomber Problem.*

*Proof.* For  $0 \leq y < x$  and  $\delta = x - y$  note that [C], that is,  $x - K(x, t) \geq y - K(y, t)$ , is equivalent to

$$K(y + \delta, t) \leq K(y, t) + \delta. \quad (25)$$

Inequality (25) holds trivially for all  $y$  such that  $K(y + \delta, t) \leq \delta$ . To handle the case  $K(y + \delta, t) > \delta$ , define

$$b(x, y) = \frac{a(x)}{a(y)} \quad \text{and} \quad G_x(y, t) = a(y)P(x - y, t),$$

the latter expression being the conditional probability of survival upon expending an amount  $y$  of ammunition at an encounter at time  $t$ , when the total amount  $x$  is available, and then proceeding optimally thereafter. With these definitions in place we obtain

$$G_{y+\delta}(v + \delta, t) = a(v + \delta)P(y - v, t) = \frac{a(v + \delta)}{a(v)} G_y(v, t) = b(v + \delta, v) G_y(v, t). \quad (26)$$

Rewriting (26) and using that  $K(x, t)$  is optimal, we have

$$\begin{aligned}
G_y(v, t) &= \frac{G_{y+\delta}(v + \delta, t)}{b(v + \delta, v)} \\
&\leq \frac{G_{y+\delta}(K(y + \delta, t), t)}{b(v + \delta, v)} \\
&= \frac{G_{y+\delta}([K(y + \delta, t) - \delta] + \delta, t)}{b(v + \delta, v)} \\
&= G_y(K(y + \delta, t) - \delta, t) \frac{b(K(y + \delta, t), K(y + \delta, t) - \delta)}{b(v + \delta, v)}, \tag{27}
\end{aligned}$$

where in the third expression we use  $K(y + \delta, t) > \delta$ , and obtain the last equality by applying (26) with  $v = K(x, t) - \delta$ .

By part 1 of Lemma 3.1  $a(y)$  is strictly  $TP_2$ , which is equivalent to  $b(x + \delta, x)$  being a strictly decreasing function of  $x$  for  $x \geq 0$ . Therefore, for  $v$  satisfying  $v < K(y + \delta, t) - \delta$ , we have

$$\frac{b(K(y + \delta, t), K(y + \delta, t) - \delta)}{b(v + \delta, v)} < 1.$$

Since  $K(x, t)$  is the minimal amount of ammunition that can be used to proceed optimally, if the smallest maximizer of the left hand side of (27) is attained at a value of  $v = K(y, t)$  satisfying  $v + \delta < K(y + \delta, t)$ , the ratio of the  $b$ -functions on the right hand side of (27) would be strictly less than 1. In this case, by (27),  $K(y + \delta, t) - \delta$  would be a better choice for  $v$  than the optimal amount  $K(y, t)$ . Note that by definition  $K(y + \delta, t) \leq y + \delta$ , so  $K(y + \delta, t) - \delta \leq y$ , and hence is a feasible quantity to expend, given ammunition  $y$ . Having reached a contradiction, we conclude

$$K(y, t) + \delta \geq K(y + \delta, t),$$

which is (25). ■

## 4 The Frail Fighter

In this section we consider the Frail Fighter, whose goal is to shoot down as many of the enemy planes as possible. Recall that if the Frail Fighter does not shoot down an encountered enemy, then he gets shot down with probability 1. In this section letting  $\otimes$  be given by case  $(F_0)$  of (11), the dynamic programming equation (5) for the Frail Fighter can be written  $\bar{N}(x, t) = \mathcal{G}(x, t, \bar{N})$  for  $\mathcal{G}(x, t, Q)$  given by (12) and  $\bar{N}(x, t) = e^t N(x, t)$ . Analogous to the previous section, in this and the next section  $a(y)$  can be interpreted as the probability that the Fighter destroys an enemy using an amount  $y$  of ammunition, although our results hold for more general  $a$ .

The following result is parallel to Lemma 3.4.

**Lemma 4.1.** *If  $e^{-t}Q(x, t)$  is nonnegative and increasing in  $x$  and  $t$  then so is  $e^{-t}\mathcal{G}(Q)$ . If  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is log-concave and  $Q$  is  $TP_2(x, t)$  then so is  $\mathcal{G}(Q)$ .*

*Proof.* It is clear that  $e^{-t}\mathcal{G}(Q)$  is nonnegative and increasing in  $x$  whenever  $e^{-t}Q$  is. Next,

$$\begin{aligned}
\frac{\partial e^{-t}\mathcal{G}(Q)}{\partial t} &= \sup_{0 \leq y \leq x} a(y)[1 + e^{-t}Q(x - y, t)] - e^{-t}\mathcal{G}(Q) \\
&= f(t) - \int_0^t f(s)e^{-(t-s)} ds \quad \text{where} \quad f(t) = \sup_{0 \leq y \leq x} a(y)[1 + e^{-t}Q(x - y, t)].
\end{aligned} \tag{28}$$



Since  $e^{-t}Q(x, t)$  is increasing in  $t$ ,  $f(t) \geq f(s)$  for all  $s \in [0, t]$ , and

$$f(t) \geq (1 - e^{-t})f(t) = \int_0^t f(t)e^{-(t-s)}ds \geq \int_0^t f(s)e^{-(t-s)}ds.$$

Hence the partial derivative (28) is nonnegative, and  $e^{-t}\mathcal{G}(Q)$  is increasing in  $t$ .

As  $Q(x, t)$  is  $\text{TP}_2$ , so is  $e^{-t}Q(x, t)$  by Lemma 3.1. Now, by Lemma 3.5,  $1 + e^{-t}Q(x, t)$  is  $\text{TP}_2(x, t)$ . As in the proof of Lemma 3.3, letting  $p > 0$  and  $I(x, v) = \mathbf{1}(0 \leq v \leq x)$ , write

$$\int_0^x a(y)^p [1 + e^{-s}Q(x - y, s)]^p dy = \int_{-\infty}^{\infty} a(x - v)^p I(x, v) [1 + e^{-s}Q(v, s)]^p dv,$$

which is  $\text{TP}_2(x, s)$ , as in the proof of Lemma 3.3. Taking  $p^{\text{th}}$  roots and letting  $p \rightarrow \infty$  we conclude that  $\sup_{0 \leq y \leq x} a(y)[1 + e^{-s}Q(x - y, s)]$ , and therefore also  $\sup_{0 \leq y \leq x} a(y)[1 + e^{-s}Q(x - y, s)]e^s$ , are  $\text{TP}_2(x, s)$ . Now

$$\int_0^t \sup_{0 \leq y \leq x} a(y)[1 + e^{-s}Q(x - y, s)]e^s ds = \int_{-\infty}^{\infty} \sup_{0 \leq y \leq x} a(y)[1 + e^{-s}Q(x - y, s)]I(s, t)e^s ds,$$

and the final claim follows. ■

The proof of Theorem 4.1, which proves the first claim of Theorem 1.4, is similar to the proof of Theorem 3.1 and is omitted.

**Theorem 4.1.** *If  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a uniformly continuous log-concave function then the solution  $\bar{N}(x, t)$  to  $\mathcal{G}(Q) = Q$  is  $\text{TP}_2$ , as is  $N(x, t) = e^{-t}\bar{N}(x, t)$ .*

The Frail Fighter's optimal policy  $K(x, t)$  satisfies

$$K(x, t) = \min\{y \in [0, x] : a(y)[1 + N(x - y, t)] = \sup_{0 \leq z \leq x} a(z)[1 + N(x - z, t)]\}.$$

Applying Lemma 3.6 with  $R(x, z, t) = a(z)[1 + N(x - z, t)]$  we obtain [A] for the Frail Fighter, thus proving the last claim of Theorem 1.4.

We now prove that [C] holds for the Frail Fighter.

**Theorem 4.2.** *If  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a strictly log-concave function, then conjecture [C] holds for the Frail Fighter.*

*Proof.* The argument is nearly identical to the proof of Theorem 3.2, and again it suffices to only consider  $y$  for which  $K(y + \delta, t) > \delta$ . Let

$$b(x, y) = \frac{a(x)}{a(y)} \quad \text{and} \quad G_x(y, t) = a(y)[1 + N(x - y, t)],$$

the latter expression being the expected 'score' for the Fighter expending  $y$ , conditional upon an encounter at time  $t$  when a total amount  $x$  is available, and then proceeding optimally thereafter. With these definitions in place we obtain

$$G_{y+\delta}(v + \delta, t) = a(v + \delta)[1 + N(y - v, t)] = \frac{a(v + \delta)}{a(v)}G_y(v, t) = b(v + \delta, v)G_y(v, t).$$

The same lines of argument of Theorem 3.2 from (26) on may now be applied to yield the conclusion. ■

## 5 The Invincible Fighter

In this section we consider an Invincible Fighter who cannot be shot down, whose goal is to shoot down as many of the enemy planes as possible. Hence, in this section we take  $\mathcal{G}$  given by (12) and  $\otimes$  by case  $(F_1)$  of (11), and the relevant equation may be written  $\bar{N}(x, t) = \mathcal{G}(x, t, \bar{N})$ . The role of the next lemma, and its proof, are parallel to that of Lemma 3.3 in Section 3.

**Lemma 5.1.** *If  $f(x)$  and  $g(x)$  are log-concave functions, then so is*

$$h(x) = \sup_{0 \leq y \leq x} [f(y)g(x-y)].$$

*If  $f(x)$  and  $g(x)$  are concave functions, then so is*

$$h(x) = \sup_{0 \leq y \leq x} [f(y) + g(x-y)].$$

*Proof.* By Lemma 3.1 part 2, the convolution of log-concave functions is log-concave. Writing

$$h(x) = \lim_{p \rightarrow \infty} \left( \int_0^x [f(y)g(x-y)]^p dy \right)^{1/p},$$

and noting that log-concavity is preserved when taking powers and under pointwise limits, we conclude that  $h(x)$  is log-concave, thus proving the first claim. To prove the second claim, write  $h(x) = \log \sup_{0 \leq y \leq x} [e^{f(y)} e^{g(x-y)}]$  and apply the first result to the log-concave functions  $e^{f(y)}$  and  $e^{g(x)}$ . ■

The proof of the following theorem uses the second part of Lemma 5.1 to show that concavity is preserved under the operation of  $(F_1)$  of (11), and then an iteration and limit argument as in the proof of Theorem 3.1. The details are omitted.

**Theorem 5.1.** *If  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a uniformly continuous concave function then the solution  $\bar{N}(x, t)$  to the equation  $\mathcal{G}(Q) = Q$ , for  $\mathcal{G}$  given in (12) for case  $(F_1)$  of (11), is concave in  $x$  for every  $t$ , as is  $N(x, t) = e^{-t} \bar{N}(x, t)$ .*

The following lemma is similar to Lemma 3.6.

**Lemma 5.2.** *If  $R(x, y)$  is a continuous function and  $R(x, y') - R(x, y)$  is increasing in  $x$  for all  $y' > y$ , then*

$$k(x) = \inf\{y : R(x, y) = \sup_{0 \leq z \leq x} R(x, z)\}$$

*is increasing in  $x$ .*

*Proof.* To reach a contradiction, assume there exist  $x < x'$  such that  $y' = k(x) > k(x') = y$ . Note then that by the definition of  $k$  we have

$$R(x, y') = R(x, k(x)) > R(x, k(x')) = R(x, y) \quad \text{and} \quad R(x', y') = R(x', k(x)) \leq R(x', k(x')) = R(x', y),$$

contradicting the assumption on  $R$ . ■

We may now provide the proof of Theorem 1.5.

*Proof of Theorem 1.5.* For fixed  $t$ , let

$$R(x, y) = a(y) + N(x - y, t).$$

Note that  $R(x, y') - R(x, y)$  being increasing in  $x$  for all  $y' > y$  is equivalent to  $N(x - y') - N(x - y)$  being increasing in  $x$  for all  $y' > y$ , which is equivalent to  $N$  being concave in  $x$ . Applying Theorem 5.1 and Lemma 5.2 completes the proof. ■

The optimal policy for the Invincible Fighter is given by

$$K(x, t) = \arg \max_{0 \leq y \leq x} [a(y) + N(x - y, t)]. \quad (29)$$

**Proposition 5.1.** *If the uniformly continuous function  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is strictly concave then the optimal policy  $K(x, t)$  for the Invincible Fighter is unique, that is, the maximum in (29) is attained uniquely.*

*Proof.* Theorem 5.1 gives that  $N(x, t)$  is concave in  $x$  for every  $t$ , and hence  $N(x - y, t)$  is concave in  $y$  for all fixed  $x$  and  $t$ . Since  $a(y)$  is strictly concave in  $y$ , so is the function  $a(y) + N(x - y, t)$ , and hence its maximum is attained uniquely. ■

The argument we have used here to show uniqueness no longer works in general in the model ( $F_u$ ) where the Fighter is vulnerable to attack, where, following (3), the function  $R(x, y)$  in the proof of Theorem 1.5 is replaced by

$$R(x, y) = a(y) + [a(y) + u(1 - a(y))]N(x - y, t).$$

In this case,  $R(x, y') - R(x, y)$  being increasing in  $x$  for  $y < y'$  is no longer equivalent to the concavity of  $N(x, t)$  in  $x$  for fixed  $t$  when  $u \neq 1$ .

We now prove that [C] holds for the Invincible Fighter.

**Theorem 5.2.** *If the uniformly continuous function  $a : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is strictly concave, then conjecture [C] holds for the Invincible Fighter.*

*Proof.* The argument is similar to the proof of Theorem 3.2. As there, it suffices to only consider  $y$  for which  $K(y + \delta, t) > \delta$ . Define

$$b(x, y) = \exp(a(x) - a(y)) \quad \text{and} \quad G_x(y, t) = \exp(a(y) + N(x - y, t)),$$

the logarithm of the latter expression being the expected ‘score’ for the Fighter expending  $y$ , conditional upon an encounter, at time  $t$ , when a total amount  $x$  is available, and then proceeding optimally thereafter. With these definitions in place we obtain

$$G_{y+\delta}(v + \delta, t) = b(v + \delta, v)G_y(v, t),$$

and the lines of argument of the proof of Theorem 3.2 may now be followed to show the desired conclusion under the condition that  $\exp(a(x))$  is strictly log-concave, that is, that  $a(x)$  is strictly concave. ■

## 6 Discussion

We have demonstrated that [A] holds for the Bomber and Frail Fighter, [B] for the Invincible Fighter, and [C] for all three, so in no model have we shown that [A] and [B] hold together. In addition, we have shown uniqueness of the optimal policy only for the Invincible Fighter. But of all the possible conjectures which can be raised in these models, the one most outstanding is that




Figure 1. The functions  $\log \bar{P}_m(x, t)$ ,  $m = 1, \dots, 10$ , for  $t = 10$ .

[B] holds for the Bomber, and naturally one may hope to use the ideas developed in Sections 2 and 3 to settle this.

The equivalence of [B] to the log-concavity of  $P(x, t)$  in  $x$  was noted in Simons and Yao (1990). Hence, parallel to the proof of [A], for a given  $Q(x, s)$  consider the function

$$R(x, t) = \int_0^t (a \otimes Q)(x, s) ds$$

where  $\otimes$  is the max-convolution operator given in case (Bo) of (11). By using results in Prékopa (1973), one can show that  $R(x, t)$  is log-concave when  $a(y)$  and  $Q(x, t)$  are log-concave, similarly as to how one shows that the  $TP_2$  is preserved. When  $R(x, t)$  is log-concave, for any nonnegative  $x_1, x_2, t_1, t_2$ , we have

$$R_{12}^2 \geq R_1 R_2 \quad \text{or} \quad R_{12} \geq (R_1 R_2)^{1/2}, \quad (30)$$

where

$$R_{12} = R\left(\frac{x_1 + x_2}{2}, \frac{t_1 + t_2}{2}\right), \quad R_1 = R(x_1, t_1) \quad \text{and} \quad R_2 = R(x_2, t_2).$$

Now to proceed as in Lemma 3.5, for  $R(x, t) + 1$  to be log-concave, we require that

$$(R_{12} + 1)^2 \geq (R_1 + 1)(R_2 + 1),$$

and by (30) it suffices that

$$2R_{12} \geq R_1 + R_2 \quad \text{or} \quad R_{12} \geq \frac{1}{2}(R_1 + R_2), \quad (31)$$




Figure 2. The functions  $\log \bar{P}_m(x, t)$ ,  $m = 1, \dots, 7$ .

that is, that  $R(x, t)$  is concave. However, unlike the argument in the proof of Lemma 3.5, we cannot conclude (31) from (30): the geometric mean of  $R_1$  and  $R_2$  in (30) is in general smaller than the arithmetic mean in (31). Moreover, numerical evidence indicates that  $R(x, t)$  may fail to be concave.

Nevertheless, as the concavity of  $R(x, t)$  is sufficient but not necessary for the log-concavity of  $R(x, t) + 1$ , log-concavity may yet be conserved in each step of the iteration, and the limiting argument pushed through as before. Taking  $a(y)$  to be given in (1) and considering the first few iterates of (10), one has  $\bar{P}_0(x, t) = 0$  and  $\bar{P}_1(x, t) = 1$ , which are indeed log-concave. Continuing, as  $a(y)$  is increasing we have  $(a \otimes 1)(x, s) = \sup_{0 \leq y \leq x} a(y) = a(x)$ , and therefore

$$\bar{P}_2(x, t) = \int_0^t (a \otimes \bar{P}_1)(x, s) ds + 1 = \int_0^t a(x) ds + 1 = ta(x) + 1.$$

Since

$$\frac{\partial^2}{\partial x^2} \log \bar{P}_2(x, t) = \frac{-t(1+t)(1-u)e^{-x}}{(ta(x) + 1)^2} \leq 0 \quad \text{for all } (x, t) \in \mathbb{R}^+ \times \mathbb{R}^+,$$

the function  $\bar{P}_2(x, t)$  is log-concave as well. Indeed, Figure 1, which gives plots of  $\log \bar{P}_m$  for  $m = 1, \dots, 10$ ,  $(x, t) \in [0, 10] \times \{10\}$ , and  $u = 0$ , suggests that the log-concave property is preserved in these further functions as well. The figure also indicates that convergence of  $\bar{P}_m(x, t)$  for the values considered is quite rapid, consistent with the exponential bound in Theorem 2.1. Expanding the range of  $t$  back toward the origin, Figure 2 shows  $\log \bar{P}_m(x, t)$  for  $m = 1, \dots, 7$  over the range  $(x, t) \in [0, 10]^2$ , and indicates that these functions are log-concave in both variables  $x$  and  $t$ .

## References

- Bartroff, J., Goldstein, L., and Samuel-Cahn, E. (2010). The spend-it-all region and small time results for the continuous Bomber Problem. *Sequential Analysis*, in press.
- Karlin, S. (1968) *Total Positivity*, Stanford University Press.
- Klinger, A. and Brown, T. A. (1968). Allocating unreliable units to random demands. In Karreman, H. F., editor, *Stochastic Optimization and Control*, number 20 in Mathematics Research Center, U.S. Army, at the University of Wisconsin, Madison, pages 173–209. Wiley, New York.
- Luenberger, D. (1969) *Optimization by vector space methods*, John Wiley and Sons, Inc.
- Prékopa, A. (1973) On logarithmic concave measures and functions. *Acta Scientiarum Mathematicarum*, 34:335–343.
- Samuel, E. (1970). On some problems in operations research. *Journal of Applied Probability*, 7:157–164.
- Schoenberg, I. J. (1951). On Pólya frequency functions, I. The totally positive functions and their Laplace transforms. *J. Analyse Math.*, 1:331-374.
- Shepp, L. A., Simons, G., and Yao, Y.-C. (1991). On a problem of ammunition rationing. *Advances in Applied Probability*, 23:624–641.
- Simons, G. and Yao, Y.-C. (1990). Some results on the bomber problem. *Advances in Applied Probability*, 22:412–432.
- Weber, R. (1985). On a problem of ammunition rationing. In *Conference Report: Stochastic Dynamic Optimization and Applications in Scheduling and Related Areas*, page 148. Universität Passau, Fakultät für Mathematik und Informatik. Abstract.