

Acarologia

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:



<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia-contact@supagro.fr



**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2020 (Volume 60): 450 €
<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2018): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

ISSN 0044-586X (print), ISSN 2107-7207 (electronic)

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme
(Labex Agro: ANR-10-LABX-0001-01)



Acarologia is under free license and distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

A new species and new record of *Agistemus Summers* (Acari: Stigmeidae) associated with *Capsicum annuum L.* var. *glabriuscum* (Solanaceae) from Northeastern of Mexico

José Irving Monjarás-Barrera^a, Guilherme Liberato Da Silva^b, Julio Cesar Chacón-Hernández^a, Onilda Santos Da Silva^c, Noeli Juarez Ferla^{b,d}, Liana Johann^b

^aUniversidad Autónoma de Tamaulipas, Instituto de Ecología Aplicada, División del Golfo 356, Colonia Libertad, C.P. 87019. Ciudad Victoria, Tamaulipas, México.

^bLaboratório de Acarologia, Tecnovates, Universidade do Vale do Taquari - Univates, Lajeado, 95914-014, Rio Grande do Sul, Brasil.

^cDepartamento de Microbiología, Imunología e Parasitología, Universidade Federal do Rio Grande do Sul. 90050-170 Porto Alegre, Rio Grande do Sul, Brasil.

^dCNPq Researcher.

Original research

ABSTRACT

Two mite species belonging to the genus *Agistemus* Summers (Acari: Stigmeidae) were collected from wild plants known as “chile piquín” (*Capsicum annuum* L. var. *glabriuscum*) in two protected natural areas, state of Tamaulipas, Mexico. *Agistemus piquinus* Monjarás-Barrera & Johann n. sp. is described and illustrated. *Agistemus brasiliensis* Matioli, Ueckermann & Oliveira, 2002 is a new record for Mexico. A key to the *Agistemus* species of Mexico is provided.

Keywords Raphignathoidea; Protected Natural Areas; Trombidiformes; chile piquín; Tamaulipas

Zoobank <http://zoobank.org/922C0CDD-E7B9-4670-8AE3-863621A79BAB>

Received 11 February 2020

Accepted 12 May 2020

Published 22 May 2020

Corresponding author

José Irving Monjarás-Barrera:
monjarasb.ji@gmail.com

Academic editor

Faraji, Farid

DOI

10.24349/acarologia/20204378

ISSN 0044-586X (print)

ISSN 2107-7207 (electronic)

© Copyright

Monjarás-Barrera J. I. et al.

Distributed under

Creative Commons CC-BY 4.0



Introduction

In several ecosystems and crops, mites of the family Stigmeidae are one of the most important predatory mites after Phytoseiidae (Santos & Laing, 1985) and are considered economically important as predators of other mites and small arthropods (Fan & Zhang, 2005; Fan & Flechtmann, 2015). Stigmeid mites have been studied as potential predators against pestiferous mites of the families Eriophyidae, Tetranychidae, Tenuipalpidae and Tarsonemidae (Gerson et al., 2003; Johann et al., 2013). Therefore, its correct identification is fundamental for management programs of these mites (Zhang, 2003).

The genus *Agistemus* Summers, 1960 comprise more than 87 described species (Fan et al., 2016; Paktinat-Saeij et al., 2017; Rehman et al., 2018); of which, seven were recorded from Mexico: *Agistemus floridanus* González-Rodríguez (González-Rodríguez, 1965), *A. longisetus* González-Rodríguez (González-Rodríguez, 1963), *A. fleschneri* Summers, *A. simplex* González-Rodríguez, *A. striolatus* González-Rodríguez (González-Rodríguez, 1965), *A. arcypaurus* González-Rodríguez and *A. terminalis* Quayle (Estrada-Venegas et al., 2002).

How to cite this article Monjarás-Barrera J. I. et al. (2020), A new species and new record of *Agistemus Summers* (Acari: Stigmeidae) associated with *Capsicum annuum* L. var. *glabriuscum* (Solanaceae) from Northeastern of Mexico. *Acarologia* 60(2): 470-480; DOI 10.24349/acarologia/20204378

The present study aims to (i) described a new species of stigmacid mites from Mexico, (ii) record of *Agistemus brasiliensis* for the first time from this country, and (iii) provide an identification key to species of *Agistemus* so far reported from Mexico.

Materials and methods

The specimens were collected on leaves of the wild plant “chile piquín” (*Capsicum annuum* L. var. *glabriuscum*) in two Protected Naturals Areas (PNA): “El Cielo” Biosphere Reserve (23°01'07" N, 99°08'54" W) and PNA “Altas Cumbres” (23°41'52" N, 99°11'04" W and 23°46'41" N, 99°12'12" W), located in the municipalities of Gómez Farias and Victoria in Tamaulipas, Mexico, respectively. The collected leaves were transported to the laboratory of Ecología de Poblaciones in the Instituto de Ecología Aplicada of Universidad Autónoma de Tamaulipas. Mite extraction followed Dhooria (2016) and slide preparations were carried out according to Jeppson *et al.*, (1975). The mites were observed under Phase Contrast (PH) and Differential Interference Contrast (DIC) microscope Zeiss AXIO Imager.Z2 in the Laboratory of Acarology of Universidade do Vale do Taquari - Univates, Lajeado, Rio Grande do Sul, Brazil.

The terminology and abbreviations are based on Grandjean (1944) and Kethley (1990). All measurements are given in micrometers (μm) and the measurements of the holotype are followed by ranges of the paratypes in parentheses.

Holotype female and three paratype males were deposited at Departamento de Entomología e Acarología, Escola Superior de Agricultura “Luiz de Queiroz”, Universidade de São Paulo (ESALQ/USP), Piracicaba, São Paulo State, Brazil. Two paratypes female were deposited at “Colección Nacional de Ácaros” (CNAC) of Instituto de Biología, Universidad Nacional Autónoma de México (UNAM) in Mexico city, Mexico.

Taxonomy

Family: Stigmeidae Oudemans, 1931

Genus: *Agistemus* Summers, 1960

Type species: *Caligonus terminalis* Quayle, 1912, by original designation. *Agistemus terminalis* (Quayle), Summers, 1960: 234.

Agistemus brasiliensis Matioli, Ueckermann & Oliveira, 2002

(Table 1)

Distribution — *Agistemus brasiliensis* was reported on *Citrus sinensis* (L.) Osbeck in São Paulo, Brazil, feeding on *Panonychus citri* (McGregor) (Tetranychidae) and *Brevipalpus phoenicis* (Geijskes) (Tenuipalpidae) (Matioli *et al.*, 2002). It is recorded from *Coffea arabica* L. and *C. canephora* P. (Mineiro *et al.*, 2010). It is the only species collected on *Vitis vinifera* L. Chardonnay and Merlot cultivar, *Vitis labrusca* L. Bordeaux cultivar and *Solanum americanum* Mill. (Johann *et al.*, 2013). *Agistemus brasiliensis* has only been recorded from two countries worldwide, Brazil (Fan *et al.*, 2019) and Mexico (present study).

Remarks — No morphological differences were observed between specimens from Mexico and those described by Matioli *et al.*, (2002). Measurements of the Mexican specimens are provided in Table 1.

Material examined — All specimens were found on leaves of *Capsicum annuum* L. var. *glabriuscum* at, Ojo de Agua, Gómez Farias Tamaulipas, Mexico; J. I. Monjarás B. leg.; two females; February 17, 2017; one female, March 3, 2017; one male, February 17, 2017; one male, October 13, 2017; one male, November 10, 2017.

***Agistemus piquinnus* Monjarás-Barrera & Johann n. sp.**

Zoobank: AC9EF2C2-8565-4982-9AEB-F1DBE9BA2D88

(Figures 1-5)

Diagnosis (female) — Prodorsal and hysterosomal shields smooth; length of seta c_1 longer than the distance between the bases of c_1 and d_1 ; the ratio $vi/vi-vi$ 2.8 (2.8-3.0); ratio c_1/c_1-c_1 2.1 (2.1-2.3); ratio: pob/eye : 3.2 (3.0-3.2). Aggenital shield not defined, bearing two pair of setae (ag_1-ag_2); seta g_1 extending beyond the base of ps_2 ; setae ps_1 and ps_2 slightly barbed and more robust than ps_3 ; femora I-IV 5-4-2-2.

Table 1 Measurements of *Agistemus brasiliensis* Matioli, Ueckermann & Oliveira, 2002 collected in Mexico.

Character measured	Female (n=3)				Male (n=3)			
	Mean	SD	Min	Max	Mean	SD	Min	Max
Body length	325	15.62	315	343	291	27	260	310
Idiosoma	259	21.2	244	283	239	20.5	215	251
Body width	206	22.9	190	232	173	16.8	154	184
vi	50	0.6	49	50	40	2.1	38	42
ve	71	3.8	68	75	54	2.1	52	56
sce	67	1.5	66	69	50	2.1	48	52
c_1	57	3.1	54	60	41	2.3	40	44
c_2	50	1.2	49	51	40	2.3	39	43
d_1	60	1.5	59	62	35	0.6	35	36
d_2	62	0	62	62	43	3	40	46
e_1	66	0.6	65	66	28	2.5	26	31
e_2	68	2.1	66	70	43	2.1	41	45
f	55	2.9	53	58	49	1	48	50
h_1	37	1.2	36	38	14	0.6	14	15
h_2	24	0.6	23	24	18	0.6	17	18
$vi-vi$	22	2.9	20	25	20	1.7	18	21
c_1-c_1	28	1.5	26	29	21	4.6	17	26
d_1-d_1	86	2.1	84	88	68	10	60	79
e_1-e_1	26	1.7	24	27	15	3.5	12	19
$f-f$	60	4.9	54	63	30	2.3	27	31
h_1-h_1	13	1	12	14	5	1	4	6
$vi/vi-vi$	2.3	0.3	2	2.5	2	0.3	1.8	2.3
c_1/c_1-c_1	2.1	0.2	1.9	2.3	2	0.4	1.5	2.4
d_1/d_1-d_1	0.7	0	0.7	0.7	1	0.1	0.5	0.6
e_1/e_1-e_1	2.5	0.2	2.4	2.7	2	0.5	1.4	2.3
$f/f-f$	0.9	0.1	0.8	1	2	0.1	1.5	1.8
h_1/h_1-h_1	2.8	0.2	2.7	3	3	0.5	2.5	3.5

SD= Standard Deviation

Description

Female

(n=3)

Dorsum (Figures 1A-B) — Length of idiosoma 323 (323-365); width of idiosoma 264 (264-295). Idiosoma broadly oval in dorso-ventral view. Prodorsal shield triangular, indented posteriorly and smooth. Prodorsal shield with 3 pairs of setae (*vi*, *ve* and *sci*), one pair of eyes 9 (9-10) in diameter and one pair of post-ocular bodies (*pob*) 29 (29-30) in diameter; humeral shields smooth, bearing *c*₂. Hysterosomal shield smooth, with five pairs of setae (*c*₁, *d*₁, *d*₂, *e*₁ and *e*₂); intercalary shields smooth, with setae *f*₁; suranal shield entire and smooth, with two pairs of setae, *h*₁ and *h*₂. Dorsal setae set on small tubercles, barbed, with blunt tips, *h*₁ and *h*₂ are distinctly shorter. Length of dorsal setae: *vi* 58 (58-60); *ve* 82 (82-85); *sci* 79 (73-79); *c*₁ 73 (73-75); *c*₂ 46 (45-50); *d*₁ 72 (70-75); *d*₂ 67 (67-73); *e*₁ 73 (73-80); *e*₂ 76 (76-80); *f*₁ 60 (60-63); *h*₁ 38 (38-40) and *h*₂ 25 (25-28). Distances between dorsal setae: *vi*-*vi* 21 (20-21); *ve*-*ve* 56 (50-56); *vi*-*ve* 19 (18-23); *ve*-*sci* 47 (43-47); *sci*-*sci* 114 (114-118); *c*₁-*c*₁ 35 (33-35); *c*₁-*d*₁ 51 (51-55); *c*₁-*c*₂ 83 (83-98); *c*₁-*d*₂ 59 (59-63); *d*₁-*d*₁ 74 (68-75); *d*₁-*d*₂ 29 (25-29); *d*₁-*e*₁ 59 (50-59); *d*₁-*e*₂ 37 (35-40); *d*₂-*d*₂ 133 (130-135); *d*₂-*e*₂ 42 (42-48); *e*₁-*e*₁ 30 (25-30); *e*₂-*e*₂ 108 (108-115); *e*₁-*f*₁ 30 (30-58); *f*₁-*f*₁ 67 (63-75); *f*₁-*h*₁ 38 (38-48); *h*₁-*h*₁ 13 (10-13); *h*₁-*h*₂ 21 (15-21); *h*₂-*h*₂ 49 (25-49). Ratios: *pob*/*eye*: 3.2 (3.0-3.2); *vi*/*vi*-*vi* 2.8 (2.8-3.0); *c*₁/*c*₁-*c*₁ 2.1 (2.1-2.3); *d*₁/*d*₁-*d*₁ 1.0 (1.0); *e*₁/*e*₁-*e*₁ 2.4 (2.4-3.2); *f*₁/*f*₁-*f*₁ 0.9 (0.8-1.0); *h*₁/*h*₁-*h*₁ 2.9 (2.9-4.0); *c*₁-*c*₁/*d*₁-*d*₁ 0.5 (0.4-0.5); *c*₁-*c*₁/*e*₁-*e*₁ 1.2 (1.2); *d*₁-*d*₁/*e*₁-*e*₁ 2.5 (2.5-2.7); *h*₁/*h*₂ 1.5 (1.4-1.5); *c*₁-*c*₁: *d*₁-*d*₁: *e*₁-*e*₁: *f*₁-*f*₁ = 1.2 (1.2): 2.5 (2.5-2.7): 1.0: 2.2 (2.2-2.7).

Venter (Figures 1C-D) — Ventral surface ornamented with striations; bearing three pairs of setae (*1a*, *3a* and *4a*). Aggenital shield not defined, bearing two pairs of setae (*ag*₁ and *ag*₂). Anogenital valves with one pair of genital (*g*₁) setae and three pairs of pseudanal setae (*ps*₁-*ps*₃). Seta *g*₁ reaching to base of seta *ps*₂. Setae *ps*₁ and *ps*₂ slightly barbed and more robust than *ps*₃. Measurements of setae: *1a* 31 (25-31); *3a* 28 (23-28); *4a* 23 (20-23); *ag*₁ 16 (13-16); *ag*₂ 15 (13-15); *g*₁ 26 (20-26); *ps*₁ 14 (13-14); *ps*₂ 16 (15-16); *ps*₃ 13 (13-15); *g*₁-*ps*₃ 29 (28-29).

Gnathosoma (Figure 2A) — Gnathosoma (including palp) 143 (143-150); subcapitulum 84 (80-84); palp 101 (100-103); chelicera 99 (93-105). Subcapitulum bearing subcapitular setae *m* 32 (30-33) and *n* 48 (48) and adoral setae *or*₁ 18 (18-19) and *or*₂ 19 (18-23). Distance *m*-*m* 44 (43-45), *n*-*n* 36 (30-36) and *m*-*n* 7 (7-8). Palp five segmented; palptrochanter without setae; palpfemur with two smooth and one barbed setae; palpgenu with one seta; palptibia with two tactile setae + one well-developed claw + one spine-like accessory claw; palptarsus with four tactile setae + one solenidion (ω), one subapical eupathidium and one distal trifurcate eupathidium.

Legs (Figures 3A-D) — Length of legs I-IV: 297 (297-325); 264 (264-293); 253 (250-275); 277 (275-290). Femur I 77 (77-85), genu I 27 (27-28), tibia I 55 (55-57), tarsus I 75 (70-83). Dorsal most seta on femur I (*dFI*) barbed 41 (41-45), longer than *h*₁, 1.1 (1.1-1.2) times length of *h*₁; dorsal most seta on genu I (*dGI*) barbed 36 (35-38). Chaetotaxy: coxae (excluding *1a*, *3a* and *4a*) 2(+1elcl)-1-2-2, trochanters 1-1-1-1, femora 5-4-2-2, genua 3(+1 κ)-1-0-0, tibiae 5(+1 φ p)-5(+1 φ p)-5(+1 φ p)-5(+1 φ p), tarsi 12(+1 ω)-9(+1 ω)-7(+1 ω)-7. Length of solenidia: ω I 27 (25-27); ω II 26 (23-26); ω III 16 (15-18).

Male

(n=3)

Dorsum (Figures 4A-B) — Length of idiosoma 266 (220-266); width of idiosoma 200 (177-200); idiosoma broadly oval in dorso-ventral view. Prodorsal shield triangular, indented posteriorly and smooth. Prodorsal shield with 3 pairs of setae (*vi*, *ve* and *sci*), one pair of eyes 8 (8-10) in diameter and one pair of post-ocular bodies (*pob*) 23 (22-23) in diameter; humeral shields smooth, bearing *c*₂. Hysterosomal shield smooth, with six pairs of setae (*c*₁, *d*₁, *d*₂, *e*₁, *e*₂, and *f*₁); suranal shield entire and smooth, with two pairs of setae, *h*₁ and *h*₂. Dorsal setae set on small tubercles, short and barbed, with blunt tips, *e*₁, *h*₁ and *h*₂ are distinctly shorter. Length




Figure 1 *Agistemus piquinnus* n. sp. (female); A – Dorsum, B – Dorsum (PH), C – Anogenital area, D – Anogenital area (DIC).

of dorsal setae: vi 45 (45-54); ve 67 (67-68); sci 63 (60-65); c_1 61 (57-61); c_2 37 (37-40); d_1 50 (47-52); d_2 57 (48-59); e_1 21 (21-28); e_2 60 (56-60); f_1 56 (54-61); h_1 13 (13-15) and h_2 19 (18-20). Distances between dorsal setae: $vi-vi$ 20 (17-20); $ve-ve$ 51 (46-51); $vi-ve$ 18 (18-20); $ve-sci$ 41 (36-41); $sci-sci$ 102 (92-102); c_1-c_1 28 (28-35); c_1-d_1 44 (41-44); c_1-c_2 78 (63-78); c_1-d_2 51 (46-51); d_1-d_1 63 (60-63); d_1-d_2 21 (19-21); d_1-e_1 41 (39-41); d_1-e_2 26 (24-28); d_2-d_2 107 (91-107); d_2-e_2 31 (29-31); e_1-e_1 23 (23); e_2-e_2 87 (81-90); e_1-f_1 15 (14-16); f_1-f_1 30 (30-33); f_1-h_1 42 (24-42); h_1-h_1 7 (5-7); h_1-h_2 11 (10-11); h_2-h_2 34 (28-34). Ratios: pob/eye : 2.9 (2.2-2.9); $vi/vi-vi$ 2.3 (2.3-3.2); c_1/c_1-c_1 1.4 (1.4); d_1/d_1-d_1 0.8 (0.8-0.9);




Figure 2 *Agistemus piquinnus* n. sp. Gnathosoma; A – Female, B – Male.

e_1/e_1-e_1 0.9 (0.9-1.2); f_1/f_1-f_1 1.9 (1.6-2.0); h_1/h_1-h_1 1.9 (1.9-3.0); c_1-c_1/d_1-d_1 0.4 (0.4-0.6); c_1-c_1/e_1-e_1 1.2 (1.2-1.5); d_1-d_1/e_1-e_1 2.7 (2.6-2.7); h_1/h_2 0.7 (0.7-0.8); $c_1-c_1/d_1-d_1: e_1-e_1: f_1-f_1$ = 1.2 (1.2-1.5); 2.7 (2.6-2.7); 1.0: 1.3 (1.3-1.4)

Venter (Figures 4C-D) — Ventral surface ornamented with striations; bearing three pairs of setae ($1a$, $3a$ and $4a$). Aggenital shield defined, with one pair of setae (g_1). Anogenital valves with three pairs of pseudanal setae (ps_{1-3}), ps_1 smaller than ps_2 and ps_3 . Measurements of setae: $1a$ 29 (28-31); $3a$ 22 (22-34); $4a$ 24 (24-25); ag_1 14 (14-15); ps_1 5 (5-10); ps_2 12 (11-12); ps_3 11 (11-12).

Gnathosoma (Figure 2B) — Gnathosoma (including palp) 127 (127-143); subcapitulum 79 (77-85); palp 93 (93-96); chelicera 95 (84-95). Subcapitulum bearing subcapitular setae m 20 (20-37) and n 42 (42-48) and adoral setae or_1 18 (16-23) and or_2 21 (20-21). Distance $m-m$ 42 (35-42), $n-n$ 31 (25-34) and $m-n$ 8 (6-8). Palp five segmented; palptrochanter without setae; palp femur with two barbed and one smooth seta; palpgenu with one seta; palptibia with two tactile setae + one well-developed claw + one spine-like accessory claw; palptarsus with four tactile setae + one solenidion (ω), one subapical eupathidium and one distal trifid eupathidium.

Legs (Figures 5A-D) — Length of legs I–IV: 288 (288-300); 240 (239-246); 219 (219-236); 247 (243-258). Femur I 76 (75-77), genu I 22 (22-24), tibia I 49 (49-51), tarsus I 69 (69-74). Dorsal most seta on femur I (dFI) barbed 49 (49-53), longer than h_1 , 3.8 (3.5-4.1) times length of h_1 ; dorsal most seta on genu I (dGI) barbed 36 (36-39). Chaetotaxy: coxae (excluding $1a$, $3a$ and $4a$) 2(+1elcl)-1-2-2, trochanters 1-1-1-1, femora 5-4-2-2, genua 3(+1 κ)-1-0-0, tibiae 5(+1 φ p)-5(+1 φ p)-5(+1 φ p)-5(+1 φ p), tarsi 12(+2 ω)-9(+2 ω)-7(+1 ω)-7(+1 ω). Length of solenidia: Tarsus I ω I 37 (34-37), ω II 26 (25-26); Tarsus II ω I 36 (32-36), ω II 32 (27-32); Tarsus III ω I 15 (15); Tarsus IV ω I 19 (19-20).

Immature stages — unknown

Remarks — The new species resembles *Agistemus longisetus* González-Rodríguez, 1963 and *A. brasiliensis* Matioli et al., 2002; but *Agistemus piquinnus* n. sp. can be distinguished




Figure 3 *Agistemus piquinnus* n. sp. (female); A – Leg I, B Leg II, C – Leg III, D – Leg IV.

from *A. longisetus* by a) lengths of median hysterosomal setae c_1 (73), d_1 (72) and e_1 (73) are shorter (vs 104, 103 and 115, respectively); b) ventral setae $1a$ (29), $3a$ (22) and $4a$ (24) are shorter (vs 41, 39 and 39, respectively) c) the ratio pob/eye (3.2), sci/pob (2.7), $vi/vi-vi$ (2.8), c_1/c_1-c_1 (2.1) and e_1/e_1-e_1 (2.4) are different (vs 4.1, 2.2, 2.6, 2.8 and 2.8, respectively); d) aggenital shield not horseshoe-shape and g_1 (26) twice longer than ps_3 (13) [vs aggenital shield in horseshoe-shape and g_1 (23) slightly longer than ps_3 (20)]. *Agistemus piquinnus n. sp.* can be distinguished from *A. brasiliensis* by a) setae vi (58), ve (82) and sci (79) are longer (vs 47, 77 and 72, respectively); b) ratio pob/eye (3.2), $vi/vi-vi$ (2.8), c_1/c_1-c_1 (2.1), d_1/d_1-d_1 (1.0), e_1/e_1-e_1 (2.4) are greater (vs 2.0, 2.5, 1.7, 0.8 and 1.9, respectively), h_1/h_1-h_1 (2.9) and h_1/h_2 (1.5) are less (vs 5.6 and 1.9, respectively); c) aggenital shield not horseshoe-shape and seta g_1 extending beyond the base of ps_2 (vs aggenital shield in horseshoe-shape and setae g_1 reach just past ps_3).

Etymology—The new species is named after the regional name of host plant “piquín”.

Type material—Holotype, 2 females paratypes and 3 paratypes males were collected on leaves of *Capsicum annuum* L. var. *glabriuscum* (Solanaceae), Ciudad Victoria, Tamaulipas, Mexico. Holotype, female on leaf of *Capsicum annuum* L. var. *glabriuscum* (Solanaceae), Cañón de la Peregrina, Ciudad Victoria, Tamaulipas. November 12, 2017. J. I. Monjarás B. leg.; two females collected on leaves of *Capsicum annuum* L. var. *glabriuscum* (Solanaceae).




Figure 4 *Agistemus piquinnus* n. sp. (male); A – Dorsum, B – Dorsum (PH), C – Anogenital area, D – Anogenital area (DIC).

Cañón de la Peregrina, Ciudad Victoria, Tamaulipas. July 10, 2019. J. I. Monjarás B. One male collected on leaf of *Capsicum annuum* L. var. *glabriuscum* (Solanaceae), Cañón de la Peregrina, Ciudad Victoria, Tamaulipas. April 28, 2017. J. I. Monjarás B. leg. Two males collected on leaves of *Capsicum annuum* L. var. *glabriuscum* (Solanaceae). Cañón de la Peregrina, Ciudad Victoria, Tamaulipas. November 12, 2017. J. I. Monjarás B. leg.




Figure 5 *Agistemus piquinnus* n. sp. (male); A – Leg I, B – Leg II, C – Leg III, D – Leg IV.

Key to females of *Agistemus* from Mexico

(modified from Paktinat-Saeij *et al.*, 2016)

1. One pair of aggenital setae (species group *fleschneri*) 2
— Two pairs of aggenital setae (species group *terminalis*) 3
2. Dorsal shield reticulated; ratios *vi/vi-vi* 2.6, *c₁/c₁-c₁* 1.0, *e₁/e₁-e₁* 1.8 *A. fleschneri* Summers
— Dorsal shield smooth; ratios *vi/vi-vi* 2.0, *c₁/c₁-c₁* 1.0, *e₁/e₁-e₁* 1.2 *A. striolatus* González-Rodríguez
3. Dorsal shield ornamented with small alveoli; ratios *vi/vi-vi* 2.6, *e₁/e₁-e₁* 2.2 *A. arcypaurus* González-Rodríguez
— Dorsal shield smooth 4
4. Setae *c₁* shorter than distance between the bases of *c₁-d₁* 5
— Setae *c₁* equal or longer than distance between the bases of *c₁-d₁* 7
5. Setae *g₁* reach to the base of setae *ps₃*; ratios *vi/vi-vi* 2.4, *c₁/c₁-c₁* 0.6 *A. simplex* González-Rodríguez

- Setae g_1 reach to the base of setae ps_2 6
6. Ratios $vi/vi-vi$ 1.0, c_1/c_1-c_1 0.6 *A. terminalis* (Quayle)
- Ratio $vi/vi-vi$ 1.5-1.7, c_1/c_1-c_1 1.5-1.6 *A. floridanus* González-Rodríguez
7. Length of medial hysterosomal setae > 100 μ m *A. longisetus* González-Rodríguez
- Length of medial hysterosomal setae < 100 μ m 8
8. Ratios pob/eye 3.1, d_1/d_1-d_1 1.0, h_1/h_2 1.4 *A. piquinnus* n. sp.
- Ratios pob/eye 2.0, d_1/d_1-d_1 0.8, h_1/h_2 1.9 *A. brasiliensis* Matoli et al.

Acknowledgements

We thank the Universidad Autónoma de Tamaulipas for supporting this research through project PFI2016-EB-07, PRODEP and CONACYT for the economic support to the doctoral stay. The authors are grateful to Universidade do Vale do Taquari - Univates for providing necessary materials during the study.

References

- Dhooria M.S. 2016. Acarine Technology. In: Dhooria M.S. (Ed) Fundamental of Applied Acarology, Springer Nature, Singapore. pp 21-39. doi:[10.1007/978-981-10-1594-6](https://doi.org/10.1007/978-981-10-1594-6)
- Estrada-Venegas E.G., Rodriguez-Navarro S., McMurry J.A. 2002. Some avocado mites from Michoacán, México. Int. J. Acarol., 28: 387-393. doi:[10.1080/01647950208684315](https://doi.org/10.1080/01647950208684315)
- Fan Q.-H., Flechtmann C.H.W. 2015. Chapter 7. Stigmeidae. In: Carrillo D., de Moraes G.J., Peña J. (Eds) Prospects for Biological Control of Plant Feeding Mites and Other Harmful Organisms, Springer, NY. pp 185-206. doi:[10.1007/978-3-319-15042-0_7](https://doi.org/10.1007/978-3-319-15042-0_7)
- Fan Q.-H., Zhang Z.-Q. 2005 Raphignathoidea (Acari: Prostigmata). Fauna of New Zealand, 52, 1-400.
- Fan Q.-H., Flechtmann C.H.W., de Moraes G.J. 2016. Annotated catalogue of Stigmeidae (Acari: Prostigmata), with a pictorial key to genera. Zootaxa, 4176: 1-199. doi:[10.11646/zootaxa.4176.1.1](https://doi.org/10.11646/zootaxa.4176.1.1)
- Fan Q.-H., Flechtmann, C.H.W., de Moraes G.J. 2019 Emendations and updates to "Annotated catalogue of Stigmeidae (Acari: Prostigmata), with a pictorial key to genera. Zootaxa, 4647(1): 088-103. doi:[10.11646/zootaxa.4647.1.9](https://doi.org/10.11646/zootaxa.4647.1.9)
- Gerson U., Smiley R., Ochoa R. 2003 Acarine pests of citrus: overview and non-chemical control. Syst. Appl. Acarol., 8: 3-12. doi:[10.11158/saa.8.1.1](https://doi.org/10.11158/saa.8.1.1)
- González-Rodríguez R.H. 1963. Four new mites of the genus *Agistemus* Summers, 1960 (Acarina: Stigmeidae). Acarología, 5: 342-350.
- González-Rodríguez R.H. 1965. A taxonomic study of the genera *Mediolata*, *Zetzellia* and *Agistemus* (Acarina: Stigmeidae). Univ. Calif. Publ. Entomol., 41: 1-64.
- Grandjean F. 1944. Observations sur les acariens de la famille des Stigmeidae. Arch. Sci. Phys. Nat., 26: 103-131.
- Jeppson L.R., Keifer H.H., Baker E.W. 1975. Mites injurious to economic plants. Berkeley: University of California Press, 641 pp.
- Johann L., Carvalho G.S., Majolo F., Ferla N.J. 2013. Stigmeid mites (Acari: Stigmeidae) from vineyards in the state of Rio Grande do Sul, Brazil. Zootaxa, 3701: 238-256. doi:[10.11646/zootaxa.3701.2.6](https://doi.org/10.11646/zootaxa.3701.2.6)
- Kethley J. 1990. Acariformes, Prostigmata (Actinedida). In: Dindal D.L. (Ed) Soil Biology Guide. John Wiley & Sons, New York, pp 667-756.
- Matioli A.L., Ueckermann E.A., Oliveira C.A.L. 2002. Some stigmeid and eupalopsellid mites from citrus orchards in Brazil (Acari: Stigmeidae and Eupalopsellidae). Int. J. Acarol., 28: 99-120. doi:[10.1080/01647950208684287](https://doi.org/10.1080/01647950208684287)
- Mineiro, J.L.C., Raga, A., Sato, M.E., Matoli, A.L., Berton, L.H.C. 2010. Mites of coffee plants (*Coffea* spp.) in State of São Paulo. Part II. Prostigmata [Ácaros de cafeiro (*Coffea* spp.) no Estado de São Paulo, Brasil. Parte II. Prostigmata]. Biota Neotrop., 10(4): 215-226. doi:[10.1590/S1676-06032010000400027](https://doi.org/10.1590/S1676-06032010000400027)
- Oudemans, A.C. 1931. Acarologische aanteekeningen CVIII. Entomol. Ber. 8(179): 251-263.
- Paktinat-Saeij S., Barroso G., Cruz W.P.D. 2017. Two new species of Stigmeidae (Acari: Trombidiformes: Rhaphignathoidea) from Brazil. Zootaxa, 4242: 372-382. doi:[10.11646/zootaxa.4242.2.9](https://doi.org/10.11646/zootaxa.4242.2.9)
- Rehman M.U., Kamran M., Alatawi F.J. 2018. Genus *Agistemus* Summers (Acari: Trombidiformes: Stigmeidae) from Saudi Arabia and a key to the world species. Syst. Appl. Acarol., 23: 1051-1072. doi:[10.11158/saa.23.6.5](https://doi.org/10.11158/saa.23.6.5)
- Santos M.A., Laing J.E. 1985. Stigmeidae predators. In: Helle W, Sabelis MW (Eds) Spider mites: their biology, natural enemies and control. V.1B. Amsterdam: Elsevier Science, pp 197-203.

- Summers, F.M. 1960. Several stigmaeid mites formerly included in *Mediolata* redescribed in *Zetzellia* Ouds and *Agistemus* new genus. Proc. Entomol. Soc. Wash., 62(4): 233-247.
- Zhang Z.-Q. 2003. Mites of Greenhouses, Identification, Biology and Control. CABI Publishing, Wallingford, UK, 244 pp. doi:10.1079/9780851995908.0000