

University of Dundee

Does additional monitoring status increase the reporting of ADRs?

Segec, A.; Slattery, J.; Morales, D. R.; Januskiene, J.; Kurz, X.; Arlett, P.

Published in:
Pharmacoepidemiology and Drug Safety

DOI:
[10.1002/pds.5174](https://doi.org/10.1002/pds.5174)

Publication date:
2020

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Segec, A., Slattery, J., Morales, D. R., Januskiene, J., Kurz, X., & Arlett, P. (2020). Does additional monitoring status increase the reporting of ADRs? An interrupted time series analysis of EudraVigilance data. *Pharmacoepidemiology and Drug Safety*. <https://doi.org/10.1002/pds.5174>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Does additional monitoring status increase the reporting of adverse drug reactions? An interrupted time series analysis of EudraVigilance data

Andrej Segec¹ | Jim Slattery¹ | Daniel R. Morales^{1,2} |
Justina Januskiene¹ | Xavier Kurz¹ | Peter Arlett^{1,3}

¹Pharmacovigilance and Epidemiology Department, European Medicines Agency, Amsterdam, The Netherlands

²Division of Population Health and Genomics, University of Dundee, Dundee, UK

³London School of Hygiene & Tropical Medicine, London, UK

Correspondence

Andrej Segec, European Medicines Agency, Domenico Scarlattilaan 6, 1083 HS Amsterdam, The Netherlands.
Email: andrej.segec@ema.europa.eu

Abstract

Purpose: To evaluate the impact of including a medicine in the list of medicinal products subject to additional monitoring (AM) on the reporting of adverse drug reactions (ADRs) in the European Economic Area (EEA).

Methods: Interrupted time series using the monthly number of EEA ADR reports in EudraVigilance during 12 months before and after the addition to AM list. The main outcome was the change (%) in reporting of ADRs with step change as the *a priori* impact model. Further time series analysis was performed using Joinpoint Regression.

Results: The analysis included 11 active substances. No significant immediate (step change) increase of reporting was identified for any product at time of addition to AM list. We identified a significant gradual increase of ADR reporting after addition to AM list (slope change) for two out of five new products—boceprevir (10% per month, 95% confidence interval (CI) 3%–18%) and denosumab-Xgeva (13% per month, 95% CI 4%–22%). No change was identified for Prolia, another denosumab-containing product not subject to AM. No significant increase was identified for any product included in the AM list due to the requirement to conduct a PASS. Conversely, a gradual decrease in reporting was identified for natalizumab (–5% per month; 95% CI –10% to –1%), rivaroxaban (–5%; –8 to –3%), and varenicline (–16%; –21 to –10%). The results were corroborated by the Joinpoint analyses, which yielded similar results.

Conclusions: We identified limited evidence that reporting of ADRs increased modestly and gradually for some new products and not for products with PASS requirement.

KEYWORDS

additional monitoring, adverse drug reactions, impact, pharmacovigilance

1 | INTRODUCTION

The views expressed in this article are the personal views of the author(s) and may not be understood or quoted as being made on behalf of or reflecting the position of the regulatory agency/agencies or organisations with which the author(s) is/are employed/affiliated.

Pharmacovigilance is the science and activities relating to the detection, assessment, understanding and prevention of adverse drug reactions (ADRs) or any other medicine-related problem.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Pharmacoepidemiology and Drug Safety* published by John Wiley & Sons Ltd.

Pharmacovigilance contributes to the protection and promotion of public health, timely access to innovative medicines and their safe and effective use after authorisation. Reporting of suspected ADRs is the cornerstone of pharmacovigilance as regular monitoring of newly received reports allows the identification of new risks and changes in known risks for medicines. Additional monitoring (AM) was introduced in Europe in 2010, following the revision of pharmacovigilance legislation (Regulation [EC] No 726/2004, Article 23), to increase the reporting of ADRs for targeted medicines.^{1,2}

Medicines falling under the mandatory scope of additional monitoring include new medicines, new biologicals, medicines authorised subject to conditions or under exceptional circumstances, and medicines with the obligation to conduct a post-authorisation safety study (PASS). Such medicines are identified with an inverted black triangle and an accompanying explanatory statement in their product information and are included in the list of medicines under additional monitoring which is updated and published monthly (AM list). Once on the list, a medicine will remain under additional monitoring for 5 years or until removed from the list by the Pharmacovigilance Risk Assessment Committee (PRAC). The black triangle allows to quickly identify medicines that are subject to additional monitoring and encourages patients and healthcare professionals to report any ADRs. At the end of 2019, 342 medicines were included in the AM list.²⁻⁴

The impact of additional monitoring as a policy intervention on the reporting of ADRs is unknown. Measuring the impact of regulatory decisions is important for all pharmacovigilance activities in order to improve existing processes.^{5,6} Consequently, we undertook this study using EudraVigilance data to investigate whether the inclusion of medicines in the AM list increases the reporting of ADRs for those medicines in Europe. The study was part of a data gathering project on the experience with additional monitoring, together with a survey of patients' and healthcare professionals' attitudes and behaviours towards reporting ADRs.^{1,7}

2 | METHODS

2.1 | Data sources

Information on the number of reported ADRs was obtained from EudraVigilance, the European database of ADR reports maintained by the European medicines agency (EMA) on behalf of the EU network, which at the end of 2018 contained over 14.5 million reports of suspected ADRs.⁸⁻¹⁰

2.2 | Exposures

Eligible medicines were identified from the December 2015 version of the AM list,³ by excluding those which did not have 12 months of authorised use prior to their addition to the list (to ensure we had a baseline of reporting for comparison) and those medicines for which their monthly reporting of ADRs from the EEA was less than 10 reports (over the 24 months of observation).

KEY POINTS

- EU legislation mandates that new medicines including new biologicals, medicines authorised subject to conditions or under exceptional circumstances and medicines with the obligation to conduct a post-authorisation safety study (PASS) are subject to additional monitoring aimed at increasing the reporting of ADRs by patients and healthcare providers.
- The real impact of additional monitoring on reporting of ADRs is currently unknown.
- Using an interrupted time series analysis of EudraVigilance ADR reporting data, we identified limited evidence that reporting of ADRs increased modestly and gradually for some new products. In contrast, reporting of ADRs did not increase (or even decreased) for products subject to AM due to the requirement to conduct a PASS.
- Further work is required to determine the effectiveness of AM as a policy intervention.

For each substance, we obtained the monthly number of post-marketing individual case reports of ADRs originating from the EEA from EudraVigilance,⁸⁻¹⁰ the EU database of ADR reports, for 12 months prior to and 12 months after inclusion of the product in the AM list (the intervention).

To account for changes in the number of patients exposed to the medicines over time, we calculated the size of at-risk population using medicine consumption/exposure data estimates obtained from Periodic Safety Update Reports¹¹ (PSUR) held by EMA for centrally-authorised products. Exposure data are normally reported as the total person-years exposed during the interval covered by the PSUR which can range from 6 months to several years. We therefore divided the person-years by the number of years covered by a PSUR period and smoothed the data over time using a piecewise linear form between the mid-point intervals. Where PSUR data were not available for the whole period under observation, but was missing only for a few months, we imputed the modelled exposure to the missing months based on the period with available data, as a continuation of that trendline. For products for which PSUR data was not available, we substituted the estimates of exposure by defined daily dose from publicly available nationwide drug consumption databases available from Denmark,¹² France,¹³ Netherlands,¹⁴ Norway,¹⁵ Sweden¹⁶ and United Kingdom,¹⁷ representing approx. 30% of the EU population.

2.3 | Analysis

Interrupted time series (ITS) is a suitable analytical method for evaluating the effect of interventions implemented at population level at a

defined time point.¹⁸ We applied segmented regression analysis to the interrupted time series data using Poisson regression and modelling the count of events per month, whilst offsetting the changes in at-risk population. To account for over-dispersion of the data we corrected the standard error by applying a scale parameter based on Pearson chi-square statistic divided by the residual degrees of freedom and we adjusted for seasonality using Fourier terms with two pairs of sine/cosine functions.^{18,19} We postulated an immediate effect (step change) as our *a priori* impact model; we also investigated a possible gradual change in trend (slope change). STATA (College Station, Texas, USA) version 15 was used for the analyses.

Further time series analysis was performed in parallel using Joinpoint Regression Program²⁰ (National Cancer Institute, Maryland, USA) version 4.5.0.1, using the Grid search method and Permutation testing. Statistical analysis using joinpoint regression identifies the time point(s) where a marked change in trend (the 'joinpoint') has occurred and estimates the regression function compared with the previously identified joinpoints. As the final number of joinpoints is established on the basis of a statistical criterion and their position is not fixed it does not require that an intervention date is pre-specified unlike interrupted time series regression.²¹ We compared the results of both statistical analyses.

3 | RESULTS

We identified 82 eligible products corresponding to 79 substances from the AM list with at least 12 months of baseline data and excluded 68 substances with low ADR reporting. The final analysis therefore contained 11 substances, five of which were included in the AM list due to new substance status (boceprevir, telaprevir, vemurafenib, fingolimod and denosumab) and six that were included because of an imposed PASS (imatinib, lenalidomide, natalizumab, rivaroxaban, valproic acid and varenicline), as detailed in Flowchart 1 and Table 1. Analyses were performed at substance level except for denosumab, for which we identified two products with different indications (Xgeva and Prolia) and discrepant AM status. Xgeva is

indicated for the prevention of skeletal related events in adults with advanced malignancies involving the bone and for the treatment of giant cell tumour of bone and is subject to AM.²² Prolia is indicated for the treatment of osteoporosis and for bone loss associated with hormone ablation in men with prostate cancer and is not subject to AM.²³ The analysis of these products was performed separately at product level with Prolia used as a control for Xgeva.


As presented in Table 2 and Figures 1 and 2, we did not identify a significant immediate effect (step change) for any studied product based on ITS analysis. We identified a significant increase of ADR reporting as a gradual change in trend after the addition to AM list (slope change) for boceprevir (10% per month, 95% CI 3%–18%) and denosumab-Xgeva (13% per month, 95% CI 4%–22%). No significant change in slope was identified for the other new substances (telaprevir, vemurafenib and fingolimod). No significant change (step or slope) was identified for imatinib, lenalidomide and valproate. Conversely, for three products included in the AM list due to the requirement to conduct a PASS, we identified a decrease in the reporting of ADRs after addition to the list (slope decrease): natalizumab (–5% per month, 95% CI –10% to –1%), rivaroxaban (–5% per month, 95% CI –8% to –3%) and varenicline (–16% per month, 95% CI –21% to –10%).

When the two denosumab-containing products were compared, the reporting for Xgeva, a product that was subject to AM, increased by 13% (95% CI 4%–22%) per month, but we detected no significant changes in reporting of ADRs for Prolia (0%, 95% CI –6% to 7%), a product that was not subject to AM.

The results were broadly comparably with the results of the Joinpoint analyses, which are presented in Figures 3 and 4, with the exception of natalizumab, for which no change was identified in the Joinpoint analysis.

4 | DISCUSSION

In our study we identified that ADR reporting increased after addition to the AM list for two out of five new products, in the order of 10%–



FLOWCHART 1 Selection of substances for the study. *Count includes valproic acid as one product and does not include denosumab-Prolia (used as control)

TABLE 1 Characteristics of medicinal products included in the study

Active substance	Product name	Indications/therapeutic area (MeSH)	Date of inclusion in the AM list	Reason for inclusion in the AM list
Boceprevir	Victrelis	Hepatitis C, chronic	April 2013	New active substance
Telaprevir	Incivo	Hepatitis C, chronic	April 2013	New active substance
Vemurafenib	Zelboraf	Melanoma	April 2013	New active substance
Fingolimod	Gilenya	Multiple sclerosis	April 2013	New active substance, PASS
Denosumab	Xgeva	<ul style="list-style-type: none"> •Fractures, bone •Neoplasm metastasis 	April 2013	New biological
	Prolia (control for Xgeva)	<ul style="list-style-type: none"> •Bone resorption •Osteoporosis, Postmenopausal 	N/A	N/A
Imatinib	Glivec	<ul style="list-style-type: none"> •Precursor cell lymphoblastic leukaemia-lymphoma •Gastrointestinal stromal tumours •Dermatofibrosarcoma •Myelodysplastic-myeloproliferative diseases •Leukaemia, myelogenous, chronic, BCR-ABL positive •Hypereosinophilic syndrome 	September 2014	PASS
Lenalidomide	Revlimid	•Multiple myeloma	June 2014	PASS
		•Lymphoma, mantle-cell		
		•Myelodysplastic Syndromes		
Natalizumab	Tysabri	Multiple sclerosis	April 2013	PASS
Rivaroxaban	Xarelto	<ul style="list-style-type: none"> •Arthroplasty, replacement •Venous thromboembolism 	July 2013	PASS
Valproic acid	Various	<ul style="list-style-type: none"> •Bipolar disorder •Epilepsy •Migraine disorders 	January 2015	PASS
Varenicline	Champix	Tobacco use cessation	April 2013	PASS

13% per month. For products included in the AM list due to the requirement for a PASS we identified either no changes or a decrease in the reporting of ADRs, in the order of -5 and -16% per time point. For denosumab, the only medicine that had two products with differing AM status, we observed a 13% increase in reporting for Xgeva, with no changes observed for Prolia (used as a control for Xgeva), which is compatible with an effect of additional monitoring. The results were broadly comparable with the results of the Joinpoint regression analysis.

This is to our knowledge the first study evaluating the impact of additional monitoring on the reporting of ADRs. Studies have previously reported that ADRs for medicines are under-reported, including for medicines subject to additional monitoring, and including serious and fatal ADRs.²⁴⁻²⁷ Such information is vital for pharmacovigilance decision making. Unequal and sometimes low awareness of additional monitoring amongst healthcare professionals was reported as potential reasons for this.²⁸ Our separate study of the knowledge of and attitudes to ADR reporting found that although 88% of respondents would report an ADR for a medicine subject to AM, identified with a black triangle, 1 in 5 reporters misunderstood the concept of AM and only 37% of those who reported an ADR for a product subject to AM were influenced by the AM status.¹ These factors could partially

TABLE 2 Summary of results—interrupted time series analysis of the monthly reporting of post-marketing ADRs from the EEA

Active substance	Step change RR (95% CI)	Slope change RR (95% CI)
Boceprevir ^a	0.59 (0.23–1.55)	1.10 (1.03–1.18)
Telaprevir ^a	0.85 (0.35–2.06)	1.02 (0.96–1.09)
Vemurafenib ^a	0.90 (0.38–2.11)	0.96 (0.89–1.02)
Fingolimod ^a	1.14 (0.74–1.75)	0.97 (0.94–1.00)
Denosumab ^a –Xgeva	0.73 (0.30–1.77)	1.13 (1.04–1.22)
Denosumab ^a –Prolia (not in the AM list, control for Xgeva)	1.43 (0.62–3.28)	1.00 (0.94–1.07)
Imatinib ^b	0.78 (0.42–1.42)	1.03 (0.99–1.08)
Lenalidomide	1.54 (0.71–3.53)	1.03 (0.98–1.10)
Natalizumab	0.96 (0.49–1.87)	0.95 (0.90–0.99)
Rivaroxaban	1.30 (0.92–1.85)	0.95 (0.92–0.97)
Valproic acid ^b	0.91 (0.50–1.67)	1.01 (0.97–1.06)
Varenicline	0.96 (0.42–2.23)	0.84 (0.79–0.90)

^aNew medicines.

^bAnalysis using exposure data from prescription databases (PSUR data not available).




FIGURE 1 Interrupted time series analysis. New products included in the study. Reporting of post-marketing ADRs from the EEA. Green vertical line: addition to AM list. Red line: predicted reporting from seasonality-adjusted model. Dashed line: de-seasonalised trend

explain the limited effect we found on increasing the reporting of ADRs.

Due to methodological limitations, the maximum number of substances that we were able to include in our study (11 of 79 substances) was limited by the availability of their baseline data or low reporting, which restricts the overall generalisability of the findings. A large number of new medicines were excluded simply because there was no comparative data available before their inclusion in the AM list. The small sample of substances that we were able to study is also

limited in terms of chemical/biological and pharmacological classes and may therefore not be universally generalisable to all medicines subject to AM. A possible solution to overcome this limitation would be to stagger future implementation of such policy interventions in various regions over time, which would serve as comparators in a step-wedge type of approach, but this may not be feasible for EU wide interventions. Alternatively, where possible, a controlled ITS design can be used, as illustrated with the denosumab example. Another approach that could be employed would be to study the




FIGURE 2 Interrupted time series analysis. Products included in the study due to the requirement to conduct a PASS. Reporting of post-marketing ADRs from the EEA. Green vertical line: addition to AM list. Red line: predicted reporting from seasonality-adjusted model. Dashed line: de-seasonalised trend

change in reporting at the time of removal of products from the AM list.

Our study focused on changes during 12 months after the addition to AM list. For products already on the market which newly acquire additional monitoring status, it may take several months or years for the medicines with the updated product information to reach patients (medicines already distributed are not re-labelled). It is therefore possible that a longer follow-up period may be needed to detect significant changes, if awareness of the AM status increases slowly and gradually. Considering that no step change was observed

in our study, this remains a possibility. Longer follow up would also allow studying whether any observed increases are sustained or temporary and thus whether immediate changes and long-term patterns differ. However, as ITS examines associations only, this would be at the risk that other interventions including regulatory interventions, may confound the interpretation over time.

Additionally, the monthly counts of ADR reports were generally low. This combined with 24 time points and unequal variability in the monthly reports, restricted the power to detect a difference, as well as the utility of seasonality adjustment. Due to the small number of




FIGURE 3 Jointpoint regression analysis—new products included in the study. Reporting of post-marketing ADRs from the EEA, addition to AM list is at time point 12 (time in months)




FIGURE 4 Joinpoint regression analysis—products included in the study due to PASS requirement. Reporting of post-marketing ADRs from the EEA, addition to AM list is at time point 12 (time in months)

substances which we were able to study as well as the low monthly ADR counts we were unable to stratify the analyses by the reason of addition to AM list, by country, or by reporter type (patients or healthcare professionals) as initially intended. Therefore although differences in ADR reporting may exist between different reporter types, as reported in our separate study on reporter attitudes and behaviours,¹ we could not examine this effect in this study. Seven of the studied substances were included in the first version of the list in April 2013 and the remaining four between 2013 and 2015. Therefore, changes in these reporting sub-categories and changes over time cannot be excluded.

The application of ITS methods to medicine use data, other than being limited by the lack of baseline data for new medicines, also requires reliable estimates of exposure data due to the often rapid changes in the size of the population at risk, as opposed to population studies. Medicine use can increase rapidly after authorisation/reimbursement and inclusion in clinical guidelines or decrease due to safety concerns or replacement with a more effective, safer or more convenient competitor. The estimates of exposure that we have relied on are often based on approximations from sales data using an expected dose and treatment duration¹¹ and can therefore be imprecise and be a source of residual confounding as well as distort important trends.

Other influences in the changes in exposure, especially if they also relate to the reporting of ADRs would be equally obscured by such calculations and cannot thus be investigated. Similarly, the absence of knowledge of other time-varying confounders and their exact timing, for example, media attention to the results of a study suggesting the risk of serious ADRs, direct healthcare professional communication, restrictions in authorised use due to ADRs, increased awareness due to changes in the international or national treatment guidelines and precautions/screening at treatment initiation or recommendations for regular testing, and so on, which may have influenced the reporting of ADRs, can further complicate the interpretation of time series analyses or lead to unmeasured confounding. Additionally, patients who are at the highest risk of an adverse outcome may no longer be treated after such regulatory actions, and thus the occurrence of ADRs decreases. We did not collect details about the lifecycle of individual medicines included in our study, and therefore couldn't investigate these factors in the current study.

Indeed, half of the substances included in our study were included in the AM list due to the obligation to conduct a PASS. The imposition of a PASS often follows the emergence of serious ADRs, concerns about medicine safety or the evaluation of benefit–risk balance by the PRAC (e.g. via a referral procedure), with consequent media attention. Such a setting is prone to confounding due to other regulatory actions and media attention possibly influencing ADR reporting and therefore the results are difficult to interpret in terms of causality. This may be one reason why we observed a fall in ADR reporting for several products added to the AM list due to an imposed PASS. However, should this observation be confirmed in future research, it may serve to inform any future discussions on the legislation governing additional monitoring.

5 | CONCLUSION

In summary, we identified limited evidence that reporting of ADRs increased modestly and gradually for some new products and did not increase for products subject to AM due to the requirement to conduct a PASS. The small number of medicines that we were able to include in our study, together with its ecological design, makes the causality of this observation difficult to establish. We suggest that, in the future it would be worthwhile to pre-specify the methods for the evaluation of policy interventions such as the introduction of additional monitoring, to help overcome the shortcomings of a retrospective evaluation. Given the limitations in our results, we would welcome suggestions from the research and regulatory communities on complementary methods that might be applied to study the impact of AM.

ETHICAL STATEMENT

The study did not involve human subjects and relied on routine data.

ACKNOWLEDGMENTS

We are grateful to the members of the EMA's Pharmacovigilance Risk Assessment Committee for their comments on the design of this study. We are also grateful to Dr Georgy Genov for helpful comments on the study outline and draft article. The work presented in this article did not receive project funding or grants.

Prior postings, sponsors and grants: Partial results were included in the European Commission report⁷ (which has since been published). DRM has received funding support from the Wellcome Trust, NIHR, Chief Scientist Office and Tenovus Scotland for research unrelated to this work.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

ORCID

Andrej Segec  <https://orcid.org/0000-0003-2881-0593>

Jim Slattery  <https://orcid.org/0000-0002-8352-9350>

Daniel R. Morales  <https://orcid.org/0000-0002-0063-8069>

Justina Januskiene  <https://orcid.org/0000-0001-6758-650X>

Xavier Kurz  <https://orcid.org/0000-0002-9838-7754>

Peter Arlett  <https://orcid.org/0000-0002-6640-0117>

REFERENCES

1. Januskiene J, Segec A, Slattery JM, et al. What are the patients' and health care professionals' understanding and behaviors towards adverse drug reaction reporting and additional monitoring? *Pharmacoepidemiol Drug Saf.* 2020 Oct 25. <https://doi.org/10.1002/pds.5162>.
2. Regulation (EC) No 726/2004 of the European Parliament and of the council of 31 March 2004 laying down Community procedures for the authorisation and supervision of medicinal products for human and veterinary use and establishing a European Medicines Agency, *Official Journal L - 136, 30/04/2004, p. 1–33.* <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex:02004R0726-20190128>. Accessed April 30, 2019.
3. European Medicines Agency. List of medicines subject to additional monitoring, EMA/245297/2013 Rev.60. <https://www.ema.europa>.

- eu/en/human-regulatory/post-authorisation/pharmacovigilance/medicines-under-additional-monitoring/list-medicines-under-additional-monitoring. Accessed January 17, 2020.
4. European Medicines Agency. Guideline on good pharmacovigilance practices Module X – additional monitoring. http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2013/04/WC500142282.pdf. Accessed April 30, 2019.
 5. Goedecke T, Morales DR, Pacurariu A, Kurz X. Measuring the impact of medicines regulatory interventions—systematic review and methodological considerations. *Br J Clin Pharmacol*. 2018;84(3):419-433.
 6. European Medicines Agency. PRAC strategy on measuring the impact of pharmacovigilance activities (Rev 1), EMA/165407/2017. http://www.ema.europa.eu/docs/en_GB/document_library/Other/2016/01/WC500199756.pdf. Accessed April 30, 2019.
 7. European Commission. European Medicines Agency and Member States joint report to the European Commission on the experience with the list of products subject to additional monitoring. https://ec.europa.eu/health/sites/health/files/files/pharmacovigilance/docs/2019_ema-ms_additional-monitoring_en.pdf. Accessed November 19, 2019.
 8. European Medicines Agency. 2018 Annual Report on EudraVigilance for the European Parliament, the Council and the Commission. https://www.ema.europa.eu/en/documents/report/2018-annual-report-eudravigilance-european-parliament-council-commission-reporting-period-1-january_en.pdf. Accessed July 18, 2019.
 9. European Medicines Agency. EudraVigilance. <https://www.ema.europa.eu/en/human-regulatory/research-development/pharmacovigilance/eudravigilance>. Accessed July 24, 2019.
 10. Postigo R, Brosch S, Slattery J, et al. EudraVigilance medicines safety database: publicly accessible data for research and public health protection. *Drug Saf*. 2018;41(7):665-675.
 11. European Medicines Agency. Guideline on good pharmacovigilance practices (GVP)—Module VII – Periodic safety update report (Rev 1). https://www.ema.europa.eu/en/documents/scientific-guideline/guideline-good-pharmacovigilance-practices-gvp-module-vii-periodic-safety-update-report_en.pdf. Accessed April 30, 2019.
 12. Register of Medicinal Product Statistics. www.medstat.dk.
 13. Health Insurance system. <http://www.ameli.fr/l-assurance-maladie/statistiques-et-publications/donnees-statistiques/medicament/medic-am/medic-am-mensuel-2016.php>.
 14. The Drug Information System of National Health Care Institute (GIP Databank). www.gipdatabank.nl.
 15. Norwegian Prescription Database (NorPD). <http://www.norpd.no>.
 16. Statistikdatabas för läkemedel. <http://www.socialstyrelsen.se/statistik/statistikdatabas/lakemedel>.
 17. National Health Service/Information Services Division Scotland. <http://www.isdscotland.org/Health-Topics/Prescribing-and-medicines/Community-Dispensing/Prescription-Cost-Analysis/>. Welsh Government: <http://gov.wales/statistics-and-research/prescriptions-dispensed-community/?tab=previous&lang=en>. National Health Service (NHS) Digital: <https://data.gov.uk/dataset/prescription-cost-analysis-england>.
 18. Lopez Bernal J, Cummins S, Gasparrini A. Interrupted time series regression for the evaluation of public health interventions: a tutorial. *Int J Epidemiol*. 2017;46(1):348-355.
 19. Bhaskaran K, Gasparrini A, Hajat S, Smeeth L, Armstrong B. Time series regression studies in environmental epidemiology. *Int J Epidemiol*. 2013;42:1187-1195.
 20. U.S. Department of Health and Human Services—National Institutes of Health—National Cancer Institute: Joinpoint. <https://surveillance.cancer.gov/joinpoint/>. Accessed July 24, 2019.
 21. Kim HJ, Fay MP, Feuer EJ, Midthune DN. Permutation tests for joinpoint regression with applications to cancer rates. *Stat Med*. 2000;19:335-351.
 22. European Medicines Agency. Xgeva European Public Assessment Report. <https://www.ema.europa.eu/en/medicines/human/EPAR/xgeva>. Accessed July 18, 2019.
 23. European Medicines Agency. Prolia European Public Assessment Report. <https://www.ema.europa.eu/en/medicines/human/EPAR/prolia>. Accessed July 18, 2019.
 24. Patrignani A, Palmieri G, Ciampini N, Moretti V, Mariani A, Racca L. Under-reporting of adverse drug reactions, a problem that also involves medicines subject to additional monitoring. Preliminary data from a single-center experience on novel oral anticoagulants. *G Ital Cardiol*. 2018;19(1):54-61.
 25. Hazell L, Shakir SA. Under-reporting of adverse drug reactions: a systematic review. *Drug Saf*. 2006;29(5):385-396.
 26. Varallo FR, Guimarães SOP, Abjaude SAR, Mastroianni PC. Causes for the underreporting of adverse drug events by health professionals: a systematic review. *Rev Esc Enferm USP*. 2014;48(4):739-747.
 27. Al Dweik R et al. Factors affecting patient reporting of adverse drug reactions: a systematic review. *Br J Clin Pharmacol*. 2017;83(4):875-883.
 28. O'Callaghan J et al. Knowledge of adverse drug reaction reporting and the pharmacovigilance of biological medicines: a survey of healthcare professionals in Ireland. *BioDrugs*. 2018;32(3):267-280.

How to cite this article: Segec A, Slattery J, Morales DR, Januskiene J, Kurz X, Arlett P. Does additional monitoring status increase the reporting of adverse drug reactions? An interrupted time series analysis of EudraVigilance data. *Pharmacoepidemiol Drug Saf*. 2020;1–10. <https://doi.org/10.1002/pds.5174>