

First-order virial expansion of short-time diffusion and sedimentation coefficients of permeable particles suspensions

Bogdan Cichocki, Maria L. Ekiel-Jeżewska, Gerhard Nägele, and Eligiusz Wajnryb

Citation: *Phys. Fluids* **23**, 083303 (2011); doi: 10.1063/1.3626196

View online: <http://dx.doi.org/10.1063/1.3626196>

View Table of Contents: <http://pof.aip.org/resource/1/PHFLE6/v23/i8>

Published by the [American Institute of Physics](http://www.aip.org).

Additional information on Phys. Fluids

Journal Homepage: <http://pof.aip.org/>

Journal Information: http://pof.aip.org/about/about_the_journal

Top downloads: http://pof.aip.org/features/most_downloaded

Information for Authors: <http://pof.aip.org/authors>

ADVERTISEMENT



**Running in Circles Looking
for the Best Science Job?**

Search hundreds of exciting
new jobs each month!

<http://careers.physicstoday.org/jobs>

physicstodayJOBS



First-order virial expansion of short-time diffusion and sedimentation coefficients of permeable particles suspensions

Bogdan Cichocki,¹ Maria L. Ekiel-Jeżewska,^{2,a)} Gerhard Nägele,³ and Eligiusz Wajnryb²

¹*Faculty of Physics, Institute of Theoretical Physics, University of Warsaw, Hoża 69, 00-681 Warsaw, Poland*

²*Institute of Fundamental Technological Research, Polish Academy of Sciences, Pawińskiego 5B, 02-106 Warsaw, Poland*

³*Institute of Complex Systems, ICS-3, Forschungszentrum Jülich, D-52425 Jülich, Germany*

(Received 23 January 2011; accepted 20 July 2011; published online 31 August 2011)

For suspensions of permeable particles, the short-time translational and rotational self-diffusion coefficients, and collective diffusion and sedimentation coefficients are evaluated theoretically. An individual particle is modeled as a uniformly permeable sphere of a given permeability, with the internal solvent flow described by the Debye-Bueche-Brinkman equation. The particles are assumed to interact non-hydrodynamically by their excluded volumes. The virial expansion of the transport properties in powers of the volume fraction is performed up to the two-particle level. The first-order virial coefficients corresponding to two-body hydrodynamic interactions are evaluated with very high accuracy by the series expansion in inverse powers of the inter-particle distance. Results are obtained and discussed for a wide range of the ratio, x , of the particle radius to the hydrodynamic screening length inside a permeable sphere. It is shown that for $x \gtrsim 10$, the virial coefficients of the transport properties are well-approximated by the hydrodynamic radius (annulus) model developed by us earlier for the effective viscosity of porous-particle suspensions.

© 2011 American Institute of Physics. [doi:10.1063/1.3626196]

I. INTRODUCTION

One of the theoretical methods to analyze transport properties in suspensions of interacting colloidal particles is the virial expansion in terms of the particle volume fraction ϕ . For suspensions of non-permeable hard spheres with stick hydrodynamic boundary conditions, virial expansion results for short-time properties are known to high numerical precision up to the three-particle level, i.e., to quadratic order in ϕ for diffusion and sedimentation coefficients,^{1,2} and to third order in ϕ for the effective viscosity.³ The concentration range of applicability of these hard-sphere virial expansion results in comparison to simulation data that has been discussed in Ref. 4. Our knowledge on virial expansion coefficients of colloidal transport properties is less developed when suspensions of solvent-permeable particles are considered. The theoretical description of their dynamics is more complicated since one needs to account for the solvent flow also inside the particles. Permeable particle systems are frequently encountered in soft matter science. Prominent examples of practical relevance, which are the subject of ongoing research, are dendrimers,^{5,6} microgel particles,⁷⁻⁹ a large variety of core-shell particles with a dry core and an outer porous layer,¹⁰⁻¹⁴ fractal aggregates,¹⁵ and star-like polymers of lower functionality.¹⁶

In a series of recent articles,¹⁷⁻²⁰ we have explored the generic effect of solvent permeability on the short-time transport using the model of uniformly permeable colloidal spheres with excluded volume interactions. This simple model is specified by two parameters only, namely, the particle volume fraction $\phi = (4\pi/3)na^3$, where n is the number

concentration and a is the particle radius, and the ratio x of the particle radius to the hydrodynamic penetration depth inside a permeable sphere. Large (low) values of x correspond to weakly (strongly) permeable particles. While the model of uniformly permeable hard spheres ignores a specific intra-particle structure, it is generic in the sense that the hydrodynamic structure of more complex porous particles can be approximately accounted for in terms of a mean permeability. In our related previous publications, using a hydrodynamic multipole simulation method of a very high accuracy,²¹ encoded in the HYDROMULTIPOLE program package,¹ we have calculated the short-time translational diffusion properties^{17,18} and the high-frequency viscosity^{19,20} of the permeable spheres model as functions of ϕ and x . These results cover the full range of permeabilities, with volume fractions extending up to the liquid-solid transition.

While the simulation results are important for the general understanding of permeability effects in concentrated systems, for practical use in experimental data evaluation and as input in long-time theories, virial expansion results based on a rigorous theoretical calculation are still strongly on demand. In fact, the knowledge of the leading-order virial coefficients can be a good starting point in deriving approximate expressions for transport properties, which may be applicable at concentrations much higher than those where the original (truncated) virial expansion result is useful. An example in case is provided in our recent derivation²⁰ of a generalized Saitô expression for the effective high-frequency viscosity η_∞ of permeable spheres, based on the second-order concentration expansion result, i.e., a Huggins coefficient calculation, of this property.

In Ref. 19, we have performed virial expansion calculations for η_∞ . We have investigated therein a simplifying hydrodynamic radius model (HRM), where a uniformly

^{a)}Electronic mail: mekiel@ippt.gov.pl.

permeable sphere of radius a is described by a spherical annulus particle with an inner hydrodynamic radius $a_{\text{eff}}(x) < a$, and unchanged excluded-volume radius a . In this annulus model (HRM), the Huggins coefficient describing two-body viscosity contributions has been evaluated and shown to be in a remarkably good agreement with the precise numerical data for porous particles, characterized by a wide range of permeabilities realized in experimental systems.

In the present article, the aforementioned theoretical work on the virial expansion of the high-frequency viscosity is generalized to short-time diffusion properties. In Sec. II, the virial expansion is performed for the translational and rotational self-diffusion coefficients D_t and D_r , respectively, the sedimentation coefficient K , and the associated collective diffusion coefficient $D_C = KD'_0/S(0)$. Here, $S(0)$ is the small-wavenumber limit of the static structure factor and D'_0 is the single-particle translational diffusion coefficient. In Sec. III, we provide highly accurate numerical values for the first-order (i.e., two-particle) virial coefficients, $\lambda_t(x)$, $\lambda_r(x)$, $\lambda_C(x)$, and $\lambda_K(x)$, of D_t , D_r , D_C , and K , respectively, in the full range of permeabilities. In Sec. IV, we also recalculate these virial coefficients approximately on the basis of the simplifying annulus model (HRM). In Sec. VI, we conclude that in the range $x \gtrsim 10$ typical of many permeable particle systems, the virial coefficients are well approximated by the annulus model.

II. THEORY

We consider a suspension made of a fluid with shear viscosity η_0 and identical porous particles of radius a . The fluid flow is characterized by Reynolds number $\text{Re} \ll 1$. Outside the particles, the fluid velocity \mathbf{v} and pressure p satisfy the Stokes equations,^{22,23}

$$\begin{aligned} \eta_0 \nabla^2 \mathbf{v}(\mathbf{r}) - \nabla p(\mathbf{r}) &= 0 \\ \nabla \cdot \mathbf{v}(\mathbf{r}) &= 0, \end{aligned} \quad (1)$$

and inside the particles, the Debye-Büchse-Brinkman (DBB) equations,^{24,25}

$$\begin{aligned} \eta_0 \nabla^2 \mathbf{v}(\mathbf{r}) - \eta_0 \kappa^2 [\mathbf{v}(\mathbf{r}) - \mathbf{u}_i(\mathbf{r})] - \nabla p(\mathbf{r}) &= 0 \\ \nabla \cdot \mathbf{v}(\mathbf{r}) &= 0, \end{aligned} \quad (2)$$

where κ^{-1} is the hydrodynamic penetration depth. The skeleton of the particle i , centered at \mathbf{r}_i , moves rigidly with the local velocity $\mathbf{u}_i(\mathbf{r}) = \mathbf{U}_i + \boldsymbol{\omega}_i \times (\mathbf{r} - \mathbf{r}_i)$, determined by the translational and rotational velocities \mathbf{U}_i and $\boldsymbol{\omega}_i$ of the particle i , respectively. The fluid velocity and stress tensor are continuous across a particle surface. The effect of the particle porosity is therefore described by the ratio x of the particle radius a to the hydrodynamic screening length κ^{-1} of the porous material inside the particle, i.e.,

$$x = \kappa a. \quad (3)$$

Owing to linearity of the Stokes and DBB equations and the boundary conditions, the particle velocity \mathbf{U}_i depends linearly on the external forces \mathbf{F}_j exerted on a particle j . In particular, for two interacting spherical particles, 1, 2, in the absence of external torques and flows,

$$\mathbf{U}_1 = \boldsymbol{\mu}_{11}^t(\mathbf{r}_1, \mathbf{r}_2) \cdot \mathbf{F}_1 + \boldsymbol{\mu}_{12}^t(\mathbf{r}_1, \mathbf{r}_2) \cdot \mathbf{F}_2, \quad (4)$$

$$\mathbf{U}_2 = \boldsymbol{\mu}_{21}^t(\mathbf{r}_1, \mathbf{r}_2) \cdot \mathbf{F}_1 + \boldsymbol{\mu}_{22}^t(\mathbf{r}_1, \mathbf{r}_2) \cdot \mathbf{F}_2. \quad (5)$$

In this paper, the two-particle translational-translational mobility matrices $\boldsymbol{\mu}_{ij}^t(\mathbf{r}_1, \mathbf{r}_2)$ are evaluated using the multipole method of solving the Stokes and DBB equations.²¹ The cluster expansion of the above mobility matrices reads,

$$\boldsymbol{\mu}_{ij}^t(\mathbf{r}_1, \mathbf{r}_2) = \mu_0^t \delta_{ij} \mathbf{1} + \boldsymbol{\mu}_{ij}^{t(2)}(\mathbf{r}), \quad (6)$$

where $\mathbf{r} = \mathbf{r}_2 - \mathbf{r}_1$ and

$$\mu_0^t = \frac{1}{4\pi\eta_0 A_{10}} \quad (7)$$

is the single porous-particle translational mobility. Here A_{10} is a single porous-particle scattering coefficient,²⁶ given explicitly in Appendix A. For a non-permeable hard sphere with stick boundary conditions, $A_{10}^{\text{hs}} = 3a/2$.

The single particle scattering coefficients $A_{l\sigma}$, with $l = 1, 2, 3, 4, \dots$ and $\sigma = 1, 2, 3, \dots$ are essential to perform the multipole expansion. They determine the corresponding multipoles of the hydrodynamic force density on a particle immersed in an ambient flow; examples are Eqs. (7) and (24). The same coefficients $A_{l\sigma}$ specify also the corresponding multipoles of the fluid velocity, reflected (scattered) by a particle immersed in a given ambient flow. This is why $A_{l\sigma}$ is called ‘‘scattering coefficients.’’ In the multipole approach, differences in the internal structure of particles (e.g., solid, liquid, gas, porous, core-shell, stick-slip) are fully accounted for by different scattering coefficients. The other parts of the multipole algorithm need not to be changed. The scattering coefficients are the matrix elements of two single-particle friction operators, \mathbf{Z}_0 and $\hat{\mathbf{Z}}_0$, which determine the hydrodynamic force density exerted by a given ambient flow on a motionless and a freely moving particle, respectively.

In the multipole expansion method, the two-particle mobility $\boldsymbol{\mu}^{(2)}(1, 2)$ (e.g., translational one, as in Eq. (6), or rotational one, as in Eq. (23)) is expressed in terms of the single-particle friction operators, $\mathbf{Z}_0(i)$ and $\hat{\mathbf{Z}}_0(i)$, with $i = 1, 2$, and the Green operator $\mathbf{G}(1, 2)$. The latter relates the flow outgoing from particle 2 and incoming on particle 1. We can write $\boldsymbol{\mu}^{(2)}$ as an infinite scattering series,

$$\begin{aligned} \boldsymbol{\mu}^{(2)} &= \boldsymbol{\mu}_0 \mathbf{Z}_0 (1 + \mathbf{G} \hat{\mathbf{Z}}_0)^{-1} \mathbf{G} \mathbf{Z}_0 \boldsymbol{\mu}_0 \\ &= \boldsymbol{\mu}_0 \mathbf{Z}_0 \mathbf{G} \mathbf{Z}_0 \boldsymbol{\mu}_0 - \boldsymbol{\mu}_0 \mathbf{Z}_0 \mathbf{G} \hat{\mathbf{Z}}_0 \mathbf{G} \mathbf{Z}_0 \boldsymbol{\mu}_0 + \dots \end{aligned} \quad (8)$$

Since the multipole matrix elements of the Green tensor \mathbf{G} scale as inverse powers of the interparticle distance r , Eq. (8) corresponds to a power series in $1/r$. Truncating the expansion at order $1/r^{1000}$, we obtain a very high precision of the mobility calculation, actually much higher than that needed for any practical applications.

In the present work, we investigate the short-time dynamics, at time scales $t \ll a^2/D'_0$, where

$$D'_0 = k_B T \mu_0^t, \quad (9)$$

is the single-particle translational diffusion coefficient, with the Boltzmann constant k_B and temperature T . On the short-time

scale, the system is described by the equilibrium particle distribution.²⁷ In the further analysis, we will need only the small-concentration limit $g_0(r)$ of the pair distribution function, where r is the interparticle distance. For particle-particle direct interactions described by a pair potential $V(r)$, this pair distribution is $g_0(r) = \exp(-V(r)/k_B T)$. For non-overlapping spheres of radius a ,

$$g_0(r) = \begin{cases} 0 & \text{for } r \leq 2a, \\ 1 & \text{for } r > 2a. \end{cases} \quad (10)$$

The first-order terms in the virial expansion of the short-time transport coefficients are obtained by averaging the corresponding two-particle mobility elements. As a result, the virial coefficients are obtained as integrals, which involve the mobility elements and $g_0(r)$.

The first-order virial expansion of the short-time translational self-diffusion coefficient has the form,

$$D_t = D_0^t(1 + \lambda_t \phi + \mathcal{O}(\phi^2)). \quad (11)$$

The coefficient λ_t is given by the relation,²⁸

$$\lambda_t = 8 \int_1^{+\infty} g_0(R) J_t(R) R^2 dR, \quad (12)$$

with $\mathbf{R} = \mathbf{r}/2a$ and

$$J_t(R) = \frac{1}{\mu_0^t} \text{Tr} \boldsymbol{\mu}_{11}^{tt(2)}(\mathbf{r}). \quad (13)$$

Here, Tr denotes the trace operation. For the sedimentation coefficient, one obtains

$$K = 1 + \lambda_K \phi + \mathcal{O}(\phi^2), \quad (14)$$

where

$$\lambda_K = \frac{2}{5a^3} A_{12} + \frac{8}{a} A_{10} \int_0^{+\infty} [g_0(R) - 1] R dR, \quad (15)$$

$$+ 8 \int_1^{+\infty} g_0(R) J_K(R) R^2 dR, \quad (16)$$

and

$$J_K(R) = \frac{1}{\mu_0^t} \text{Tr} \left[\boldsymbol{\mu}_{11}^{tt(2)}(\mathbf{r}) + \boldsymbol{\mu}_{12}^{tt(2)}(\mathbf{r}) - \mathbf{T}_0(\mathbf{r}) \right]. \quad (17)$$

In the above expression,

$$\mathbf{T}_0(\mathbf{r}) = \frac{\mathbf{1} + \hat{\mathbf{r}}\hat{\mathbf{r}}}{8\pi\eta_0 r} \quad (18)$$

is the Oseen tensor and $\hat{\mathbf{r}} = \mathbf{r}/r$.

The scattering coefficient A_{12} for a porous particle²⁶ is given explicitly in Appendix A. For a non-permeable hard sphere with the stick boundary conditions, $A_{12}^{\text{hs}} = 5a^3/2$.

The collective diffusion coefficient is given by

$$D_C = D_0^t K / S(0), \quad (19)$$

where $S(0)$ is the zero-wavenumber limit of the static structure factor, $S(0) = \lim_{q \rightarrow 0} S(q)$. The first-order virial expansion of Eq. (19) has the form,

$$D_C = D_0^t(1 + \lambda_C \phi + \mathcal{O}(\phi^2)). \quad (20)$$

For the non-overlapping spheres,²⁷

$$S(0) = 1 - 8\phi + \mathcal{O}(\phi^2). \quad (21)$$

In this case,

$$\lambda_C = \lambda_K + 8. \quad (22)$$

The relation (22) follows from Eqs. (14), (19), and (20).

We proceed by analyzing the short-time rotational self-diffusion coefficient. In the absence of external forces and flows, the two-particle rotational-rotational mobility matrices $\boldsymbol{\mu}_{ij}^{rr}(\mathbf{r}_1, \mathbf{r}_2)$ satisfy the relation analogical to Eqs. (4)–(5), with the translational velocities replaced by the rotational ones, and the forces replaced by the torques. The two-particle cluster expansion now reads,

$$\boldsymbol{\mu}_{ij}^{rr}(\mathbf{r}_1, \mathbf{r}_2) = \mu_0^r \delta_{ij} \mathbf{1} + \boldsymbol{\mu}_{ij}^{rr(2)}(\mathbf{r}), \quad (23)$$

with

$$\mu_0^r = \frac{1}{8\pi\eta_0 A_{11}}. \quad (24)$$

The scattering coefficient A_{11} for a porous particle²⁶ is given in Appendix A. For a non-permeable hard sphere with the stick boundary conditions, $A_{11}^{\text{hs}} = a^3$.

The virial expansion of the rotational self-diffusion coefficient is

$$D_r = D_0^r(1 + \lambda_r \phi + \mathcal{O}(\phi^2)), \quad (25)$$

where

$$D_0^r = k_B T \mu_0^r, \quad (26)$$

and

$$\lambda_r = 8 \int_1^{+\infty} g_0(R) J_r(R) R^2 dR, \quad (27)$$

with

$$J_r(R) = \frac{1}{\mu_0^r} \text{Tr} \boldsymbol{\mu}_{11}^{rr(2)}(\mathbf{r}). \quad (28)$$

In the limit $x \rightarrow \infty$, the first-order virial coefficients for a suspension of hard non-permeable spheres are recovered, and the Batchelor's values of the first-order virial coefficients, $\lambda(x) \rightarrow \lambda^{\text{hs}}$.

In the limit $x \rightarrow 0$, the scattering coefficients scale as $\sim x^2$, and as a result, $D_0 \sim 1/x^2$, and $\lambda \rightarrow 0$, with $\lambda_C \sim x^2$, and $\lambda_t, \lambda_r \sim x^4$.

III. RESULTS

To evaluate the first order virial coefficients $\lambda = \lambda_t, \lambda_K, \lambda_r$, we calculate two-particle mobility matrix elements, performing a series expansion in powers of $1/r$ up to the order 1000, as described in Sec. II. Integration with respect to the

TABLE I. First virial coefficients λ_t , λ_K , and λ_r for the short-time translational self-diffusion, sedimentation, and rotational self-diffusion, respectively. The limit $x \rightarrow \infty$ corresponds to hard non-permeable spheres.

x	λ_t	λ_K	λ_r
3	-0.2497	-3.4451	-0.03257
4	-0.4159	-4.1066	-0.06336
5	-0.5692	-4.5539	-0.09682
6	-0.7021	-4.8722	-0.12956
7	-0.8149	-5.1084	-0.16012
8	-0.9102	-5.2898	-0.18802
9	-0.9909	-5.4328	-0.21327
10	-1.0598	-5.5480	-0.23606
11	-1.1190	-5.6426	-0.25662
13	-1.2151	-5.7884	-0.29208
16	-1.3202	-5.9380	-0.33426
18	-1.3730	-6.0095	-0.35692
20	-1.4161	-6.0662	-0.37628
30	-1.5499	-6.2335	-0.44202
40	-1.6190	-6.3149	-0.48007
50	-1.6610	-6.3628	-0.50497
65	-1.7001	-6.4064	-0.52959
100	-1.7460	-6.4563	-0.56075
∞	-1.8315	-6.5465	-0.63102

particle positions has been performed analytically term by term, using the expressions given in Sec. II. The virial coefficients λ have been evaluated for a wide range of x . Selected results are listed in Table I. All displayed digits are significant (accuracy is discussed in Appendix C).

In Sec. VI, the dependence of the first-order virial coefficients λ on the permeability parameter $1/x$ will be discussed and plotted. Tables of values presented here can be used to construct approximate description of suspensions made of permeable particles with different internal structures.

IV. ANNULUS MODEL

The annulus model is another simplification, which can be applied to describe the dynamics of suspensions made of soft particles having a complex internal structure. The annulus model is alternative to the model of uniformly permeable particles.

In the annulus model,³¹ a particle suspended in a viscous fluid is characterized by two radii, $a_<$ and $a_>$. Its hydrodynamic interactions are governed by the smaller radius $a_<$. In addition, there exist also direct pair interactions. Two particles cannot come too close to each other, with the no-overlap radius equal to $a_>$. For such a model, the first-order virial expansion of transport coefficients has been performed with respect to the volume fraction $\phi_> = (4\pi/3)na_>^3$. The corresponding first-order virial coefficients $\lambda^A = \lambda_t^A, \lambda_K^A, \lambda_r^A$ have been evaluated as functions of ϵ , where

$$\epsilon = \frac{a_> - a_<}{a_<}. \quad (29)$$

The method used to determine λ^A is described in Appendix B. The calculated values are listed in Table II.

In Sec. VI, the dependence of the annulus first-order virial coefficients λ^A on the permeability parameter $1/x$ will be

TABLE II. First-order virial coefficients λ_t^A , λ_K^A , and λ_r^A for the short-time translational self-diffusion, sedimentation, and rotational self-diffusion, respectively, for a suspension of the annulus particles.

ϵ	λ_t^A	λ_K^A	λ_r^A
0.00	-1.8315	-6.5464	-0.63055
0.01	-1.7523	-6.4601	-0.56666
0.02	-1.6793	-6.3769	-0.51671
0.03	-1.6109	-6.2962	-0.47417
0.04	-1.5466	-6.2179	-0.43699
0.05	-1.4860	-6.1419	-0.40402
0.06	-1.4286	-6.0680	-0.37451
0.07	-1.3743	-5.9962	-0.34791
0.08	-1.3228	-5.9263	-0.32381
0.09	-1.2739	-5.8582	-0.30189
0.10	-1.2274	-5.7918	-0.28187
0.11	-1.1832	-5.7272	-0.26354
0.13	-1.1008	-5.6027	-0.23122
0.18	-0.9253	-5.3166	-0.16974
0.24	-0.7595	-5.0135	-0.12034
0.31	-0.6111	-4.7051	-0.08296
0.45	-0.4093	-4.1986	-0.04242
0.66	-0.2401	-3.6278	-0.01775

discussed, plotted, and compared with predictions of the model of uniformly porous particles. Tables of the annulus values presented here can be used to construct approximate description of suspensions made of permeable particles with different internal structures.

V. DISCUSSION

A. Comparison with the annulus model

Now we are going to compare the first-order virial coefficients, calculated in Sec. III for porous particles, with the corresponding results, obtained in Sec. IV for the annulus model, also called HRM. A similar comparison has been done in Ref. 19 for the effective viscosity. The key concept in this procedure is the hydrodynamic radius of a porous particle. For the translational diffusion (self-diffusion and sedimentation), the hydrodynamic radius a_{eff}^t is obtained from the single-particle translational diffusion coefficient, with the use of the relation,

$$D_0^t = \frac{k_B T}{6\pi\eta_0 a_{\text{eff}}^t}. \quad (30)$$

The dependence of a_{eff}^t on the porosity parameter x follows from Eqs. (7) and (9), which determine the translational diffusion coefficient of a single porous particle,^{24,25,32} and Eq. (A1), which specifies the scattering coefficient A_{10} . Explicitly,

$$a_{\text{eff}}^t(x) = a \frac{2x^2(x - \tanh(x))}{2x^3 + 3(x - \tanh(x))}. \quad (31)$$

For the rotational self-diffusion, the hydrodynamic radius a_{eff}^r is obtained from the rotational diffusion coefficient of a single porous particle,^{32,33} with the use of the relation,

$$D_0^r = \frac{k_B T}{8\pi\eta_0 (a_{\text{eff}}^r)^3}. \quad (32)$$




FIG. 1. The annulus (or hydrodynamic radius) model of a porous particle.

The dependence of a_{eff}^r on the porosity parameter x follows from Eqs. (24) and (26), with the scattering coefficient A_{11} given by Eq. (A2). As a result,

$$a_{\text{eff}}^r(x) = a \left[1 + \frac{3}{x^2} - \frac{3 \coth(x)}{x} \right]^{1/3}. \quad (33)$$

For large x ,

$$a_{\text{eff}}^r(x) = a_{\text{eff}}^r(x) + \mathcal{O}(1/x^2). \quad (34)$$

A porous particle of radius a and the porosity parameter x is modeled as an annulus particle, see Fig. 1, by matching its geometrical radius a to the annulus no-overlap radius $a_{>} = a$. The smaller annulus radius $a_{<} = a_{\text{eff}}$ is determined by the effective hydrodynamic radius a_{eff} , given in Eqs. (31)–(33).

In this way, the annulus parameter ϵ , defined in Eq. (29), becomes the following function of x :

$$\epsilon(x) = \frac{a - a_{\text{eff}}(x)}{a_{\text{eff}}(x)}, \quad (35)$$

with $a_{\text{eff}}(x)$ determined by Eqs. (31) and (33).

In Figs. 2–4, the annulus coefficients $\lambda^A(\epsilon)$, with $\epsilon(x)$ specified by Eq. (35) (dashed lines), are plotted together with the porous-particle virial coefficients $\lambda(x)$ (solid lines).

As shown in Fig. 2, for a low permeability, the coefficient $\lambda_K \approx \lambda_K^{\text{hs}} + 10/x$ is approximately a linear function of $1/x$. The coefficients λ_t and λ_r as functions of $1/x$ are shown in Figs. 3 and 4, respectively. In the plots, $1/x \leq 0.1$, i.e., for $x \geq 10$. For




FIG. 2. (Color online) Two-particle sedimentation virial coefficient $\lambda_K(x)$. Our precise results for porous particles (solid line) are well-approximated by the annulus model (dashed line).




FIG. 3. (Color online) Two-particle translational self-diffusion virial coefficient $\lambda_t(x)$. Our precise results for porous particles (solid line) are well-approximated by the annulus model (dashed line).

$1/x = 0$, i.e., for $x = \infty$, the limit of a non-permeable hard sphere with radius a is recovered, $\lambda^{\text{hs}} = \lambda(\infty)$.

The annulus model slightly underestimates the accurate values of the first-order virial coefficients for suspensions of uniformly permeable particles, and this very small difference slightly increases when x decreases (when the particles become more permeable).

For the sedimentation coefficient, the annulus model is accurate, with a half-percent relative accuracy already at $x = 20$ and a reasonable 3% precision at $x = 5$. For the translational self-diffusion, the annulus model is less accurate, but still it gives only a 2% error for $x = 20$, a 5% error for $x = 10$, and a 7% error for $x = 5$. The least accurate is the annulus prediction for the rotational self-diffusion, with a 5% error for $x = 20$ and a 11% error for $x = 10$.

Summarizing, the annulus model (HRM) approximates well with the first virial coefficients of porous particles suspensions, in the range of intermediate and small particle permeability (i.e., for moderate and large values of x).




FIG. 4. (Color online) Two-particle rotational self-diffusion virial coefficient $\lambda_r(x)$. Our precise results for porous particles (solid line) are well-approximated by the annulus model (dashed line).

B. Comparison with other results for permeable particles suspensions

In this section, we compare our first-order virial approximation for diffusion and sedimentation of porous particles suspensions, given by Eqs. (11), (14), (20), and (25), with the corresponding values evaluated by other methods, and with the accurate data obtained from numerical simulations of the same system, what allows to estimate the range of validity of the first-order virial approximation.

In Fig. 5, our first-order virial expansion of the short-time translational self-diffusion coefficient of a porous-particle suspension (dashed lines) is compared with the accurate simulation results performed in Ref. 17 for a porous-particle suspension in the whole range of volume fractions $\phi \leq 0.45$ (solid lines, which interpolate the simulation points indicated by symbols). The first order virial expansion, see Eq. (11), can be used as an accurate approximation of D_t in a very wide range of volume fractions, even for relatively large values of x (i.e., low permeabilities).

The small difference between the simulations and the first virial expansion has a tendency to increase when the volume fraction is increased. The reason is that in this case, the two-particle hydrodynamic interactions, taken into account in the first-order virial expansion, are not sufficient to describe very accurately the whole system, in which the N -particle hydrodynamic interactions, with $N \geq 3$, become more and more important.

Also, it is intuitive that at larger volume fractions, the first-order virial approximation is closer to the accurate value, when x is smaller, i.e., for a larger permeability—in this case the effect of many-body hydrodynamic interactions is smaller.

In contrast to the translational self-diffusion, values of the sedimentation coefficient K differ significantly from the first-order virial estimation already at rather small volume fractions, see Ref. 17. Our values of $\lambda_K(\infty)$ reproduce with a higher accuracy, the classic Batchelor's result,²⁹ for non-permeable hard spheres. For uniformly porous particles, the




FIG. 5. (Color online) Translational self-diffusion coefficient D_t for a suspension of porous particles. Symbols connected by splines (solid lines) represent accurate simulation results from Ref. 17. Dashed straight lines represent first-order virial expansion calculated in this work.

boundary collocation method was applied by Chen and Cai³⁰ to evaluate $\lambda_K = -3.46, -5.50, -6.23, -6.44$ for $x^2 = 10^\alpha$, with $\alpha = 1, 2, 3, 4$, respectively. Comparing their results with our very accurate values, $\lambda_K = -3.5723, -5.5480, -6.2504, -6.4563$, we conclude that the uncertainty of their results is decreasing from 3% at $\alpha = 1$ to 0.3% at $\alpha = 4$. The accuracy of the boundary collocation method is worse at smaller values of x , i.e., for larger permeabilities.

VI. CONCLUSIONS

In this paper, the short-time diffusion properties of dilute suspensions of uniformly porous spherical particles have been investigated. The first-order virial coefficients λ of the diffusion and sedimentation coefficients have been evaluated as functions of the permeability parameter x , equal to the ratio of the particle radius to the hydrodynamic screening length. In the limit of $x \rightarrow \infty$, the hard non-permeable limit is recovered, with the Batchelor's values of the first-order virial coefficients.

It is worthwhile to point out that values of the first-order virial coefficients calculated in this work are essential also for concentrated suspensions of uniformly porous particles. As it has been recently shown in Ref. 34, the first-order virial coefficients provide such a significant information about suspension of permeable particles, that they are sufficient to determine very accurately the translational and rotational self-diffusion in the whole range of volume fractions up to $\phi = 0.45$, based on the known results for a non-permeable hard-sphere suspension of the same volume fraction. A simple and accurate (but not straightforward) scaling has been found and described in detail in Ref. 34. Therefore, the first-order virial coefficients evaluated here are of fundamental importance for the self-diffusion in concentrated systems of permeable particles.

In this work, it has been checked that values of the first-order virial coefficients λ are well-approximated by the annulus (hydrodynamic radius) model, if the parameter x is sufficiently large, i.e., the permeability is sufficiently low. Systematically, the annulus approximation slightly underestimates the virial coefficients of porous particle suspensions. For rotational diffusion, a reasonable 5% accuracy of this model is reached at $x \gtrsim 20$. For translational diffusion (collective and self), the comparable or even better 3%–5% precision is obtained already for $x \gtrsim 10$. For the sedimentation coefficient, the accuracy is even higher (a 3% precision already at $x = 5$), owing to much larger absolute values of $\lambda_K = 8 + \lambda_C$.

The annulus model is expected to work well also for suspensions of uniformly porous particles at larger volume fractions, if the permeability parameter x is sufficiently large. The accuracy of this approximation at larger volume fractions will be investigated in a separate publication.

The annulus model is an approximate model of real suspensions of particles, which can exhibit a complex non-uniform internal structure. The advantage of the annulus model is its simplicity—there is only one parameter, i.e., the inner (hydrodynamic) radius, which can be easily determined from experiments with individual particles. Here we provide

tables of values for the virial coefficients following from this model, and we check that the model is reasonably accurate for the special case of a suspension made of uniformly permeable (porous) particles. The tables of the annulus virial coefficients presented in this work can be used also in a wide range of other contexts, to estimate diffusion and shear viscosity coefficients of suspensions made of particles characterized by a different internal structure. The core-shell is an example. Comparison with experiments, polydispersity, non-uniform porosity, core-shell particles, and relation to practical aspects are the subjects of future studies.

ACKNOWLEDGMENTS

M.L.E.J. and E.W. were supported in part by the Polish Ministry of Science and Higher Education Grant N N501 156538. G.N. thanks the Deutsche Forschungsgemeinschaft (SFB-TR6, project B2) for financial support.

APPENDIX A: SCATTERING COEFFICIENTS

The scattering coefficients for a uniformly permeable sphere with $x = \kappa a$, where a is the sphere radius and $1/\kappa$ is the hydrodynamic penetration depth, have the form,³²

$$A_{l0} = \frac{(2l+1)g_l(x)}{2g_{l-2}(x)} \left[1 + \frac{l(2l-1)(2l+1)g_l(x)}{(l+1)x^2 g_{l-2}(x)} \right]^{-1} a^{2l-1}, \quad (\text{A1})$$

$$A_{l1} = \frac{g_{l+1}(x)}{g_{l-1}(x)} a^{2l+1}, \quad (\text{A2})$$

$$A_{l2} = \left[\frac{2l+3}{2l-1} + \frac{2(2l+1)(2l+3)}{(l+1)x^2} \right] a^2 A_{l0} - \frac{2l+3}{2l-1} a^{2l+1}, \quad (\text{A3})$$

$$B_{l2} = \left[1 + \frac{2(2l-1)(2l+1)}{(l+1)x^2} \right] a^2 A_{l2} - a^{2l+3}, \quad (\text{A4})$$

where $l = 1, 2, \dots$ and $g_l(x) = \sqrt{\pi/2} x^{l+1/2}(x)$ is the modified spherical Bessel function of the first kind.

APPENDIX B: THE ANNULUS (HYDRODYNAMIC RADIUS) MODEL

For a suspension of particles described by the annulus (hydrodynamic radius) model, the virial coefficients λ^A are functions of the model parameter ϵ , defined by Eq. (29) and listed in Table II. In this Appendix, we explain how these values have been evaluated.

The first-order virial expansion of D_t/D_0^t , D_C/D_0^t , K , and D_r/D_0^r , can be performed with the use of $\phi_< = (4\pi/3)na_<^3$, or $\phi_>$ defined by the analogical expression,

$$1 + \lambda^A(\epsilon)\phi_> + \mathcal{O}(\phi_>^2) = 1 + \bar{\lambda}^A(\epsilon)\phi_< + \mathcal{O}(\phi_<^2). \quad (\text{B1})$$

Therefore,

$$\lambda^A(\epsilon) = \frac{\bar{\lambda}^A(\epsilon)}{(1+\epsilon)^3}. \quad (\text{B2})$$

To evaluate $\bar{\lambda}^A(\epsilon)$, we now introduce the dimensionless inter-particle distance as $R = r/2a_<$, and we replace the pair distribution function from Eq. (10) by the following expression:

$$g_0(R) = \begin{cases} 0 & \text{for } R \leq 1 + \epsilon, \\ 1 & \text{for } R > 1 + \epsilon. \end{cases} \quad (\text{B3})$$

which corresponds to the no-overlap condition at a larger radius $a_>$. Then, we apply the Eqs. (12), (16), and (27), taken in the non-permeable hard-sphere limit, $x = \infty$. We obtain the following expressions:

$$\begin{aligned} \bar{\lambda}_t^A(\epsilon) &= 8 \int_{1+\epsilon}^{+\infty} J_l(R) R^2 dR \\ &= \lambda_t^{\text{hs}} - 8 \int_1^{1+\epsilon} J_l(R) R^2 dR, \end{aligned} \quad (\text{B4})$$

$$\begin{aligned} \bar{\lambda}_K^A(\epsilon) &= \frac{2}{5a_<^3} A_{12}^{\text{hs}} - \frac{4}{a_<} A_{10}^{\text{hs}} (1+\epsilon)^2 \\ &\quad + 8 \int_{1+\epsilon}^{+\infty} J_K(R) R^2 dR \\ &= \lambda_K^{\text{hs}} - 8 \int_1^{1+\epsilon} \bar{J}_K(R) R^2 dR, \end{aligned} \quad (\text{B5})$$

$$\begin{aligned} \bar{\lambda}_r^A(\epsilon) &= 8 \int_{1+\epsilon}^{+\infty} J_r(R) R^2 dR \\ &= \lambda_r^{\text{hs}} - 8 \int_1^{1+\epsilon} J_r(R) R^2 dR, \end{aligned} \quad (\text{B6})$$

where

$$\bar{J}_K(R) = \frac{1}{\mu_0^t} \text{Tr}[\mu_{11}^{t(2)}(\mathbf{R}) + \mu_{12}^{t(2)}(\mathbf{R})]. \quad (\text{B7})$$

In this Appendix, all the mobility coefficients, the associated functions J , and the superscript ^{hs}, refer to the hard spheres with the stick boundary conditions.

The above formulas have been applied to compute the functions $\lambda_t^A(\epsilon)$, $\lambda_K^A(\epsilon)$, and $\lambda_r^A(\epsilon)$, listed in Table II. The functions $\bar{\lambda}_t^A(\epsilon)$, $\bar{\lambda}_K^A(\epsilon)$, and $\bar{\lambda}_r^A(\epsilon)$ will be also listed in our future publication, where they are applied to model diffusion and viscosity of particles with a hard solid core and a thin porous shell.

APPENDIX C: ACCURACY OF THE VIRIAL COEFFICIENTS

The virial coefficients λ for uniformly permeable particles are calculated as a series expansion in powers $1/r^k$, with $k = 1, \dots, 1000$. Accuracy of this expansion has been estimated by calculating $\lambda(n)$ as the sum of all the terms $k = 1, \dots, n$. Then, the relative error of this sum has been evaluated,

$$\Delta\lambda(n) = \left| \frac{\lambda(n) - \lambda(1000)}{\lambda(1000)} \right|. \quad (\text{C1})$$

The relative error $\Delta\lambda(n)$ is plotted in Fig. 6 as a function of n . Here, we took the largest value of $x = 100$, since the larger value of x , the slower is the convergence rate. For a smaller x , the error is smaller.




FIG. 6. (Color online) Estimation of the relative uncertainty $\Delta\lambda(n)$ of the first-order (two-particle) virial coefficients as a function of the truncation parameter n of the power expansion in the inverse interparticle distance. $\Delta\lambda$ is defined in Eq. (C1), and t , r , and K refer to the translational and rotational self-diffusion and sedimentation, respectively.

Concluding, the accuracy of the first-order virial coefficient for suspensions of permeable particles is very high, exceeding significantly precision obtained usually in experiments.

- ¹B. Cichocki, M. L. Ekiel-Jeżewska, and E. Wajnryb, "Lubrication corrections for three-particle contribution to short-time self-diffusion coefficients in colloidal dispersions," *J. Chem. Phys.* **111**, 3265 (1999).
- ²B. Cichocki, M. L. Ekiel-Jeżewska, P. Szymczak, and E. Wajnryb, "Three-particle contribution to sedimentation and collective diffusion in hard-sphere suspensions," *J. Chem. Phys.* **117**, 1231 (2002).
- ³B. Cichocki, M. L. Ekiel-Jeżewska, and E. Wajnryb, "Three-particle contribution to effective viscosity of hard-sphere suspensions," *J. Chem. Phys.* **119**, 606 (2003).
- ⁴A. J. Banchio and G. Nägele, "Short-time transport properties in dense suspensions: From neutral to charge-stabilized colloidal spheres," *J. Chem. Phys.* **128**, 104903 (2008).
- ⁵C. N. Likos, S. Rosenfeldt, N. Dingenouts, M. Ballauff, P. Lindner, N. Werner, and F. Vögtle, "Gaussian effective interaction between flexible dendrimers of fourth generation: A theoretical and experimental study," *J. Chem. Phys.* **117**, 1869 (2002).
- ⁶S. Huißmann, A. Wynveen, C. N. Likos, and R. Blaak, "The effects of pH, salt and bond stiffness on charged dendrimers," *J. Phys. Condens. Matter* **22**, 232101 (2010).
- ⁷S. Pyett and W. Richtering, "Structures and dynamics of thermosensitive microgel suspensions studied with three-dimensional cross-correlated light scattering," *J. Chem. Phys.* **122**, 034709 (2005).
- ⁸T. Eckert and W. Richtering, "Thermodynamic and hydrodynamic interaction in concentrated microgel suspensions: Hard or soft sphere behavior?" *J. Chem. Phys.* **129**, 124902 (2008).
- ⁹C. A. Coutinho, R. K. Harrinath, and V. K. Gupta, "Settling characteristics of composites of PNIPAM microgels and TiO₂ nanoparticles," *Colloids Surf., A* **318**, 111 (2008).
- ¹⁰M. H. G. Duits, P. A. Nommensen, D. van den Ende, and J. Mellema, "High frequency elastic modulus of hairy particle dispersions in relation to their microstructure," *Colloids Surf., A* **183–185**, 335 (2001).

- ¹¹G. Petekidis, J. Gapiński, P. Seymour, J. S. van Duijneveldt, D. Vlassopoulos, and G. Fytas, "Dynamics of core-shell particles in concentrated suspensions," *Phys. Rev. E* **69**, 042401 (2004).
- ¹²M. Zackrisson, A. Stradner, P. Schurtenberger, and J. Bergenholtz, "Small-angle neutron scattering on a coreshell colloidal system: A contrast-variation study," *Langmuir* **21**, 10835 (2005).
- ¹³Z. Adamczyk, B. Jachimska, and M. Kolasnińska, "Structure of colloid silica determined by viscosity measurements," *J. Colloid Interface Sci.* **273**, 668 (2004).
- ¹⁴J. Masliyah, G. Neale, K. Malysa, and T. van de Ven, "Creeping flow over a composite sphere: Solid core with porous shell," *Chem. Eng. Sci.* **42**, 245 (1987).
- ¹⁵W. van Saarloos, "On the hydrodynamic radius of fractal aggregates," *Physica A* **147**, 280 (1987).
- ¹⁶C. N. Likos, H. Löwen, M. Watzlawek, B. Abbas, O. Jucknischke, J. Allgaier, and D. Richter, "Star polymers viewed as ultrasoft colloidal particles," *Phys. Rev. Lett.* **80**, 4450 (1998).
- ¹⁷G. C. Abade, B. Cichocki, M. L. Ekiel-Jeżewska, G. Nägele, and E. Wajnryb, "Short-time dynamics of permeable particles in concentrated suspensions," *J. Chem. Phys.* **132**, 014503 (2010).
- ¹⁸G. C. Abade, B. Cichocki, M. L. Ekiel-Jeżewska, G. Nägele, and E. Wajnryb, "Dynamics of permeable particles in concentrated suspensions," *Phys. Rev. E* **81**, 020404(R) (2010).
- ¹⁹G. C. Abade, B. Cichocki, M. L. Ekiel-Jeżewska, G. Nägele, and E. Wajnryb, "High-frequency viscosity of concentrated porous particles suspensions," *J. Chem. Phys.* **133**, 084906 (2010).
- ²⁰G. C. Abade, B. Cichocki, M. L. Ekiel-Jeżewska, G. Nägele, and E. Wajnryb, "High-frequency viscosity and generalized Stokes-Einstein relations in dense suspensions of porous particles," *J. Phys. Condens. Matter* **22**, 322101 (2010).
- ²¹B. Cichocki, B. U. Felderhof, K. Hinsen, E. Wajnryb, and J. Bławdziewicz, "Friction and mobility of many spheres in Stokes flow," *J. Chem. Phys.* **100**, 3780 (1994).
- ²²S. Kim and S. J. Karrila, *Microhydrodynamics. Principles and Selected Applications* (Butterworth-Heinemann, London, 1991).
- ²³J. Happel and H. Brenner, *Low Reynolds Number Hydrodynamics* (Martinus Nijhoff, Dordrecht, 1986).
- ²⁴H. C. Brinkman, "A calculation of the viscous force exerted by a flowing fluid on a dense swarm of particles," *Appl. Sci. Res. Sect. A* **1**, 27 (1947).
- ²⁵P. Debye and A. M. Bueche, "Intrinsic viscosity, diffusion, and sedimentation rate of polymers in solution," *J. Chem. Phys.* **16**, 573 (1948).
- ²⁶B. Cichocki, B. U. Felderhof, and R. Schmitz, "Hydrodynamic interactions between two spherical particles," *PhysicoChem. Hydrodyn.* **10**, 383 (1988).
- ²⁷J. P. Hansen and I. R. McDonald, *Theory of Simple Liquids* (Academic, London, 1986).
- ²⁸B. Cichocki and B. U. Felderhof, "Short-time diffusion coefficients and high frequency viscosity of dilute suspensions of spherical Brownian particles," *J. Chem. Phys.* **89**, 1049 (1988).
- ²⁹G. K. Batchelor, "Sedimentation in a dilute dispersion of spheres," *J. Fluid Mech.* **52**, 245 (1972).
- ³⁰S. B. Chen and A. Cai, "Hydrodynamic interactions and mean settling velocity of porous particles in a dilute suspension," *J. Colloid Interface Sci.* **217**, 328 (1999).
- ³¹B. Cichocki and B. U. Felderhof, "Diffusion of Brownian particles with hydrodynamic interaction and hard core repulsion," *J. Chem. Phys.* **94**, 556 (1991).
- ³²R. Reuland, B. U. Felderhof, and R. B. Jones, "Hydrodynamic interaction of two spherically symmetric polymers," *Physica A* **93**, 465 (1978).
- ³³B. U. Felderhof and J. M. Deutch, "Frictional properties of dilute polymer solutions. I. Rotational friction coefficient," *J. Chem. Phys.* **62**, 2391 (1975).
- ³⁴G. Abade, B. Cichocki, M. L. Ekiel-Jeżewska, G. Nägele, and E. Wajnryb, "Rotational and translational self-diffusion in concentrated suspensions of permeable particles," *J. Chem. Phys.* **134**, 244903 (2011).