

1 **Using animal-mounted sensor technology and machine learning to predict**
2 **time-to-calving in beef and dairy cows**

3 G.A. Miller¹, M. Mitchell², Z.E. Barker³, K. Giebel³, E.A. Codling⁴, J.R. Amory³, C.
4 Michie⁵, C. Davidson⁵, C. Tachtatzis⁵, I. Andonovic⁵ and C.-A. Duthie¹

5 ¹ *Future Farming Systems, SRUC, Peter Wilson Building, West Mains Road, King's*
6 *Buildings, Edinburgh, EH9 3JG, UK*

7 ² *Animal and Veterinary Science, SRUC, Peter Wilson Building, West Mains Road,*
8 *King's Buildings, Edinburgh, EH9 3JG, UK*

9 ³ *Writtle University College, Lordship Road, Writtle, Chelmsford, CM1 3RR, UK*

10 ⁴ *Department of Mathematical Sciences, University of Essex, Wivenhoe Park,*
11 *Colchester, CO4 3SQ, UK*

12 ⁵ *University of Strathclyde, Glasgow, Royal College Building, 204 George Street,*
13 *Glasgow, G1 1XW, UK*

14 Corresponding author: Dr Gemma Miller. Email: gemma.miller@sruc.ac.uk

15 Running head: Predict calving with sensors and machine learning

16 **Abstract**

17 Worldwide, there is a trend towards increased herd sizes and the animal to
18 stockman ratio is increasing within the beef and dairy sectors, thus the time available
19 to monitoring individual animals is reducing. The behaviour of cows is known to
20 change in the hours prior to parturition, e.g. less time ruminating and eating, and
21 increased activity level and tail raise events. These behaviours can be monitored
22 non-invasively using animal mounted sensors. Thus behavioural traits are ideal
23 variables for the prediction of calving. This study explored the potential of two sensor
24 technologies for their capabilities in predicting when calf expulsion should be
25 expected. Two trials were conducted at separate locations: i) beef cows (n = 144)
26 and (ii) dairy cows (n = 110). Two sensors were deployed on each cow: 1) Afimilk
27 Silent Herdsman (SHM) collars monitoring time spent ruminating (RUM), eating
28 (EAT) and the relative activity level (ACT) of the cow and 2) tail mounted Axivity
29 accelerometers to detect tail-raise events (TAIL). The exact time the calf was
30 expelled from the cow was determined by viewing closed-circuit television camera
31 footage. Machine learning random forest (RF) algorithms were developed to predict
32 the when calf expulsion should be expected using single sensor variables and by
33 integrating multiple sensor data-streams. The performance of the models were
34 tested by the Matthew's Correlation Coefficient (MCC), the area under the curve
35 (AUC) and the sensitivity (Se) and specificity (Sp) of predictions. The TAIL model
36 was slightly better at predicting calving within a five hour window for beef cows (MCC
37 = 0.31) than for dairy cows (MCC = 0.29). The TAIL+RUM+EAT models were equally
38 as good at predicting calving within a five hour window for beef and dairy cows (MCC
39 = 0.32 for both models). Combining data-streams from SHM and tail sensors did not
40 substantially improve model performance over tail sensors alone therefore hour-by-

41 hour algorithms for the prediction of the time of calf expulsion were developed using
42 tail sensor data. Optimal classification occurred at two hours prior to calving for both
43 beef (MCC = 0.29) and dairy cows (MCC = 0.25). This study has shown that tail
44 sensors alone are adequate for the prediction of parturition and that the optimal time
45 for prediction is two hours before expulsion of the calf.

46

47 Keywords: precision livestock farming, parturition, bovine, machine learning, sensors

48

49 Implications: The availability of alerts to when beef and dairy cows are expected to
50 deliver a calf will enable farmers to more effectively manage their time and to
51 intervene in a timely manner where necessary,, thus optimising the economic and
52 production efficiency of their business.

53 **Introduction**

54 There is a global trend towards increased herd sizes. For instance, in the UK, the
55 average dairy herd size has increased 2.7% since 2014 and the average beef herd
56 size by 1.2% (AHDB, 2018). If available labour does not increase in line with herd
57 size this can result in the cow to stockman ratio increasing and less time available for
58 monitoring of individual animal. In order to optimise the production efficiency of the
59 UK livestock sector there is a requirement for the development and use of cost-
60 effective animal monitoring solutions to inform on the health and productive status of
61 individual animals.

62 Dystocia is a considerable problem within beef and dairy systems. Internationally,
63 the prevalence of dystocia in dairy cows typically varies between 2 and 7% of
64 calvings, but is as high as 14% in the USA (Mee, 2008). In the UK, 6.9% of dairy
65 heifers experience serious difficulties during calving (Raumph and Faust, 2006).
66 Reports of assisted calvings range from 10 – 50% (Mee, 2008), with primiparous
67 cows more commonly experiencing difficulties (Lombard *et al*, 2007). In the beef
68 sector, between 1 and 8% of cows experience difficult calvings, require surgical
69 intervention or have stillbirths (Nix *et al* 1998; Phocas and Laloë, 2003; Eriksson *et*
70 *al*, 2004; De Amicis *et al*, 2018).

71 The costs associated with mild and severe cases of dystocia in the dairy sector are
72 estimated at between £110 and £400 due to milk loss (McGuirk *et al*, 2007).

73 Dystocia can lead to increased days open, increased numbers of services,
74 premature culling and poor calf health, performance and survival (McGuirk *et al*,
75 2007; López de Maturana *et al*, 2007; Lombard *et al*, 2007; Gaafar *et al*, 2011;
76 Barrier *et al*, 2013). Thus the development of methods to automatically predict the

77 onset of parturition and identify problematic calvings is important to facilitate timely
78 and appropriate interventions to prevent the losses associated with dystocia.

79 A number of physiological and behavioural changes occur around calving which offer
80 opportunities to predict the onset of parturition. Characterisation of maternal
81 hormonal profiles is able to predict calving times with limited accuracy (Shah *et al*,
82 2006) and the process is invasive and retrospective. Reductions in body temperature
83 occur on the day of calving and can be used to predict parturition onset within a 24
84 hour window, but variations in temperature change between individual animals limit
85 the predictive power of temperature alone (Saint-Dizier and Chastant-Maillard,
86 2015). Behavioural indicators, such as lying and standing, eating and rumination
87 (Kovács, *et al*, 2016) patterns, social behaviour and tail raising events are known to
88 change in the 24 hours prior to calving (Huzzey *et al*, 2005; Miedema *et al*, 2011a,b;
89 Jensen, 2012). Advances in animal mounted sensors capable of monitoring these
90 behaviours provides the opportunity to develop an automated system for prediction
91 of parturition.

92 The present study utilised two non-invasive animal mounted sensors: a near to
93 market tail mounted sensor to monitor tail raising behaviour, and an on the market
94 neck mounted sensor to monitor eating and rumination behaviour as well as a
95 relative level of activity. The objectives were to determine if variables recorded using
96 existing technologies could be used to develop algorithms to predict when calf
97 expulsion should be expected to occur, and if combining sensors could improve the
98 prediction. The hypothesis was that variables reported from existing technologies
99 could be used to develop algorithms to predict time to calf expulsion in both beef and
100 dairy cows.

101 **Methods**

102 *Ethics statement*

103 The animal trials described below were approved by the Animal Experiment
104 Committee of SRUC and were conducted in accordance with the requirements of the
105 UK Animals (Scientific Procedures) Act 1986.

106 *Animals*

107 Two studies were conducted, one with beef cows at the Beef and Sheep Research
108 Centre at Scotland's Rural College (SRUC), UK, and one at a commercial dairy farm
109 in Essex, UK. In the beef trial, a total of 144 pregnant spring-calving cows which
110 calved between March and June 2017 were monitored. The animals were a mixture
111 of breeds (51 Limousin sired; 59 Aberdeen Angus sired, 34 Luing), with 78, 54 and
112 12 calving to the first, second and third artificial insemination (AI) respectively. At the
113 beginning of the trial the average liveweight was 662 ± 91 kg and the average body
114 condition score was 2.8 ± 0.3 (using the system described in Lowman *et al*, 1976).
115 Cows ranged in age from 2-16 years and parity number from 0-13. Cows were
116 allocated to one of two group-housed straw-bedded pens prior to calving (Pen 1:
117 32m x 6.4m housing up to 24 cattle; Pen 2: 27.4m x 6.4m housing up to 20 cattle).
118 Animals entered the study based on anticipated date of calving, with those calving to
119 the first AI entering the trial first. Throughout the study, all beef cows were fed a total
120 mixed ration comprising of (per head/day on a fresh weight basis) whole crop barley
121 silage (27.7%), grass silage (41.0%), barley straw (25.6%), maize dark grains (5.1%)
122 and minerals (0.6%).

123 In the dairy trial, a total of 110 Holstein Friesian dairy cows which calved between
124 July and October 2017 were monitored. Cows ranged in age from 1-10 years and

125 parity ranged from 0-6. All dairy cows were served using AI and estimated calving
126 dates were available from the Cattle Information Service records. Cows were housed
127 in a 41 cubicle dry-cow shed (30m x 12m) from 14 or more days pre-calving, where
128 they remained loose housed until showing signs of calving (determined visually by
129 the farm staff). At which point they were moved to a smaller (6m x 10m) loose straw
130 bedded yard for calving and until approximately 24 hours post calving. Cows were
131 fed a dietary cation-anion balanced total mixed ration which was delivered once a
132 day at approximately 9am. To allow scraping and bedding up cows were removed
133 from the cubicle house once a day and held in the adjacent collecting yard (10-
134 11am).

135 *Experimental design and sensors*

136 All cows in both studies were fitted with two sensors, and data collection was started
137 immediately:

138 1. Silent Herdsman (SHM) collars (Afimilk Ltd., Israel), neck mounted
139 accelerometers originally designed to detect oestrus based on cow activity,
140 rumination and eating patterns. Data from the collars was downloaded to a base
141 station in real time and classified into behaviours by proprietary algorithms (hourly
142 eating and rumination and relative activity per 1.5 hours).

143 2. Tail mounted tri-axial accelerometers (TTA) (AX3 3-Axis logging accelerometer,
144 Axivity, Newcastle upon Tyne, UK) measuring acceleration at a frequency of 12.5
145 Hz. The TTAs have an internal battery which is rechargeable. Data is downloaded
146 manually to a computer in comma separated values format. Previous work from
147 SRUC and the University of Edinburgh has characterised tail-raise signatures and
148 demonstrated that this information may be important to predict time-to-calving during

149 the immediate pre-calving period. The TTAs were housed in synthetic pouches and
150 mounted on cow tails using hook and loop straps (Figure 1). The angle of the tail at
151 any point in time can be determined by calculating the pitch of the TTA (Figure 1). An
152 approximation to this is obtained from the magnitude of the gravitational acceleration
153 measured on the x-axis of the TTA:

$$154 \quad Acc_x = g \sin(\theta)$$

155 where θ is the angle of the TTA orientation with respect to gravity (Figure 1). Using
156 this approach, the orientation of the TTA was determined for a period of 10 minutes
157 following attachment, thereafter deviations of more than 20° from this position were
158 deemed to be when the tail was in a raised position.

159 Continuous 24 hour video data was collected for the duration of the calving period.
160 Twenty five cameras were mounted above the beef calving pens and footage
161 recorded continuously using GeoVision software (EZCCTV, Letchworth, UK). In the
162 dairy study 2 closed-circuit TV cameras were installed at positions which ensured
163 that there was full coverage of the calving pen. Closed-circuit TV videos were
164 manually reviewed to ascertain the exact time of calf expulsion (calf completely
165 expelled from the cow) for each cow.

166 *Data Analysis*

167 The SHM collars use proprietary algorithms to convert raw accelerometer data into
168 minutes per hour spent eating (EAT), minutes per hour spent ruminating (RUM) and
169 a relative numeric level of activity per 1.5 hours (ACT). Raw TTA data was
170 expressed as minutes per hour with the tail in a raised position (TAIL).

171 For the development of the prediction models, sensor variables (TAIL, RUM, EAT
172 and ACT) were combined with non-sensor variables. The non-sensor variables used
173 in the beef models were as follows: time of day, parity, breed, weight at beginning of
174 trial (kg), body condition score at beginning of trial, age (years) and AI status
175 (conceived on the first, second or third AI). For dairy cows the variables were: time of
176 day, parity (multiparous or primiparous), number of lactations and age.

177 The hour in which a calf was completely expelled from the cow was deemed 'hour 0'
178 for that cow and all previous data points were assigned a value according to number
179 of hours relative to hour 0. For each sensor variable, only animals which had at least
180 the 48 hours prior to calf expulsion recorded were included, and all data up to 196
181 hours (one week) was considered.

182 The data from individual sensor variables were plotted to visually assess changes in
183 behaviour in the week prior to calving. The five hours prior to calving was statistically
184 compared to a control period which was the corresponding five hour period 24 hours
185 before using a Wilcoxon signed-rank test. The data was then randomly divided into
186 training and validation data sub-sets (70:30) with no animal allowed to be in both the
187 training and validation sub-sets.

188 Random forest (RF) models were developed to predict when an animal was within 5
189 hours of calving using single variables and then combined variables. Random forest
190 classifiers are ensemble machine learning algorithms which are considered to be
191 more accurate than single classifiers, and more robust to noise (Agjee *et al*, 2018).

192 Ensemble algorithms construct a set of independent classifier models (decision
193 trees), with each model having a 'vote' on how to classify each new data point. RFs
194 were developed for each individual sensor variable (TAIL, RUM, EAT and ACT), and

195 then for multiple sensor variables, and finally - for the best model - hourly time points
196 leading up to calving. The algorithm creates i bootstrapped samples from the training
197 data sub-set, where i is the number of independent decision trees (ntree). A decision
198 tree is then fitted to each bootstrap sample. To overcome the unbalanced nature of
199 the data (fewer target time points than non-target) the bootstrapping, resampling
200 during parameter tuning and model evaluation were down sampled i.e. if there were
201 100 time points of interest then only 100 other data points were included. Each tree
202 was then tested with the out-of-bag (oob) data points. At each branch in each
203 decision tree, only a random subset of variables are considered (mtry), this
204 parameter and ntree were optimised during tuning of the algorithm. All possible
205 values of mtry were tested and ntree was increased (by 500 trees) until increasing
206 the number of trees further no longer reduced the model error (i.e. the oob error
207 stabilised).

208 The binary class variable 'calving' and the model predictions (class probabilities)
209 were used to create Receiver Operator Characteristic (ROC) curves and to estimate
210 the area under the ROC curve (AUC). Based on the ROC curves, a threshold for the
211 probability that a cow was within 5 hours of calving was chosen that resembled the
212 optimum balance between sensitivity (true positives divided by true positives plus
213 false negatives) and specificity (true negatives divided by true negatives plus the
214 false positives). The Matthew's Correlation Coefficient (MCC) was also calculated.
215 The MCC is a metric which assesses the performance of a binary classifier and is
216 less sensitive to imbalanced data sets (such as the test sub-sets in this case) and is
217 calculated using the following equation:

$$218 \quad MCC = \frac{TP \times TN - FP \times FN}{\sqrt{(TP + FP)(TP + FN)(TN + FP)(TN + FN)}}$$

219 Where TP = true positive, TN = true negative, FP = false positive and FN = false
220 negative. These values were derived from the optimum model identified by the ROC
221 curve. MCC values are between -1 and +1, with +1 being a perfect classifier, 0 being
222 no better than random and -1 being completely inversed classification.

223 All data analyses were undertaken in R (version 3.4.1, R core team, 2017) using the
224 dplyr (Wickham *et al*, 2018), caret (Kuhn, 2018) and pROC (Robin *et al*, 2011)
225 packages.

226 **Results**

227 *Data inclusion*

228 Table 1 gives a summary of the success of data capture for the tail sensors and
229 SHM collars in the beef and dairy trials, and the reasons for excluding animals from
230 the data analysis. Supplementary Table 1 shows how the number of animals
231 included in the analysis changed with hours prior to calving. For the beef trial, a total
232 of 124 animals were included in the eating/rumination dataset, 112 in the activity
233 dataset and 75 in the tail sensor dataset. The corresponding numbers for the dairy
234 animals were 81, 101 and 53, respectively. The data capture from the tail sensors
235 was lower than would be practical for a commercial system. This is due to the fact
236 that the sensors were designed for data gathering purposes and have not been
237 protected sufficiently robustly for commercial deployment. As a consequence there
238 were significant numbers failures. This can be readily addressed through revision of
239 the mechanical housing.

240

241 *Changes in behaviour measured by animal mounted sensors*

242 *Tail raising*

243 Mean time spent with the tail in a raised position per hour in the week prior to calving
244 was 2.1 ± 0.04 min/hr in beef cows (Figure 2a) and 3.2 ± 0.07 min/hr for dairy cows
245 (Figure 2b). In the five hours prior to calving time spent with the tail raised was
246 significantly higher than in the control period for both beef (increase from 4.7 ± 0.80
247 to 22.8 ± 1.66 min/hr, $p < 0.01$) and dairy cows (increase from 6.6 ± 1.29 to $26.2 \pm$
248 2.48 min/hr, $p < 0.01$).

249 *Time spent ruminating*

250 In the week prior to calving, the mean time spent ruminating by beef cows was 21.9
251 ± 0.12 min/hr (Figure 3a). Time spent ruminating decreased significantly in the five
252 hours prior to calving compared to the control period (from 23.8 ± 0.67 to 12.0 ± 0.59
253 min/hr, $p < 0.001$). For dairy cows the mean time spent ruminating in the week prior
254 to calving was 16.6 ± 0.10 min/hr (Figure 3b). Time spent ruminating decreased
255 significantly in the five hours prior to calving when compared to the control period
256 (from 14.9 ± 0.73 to 8.8 ± 0.73 min/hr, $p < 0.001$).

257 *Time spent eating*

258 The mean time spent eating by beef cows was 21.1 ± 0.15 min/hr (Figure 4a) in the
259 week prior to calving. During the control period, mean time spent eating was $19.1 \pm$
260 0.76 min/hr, which increased significantly in the five hours prior to calving ($23.0 \pm$
261 0.74 min/hr, $p < 0.001$.. For dairy cows the mean time spent eating in the week prior
262 to calving was 19 ± 0.1 min/hr (Figure 4b). The five hours prior to calving was $24 \pm$
263 0.9 min/hr, which was significantly higher ($p < 0.05$) than the control period (22 ± 1.0
264 min/hr).

265 *Relative activity level*

266 In the week prior to calving, the mean relative activity by beef cows was 4.2 ± 0.06
267 (Figure 5a). Relative activity significantly increased compared to the control period in
268 the five hours prior to calving (from 5.9 ± 0.54 to 13.6 ± 1.12 , $p < 0.01$). For dairy
269 cows the mean relative activity was 2.9 ± 0.04 in the week prior to calving (Figure
270 5b). There was also a significant increase in relative activity in the five hours prior to
271 calving compared to the control period in dairy cows (from 4.3 ± 0.53 to 9.1 ± 0.81).

272 *Predictive models*

273 The model performance statistics for individual and integrated sensor variables are
274 shown in Table 2. Note that one integrated sensor model contains ACT and the other
275 does not. This is due to the difference in data reporting resolution between TAIL,
276 RUM and EAT (per hour) and ACT (per 1.5 hours). Data streams had to be
277 aggregated into 3 hour blocks to resolve the differences in resolution without making
278 the assumption that behaviours were being displayed evenly throughout the reported
279 time periods. The TAIL and TAIL+RUM+EAT models were found to be the most
280 robust models in both the beef and dairy cow data sets. The TAIL model was slightly
281 better at predicting calving within a five hour window for beef cows (MCC = 0.31)
282 than for dairy cows (MCC = 0.29). The TAIL+RUM+EAT models were equally as
283 good at predicting calving within a five hour window for beef and dairy cows (MCC =
284 0.32 for both models).

285 Variables recorded by the SHM collars alone (RUM, EAT and ACT) were not good
286 predictors of onset of parturition, the RUM and EAT variables being the worst
287 performing in both beef (MCC of 0.13 and 0.15 for RUM and EAT, respectively) and
288 dairy cows (MCC of 0.12 and 0.09 for RUM and EAT, respectively). Combining these

289 variables resulted in a poorer performing model (MCC = 0.07), likely due to the lower
290 resolution of data.

291 When assessing the relative importance of the sensor variables (calculated by
292 determining the drop in prediction accuracy after shuffling the values of a given
293 predictor variable in the oob samples, rendering them random and with no predictive
294 power – data not shown) within the TAIL+RUM+EAT dairy model, the TAIL variable
295 was by far the most important. Scaled (0-100, with 0 being redundant and 100 is the
296 most important) importance for TAIL was 100 in both, with RUM and EAT models
297 having substantially less influence (scaled importance of 22.1 and 21.7, respectively
298 for beef cows and 26.2 and 29.1 for dairy cows).

299 *Predicting time to calving*

300 As TAIL was identified as the most important sensor variable for prediction of
301 parturition, and as a one sensor system is more desirable than a multiple sensor
302 system, it was selected to develop models for prediction of discreet time points prior
303 to calf expulsion. Model parameters and performance metrics are shown for hours 0-
304 12 prior to calving in Table 3. Within the beef cows, the predictive performance of
305 TAIL increases sharply after four hours prior to calf expulsion (MCC increases from
306 0.07 at four hours prior to 0.17 at three hours prior). A similar sharp increase was
307 observed in the dairy cows (MCC increased from 0.06 four hours prior to calf
308 expulsion to 0.14 at three hours prior to calf expulsion).

309 Discussion

310 *Changes in cow behaviour prior to calf expulsion*

311 The changes in rumination behaviour observed in this study are in line with those
312 found in previous studies. Reductions in rumination time Soriani *et al* (2012) found
313 reductions in rumination time of between 38-50% in Italian Friesian cows on the day
314 they calved. Calamari *et al* (2014) observed a 30% drop in rumination time on the
315 day of calving compared to the dry period, also in Italian Friesian cows. Büchel and
316 Sundrum (2014) detected an average 27% decrease in the six hours prior to Holstein
317 cows calving. Pahl *et al* (2014) found significant differences between time spent
318 ruminating in the four hours prior to calf expulsion compared to a reference period for
319 dairy cows; Braun *et al* (2014) reported a reduction in rumination time of 45% on the
320 day of parturition in Swiss Braunvieh cows.

321 An increase in tail raising behaviour, particularly in the two hours prior to calving in
322 dairy cows has also been observed previously (Miedema *et al*, 2011a,b; Jensen,
323 2012).

324 The beef cows displayed a sharp increase in the EAT variable in the hour prior to
325 calf expulsion and in the hour in which the calf was born which was not observed in
326 the dairy cows. This is contrary to other studies which report decreases when
327 measurements were made by visual observation (Miedema *et al*, 2011a), by
328 recording the time the cow spends with its head in a feed bin (Braun *et al*, 2014;
329 Büchel and Sundrum, 2014). This can be explained by the inability of the SHM collar
330 to distinguish between neck movement characteristic of eating and behaviours which
331 result in similar neck motion e.g. grooming and licking. For the beef cows, the hour in
332 which the calf was born includes the whole hour, regardless of when the cow calved

333 within that hour – e.g. if the cow calved at quarter past the hour, the next 45 minutes
334 are also included. The apparent observed increase in eating may actually be
335 misclassification of licking behaviour, this behaviour has been shown to peak in the
336 hour proceeding birth of the calf (Jensen, 2012). The same trend was not observed
337 in the dairy cows as their collars were removed directly after calving. In the hour prior
338 to calf expulsion it is possible that the cow is displaying ground licking or nesting
339 behaviours (Miedema *et al*, 2011a).

340 Activity levels are known to increase in cows in the hours prior to calf expulsion when
341 measured by visual observations (Miedema *et al*, 2011a,b) and leg mounted
342 accelerometers (Titler *et al*, 2015). In this study, neck mounted accelerometers
343 detected an increase in activity prior to calf expulsion, particularly in the final two
344 hours. Clark *et al* (2015) did not detect any increase in activity prior to calf expulsion
345 in dairy cows using similar neck mounted accelerometers. As different animal
346 mounted sensors have different algorithms to define behaviours, and have
347 undergone different validation exercises it may be expected that there will be
348 substantial differences in behavioural measurements between them.

349 There are no studies which use animal mounted sensors to detect changes in
350 rumination time, eating time, relative activity and tail raising prior to calf expulsion in
351 suckler beef cows. This study has shown that patterns of behaviours at onset of
352 parturition are very similar in suckler beef and dairy cows.

353 *Prediction of parturition*

354 Interest in developing real-time predictive models to alert farmers to when cows will
355 calve using animal mounted sensors is increasing. The majority of published studies
356 using sensors to monitor various behaviours have been on dairy cows. Some studies

357 simply use threshold changes in behaviours to define the onset of parturition. Titler
358 *et al* (2015) were able to predict parturition on average 6 hours in advance by a 50%
359 increase in activity. Krieger *et al* (2018) used threshold values for frequency and
360 duration of tail raise events to predict parturition in five cows and detected calving
361 between 6 and 121 minutes prior to expulsion of the calf. In reality, the results of
362 Krieger *et al* (2018) are similar to those found here, where sharp increases in
363 predictive accuracy of algorithms were observed one to two hours prior to calf
364 expulsion in hour-by-hour models.

365 A variety of multi-sensor systems have been used to integrate data streams
366 monitoring different behaviours. Rutten *et al* (2017) achieved a very low false
367 positive rate of 1% within three hours of calf expulsion using two sensors to measure
368 activity level, rumination time, feeding time and temperature; however the sensitivity
369 was only 42.4%. Borchers *et al* (2017) were able to predict parturition eight hours
370 prior to calf expulsion with a sensitivity of 82.8% and a specificity of 80.4% using two
371 sensors (neck mounted for rumination time and leg mounted for time spent standing
372 or lying and step count). Ouellet *et al* (2016) achieved sensitivity of 77% and
373 specificity of 77% within a 24 hour window using three sensors to record four
374 variables (vaginal temperature, rumination time, lying time and lying bouts). In the
375 present study, similar results were achieved with a single sensor system (TTA:
376 sensitivity = 78.6%, specificity = 83.5% for dairy cows). Single sensor systems may
377 be more attractive to industry in terms of the financial outlay required and may
378 encourage greater industry uptake.

379 *Conclusions*

380 In this study it was possible to predict, with reasonable accuracy, when beef or dairy
381 cows were within five hours of calf expulsion using animal mounted technologies. Of

382 the variables measured by the sensors used in this study, time spent with the tail in a
383 raised position was found to be the best predictor of parturition, and had optimal
384 predictive power at two hours prior to calf expulsion.

385 *Acknowledgements*

386 The authors gratefully acknowledge NERC and BBSRC for funding through the
387 Sustainable Agriculture Research and Innovation Club (SARIC). SRUC are funded
388 by the Scottish Government through the Strategic Research programme of the
389 Scottish Government's Rural and Environment Science and Analytical Services
390 Division (RESAS). Thanks to the commercial dairy farm for their assistance and
391 cooperation and to the technical team at SRUCs Beef Research Centre.

392 **References**

393 Agriculture and Horticulture Development Board (AHDB), Beef and Lamb (2018) AHDB UK
394 cattle yearbook 2018.

395 Agjee, N.H., Mutanga, O., Peerbhay, K. and Ismail, R. (2018) The impact of simulated
396 spectral noise on random forest and oblique random forest classification performance.
397 *Journal of Spectroscopy* doi.org/10.1155/2018/8316918

398 Barrier, A.C., Haskell, M.J., Birch, S., Bagnall, A., Bell, D.J., Dickinson, J., Macrae, A.I. and
399 Dwyer, C.M. (2013) The impact of dystocia on dairy calf health, welfare, performance and
400 survival. *The Veterinary Journal* 195:86-90

401 Borchers, M.R., Chang, Y.M., Proudfoot, K.L., Wadsworth, B.A., Stone, A.E. and Bewley,
402 J.M. (2017) Machine-learning-based calving prediction from activity, lying, and ruminating
403 behaviors in dairy cattle. *Journal of Dairy Science* 100:5664-5674

404 Braun, U. Tschoner, T. and Hässig, M. (2014) Evaluation of eating and rumination behaviour
405 using a noseband pressure sensor in cows during the peripartum period. *BMC Veterinary*
406 *Research* 10:195

407 Büchel, S. and Sundrum, A. (2014) Decrease in rumination time as an indicator of the onset
408 of calving. *Journal of Dairy Science* 97:3120-3127

409 Calamari, L., Soriani, N., Panella, G., Petrera, F., Minuti, A. and Trevisi, E. (2014)
410 Rumination time around calving: An early signal to detect cows at greater risk of disease.
411 *Journal of Dairy Science* 97:3635-3647

412 Clark, C.E.F., Lyons, N.A., Millapan, L., Talukder, S., Cronin, G.M., Kerrisk, K.L. and Garcia,
413 S.C. (2015) Rumination and activity levels as predictors of calving for dairy cows. *Animal*
414 9:691-695

415 De Amicis, I., Veronesi, M.C., Robbe, D., Gloria, A., Carluccio, A. (2018) Prevalence,
416 causes, resolution and consequences of bovine dystocia in Italy. *Theriogenology* 1007:104-
417 108

418 Eriksson, S., Näsholm, A., Johansson, K. and Philipsson, J. (2004) Genetic parameters for
419 calving difficulty, stillbirth, and birth weight for Hereford and Charolais at first and later
420 parities. *Journal of Animal Science* 82:375-383

421 Gaafar, H.M.A., Shamiah, M., Abu El-Hamd, M.A., Shitta, A.A. and Tag El-Din, M.A. (2011)
422 Dystocia in Friesian cows and its effects on postpartum reproductive performance and milk
423 production. *Tropical Animal Health and Production* 43:229-234

424 Huzzey, J.M., von Keyserlingk, M.A.G. and Weary, D.M. (2005) Changes in feeding, drinking
425 and standing behaviour of dairy cows during the transition period. *Journal of Dairy Science*
426 88:2454-2461

427 Jensen, M.B. (2012) Behaviour around the time of calving in dairy cows. *Applied Animal*
428 *Behaviour Science* 139:195-202

429 Kuhn, M. Contributions from Jed Wing, Steve Weston, Andre Williams, Chris Keefer, Allan
430 Engelhardt, Tony Cooper, Zachary Mayer, Brenton Kenkel, the R Core Team, Michael
431 Benesty, Reynald Lescarbeau, Andrew Ziem, Luca Scrucca, Yuan Tang, Can Candan and
432 Tyler Hunt (2018) caret: Classification and Regression Training. R package version 6.0-80.
433 <https://CRAN.R-project.org/package=caret>

434 Kovács, L., Kézér, F.L., Ruff, F. and Szenci, O. (2016) Rumination time and reticuloruminal
435 temperature as possible predictors of dystocia in dairy cows. *Journal of Dairy Science*
436 100:1568-1579

437 Krieger, S., Sattlecker, G., Kicking, F., Auer, W., Drillich, M. and Iwersen, M. (2018)
438 Prediction of calving in dairy cows using a tail-mounted tri-axial accelerometer: A pilot study.
439 *Biosystems Engineering* 173:79-84

440 Lombard, J.E., Garry, F.B., Tomlinson, S.M. and Garber, L.P. (2007) Impacts of dystocia on
441 health and survival of dairy calves. *Journal of Dairy Science* 90:1751-1760

442 López de Maturana, E., Legarra, A., Varona, L. and Ugarte, E. (2007) Analysis of fertility and
443 dystocia in Holsteins using recursive models to handle censored and categorical data.
444 *Journal of Dairy Science* 90:2012-2024

445 Lowman, B.G., Scott, N., Somerville, S. (1973) Condition scoring of cattle. Bulletin
446 No. 6. Edinburgh (United Kingdom): East of Scotland College of Agriculture. McGuirk,
447 B.J., Forsyth, R. and Dobson, H. (2007) Economic cost of difficult calvings in the United
448 Kingdom dairy herd. *Veterinary Record* 161:685-687

449 Mee, J.F. (2008) Prevalence and risk for dystocia in dairy cattle: A review. *The Veterinary*
450 *Journal* 176:93-101

451 Miedema, H.M., Cockram, M.S., Dwyer, C.M. and Macrae, A.I. (2011a) Changes in the
452 behaviour of dairy cows during the 24h before normal calving compared to behaviour during
453 late pregnancy. *Applied Animal Behaviour Science* 131:8-14

454 Miedema, H.M., Cockram, M.S., Dwyer, C.M. and Macrae, A.I. (2011b) Behavioural
455 predictors of the start of normal and dystocic calving in dairy cows and heifers. *Applied*
456 *Animal Behaviour Science* 132:14–19

457 Nix, J.M., Spitzer, J.C., Grimes, L.W., Burns, G.L. and Plyler, B.B. (1998) A retrospective
458 analysis of factors contributing to calf mortality and dystocia in beef cattle. *Theriogenology*
459 49:1515-1523

460 Ouellet, V., Vasseur, E., Heuwieser, W., Burfeind, O., Maldague, X. and Charbonneau, E.
461 (2016) Evaluation of calving indicators measured by automated monitoring devices to predict
462 the onset of calving in Holstein dairy cows. *Journal of Dairy Science* 99:1539-1548

463 Pahl, C., Hartung, E., Grothmann, A. and Mahlkow-Nerge, K. (2014) Rumination activity of
464 dairy cows in the 24 hours before and after calving. *Journal of Dairy Science* 97:6935-6941

465 Phocas, F. and Laloë, D. (2003) Evaluation of genetic parameters for calving difficulty in
466 beef cattle. *Journal of Animal Science* 81:933-938

467 R Core Team (2017) R: A language and environment for statistical computing. R Foundation
468 for Statistical Computing, Vienna, Austria. <https://www.R-project.org>

469 Robin, X., Turck, N., Hainard, A., Tiberti, N., Lisacek, F., Sanchez, J. and Muller, M. (2011)
470 pROC: an open-source package for R and S+ to analyse and compare ROC curves. *BMC*
471 *Bioinformatics*, 12, p.77. DOI: 10.1186/1471-2105-12-77

472 Rumph, J.M., Faust, M.A. (2006) Genetic analysis of calving ease in Holsteins in the U.K.
473 based on data from heifers and cows. *Abstracts Book of the 8th World Congress on*

474 *Genetics Applied to Livestock Production*, August 13–18, Belo Horizonte, MG, Brazil,
475 Abstract 1– 25. p. 11.

476 Rutten, C.J., Kamphuis, C., Hogeveen, H., Huijps, K., Nielen, M. and Steeneveld, W. (2017)
477 Sensor data on cow activity, rumination, and ear temperature improve prediction of the start
478 of calving in dairy cows. *Computers and Electronics in Agriculture* 132:108-118

479 Saint-Dizier, M. and Chastant-Maillard, S. (2015) Methods and on-farm devices to predict
480 calving time in cattle. *The Veterinary Journal* 205:349-356

481 Schirmann, K., Chapinel, N., Weary, D.M., Vickers, L. and von Keyserlingk, M.A.G. (2013)
482 Rumination and feeding behaviour before and after calving in dairy cows. *Journal of Dairy*
483 *Science* 96:7088-7092

484 Schuenemann, G.M., Nieto, I., Bas, S., Galvão, K.N. and Workman, J. (2011) Assessment of
485 calving progress and reference times for obstetric intervention during dystocia in Holstein
486 dairy cows. *Journal of Dairy Science* 94:5494-5501

487 Shah, K.D., Nakao, T. and Kubota, H. (2006) Plasma estrone sulphate (E₁S) and estradiol-
488 17β (E₂β) profiles during pregnancy and their relationship with the relaxation of sacrosciatic
489 ligament, and prediction of calving time in Holstein-Fresian cattle. *Animal Reproduction*
490 *Science* 95:38-53

491 Soriani, N., Trevisi, E. and Calamari, L. (2012) Relationships between rumination time,
492 metabolic conditions, and health status in dairy cows during the transition period. *Journal of*
493 *Animal Science* 90:4544-4554

494 Titler, M., Maquivar, M.G., Bas, S., Rajala-Schultz, P.J., Gordon, E., McCullough, K. and
495 Federico, P. (2015) Prediction of parturition in Holstein dairy cattle using electronic data
496 loggers. *Journal of Dairy Science* 98:5304-5312

497 Wickham, H., Francois, R., Henry, L. and Muller, K. (2018) dplyr: A grammar of data
498 manipulation. R package version 0.7.6. [https:// CRAN.R-project.org/package=dplyr](https://CRAN.R-project.org/package=dplyr)

499

500 Table 1: Success of data recording for SHM collars and tail sensors on beef and
 501 dairy cows

	Beef			Dairy		
	Eating / Rumination	Activity	Tail raise	Eating / Rumination	Activity	Tail raise
Total animals	144	144	144	110	110	110
Successful recording	137	128	93	85	103	55
Not attached	-	-	3	-	-	2
No calving time	9	9	9	-	-	-
Less than 48 hours	4	15	3	4	2	0
Animals in analysis	124	111	75	81	101	53

502

503

504 Table 2: Model parameter tuning and performance statistics for single and combined
505 sensor variable random forest models. mtry = number of variables used at each split
506 in each independent decision tree, ntree = number of independent decision trees
507 oob error = out of bag error, AUC = area under the curve, CI = confidence interval,
508 Se = sensitivity, Sp = specificity, MCC = Matthew's Correlation Coefficient, TAIL =
509 number of tail raise events per hour, EAT = time spent eating per hour (minutes),
510 RUM = time spent ruminating per hour (minutes), ACT = relative level of activity per
511 1.5 hours (minutes).

	mtry	ntree	oob error	AUC (95% CI)	Sensitivity (%)	Specificity (%)	MCC
Beef							
TAIL	3	1000	0.187	86.7 (83.1, 90.4)	76.1	83.3	0.31
RUM	4	2500	0.376	69.5 (65.1, 73.9)	69.6	62.3	0.13
EAT	4	2500	0.386	71.7 (67.5, 75.9)	63.8	70.2	0.15
ACT	3	2500	0.296	78.1 (73.8, 82.4)	70.9	71.5	0.18
TAIL+RUM+EAT	2	2500	0.187	86.7 (83.1, 90.3)	75.4	84.6	0.32
RUM+EAT+ACT	5	2500	0.526	46.7 (55.3, 62.5)	62.5	55.3	0.07
TAIL+RUM+EAT+ACT	6	1500	0.526	72.9 (60.5, 85.3)	81.3	69.7	0.22
Dairy							
TAIL	2	2000	0.267	87.9 (81.5, 90.1)	78.6	83.5	0.29
RUM	1	1000	0.491	64.0 (58.5, 69.5)	69.8	59.3	0.12
EAT	3	500	0.463	62.4 (56.4, 68.5)	59.3	61.7	0.09
ACT	5	2000	0.421	68.2 (63.7, 72.7)	66.7	62.3	0.11
TAIL+RUM+EAT	3	2000	0.226	85.2 (80.5, 89.8)	76.7	85.1	0.32
RUM+EAT+ACT	4	1500	0.345	51.4 (68.8, 75.0)	75.0	68.8	0.18
TAIL+RUM+EAT+ACT	5	1000	0.242	86.9 (78.8, 95.1)	79.2	81.3	0.30

512 ¹ ACT models have a 1.5 hour time step due to the resolution of data collection for
513 this sensor variable.

514 ² Combined models containing ACT have a 3 hour time step to resolve differences in
515 the resolution of data collection between ACT and other sensor variables.

516

517 Table 3: Model parameter tuning and performance statistics for random forest
 518 models using number of tail raise events to predict parturition at discreet time points
 519 prior to calf expulsion. Mtry = number of variables used at each split in each tree,
 520 ntree = number of independent decision trees, oob error = out of bag error, AUC =
 521 area under the curve, Se = sensitivity, Sp = specificity, MCC = Matthew's Correlation
 522 Coefficient

Hours prior to calf expulsion	mtry	ntree	oob error	AUC	Se (%)	Sp (%)	MCC
Beef							
0	6	2000	0.14	88.5 (79.9, 97.1)	79.2	93.3	0.25
1	8	500	0.11	89.8 (80.0, 99.6)	90.9	90.9	0.23
2	6	2000	0.23	95.4 (92.2, 98.6)	91.3	93.5	0.29
3	6	1000	0.25	84.1 (74.6, 93.7)	78.3	87.0	0.17
4	8	2500	0.32	59.2 (45.4, 73.1)	47.8	82.2	0.07
5	8	1000	0.54	47.8 (35.7, 59.9)	52.2	53.9	0.01
6	6	2000	0.51	56.4 (44.9, 67.9)	53.1	70.5	0.05
7	8	1500	0.57	57.6 (44.1, 71.0)	68.4	60.8	0.05
8	7	1500	0.59	53.8 (40.6, 67.1)	57.9	58.1	0.03
9	7	2500	0.52	54.2 (43.1, 65.3)	57.7	51.1	0.02
10	8	500	0.44	63.4 (50.8, 69.7)	63.2	64.2	0.05
11	6	2000	0.64	59.5 (49.3, 69.7)	62.5	56.4	0.03
12	8	2500	0.69	65.3 (52.1, 78.5)	55.6	66.5	0.04
Dairy							
0	5	500	0.21	88.2 (71.9, 100)	87.5	89.7	0.16
1	5	1500	0.13	93.2 (88.5, 97.9)	81.3	89.7	0.20
2	5	2500	0.34	92.0 (86.0, 98.0)	86.7	92.4	0.25
3	4	1500	0.31	85.4 (75.5, 95.3)	70.0	90.3	0.14
4	2	1500	0.59	68.3 (48.6, 87.9)	88.9	54.1	0.06
5	3	1000	0.50	56.4 (38.2, 74.7)	58.3	61.4	0.03
6	5	1500	0.58	65.5 (51.8, 79.1)	80.0	59.0	0.06
7	1	2000	0.68	56.9 (43.7, 70.0)	50.0	61.2	0.02
8	5	500	0.83	54.5 (38.6, 70.4)	61.1	55.6	0.03
9	5	500	0.60	58.8 (41.8, 75.8)	71.4	54.1	0.04
10	5	500	0.48	57.5 (42.3, 72.8)	47.4	69.3	0.04
11	5	1500	0.42	52.7 (38.0, 67.4)	71.4	41.4	0.02
12	5	1000	0.56	50.2 (34.6, 65.9)	72.7	40.2	0.02

524

525 Figure 1: Tail mounted tri-axial accelerometer (TTA) attachment and orientation

526 Figure 2: Average number of tail raises per hour one week prior to calf expulsion for

527 a) beef and b) dairy cows.

528 Figure 3: Average time spent ruminating (minutes per hour) one week prior to calf

529 expulsion for a) beef and b) dairy cows.

530 Figure 4: Average time spent eating (minutes per hour) one week prior to calf



531 expulsion for a) beef and b) dairy cows.

532 Figure 5: Average relative activity (per hour) one week prior to calf expulsion for a)

533 beef and b) dairy cows.


534

535 Figure 1: Tail mounted tri-axial accelerometer (TTA) attachment and orientation




536

537 Figure 2: Average number of tail raises per hour one week prior to calving for a) beef
538 and b) dairy cows.



539

540 Figure 3: Average time spent ruminating (minutes per hour) one week prior to calving
541 for a) beef and b) dairy cows.



542

543

544

545

546


547

548

549

550 Figure 4: Average time spent eating (minutes per hour) one week prior to calving for

551 a) beef and b) dairy cows



552

553

554

555

556


557

558

559 Figure 5: Average relative activity (per hour) one week prior to calving for a) beef and

560 b) dairy cows.

561



562

563

564

565