


ACCEPTED MANUSCRIPT

The Prediction of Moment of Inertia of Rotating Nuclei

Abdurahim Okhunov¹ , Mohd Kh. M. Abu El Sheikh², Hasam Abu Kassim³ and
M. H. H. Handaker⁴

Accepted Manuscript online 20 August 2020 • © 2020 Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

What is an Accepted Manuscript?

aaokhunov@gmail.com

¹ Science in Engineering, International Islamic University Malaysia, IIUM, P.O. Box 10, Jalan Gombak, Kuala Lumpur, Kuala Lumpur, Selangor, 53100, MALAYSIA

² Physics, University of Malaya, Jalan Universiti, Kuala Lumpur, Wilayah Persekutuan, 50603, MALAYSIA

³ Physics, Universiti Malaya, Kuala Lumpur, Malaysia, Kuala Lumpur, Wilayah Persekutuan, 50603, MALAYSIA

⁴ Centre for Biomedical Physics, School of Healthcare and Medical Sciences,, Sunway University, Selangor, Bandar Sunway, Selangor, 47500, MALAYSIA

Abdurahim Okhunov  <https://orcid.org/0000-0001-8487-5506>

Received 23 May 2020

Revised 9 July 2020

Accepted 24 July 2020

Accepted Manuscript online 20 August 2020

<https://doi.org/10.1088/1674-1137/abb0dd>

Abstract

In this paper the mathematical expression which is given by Bohr for the moment of inertia of even-even nuclei on the basis of the hydrodynamical model is modified. This modification is on the kinetic energy of the surface oscillations including the second and third terms of R - R expansion as well as the first term which was already carried out by Bohr. Therefore, this work can be considered

the continuation and support of the hydrodynamic model of Bohr. This procedure results in a Bohr formula to be multiplied by a factor which depends on the deformation parameter. Bohr (modified) formula is examined by applying it on axially symmetric even-even nuclei with atomic mass ranged between 150 and 190 as well as to some triaxial symmetry nuclei. The modification of Bohr's formula are discussed including the information on how stable this modification with including second and third terms of R^2 - expansion of Bohr's formula. The results of calculation are compared with the experimental data and the results of Bohr, based earlier. The obtained results are in a good agreement with experimental data by describing almost 0.7% and better than that of the unmodified ones.

Export citation and abstract

[BibTeX](#)

[RIS](#)



As the Version of Record of this article is going to be/has been published on a subscription basis, this Accepted Manuscript will be available for reuse under a CC BY-NC-ND 3.0 licence after a 12 month embargo period.

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permission may be required. All third party content is fully copyright protected, unless specifically stated otherwise in the figure caption of the Version of Record.