



## CKJ REVIEW

# Chronic kidney disease in children

Francesca Becherucci<sup>1,\*</sup>, Rosa Maria Roperto<sup>1,\*</sup>, Marco Materassi<sup>1</sup>  
and Paola Romagnani<sup>1,2</sup>

<sup>1</sup>Nephrology and Dialysis Unit, Meyer Children's Hospital, Florence, Italy and <sup>2</sup>Department of Biomedical Experimental and Clinical Sciences 'Mario Serio', University of Florence, Florence, Italy

Correspondence and offprint requests to: Paola Romagnani; E-mail: p.romagnani@dfc.unifi.it, paola.romagnani@meyer.it

\*F.B. and R.M.R. equally contributed to this work.

## Abstract

Chronic kidney disease (CKD) is a major health problem worldwide. Although relatively uncommon in children, it can be a devastating illness with many long-term consequences. CKD presents unique features in childhood and may be considered, at least in part, as a stand-alone nosologic entity. Moreover, some typical features of paediatric CKD, such as the disease aetiology or cardiovascular complications, will not only influence the child's health, but also have long-term impact on the life of the adult that they will become. In this review we will focus on the unique issues of paediatric CKD, in terms of aetiology, clinical features and treatment. In addition, we will discuss factors related to CKD that start during childhood and require appropriate treatments in order to optimize health outcomes and transition to nephrologist management in adult life.

**Key words:** chronic renal failure, chronic renal insufficiency, CKD, ESRD, paediatrics

## Introduction

Chronic kidney disease (CKD) is a major health problem worldwide with increasing incidence and prevalence that is threatening to bring on the onset of a real 'epidemic' [1–5]. Independent of the initial cause, CKD is a clinical syndrome characterized by a gradual loss of kidney function over time [6]. In particular, the Kidney Disease: Improving Global Outcomes (KDIGO) guidelines have defined CKD as abnormalities of kidney structure or function, present for more than 3 months, with implications to health [6]. This definition has been formulated for the adult population, where CKD is a common and well-known health problem, but the KDIGO guidelines for definition and staging are not fully applicable to the paediatric population [6]. Indeed, paediatric CKD, while sharing the basic physiopathologic mechanisms with the same disease in the adult population, could be in some ways considered a stand-alone nosologic entity. Childhood CKD presents clinical features that are specific and totally

peculiar to the paediatric age, such as the impact of the disease on growth. In addition, some of the typical characteristics of paediatric CKD, such as the aetiology or cardiovascular complications, represent variables, not only influencing the health of the patient during childhood, but also having an impact on the life of the adult that this child will become. This impact is often under-recognized but should not be neglected. Moreover, CKD has a great psychosocial impact, both on the patient and his family. The parents not only have to fulfil the role of parents, but also take on many tasks we normally associate with nurses and doctors. Therefore, we must be aware that the increasing survival of paediatric patients with CKD, due to the improvement in the clinical and therapeutic management, will lead to a large number of affected adults facing problems that are specific to CKD that have started during childhood.

In this review, we will focus not only on the unique issues concerning paediatric CKD, but especially on those factors related to

Received: February 11, 2016. Accepted: May 4, 2016

© The Author 2016. Published by Oxford University Press on behalf of ERA-EDTA.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

CKD that start during childhood and require appropriate management to optimize health outcomes of the patient.

## Epidemiology of CKD in children

According to the KDIGO guidelines, CKD is identified by the presence of kidney damage, either structural or functional, or by a decline in glomerular filtration rate (GFR) below 60 mL/min/1.73 m<sup>2</sup> of body surface area for more than 3 months [6]. Therefore, the term CKD defines renal dysfunction as a *continuum*, rather than a discrete change in renal function, either in children or in adults [7]. This makes the epidemiology of CKD very difficult to study. Moreover, epidemiologic data on CKD may underestimate its real incidence and prevalence since CKD is often clinically asymptomatic, especially in earlier stages [8]. This is in part the result of the historical absence of a common definition of CKD and of a well-defined classification of its severity [9], that have recently, at least in part, been overcome by the introduction of the KDIGO guidelines [6, 9]. For all these reasons, in the majority of studies, estimates of CKD take into account patients with moderate to severe CKD or end-stage renal disease (ESRD) and are not population-based in nature [8, 9]. In addition, childhood CKD registries are usually limited by being restricted to small reference populations [10]. Despite these limitations, the paediatric incidence of CKD in Europe is reported to be around 11–12 per million of age-related population (pmarp) for stages 3–5, while the prevalence is ~55–60 pmarp (Figure 1, top) [11, 12, 13, 17]. Comparable amounts are reported in population-based registries of other western countries [14], although no precise data on the incidence and prevalence of pre-terminal CKD are available for the majority of them (Figure 1, top) [9]. Some differences could emerge if different age groups are analysed [10, 18]. The incidence of paediatric CKD rose slowly during the 1980s, then marginally until the first decade of the 21st century [19]. At the same time, the prevalence of the disease has significantly increased since survival and treatment of CKD have markedly improved [20]. Specific reports on CKD epidemiology in children have been focused on patients with ESRD requiring renal replacement therapy (RRT). The median incidence of RRT in children <20 years old is ~9 pmarp worldwide, whereas the prevalence is reported as ~65 pmarp (Figure 1, bottom) [9, 13, 21]. Moreover, higher values for incidence and prevalence have been reported in the USA, probably because RRT is started earlier, at higher levels of GFR, in comparison with other developed countries [15]. In any case, data coming from epidemiological studies in adults provide the dramatic evidence that ESRD represents the ‘tip of the iceberg’ of CKD and suggest that the number patients with earlier stages of the disease are likely to exceed those reaching ESRD by as much as 50 times [22]. The same consideration could probably be applied to the paediatric population, where CKD has only recently been recognized as a non-marginal issue. Notably, 80% of RRT in children is performed in Europe, Japan and North America, where the cost of these extremely expensive treatments can be afforded [9]. As a consequence, the real impact of CKD in children in developing countries is a long way from being clarified, especially in those countries where the healthcare resource allocation to RRT is inadequate or RRT is not available, and children affected by CKD frequently die [9, 23, 24].

The incidence and prevalence of CKD is greater in males than females because of the higher frequency of congenital abnormalities of the kidney and urinary tract (CAKUT) in males [11]. Finally, race is another factor specifically affecting the epidemiology of CKD. In particular, in North America, the incidence of CKD is two to three times higher in African-American children

compared with Caucasian children, irrespective of gender [14], whereas in Australia and New Zealand, the risk of ESRD is greater in indigenous children (e.g. Aborigines and Maoris) than in the remainder of the paediatric population [25].

## Aetiology of CKD in children

Primary causes of CKD in children significantly differ from those that are responsible for the adult onset of the disease. In fact, the main aetiological factors of CKD in children are represented by CAKUT, steroid-resistant nephrotic syndrome (SRNS), chronic glomerulonephritis (e.g. lupus nephritis, Alport syndrome) and renal ciliopathies, that account for approximately 49.1, 10.4, 8.1 and 5.3% of cases, respectively [11, 26, 27] and for more than 70% of all paediatric CKD cases when considered together, as recently reported (Table 1) [26]. Less common causes of CKD in children include thrombotic microangiopathies (especially atypical haemolytic uraemic syndrome), nephrolithiasis/nephrocalcinosis, Wilms tumour, infectious and interstitial diseases, and others (Table 1) [26]. While structural causes (e.g. renal hypoplasia or posterior urethral valves) clearly predominate in younger patients, the incidence of glomerulonephritis increases in those >12 years old (Figure 2) [9, 26]. However, minor reductions in nephron numbers that are seen in low-birth weight and small for gestational age newborns are now emerging as important predisposing factors to CKD and will come to represent an important issue for nephrologists as the number of premature children continues to grow [28–31]. These conditions, together with the exploding burden of paediatric obesity [32, 33], are probably destined to significantly change the relative distribution of the causes of CKD.

Interestingly enough, if analysis of the causes is limited to the population of children that have already reached ESRD, the relative percentage of glomerular diseases increases (approximately doubling), whereas that of CAKUT decreases from around 50 to 39.5%, underscoring the discrepancy between the rate of progression of these two entities (Table 1). Indeed, congenital malformative disorders are characterized by a slower progression towards ESRD in comparison with glomerular diseases so that, as mentioned before, the relative proportion of glomerular diseases increases in groups of patients with more advanced stages of CKD (Table 1) [9]. Somewhat different are the data from the Japanese National registry and the Australia and New Zealand Dialysis and Transplant (ANZDATA) registry, which reported glomerulonephritis to be the most common cause of ESRD in children and adolescents [16, 34]. Finally, although information on the aetiology of ESRD from less-developed countries is almost unavailable mainly due to the absence of renal registries, it is reasonable to state that the burden of glomerulonephritis secondary to infectious diseases (such as hepatitis C, tuberculosis, HIV) is predominant and still far from being under control [9].

The recent advent of massive-parallel sequencing technologies (also referred to as next-generation sequencing, NGS) has provided one of the most interesting and substantial clues in unravelling the aetiology of early-onset CKD. In particular, studies performed over the past few years have demonstrated that a significant proportion of cases of CKD manifesting before 25 years of age can be defined as monogenic. In other words, a single gene can be detected as the cause of the disease in ~20% of early-onset patients [26]. Nowadays, more than 200 genes are clearly recognized as causative of the most common aetiological categories of CKD in children (CAKUT, SRNS, chronic glomerulonephritis and ciliopathies) [26, 35, 36]. NGS technology presents the striking advantage of allowing us to simultaneously study an elevated number of genes in a single run of sequencing, saving




Fig. 1. Estimated prevalence of CKD (top) and ESRD (bottom) in children worldwide. Data are collected by NAPRTCS, the Italian registry, USRDS, ESPN/ERA-EDTA registry, ANZDATA and the Japanese registry [9, 11–16]. Incidence and prevalence are reported as number of patients per million of age-related population (pmarp) per year and number of patients pmarp, respectively. Data from the ESPN/ERA-EDTA registry are reported on the basis of the contribution to the European registry of each single country, as available at [www.espn-reg.org/index.jsp](http://www.espn-reg.org/index.jsp). CKD, chronic kidney disease; ESRD, end-stage renal disease.

both time and costs while being extremely highly informative. Therefore, by selecting an appropriate panel of genes to sequence on the basis of the clinical phenotype of the patient or on a precise diagnostic suspicion, it is possible to address specific aetiological questions in one-fifth of children with early-onset CKD [26]. In addition, large population-based genetic studies (e.g. genome-wide association studies) are revealing that the genetic background of patients with CKD is probably much more complex than what was previously expected. Indeed, besides clearly disease-causing genes, that are by themselves responsible for disease determination, a number of other genes are now recognized as playing an important role [26, 37]. The best known example is represented by *APOL1*, whose variants confer a

considerably higher risk of developing focal segmental glomerulosclerosis and CKD progression [37–40].


These findings have substantial implications, either for the single patient or for more generalized considerations. First of all, patients with a recognized genetic cause of paediatric onset of CKD might benefit from specific therapies or from the avoidance of ineffective and even potentially health threatening ones (e.g. immunosuppressive drugs in patients with genetic forms of SRNS) [41, 42]. In addition, molecular diagnostics enable prenatal testing in siblings of affected individuals and genetic counselling to the family, and may be of great help in assessing a patient prognosis. Finally, the categorization of disease entities by means of genetic testing is fundamental in assuring that the

**Table 1.** Frequency of different diagnostic groups as causes of CKD and ESRD in children

	Frequency as cause of CKD [12, 13, 26]	Aetiology	Proportion of cases of CKD determined by specific diagnostic sub-groups [26]	Frequency as cause of ESRD [12–15, 19]
Glomerular diseases	6.8–20.5%	SRNS	10.4%	15.2–24.3%
		Glomerulonephritis	8.1%	
		Thrombotic microangiopathies (aHUS)	2.0%	
Structural and other	56–57.6%	CAKUT	49.1%	38.3–39.5%
		Ciliopathies	5.3%	
		Nephrolithiasis, nephrocalcinosis	1.6%	

The distribution of the frequency shows that different aetiological groups are differentially responsible for CKD and ESRD, mainly because of the different rate of progression towards ESRD of the different diagnostic categories (e.g. glomerular versus structural).

CKD, chronic kidney disease; ESRD, end-stage renal disease; CAKUT, congenital abnormalities of kidney and urinary tract; SRNS, steroid-resistant nephrotic syndrome; aHUS, atypical haemolytic uraemic syndrome.



**Fig. 2.** Impact of different causes of CKD in children among age groups. The graph shows the variation of the impact of different diagnostic groups in determining CKD over time, highlighting how glomerular diseases significantly increase in older children, while structural disorders are more common as causes of CKD in infants and younger children. CKD, chronic kidney disease; FSGS, focal segmental glomerulosclerosis; GN, glomerulonephritis; yrs, years.

analysis of data from clinical research and pharmacological trials is reliable.

## Clinical features of CKD in children and future implications

### Growth impairment

Growth impairment is a common and perhaps the most visible complication of CKD in children [43–45]. The degree of growth impairment increases as GFR declines, even though a significant decrease in growth was seen at all levels of kidney function [43, 44]. The 2006 North American Pediatric Transplant Cooperative Study carried out on over 5000 children, showed that over 35% of children with CKD had a height less than the third percentile or a median height standard deviation score (HtSDS) less than  $-1.88$ . The same study found a correlation between GFR and HtSDS with, respectively,  $-3.2$ ,  $-1.9$ ,  $-1.5$  and  $-0.9$  for GFR  $<10$ ,  $10$ – $25$ ,  $25$ – $50$  and  $>50$  mL/min/1.73 m<sup>2</sup> [27, 45]. Even more striking is the correlation between growth impairment and age at the time of enrolment. The average HtSDS in infants (age 0–2 years)

and young children was  $-2.33$  and  $-1.65$ , respectively, whereas it was only  $-0.93$  for adolescents [27]. This is not unexpected, considering that one-third of total growth occurs in the first 2 years of life, so infants with CKD are at a great risk of severe growth retardation with a serious long-term impact on final height [21, 46, 47]. In children with CKD the risk factors that contribute to impaired growth include: malnutrition, metabolic acidosis, mineral and bone disorders, anaemia, and fluid and electrolyte abnormalities [48–51]. However, especially after infancy and early childhood, growth failure is mainly due to disturbances in growth hormone (GH) metabolism and its main mediator, insulin-like growth factor-I (IGF-I) [52, 53]. In fact, in infants and young children, growth is principally dependent on nutrition, which has a much greater impact on growth than the GH-IGF-I axis [52]. Therefore, inadequate nutrition (due to anorexia or vomiting) appears to be the most important factor contributing to growth impairment at that age and maximizing caloric intake to at least 80% of requirements has been found to effectively improve growth in children who developed CKD as infants [52, 53]. Treatment over 2 years with recombinant human growth hormone (rhGH) has been shown to be effective without any major adverse effects [45, 54–56]. A consensus paper on the use of rhGH in CKD recommends that all children with HtSDS  $<3$ rd percentile or with height velocity standard deviation score  $<-2$  SD should be treated with rhGH after metabolic and nutritional abnormalities have been corrected [57]. Furthermore, the 2005 Kidney Disease Outcomes Quality Initiative (KDOQI) Clinical Practice Guidelines for Bone Metabolism and Disease in Children With Chronic Kidney Disease suggest avoiding rhGH therapy in children with poorly controlled mineral and bone disease [58]. In summary, even though rhGH therapy is unavoidable in most cases, an effective management of growth impairment in children with CKD must take into account all the nutritional and metabolic aspects of this disease.

### Chronic kidney disease–mineral and bone disorder

Chronic kidney disease–mineral and bone disorder (CKD-MBD) is a systemic disorder of mineral and bone metabolism due to CKD that is defined by the presence of one or a combination of the following findings: abnormalities in calcium, phosphorus, parathyroid hormone (PTH) or vitamin D metabolism; abnormalities in bone histology, linear growth, or strength; vascular or other soft tissue calcifications [59]. Renal osteodystrophy is an aspect of CKD-MBD that refers only to bone pathology. A prompt and effective management of mineral and bone disorders of CKD during childhood is of utmost importance. In fact, changes in calcium and phosphorus metabolism can significantly alter bone

remodelling and somatic growth. Optimization of bone health, growth and final adult height must be a focus of CKD management in children [59, 60]. Furthermore, paediatric nephrologists must be aware that an effective treatment of CKD-MBD affects the progression of cardiovascular disease, as phosphate is also a strong vascular toxin either in its own right or through its effect on PTH and fibroblast growth factor 23 [61, 62]. Despite international guidelines for the management of CKD-MBD [58], many patients still have a poorly controlled mineral metabolism, especially in the later stages of CKD. This is shown in a report of data collected by the International Pediatric Peritoneal Dialysis Network on 900 children worldwide, where PTH levels were over five times above the upper limit of normal values in ~50% of the patients. The highest levels were associated with higher phosphate and lower calcium levels. Phosphate control begins with dietary restriction. Thus, it is advisable for nephrologists to work closely with a specialized dietician from the very early stages of CKD. However, dietary restriction is very rarely adequate and phosphate binders become necessary even earlier than in adult patients [62–64], due especially to the unpleasant taste of this medication and the need for its ingestion at every meal. In general, the goal of therapy is to normalize mineral metabolism with the aim of improving growth and bone strength and, at the same time, reducing bone deformities and minimizing the progression of extra-skeletal calcification [65, 66].

### Anaemia

Anaemia is a common complication in children with CKD causing many adverse clinical consequences, including poor quality of life, depressed neurocognitive ability, reduced exercise capacity and progression of cardiovascular risk factors, such as left ventricular hypertrophy (LVH) [67–70]. In adult patients with CKD, the diagnosis of anaemia is made and further evaluation warranted when haemoglobin concentration is <13.5 g/dL in men and <12 g/dL in women [71]. On the other hand, the diagnosis of anaemia in children with CKD is not as straightforward. The National Kidney Foundation KDOQI (NFK-KDOQI) clinical practice guidelines use reference data from National Health and Nutrition Examination Survey (NHANES) III to define normal values in the paediatric population and recommend initiating an evaluation for anaemia when haemoglobin levels fall below the age-specific and sex-specific 5th percentile value [71–73]. Anaemia increases in prevalence with advancing stages of CKD. Data from the North American Pediatric Renal Trials and Collaborative Studies (NAPRTCS) show that the prevalence of anaemia in children is 73% at CKD stage III, 87% at stage IV and >93% at stage V [72, 74]. Anaemia of CKD is the result of many interacting factors, but decreased production of erythropoietin by the unhealthy kidney and iron dysregulation (including iron deficiency and iron-restricted erythropoiesis) are the primary defects [75–78]. Treatment with recombinant human erythropoietin (rHuEPO) is safe and effective, both in children with conservatively treated CKD and in those on maintenance dialysis [79, 80]. As in adults, the goal of this treatment is to achieve target haemoglobin levels of approximately 11 g/dL or slightly greater. Evidence shows that, both in adult and children, haemoglobin levels >13 g/dL are not associated with improved patient outcomes (including lower mortality, less frequent hospitalization, and less severe LVH) [71]. Interestingly enough, the dosing requirements of rHuEPO usually differ markedly between children and adults. Data from NAPRTCS show that, to achieve and maintain target haemoglobin levels, young children require higher rHuEPO doses than adults, ranging from 275 U/kg to 350 U/kg per week

for infants, to 200–250 U/kg per week for older children [81, 82]. The underlying mechanism related to the need for such high rHuEPO doses has not yet been fully understood, but is probably due to a greater amount of non-hematopoietic erythropoietin binding sites (e.g. kidney, endothelium, brain, heart, skeletal muscle and retinal cells) in children, which decreases the bio-availability of the drug at its therapeutic sites [72, 83]. Supplemental iron therapy (either oral or intravenous) is also necessary for the treatment of anaemia in children with CKD. However, normal or above-normal ferritin levels in CKD, as in many other chronic diseases, could be a marker of inflammation and may not reflect the total iron body stores [84–86].

### Hypertension

Unlike many of the complications of CKD, hypertension can be present from the earliest stages of the disease and its prevalence increases as GFR progressively declines [87, 88]. A recent work by the Chronic Kidney Disease in Children (CKiD) study group showed that hypertension was present in 54% of participants at the time of enrolment and, even more strikingly, 48% of the children had high blood pressure (BP) levels despite the use of antihypertensive medications, which rarely included renin-angiotensin-aldosterone system inhibitors (RAAS-I). Interestingly enough, when BP was measured with a 24-h ambulatory BP monitoring (ABPM), children with CKD showed higher systolic and diastolic variability and lower heart rate variability compared with children without hypertension with CKD. These factors represent potential precursors for cardiovascular morbidity in adults [89]. Moreover, 38% of the CKiD cohort had so-called *masked hypertension* (normal office BP but elevated ambulatory BP), which is another known risk factor for LVH [88, 89]. Studies performed in adults have clearly demonstrated that an effective control of BP reduces not only cardiovascular morbidity and mortality, but also the rate of progression of CKD [89–91]. Similarly, the *renoprotective* effect of RAAS-I, especially for proteinuric CKD patients, is now considered an unquestionable fact [91]. In the paediatric population, the ESCAPE trial of 385 children with CKD showed that patients randomly assigned to intensified BP control (BP <50th percentile) had a 35% relative risk reduction in reaching the primary endpoint of a decline of 50% in the GFR or ESRD compared with those in the conventional BP control group (BP 50th–90th percentile). All patients were treated with ramipril and, when needed, other antihypertensive medications that did not target the renin-angiotensin system were added in order to achieve targeted BP control [90, 92].

In summary, data from CKiD and other studies show that underdiagnosis and inadequate control of BP occurs in children with CKD. To improve the recognition of hypertension in paediatric CKD patients, a 24-h ABPM monitoring should be performed whenever possible and the use of RAAS-I should be part of an effective antihypertensive medication management, especially in children with proteinuric disease.

### Cardiovascular complications and death

It is well known that adults with CKD have significantly increased rates of cardiovascular morbidity and mortality compared with the general population [61, 93, 94]. However, increased cardiovascular risk is not unique to adults with CKD and several reports confirm that cardiovascular disease (CVD) is the leading cause of death also in the paediatric CKD population, with a risk 1000 times higher in the ESRD group compared with the age-matched non-CKD population [87, 95, 96]. The American Heart

Association's guidelines for cardiovascular risk reduction in high-risk paediatric patients classified children with CKD in the highest risk group for the development of CVD, alongside individuals with homozygous familial hypercholesterolaemia, diabetes mellitus type 1, heart transplantation or coronary aneurisms due to Kawasaki disease [97].

Epidemiological and clinical studies have provided evidence that cardiovascular anomalies begin early in the course of renal failure, irrespective of the age of onset, and rapidly progress when dialysis is initiated [95, 98]. CVD in the CKD population ensues from a combination of traditional (e.g. hypertension, dyslipidaemia, abnormal glucose metabolism and obesity) and CKD-related risk factors (e.g. increased calcium-phosphorus product, hyperparathyroidism and anaemia) [87]. As CKD and dialysis are relatively uncommon in childhood, large multi-centre and longitudinal studies are difficult to perform. Consequently, establishing and predicting the cardiovascular risk in this population is even more difficult [95, 99]. What is well known is that the cardiovascular causes of mortality are slightly different in children with CKD compared with adults with CKD. Adult cardiovascular deaths are mainly determined by coronary artery disease and congestive heart failure, while the leading causes of cardiac death in children with CKD are arrhythmias, valve diseases, cardiomyopathy and cardiac arrest [61, 94]. The difference between the two populations may in part be attributed to the lower prevalence of classic risk factors for atherosclerosis in children with CKD.

From a pathophysiologic point of view, all the cardiovascular abnormalities that occur in adults with CKD are also present, to some extent, in children with CKD. As in adults, endothelial dysfunction appears early in the course of renal disease and has been observed in children with CKD undergoing conservative therapy as well as in children on dialysis [99, 100]. Arterial

stiffening due to intimal calcification is commonly found in older patients with ESRD and is associated with classic risk factors for atherosclerosis, such as age, diabetes mellitus, smoking, high low-density lipoprotein cholesterol levels and inflammation [99]. On the other hand, diffuse and non-occlusive arterial stiffening found in children and young adults with ESRD is more often due to medial calcification and is strongly associated with uraemia-related specific factors, such as hypertension, long-term dialysis and high serum phosphate levels [99–101]. Furthermore, most studies show that LVH is the most common cardiac abnormality in children with CKD, and it develops even when CKD is mild and progresses as kidney function declines [87, 102]. This remodelling causes firstly a predominantly diastolic dysfunction and ultimately leads to systolic dysfunction and cardiac failure. LVH influences also the conductive properties of the myocardium and exacerbates the risk of dangerous arrhythmias [96, 103].

In conclusion, even though classic risk factors for atherosclerosis are less prevalent in children than in adults with CKD, markers of subclinical cardiovascular damage are present also in paediatric patients [99, 104]. Several modifiable risk factors, including hyperphosphataemia, hyperparathyroidism, anaemia and hypertension, independently predict the presence of cardiovascular abnormalities in these cases. An effective control of these non-traditional risk factors of CVD could improve the survival and the future global health of these patients.

### Concluding remarks

CKD is a sly disease. Although relatively uncommon in children, CKD can be a devastating illness with many long-term consequences (Figure 3). In fact, the mortality rate for children with ESRD receiving dialysis therapy is 30–150 times higher than in the general paediatric population and the life expectancy for a




Fig. 3. Clinical complications of CKD: a double perspective. The picture shows the correspondence between clinical features and complications of CKD with onset during childhood (left, top) and the relative consequences in adult life (right, top). On the other hand, clinical and laboratory findings of kidney disease in an adult (right, bottom) may find an explanation in kidney functional and/or structural abnormalities that already existed during infancy and childhood (left, bottom) but that may have been missed or underdiagnosed because of being clinically silent. Therefore, nephrologists, should have a *global vision* of their patients, regardless of whether the patient with CKD is a child or an adult: the first with a look towards the future, the other to the past. To underline this aspect, each box on the left side of the picture corresponds to one on the right side, as highlighted by the colour code. CKD, chronic kidney disease; GH-IGF-I, growth hormone and insulin-like growth factor I; LVH, left ventricular hypertension; CKD-MBD, chronic kidney disease–mineral and bone disorder; CV, cardiovascular; FSGS, focal segmental glomerulosclerosis.

child on dialysis is ~50 years less than a healthy child [9, 19, 105]. Kidney transplantation is characterized by a significant improvement in prognosis and is the best therapeutic option for children with ESRD. However, most of the complications of this clinical syndrome have consequences on the patients' health well before kidney function is irreversibly lost, even when it is maintained stable over time with conservative therapy.

In addition, despite similarities to the adult disease, CKD in children presents unique features and challenges that are not usually faced by adult patients and that make paediatric CKD a stand-alone nosologic entity. Nevertheless, paediatric nephrologists should be aware that complications in childhood CKD will have consequences well beyond paediatric age and influence outcomes of affected young adults with CKD (Figure 3). On the other hand, nephrologists who take care of young adults with CKD or adults with childhood CKD should understand the unique characteristics that CKD presents in children, especially the aetiology, in order to significantly ameliorate their patients' care (Figure 3).

In summary, nephrologists, whether caring for children or for adults with CKD, should have a *global vision* of their patients: the first with a look towards the future, the other to the past.

## Acknowledgements

This article is supported by funding from the Meyer Children's Hospital Foundation and from the Fondazione AMARTI to P.R.

## Conflict of interest statement

The results presented in this review have not been published previously in whole or part. All the authors declare no conflict of interests.

## References

- Lysaght MJ. Maintenance dialysis population dynamics: current trends and long-term implications. *J Am Soc Nephrol* 2002; 13: 37–40
- United States Renal Data System. *USRDS 2013 Annual Data Report: Atlas of Chronic Kidney Disease and End-Stage Renal Disease in the United States*. National Institutes of Health. Bethesda, MD: National Institute of Diabetes and Digestive and Kidney Diseases, 2013
- Schaefer B, Wühl E. Educational paper: progression in chronic kidney disease and prevention strategies. *Eur J Pediatr* 2012; 171: 1579–1588
- Schieppati A, Remuzzi G. Chronic renal diseases as a public health problem: epidemiology, social, and economic implications. *Kidney Int Suppl* 2005; 98: S7–S10
- Brück K, Stel VS, Fraser S et al. Translational research in nephrology: chronic kidney disease prevention and public health. *Clin Kidney J* 2015; 8: 647–655
- Kidney Disease: Improving Global Outcomes (KDIGO) CKD work group. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int Suppl* 2013; 3: 1–150
- Kidney Disease: Improving Global Outcomes (KDIGO) Glomerulonephritis Work Group. KDIGO clinical practice guideline for glomerulonephritis. *Kidney Int Suppl* 2012; 2: 139–274
- Wong CS, Warady BA. Epidemiology, etiology, and course of chronic kidney disease in children. 2015, [www.uptodate.com](http://www.uptodate.com) (February 2016, date last accessed)
- Warady BA, Chadha V. Chronic kidney disease in children: the global perspective. *Pediatr Nephrol* 2007; 22: 1999–2009
- Esbjörner E, Berg U, Hansson S. Epidemiology of chronic renal failure in children: a report from Sweden 1986–1994. Swedish Pediatric Nephrology Association. *Pediatr Nephrol* 1997; 11: 438
- Harambat J, van Stralen KJ, Kim JJ et al. Epidemiology of chronic kidney disease in children. *Pediatr Nephrol* 2012; 27: 363–373
- Ardissino G, Daccò V, Testa S et al. Epidemiology of chronic renal failure in children: data from the Italkid project. *Pediatrics* 2003; 111 (4 Pt 1): e382
- ESPN/ERA-EDTA registry. European registry for children on renal replacement therapy. [www.espn-reg.org/index.jsp](http://www.espn-reg.org/index.jsp) (February 2016, date last accessed)
- NAPRTCS: 2008 Annual Report. Rockville, MD: EMMES, 2008
- Saran R, Li Y, Robinson B et al. US Renal Data System 2014 annual data report: epidemiology of kidney disease in the United States. *Am J Kidney Dis* 2015; 66 (1 Suppl 1): S7, S1
- Australia and New Zealand Dialysis and Transplant Registry. *The 28th annual report. 2005 report-data to 2004*. 2005. <http://www.anzdata.org> (February 2016, date last accessed)
- Wong CJ, Moxey-Mims M, Jerry-Fluker J et al. CKiD (CKD in children) perspective cohort study: a review of current findings. *Am J Kidney Dis* 2012; 60: 1002–1011
- Lagomarsimo E, Valenzuela A, Cavagnaro F et al. Chronic renal failure in pediatrics 1996. Chilean survey. *Pediatr Nephrol* 1999; 13: 288
- U.S. renal data system, USRDS 2005. *Annual data report: Atlas of end-stage renal disease in the United States*, National Institutes of Health. Bethesda, MD: National Institute of Diabetes and Digestive and Kidney Diseases, 2005
- Baum M. Overview of chronic kidney disease in children. *Curr Opin Pediatr* 2010; 22: 158–160
- Harambat J, Bonthuis M, van Stralen KJ et al. Adult height in patients with advanced CKD requiring renal replacement therapy during childhood. *Clin J Am Soc Nephrol* 2014; 9: 92–99
- Coresh J, Astor BC, Greene T et al. Prevalence of chronic kidney disease and decreased kidney function in the adult US population: third National Health and Nutrition Examination Survey. *Am J Kidney Dis* 2003; 41: 1–12
- De Vecchi AF, Dratwa M, Wiedmann ME. Healthcare systems and end-stage renal disease: an international review—costs and reimbursement of ESRD therapies. *N Eng J Med* 1999; 14: 31–41
- Moosa MR, Kidd M. The dangers of rationing dialysis treatment: the dilemma facing a developing country. *Kidney Int* 2006; 70: 1107–1114
- Hoy WE. Renal disease in Australian aboriginals. *Med J Aust* 1996; 165: 126
- Vivante A, Hildebrandt F. Exploring the genetic basis of early-onset chronic kidney disease. *Nat Rev Nephrol* 2016; 12: 133–146
- Smith JM, Stablein DM, Munoz R et al. Contributions of the Transplant Registry: The 2006 Annual Report of the North American Pediatric Renal Trials and Collaborative Studies (NAPRTCS). *Pediatr Transplant* 2007; 11: 366–373
- Cain JE, Di Giovanni V, Smeeton J et al. Genetics of renal hypoplasia: insights into the mechanisms controlling nephron endowment. *Pediatr Res* 2010; 68: 91–98
- Becherucci F, Lazzeri E, Lasagni L et al. Renal progenitors and childhood: from development to disorders. *Pediatr Nephrol* 2014; 29: 711–719

30. Black MJ, Sutherland MR, Gubhaju L et al. When birth comes early: effects on nephrogenesis. *Nephrology* 2013; 18: 180
31. Schreuder MF. Safety in glomerular numbers. *Pediatr Nephrol* 2012; 27: 1881
32. Ding W, Cheung WW, Mak RH. Impact of obesity on kidney function and blood pressure in children. *World J Nephrol* 2015; 4: 223–229
33. Tullus K. Is there an obesity-related epidemic of CKD starting already in childhood? *Nephrol Dial Transplant* 2013; 28 (Suppl 4): iv114–iv116
34. Hattori S, Yosioka K, Honda M et al. The 1998 report of the Japanese National Registry data on pediatric end-stage renal disease patients. *Pediatr Nephrol* 2002; 17: 456–461
35. Devuyst O, Knoers NV, Remuzzi G et al. Rare inherited kidney diseases: challenges, opportunities, and perspectives. *Lancet* 2014; 383: 1844–1859
36. Hildebrandt F. Genetic kidney diseases. *Lancet* 2010; 375: 1287–1295
37. Gupta J, Kanetsky PA, Wuttke M et al. Genome-wide association studies in pediatric chronic kidney disease. *Pediatr Nephrol* 2015 (epub ahead of print)
38. Dummer PD, Limou S, Rosenberg AZ et al. APOL1 kidney disease risk variants: an evolving landscape. *Semin Nephrol* 2015; 35: 222–236
39. Genovese G, Friedman DJ, Ross MD et al. Association of trypanolytic Apol1 variants with kidney disease in African Americans. *Science* 2010; 329: 841–845
40. Parsa A, Kao WH, Xie D et al. APOL1 risk variants, race, and progression of chronic kidney disease. *N Engl J Med* 2013; 369: 2183–2196
41. Giglio S, Provenzano A, Mazzinghi B et al. Heterogeneous genetic alterations in sporadic nephrotic syndrome associate with resistance to immunosuppression. *J Am Soc Nephrol* 2015; 26: 230–236
42. Büscher AK, Beck BB, Melk A et al. Rapid response to cyclosporin A and favorable renal outcome in nongenetic versus genetic steroid-resistant nephrotic syndrome. *Clin J Am Soc Nephrol* 2014; 11: 245–253
43. Seikaly MG, Salhab N, Gipson D et al. Stature in children with chronic kidney disease: analysis of NAPRTCS database. *Pediatr Nephrol* 2006; 21: 793–799
44. Rodig NM, McDermott KC, Schneider MF et al. Growth in children with chronic kidney disease: a report from the chronic kidney disease in children study. *Pediatr Nephrol* 2014; 29: 1987–1995
45. Rees L. Growth hormone therapy in children with CKD after more than two decades of practice. *Pediatr Nephrol* 2015 (epub ahead of print).
46. André JL, Bourquard R, Guillemin F et al. Final height in children with chronic renal failure who have not received growth hormone. *Pediatr Nephrol* 2003; 18: 685–691
47. Mekahli D, Shaw V, Ledermann SE et al. Long-term outcome of infants with severe chronic kidney disease. *Clin J Am Soc Nephrol* 2010; 5: 10–17
48. Gat-Yablonski G, Phillip M. Nutritionally-induced catch-up growth. *Nutrients* 2015; 7: 517–551
49. Kraut JA, Madias NE. Consequences and therapy of the metabolic acidosis of chronic kidney disease. *Pediatr Nephrol* 2011; 26: 19–28
50. Farquharson C, Ahmed SF. Inflammation and linear bone growth: the inhibitory role of SOCS2 on GH/IGF-1 signalling. *Pediatr Nephrol* 2013; 28: 547–556
51. Sanchez CP, Salusky IB, Kuizon BD et al. Growth of long bones in renal failure: roles of hyperparathyroidism, growth hormone and calcitriol. *Kidney Int* 1998; 54: 1879–1887
52. Rees L, Mak RH. Nutrition and growth in children with chronic kidney disease. *Nat Rev Nephrol* 2011; 7: 615–623
53. Mak RH, Cheung W, Cone RD et al. Orexigenic and anorexigenic mechanisms in the control of nutrition in chronic kidney disease. *Pediatr Nephrol* 2005; 20: 427–431
54. Hodson EM, Willis NS, Craig JC. Growth hormone for children with chronic kidney disease. *Cochrane Database Syst Rev* 2012; 2: CD003264
55. Gil S, Vaiani E, Guercio G et al. Effectiveness of rhGH treatment on final height of renal-transplant recipients in childhood. *Pediatr Nephrol* 2012; 27: 1005–1009
56. Fine RN, Stablein D. Long-term use of recombinant human growth hormone in pediatric allograft recipients: a report of the NAPRTCS transplant registry. *Pediatr Nephrol* 2005; 20: 404–408
57. Mahan JD, Warady BA, Consensus Committee. Assessment and treatment of short stature in pediatric patients with chronic kidney disease: a consensus statement. *Pediatr Nephrol* 2006; 21: 917–930
58. Kidney Disease: Improving Global Outcomes (KDIGO) CKD-MBD Work Group. KDIGO clinical practice guideline for the diagnosis, evaluation, prevention, and treatment of chronic kidney disease–mineral and bone disorder (CKD-MBD). *Kidney Int* 2009; 76 (Suppl 113): S1–S130
59. Wesseling-Perry K, Salusky IB. Chronic kidney disease: mineral and bone disorder in children. *Semin Nephrol* 2013; 33: 169–179
60. Wesseling-Perry K. Bone disease in pediatric chronic kidney disease. *Pediatr Nephrol* 2013; 28: 569–576
61. Gansevoort RT, Correa-Rotter R, Hemmelgarn BR et al. Chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. *Lancet* 2013; 382: 339–352
62. Rees L, Shroff R. The demise of calcium-based phosphate binders—is this appropriate for children? *Pediatr Nephrol* 2015; 30: 2061–2071
63. Rees L, Azocar M, Borzych D et al. Growth in very young children undergoing chronic peritoneal dialysis. *J Am Soc Nephrol* 2011; 22: 2303–2312
64. Klaus G, Watson A, Edefonti A et al. Prevention and treatment of renal osteodystrophy in children on chronic renal failure: European guidelines. *Pediatr Nephrol* 2006; 21: 151–159
65. Block GA, Spiegel DM, Ehrlich J et al. Effects of sevelamer and calcium on coronary artery calcification in patients new to hemodialysis. *Kidney Int* 2005; 68: 1815–1824
66. Di Iorio B, Molony D, Bell C et al. Sevelamer versus calcium carbonate in incident hemodialysis patients: results of an open-label 24-month randomized clinical trial. *Am J Kidney Dis* 2013; 62: 771–778
67. Mitsnefes MM, Kimball TR, Kartal J et al. Progression of left ventricular hypertrophy in children with early chronic kidney disease: 2-year follow-up study. *J Pediatr* 2006; 149: 671–675
68. Kurella Tamura M, Vittinghoff E, Yang J et al. Anemia and risk for cognitive decline in chronic kidney disease. *BMC Nephrol* 2016; 17: 13
69. Dahlinghaus EK, Neu AM, Atkinson MA et al. Hemoglobin level and risk of hospitalization and mortality in children on peritoneal dialysis. *Pediatr Nephrol* 2014; 29: 2387–2394



70. Gerson A, Hwang W, Fiorenza J et al. Anemia and health-related quality of life in adolescents with chronic kidney disease. *Am J Kidney Dis* 2004; 44: 1017–1023
71. KDOQI: National Kidney Foundation. KDOQI clinical practice guidelines and clinical practice recommendations for anemia in chronic kidney disease. *Am J Kidney Dis* 2006; 47 (Suppl 3): S11–S145
72. Atkinson MA, Furth SL. Anemia in children with chronic kidney disease. *Nat Rev Nephrol* 2011; 7: 635–641
73. Keithi-Reddy SR, Singh AK. Hemoglobin target in chronic kidney disease: a pediatric perspective. *Pediatr Nephrol* 2009; 24: 431–434
74. Atkinson MA, Martz K, Warady BA et al. Risk for anemia in pediatric chronic kidney disease patients: a report of NAPRTCS. *Pediatr Nephrol* 2010; 25: 1699–1706
75. Ratcliffe LE, Thomas W, Glen J et al. Diagnosis and management of iron deficiency in CKD: a summary of the NICE guideline recommendations and their rationale. *Am J Kidney Dis* 2016; 67: 548–558
76. Fraenkel PG. Understanding anemia of chronic disease. *Hematology Am Soc Hematol Educ Program* 2015; 2015: 14–18
77. Goodnough LT, Nemeth E, Ganz T. Detection, evaluation, and management of iron-restricted erythropoiesis. *Blood* 2010; 116: 754–4761
78. Nangaku M, Eckardt KU. Pathogenesis of renal anemia. *Semin Nephrol* 2006; 26: 261–268
79. Warady BA, Silverstein DM. Management of anemia with erythropoietic-stimulating agents in children with chronic kidney disease. *Pediatr Nephrol* 2014; 29: 1493–1505
80. Boehm M. Early erythropoietin therapy is associated with improved growth in children with chronic kidney disease. *Pediatr Nephrol* 2007; 22: 1189–1193
81. Port RE, Mehls O. Erythropoietin dosing in children with chronic kidney disease: based on body size or on hemoglobin deficit? *Pediatr Nephrol* 2009; 24: 435–437
82. Port RE, Kiepe D, Van Guilder M et al. Recombinant human erythropoietin for the treatment of renal anaemia in children: no justification for bodyweight-adjusted dosage. *Clin Pharmacokinet* 2004; 43: 57–70
83. Uemura O, Hattori M, Hataya H et al. Pharmacokinetics of darbepoetin alfa after single, intravenous or subcutaneous administration in Japanese pediatric patients with chronic kidney disease. *Clin Exp Nephrol* 2014; 18: 932–938
84. Kalantar-Zadeh K, Kalantar-Zadeh K, Lee GH. The fascinating but deceptive ferritin: to measure it or not to measure it in chronic kidney disease? *Clin J Am Soc Nephrol* 2006; 1 (Suppl 1): S9–S18
85. Morgan HE, Holt RC, Jones CA et al. Intravenous iron treatment in paediatric chronic kidney disease patients not on erythropoietin. *Pediatr Nephrol* 2007; 22: 1963–1965
86. Albaramki J, Hodson EM, Craig JC et al. Parenteral versus oral iron therapy for adults and children with chronic kidney disease. *Cochrane Database Syst Rev* 2012; 1: CD007857
87. Mitsnefes MM. Cardiovascular disease in children with chronic kidney disease. *J Am Soc Nephrol* 2012; 23: 578–585
88. Flynn JT, Mitsnefes M, Pierce C et al. Blood pressure in children with chronic kidney disease: a report from the Chronic Kidney Disease in Children study. *Hypertension* 2008; 52: 631–637
89. Mitsnefes M, Flynn J, Cohn S et al. Masked hypertension associates with left ventricular hypertrophy in children with CKD. *J Am Soc Nephrol* 2010; 21: 137
90. VanDeVoorde RG, Misnefes MM. Hypertension and CKD. *Adv Chronic Kidney Dis* 2011; 18: 355–361
91. Hsu TW, Liu JS, Hung SC et al. Renoprotective effect of renin-angiotensin-aldosterone system blockade in patients with predialysis advanced chronic kidney disease, hypertension, and anemia. *JAMA Intern Med* 2014; 174: 347–354
92. Wühl E, Trivelli A, Picca S et al. Strict blood-pressure control and progression of renal failure in children. *N Engl J Med* 2009; 361: 1639–1650
93. Sarnak MJ, Levey AS, Schoolwerth AC et al. Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. *Hypertension* 2003; 42: 1050–1065
94. Anavekar NS, Pfeffer MA. Cardiovascular risk in chronic kidney disease. *Kidney Int Suppl* 2004; 92: S11–S15
95. Shroff R, Dégi A, Kerti A et al. Cardiovascular risk assessment in children with chronic kidney disease. *Pediatr Nephrol* 2013; 28: 875–884
96. Safder O, Al sharif S, Kari JA. Pediatric CKD and cardiovascular disease. *Cardiovasc Hematol Disord Drug Targets* 2014; 14: 177–184
97. Kavey RE, Allada V, Daniels SR et al. Cardiovascular risk reduction in high-risk pediatric patients: a scientific statement from the American Heart Association Expert Panel on Population and Prevention Science; the Councils on Cardiovascular Disease in the Young, Epidemiology and Prevention, Nutrition, Physical Activity and Metabolism, High Blood Pressure Research, Cardiovascular Nursing, and the Kidney in Heart Disease; and the Interdisciplinary Working Group on Quality of Care and Outcomes Research: endorsed by the American Academy of Pediatrics. *Circulation* 2006; 114: 2710–2738
98. Paoli S, Mitsnefes MM. Coronary artery calcification and cardiovascular disease in children with chronic kidney disease. *Curr Opin Pediatr* 2014; 26: 193–197
99. Lilien MR, Groothoff JW. Cardiovascular disease in children with CKD or ESRD. *Nat Rev Nephrol* 2009; 5: 229–235
100. Tripepi G, Mallamaci F, Zoccali C. Inflammation markers, adhesion molecules, and all-cause and cardiovascular mortality in patients with ESRD: searching for the best risk marker by multivariate modeling. *J Am Soc Nephrol* 2005; 16 (Suppl 1): S83–S88
101. Briet M, Boutouyrie P, Laurent S et al. Arterial stiffness and pulse pressure in CKD and ESRD. *Kidney Int* 2012; 2: 388–400
102. Matteucci MC, Wühl E, Picca S et al. Left ventricular geometry in children with mild to moderate chronic renal insufficiency. *J Am Soc Nephrol* 2006; 17: 218–226
103. Matsushita K, Sang Y, Ballew SH et al. Subclinical atherosclerosis measures for cardiovascular prediction in CKD. *J Am Soc Nephrol* 2015; 26: 439–447
104. Yao Q, Pecoits-Filho R, Lindholm B et al. Traditional and non-traditional risk factors as contributors to atherosclerotic cardiovascular disease in end-stage renal disease. *Scand J Urol Nephrol* 2004; 38: 405–416
105. McDonald SP, Craig JC. Long-term survival of children with end-stage renal disease. *N Engl J Med* 2004; 350: 2654–2662