

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Assessing outcomes after adrenalectomy for unilateral primary aldosteronism

### This is the author's manuscript

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1717651> since 2019-11-25T23:46:21Z

*Published version:*

DOI:10.1016/j.surg.2019.06.027

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Dear Editor,

We read with interest the article “Clinical outcomes after surgery for primary aldosteronism: evaluation of the PASO-investigators’ consensus criteria within a worldwide cohort of patients” by Vorselaars and colleagues.<sup>1</sup> From their retrospective study the authors concluded that 16% of their patients were incorrectly or debatably classified using the criteria of the PASO consensus.<sup>2</sup> The authors suggested that this was mainly due to using a cut-off of  $\geq 20$  mmHg to classify a clinically relevant change in systolic blood pressure. In addition, the authors suggested that use of percentages instead of absolute values in change of antihypertensive medication (daily defined doses) contributed to this incorrect classification.

We would like to clarify a number of points:

- 1.** The use of office blood pressure measurements are notorious for their variability. This variability may even be exaggerated in the study because blood pressure (as main outcome parameter) was not measured in a standardized way. This is clearly illustrated by the fact that in 27% of the patients the preoperative blood pressure level was based on only one blood pressure value and this was increased to 50% of patients for assessment of postoperative blood pressure. The reliability and accuracy of the changes in blood pressure in this study are therefore questionable.
- 2.** Related to the previous point, a cut-off change in systolic blood pressure of 10 mmHg, as recommended by the authors, cannot be considered sufficiently robust to estimate partial clinical success. The cut-off level of 20 mmHg was selected from the differences between blood pressure levels that define blood pressure categories of hypertension<sup>3</sup> and used by the PASO group of experts to define criteria for clinical outcomes with the structured Delphi method. The authors argue that 10 mmHg is more appropriate than 20 mmHg, based on the associated decrease in cardiovascular morbidity and mortality from studies in patients with primary hypertension. However, data from primary hypertensives cannot be translated to patients with primary aldosteronism in whom a complete biochemical response to treatment is essential<sup>4</sup> (which was not assessed by Vorselaars et al.<sup>1</sup>) and who have an increased cardiovascular risk compared with patients with primary hypertension.<sup>5</sup>
- 3.** The PASO consensus used an absolute value of  $\geq 0.5$  DDD to define a relevant change in medication and not a 50% change. Thus, the authors have incorrectly applied the PASO criteria to evaluate outcomes. According to table 4, 15 of their 26 patients had a decrease in DDD of  $>0.5$  DDD, so these 15 patients should be classified as partial and not as absent clinical success.
- 4.** Finally, the heterogeneous approach for the identification of surgically-treatable patients (with reliance on nonfunctional imaging in 36% of patients) and the considerable deviation from the PASO recommendations for the timeframe of follow up in 68% of patients underlines further the inappropriateness of comparing data from Vorselaars et al.<sup>1</sup> with the PASO study.<sup>2</sup>

*Tracy Ann Williams PhD<sup>1,2</sup>*  
*Jacques WM Lenders MD PhD<sup>3,4</sup>*  
*Martin Reincke MD<sup>1</sup>*

*<sup>1</sup>Medizinische Klinik und Poliklinik IV, Klinikum der Universität, LMU München, Munich, Germany and Department of Medicine*

*<sup>2</sup>Division of Internal Medicine and Hypertension, Department of Medical Sciences, University of Turin, Turin, Italy*

<sup>3</sup>Radboud University Medical Center, Nijmegen, The Netherlands

<sup>4</sup>Department of Medicine III, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany

Corresponding author TAW: Email: [Tracy.Williams@med.uni-muenchen.de](mailto:Tracy.Williams@med.uni-muenchen.de)

### Disclosure statement

The authors declare no conflicts of interest

### Funding source

The work of the authors is supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No [694913] to M Reincke) and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Projektnummer: 314061271-TRR 205 to TA Williams, JWM Lenders and M Reincke.

### References

- 1) Vorselaars WMCM, Beek DV, Postma EL, Spiering W, Borel Rinkes IHM, Valk GD, Vriens MR; International CONNsorium study group. Clinical outcomes after surgery for primary aldosteronism: Evaluation of the PASO-investigators' consensus criteria within a worldwide cohort of patients. *Surgery*. 2019 Apr 30. pii: S0039-6060(19)30063-7. doi: 10.1016/j.surg.2019.01.031
- 2) Williams TA, Lenders JWM, Mulatero P, Burrello J, Rottenkolber M, Adolf C, Satoh F, Amar L, Quinkler M, Deinum J, Beuschlein F, Kitamoto KK, Pham U, Morimoto R, Umakoshi H, Prejbisz A, Kocjan T, Naruse M, Stowasser M, Nishikawa T, Young WF Jr, Gomez-Sanchez CE, Funder JW, Reincke M; Primary Aldosteronism Surgery Outcome (PASO) investigators. Outcomes after adrenalectomy for unilateral primary aldosteronism: an international consensus on outcome measures and analysis of remission rates in an international cohort. *Lancet Diabetes Endocrinol*. 2017 Sep;5(9):689-699. doi: 10.1016/S2213-8587(17)30135-3
- 3) Mancia G, Fagard R, Narkiewicz K, Redón J, Zanchetti A, Böhm M, Christiaens T, Cifkova R, De Backer G, Dominiczak A, Galderisi M, Grobbee DE, Jaarsma T, Kirchhof P, Kjeldsen SE, Laurent S, Manolis AJ, Nilsson PM, Ruilope LM, Schmieder RE, Sirnes PA, Sleight P, Viigimaa M, Waeber B, Zannad F; Task Force Members. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of

Cardiology (ESC). J Hypertens. 2013 Jul;31(7):1281-357. doi: 10.1097/01.hjh.0000431740.32696

- 4) Hundemer GL, Curhan GC, Yozamp N, Wang M, Vaidya A. Cardiometabolic outcomes and mortality in medically treated primary aldosteronism: a retrospective cohort study. *Lancet Diabetes Endocrinol.* 2018 Jan;6(1):51-59. doi: 10.1016/S2213-8587(17)30367-4.
- 5) Monticone S, D'Ascenzo F, Moretti C, Williams TA, Veglio F, Gaita F, Mulatero P. Cardiovascular events and target organ damage in primary aldosteronism compared with essential hypertension: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol.* 2018 Jan;6(1):41-50. doi: 10.1016/S2213-8587(17)30319-4