

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Sign-changing bubble-tower solutions to fractional semilinear elliptic problems

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1725912> since 2020-01-30T13:13:02Z

Published version:

DOI:10.3934/dcds.2019268

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

SIGN-CHANGING BUBBLE-TOWER SOLUTIONS TO FRACTIONAL SEMILINEAR ELLIPTIC PROBLEMS

GABRIELE CORA

Dipartimento di Matematica "G. Peano"
Università di Torino
via Carlo Alberto 10 – 10123 Torino, Italy

ALESSANDRO IACOPETTI

Dipartimento di Matematica
Università di Roma "La Sapienza"
P.le Aldo Moro 5 – 00185 Roma, Italy

ABSTRACT. We study the asymptotic and qualitative properties of least energy radial sign-changing solutions to fractional semilinear elliptic problems of the form

$$\begin{cases} (-\Delta)^s u = |u|^{2_s^* - 2 - \varepsilon} u & \text{in } B_R, \\ u = 0 & \text{in } \mathbb{R}^n \setminus B_R, \end{cases}$$

where $s \in (0, 1)$, $(-\Delta)^s$ is the s -Laplacian, B_R is a ball of \mathbb{R}^n , $2_s^* := \frac{2n}{n-2s}$ is the critical Sobolev exponent and $\varepsilon > 0$ is a small parameter. We prove that such solutions have the limit profile of a “tower of bubbles”, as $\varepsilon \rightarrow 0^+$, i.e. the positive and negative parts concentrate at the same point with different concentration speeds. Moreover, we provide information about the nodal set of these solutions.

1. INTRODUCTION

Let $s \in (0, 1)$, let $n \in \mathbb{N}$ be such that $n > 2s$ and let $\Omega \subset \mathbb{R}^n$ be a bounded smooth domain. Consider the following non-local elliptic problem

$$\begin{cases} (-\Delta)^s u = f(u) & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases} \quad (1.1)$$

where $(-\Delta)^s$ is the s -Laplacian, $f(u) = |u|^{2_s^* - 2 - \varepsilon} u$ or $f(u) = \varepsilon u + |u|^{2_s^* - 2} u$ for $n > 6s$, $\varepsilon > 0$ is a small parameter and $2_s^* := \frac{2n}{n-2s}$ is the critical exponent for the fractional Sobolev embedding.

In the recent paper [7] the authors studied the asymptotic properties of least energy positive solutions to Problem (1.1), i.e. positive solutions u_ε such that $\|u_\varepsilon\|_s^2 \rightarrow S_s^{\frac{n}{2s}}$, as $\varepsilon \rightarrow 0^+$, where $\|\cdot\|_s$ is the standard seminorm in $H^s(\mathbb{R}^n)$ and S_s is the best fractional Sobolev constant. They proved, in the case of the spectral fractional Laplacian, that such solutions concentrate and blow-up at some point $x_0 \in \Omega$, providing also information about the blow-up speed with respect to ε . Their result is hence the fractional counterpart of the classical results of Han and Rey (see [12, 23]) for the Laplacian.

2010 *Mathematics Subject Classification.* Primary: 35J61, 35B05; Secondary: 35B40, 35B44.

Key words and phrases. Fractional semilinear elliptic equations, critical exponent, nodal regions, sign-changing radial solutions, asymptotic behavior.

Acknowledgements. Research partially supported by the project ERC Advanced Grant 2013 n. 339958 Complex Patterns for Strongly Interacting Dynamical Systems COMPAT, and by Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

Motivated by that, it is natural to ask whether is possible or not to extend to the fractional framework analogous results about the asymptotic behavior of least energy sign-changing solutions to almost critical and critical semilinear elliptic problems for the Laplacian (see [2, 3, 14, 15, 16, 22]).

At first glance the answer seems to be positive, but differently from the case of constant-sign solutions, several difficulties arise when studying the qualitative properties of sign-changing solutions. Indeed, in view of the non-local interactions between the nodal components, we cannot take benefit from the fractional moving plane method (see [5]), and the strong maximum principle does not work properly (see [8, Sect. 1]). Moreover, when considering least energy sign-changing solutions, i.e. sign-changing solutions u_ε to (1.1) such that $\|u_\varepsilon\|_s^2 \rightarrow 2S_s^{\frac{n}{2s}}$, as $\varepsilon \rightarrow 0^+$, we cannot establish by mere energetic arguments, neither by a Morse-index approach, the number of nodal components. In the local case it is well known that they possess exactly two nodal regions, since each nodal component carries the energy $S_1^{\frac{n}{2}}$ (see [2, 3]). In the fractional case we can only say that both the positive and the negative part globally carry the same energy $S_s^{\frac{n}{2s}}$, when $\varepsilon \rightarrow 0^+$, but this does not hold true in general for each individual nodal component and causes many troubles when performing the asymptotic analysis.

In our contribution [8] we tackled the case of least energy radial sign-changing solutions to Problem (1.1) in a ball, when $f(u) = \varepsilon u + |u|^{2_s^*-2}u$ is the critical nonlinearity and $n > 6s$. In the spirit of the pioneering papers [10, 11], we showed that these solutions change sign at most twice and exactly once when s is close to 1. Moreover, when $s > \frac{1}{2}$, we proved that they behave like a tower of two bubbles as $\varepsilon \rightarrow 0^+$, namely, the positive and the negative part blow-up and concentrate at the same point (which is the center of the ball) with different speeds. Nevertheless, we needed to assume that these solutions change sign exactly once to determine which one between the positive and the negative part blew-up faster (see [8, Sect.1]).

We point out that for $2s < n \leq 6s$, according to a classical result of Atkinson, Brezis, and Peletier (see [1]), radial sign-changing solutions in a ball may not exist when $\varepsilon > 0$ is close to zero, while they do exist for $n > 6s$ (see [8, Theorem 3.7]).

In this paper we consider slightly subcritical nonlinearities $f(u) = |u|^{2_s^*-2-\varepsilon}u$, and we extend the results of [8] to all $s \in (0, 1)$ without any extra assumption. The same proofs work also in the case of critical nonlinearities with minor modifications. The main result of our paper is the following:

Theorem 1.1. *Let $s \in (0, 1)$ and let $n > 2s$. Let $(u_\varepsilon)_\varepsilon$ be a family of least energy radial sign-changing solutions to*

$$\begin{cases} (-\Delta)^s u = |u|^{2_s^*-2-\varepsilon}u & \text{in } B_R, \\ u = 0 & \text{in } \mathbb{R}^n \setminus B_R, \end{cases} \quad (1.2)$$

where B_R is the euclidean ball of radius $R > 0$ centered at the origin. Assume without loss of generality that $u_\varepsilon(0) > 0$ and set $M_\varepsilon^\pm := |u_\varepsilon^\pm|_\infty$. Then, as $\varepsilon \rightarrow 0^+$ it holds that:

- (i) $M_\varepsilon^\pm \rightarrow +\infty$,
- (ii) $\frac{M_\varepsilon^+}{M_\varepsilon^-} \rightarrow +\infty$,
- (iii) $|x_\varepsilon| \rightarrow 0$, where $x_\varepsilon \in B_R$ is any point such that $u_\varepsilon(x_\varepsilon) = M_\varepsilon^+$,
- (iv) the rescaled function

$$\tilde{u}_\varepsilon(x) := \frac{1}{M_\varepsilon^+} u_\varepsilon \left(\frac{x}{(M_\varepsilon^+)^{\frac{2}{n-2s-\frac{\varepsilon}{2s}}}} \right), \quad x \in \mathbb{R}^n,$$

converges in $C_{loc}^{0,\alpha}(\mathbb{R}^n)$, for some $\alpha \in (0, 1)$, to the fractional standard bubble $U_{0,\mu}$ in \mathbb{R}^n centered at the origin and such that $U_{0,\mu}(0) = 1$,
(v) if $s \in (\frac{1}{2}, 1)$ then $|y_\varepsilon| \rightarrow 0$, where $y_\varepsilon \in B_R$ is any point such that $|u_\varepsilon(y_\varepsilon)| = M_\varepsilon^-$.

Theorem 1.1 establishes the first existence result of sign-changing bubble-tower solutions for non-local semilinear elliptic problems driven by the s -Laplacian, when $s > \frac{1}{2}$. For $s \in (0, \frac{1}{2}]$ we still get that the positive and the negative part blow-up with different speeds, but for the negative part we cannot provide any information about its concentration point. From a technical point of view (see the proof of Lemma 4.3) this is due to the fractional Strauss inequality for radial functions, namely

$$\sup_{x \in \mathbb{R}^n \setminus \{0\}} |x|^{\frac{n-2s}{2}} |u(x)| \leq K_{n,s} \|u\|_s^2, \quad (1.3)$$

where $K_{n,s}$ is an explicit positive constant depending only on n, s . Indeed, as pointed out in [6, Remark 2, Remark 4], (1.3) does not hold when $s \in (0, \frac{1}{2}]$. We also stress that in view of the non-local nature of our problem the positive and negative parts are not, in general, sub or super solutions to Problem (1.1) in their domain of definition, so it seems quite hard to overcome this difficulty by applying scaling arguments to $u_\varepsilon^+, u_\varepsilon^-$ separately.

On the other hand, as proved in [3] for the Laplacian, if the blow-up speeds of $u_\varepsilon^+, u_\varepsilon^-$ are comparable then they must concentrate at two separate points. Therefore, in view of (ii), we believe that also for $s \in (0, \frac{1}{2}]$ the negative part concentrates at the center of the ball. We plan to investigate this question in separate paper. In addition, we think that, as done in [22] for the Laplacian, by using a Lyapunov–Schmidt reduction method it should be possible to construct sign-changing bubble-tower solutions in general bounded domains, for all $s \in (0, 1)$.

We point out that, thanks to (ii) and (iii), any global maximum point is close to the origin, when $\varepsilon > 0$ is sufficiently small. Moreover, in Lemma 4.6 we specify that any such a point belongs to the nodal component containing the origin and blows-up faster than any other extremal value achieved in the other nodal components, independently on the number of sign-changes. In the local case, by using ODE techniques, it is well known that the global maximum point is the origin and the absolute values of the extrema are ordered in a radially decreasing way. Our result allows to recover these properties, at least asymptotically, via PDE-only arguments.

In the second part of this work we study the nodal set of least energy radial sign-changing solutions to (1.2). We remark that, if u_ε is a nodal solution to (1.2) and $u_\varepsilon \geq 0$ in a subdomain $D \subset B_R$, the fractional strong maximum principle does not ensure, in general, that $u_\varepsilon > 0$ in D (see [4, Remark 4.2] and [8, Sect. 1]). In addition [19, Theorem 1.4] only grants that u_ε does not vanish in a set of positive measure. Nevertheless, combining the results of [8] with a new argument based on energy and regularity estimates, we show that for any $s \in (0, 1)$ least energy radial sign-changing solutions to (1.2) vanish only where a change of sign occurs (see Lemma 4.5, Lemma 5.2).

Finally, in Theorem 5.8 we prove that for any $s_0 \in (0, 1)$, if there exists a $L^2(B_R)$ -continuous family $\mathcal{A} = \{u_{\varepsilon,s}\}_{s \in [s_0, 1]}$ of least energy nodal radial solutions to (1.2), then every element of the family changes sign exactly once, provided that $\varepsilon > 0$ is small enough. The key ingredients of the proof are the estimates contained in [24, Theorem 1.2], and the continuity of the map $s \mapsto C_{\mathcal{M}^r(B_R)}(s, \varepsilon)$, where $C_{\mathcal{M}^r(B_R)}(s, \varepsilon)$ is the infimum of the energy over the nodal Nehari set, which is a new result of its own interest (see Proposition 5.6).

The outline of the paper is the following: in Section 2 we fix the notation and we recall some known results about the existence of sign-changing solutions to (1.2), in Section 3 we study the asymptotic behavior, as $\varepsilon \rightarrow 0^+$, of the energy levels $C_{\mathcal{M}(\Omega)}(s, \varepsilon)$ in generic bounded domains. In Section 4 we prove Theorem 1.1. Finally in Section 5 we analyze the nodal set of least energy radial sign-changing solutions to (1.2) and we prove Theorem 5.8.

2. NOTATION AND PRELIMINARY RESULTS

In this section we recall some definitions and known facts that will be used in this work.

2.1. Functional setting, standard bubbles. In this paper $(-\Delta)^s$ stands for the (restricted) s -Laplacian operator, which is formally defined as

$$(-\Delta)^s u(x) := C_{n,s} P.V. \int_{\mathbb{R}^n} \frac{u(x) - u(y)}{|x - y|^{n+2s}} dy = C_{n,s} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}^n \setminus B_\varepsilon(x)} \frac{u(x) - u(y)}{|x - y|^{n+2s}} dy,$$

where the constant $C_{n,s}$ is given by

$$C_{n,s} := \frac{2^{2s} \Gamma\left(\frac{n}{2} + s\right)}{\pi^{\frac{n}{2}} |\Gamma(-s)|}.$$

Let $s \in (0, 1)$ and let $n > 2s$. For a given smooth bounded domain $\Omega \subset \mathbb{R}^n$, we consider as a working functional space the Sobolev space

$$X_0^s(\Omega) := \{u \in H^s(\mathbb{R}^n) ; u = 0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega\},$$

endowed with the norm

$$\|u\|_s^2 := \frac{C_{n,s}}{2} \int_{\mathbb{R}^{2n}} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy,$$

and whose associated scalar product is

$$(u, v)_s := \frac{C_{n,s}}{2} \int_{\mathbb{R}^{2n}} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{n+2s}} dx dy.$$

The Sobolev space $\mathcal{D}^s(\mathbb{R}^n)$ is defined as the completion of $C_0^\infty(\mathbb{R}^n)$ with respect to the above norm. By the fractional Sobolev embedding theorem it holds that $\mathcal{D}^s(\mathbb{R}^n) \hookrightarrow L^{2_s^*}(\mathbb{R}^n)$ and $X_0^s(\Omega) \hookrightarrow L^p(\Omega)$ for all $p \in [1, 2_s^*]$, where $2_s^* = \frac{2n}{n-2s}$. The previous embeddings are continuous, and the second one is compact when $p \in [1, 2_s^*)$. The best Sobolev constant is characterized as

$$S_s := \inf_{v \in \mathcal{D}^s(\mathbb{R}^n) \setminus \{0\}} \frac{\|v\|_s^2}{|v|_{2_s^*}^2},$$

where $|\cdot|_p$ denotes the usual L^p -norm, for $p \in [1, \infty]$. To simplify the notation we will not specify the domain of integration in $|\cdot|_p$, but it will be always clear from the context that it is either \mathbb{R}^n , or a fixed bounded domain Ω , or a family of bounded domains when considering rescaled functions. The value of S_s is explicitly known (see [9]), it depends continuously on $s \in [0, 1]$, and it is achieved exactly by the family

$$\mathcal{U}_{\mu, x_0, k}(x) := k \left(\frac{\mu}{\mu^2 + |x - x_0|^2} \right)^{\frac{n-2s}{2}}, \quad \mu > 0, x_0 \in \mathbb{R}^n, k \in \mathbb{R}.$$

If we choose $k = b_{n,s}$, where

$$b_{n,s} := 2^{\frac{n-2s}{2}} \left(\frac{\Gamma\left(\frac{n+2s}{2}\right)}{\Gamma\left(\frac{n-2s}{2}\right)} \right)^{\frac{n-2s}{4s}}, \quad (2.1)$$

then the functions

$$U_{x_0, \mu}(x) := b_{n,s} \left(\frac{\mu}{\mu^2 + |x - x_0|^2} \right)^{\frac{n-2s}{2}}, \quad (2.2)$$

also known as ‘‘standard fractional bubbles’’, satisfy

$$(-\Delta)^s U_{x_0, \mu} = U_{x_0, \mu}^{2_s^* - 1} \quad \text{in } \mathbb{R}^n \quad (2.3)$$

for all $\mu > 0, x_0 \in \mathbb{R}^n$ and

$$\|U_{x_0, \mu}\|_s^2 = |U_{x_0, \mu}|_{2_s^*}^{2_s^*}.$$

2.2. Existence of constant-sign and sign-changing solutions. Let $\Omega \subset \mathbb{R}^n$ be a smooth bounded domain and consider the problem

$$\begin{cases} (-\Delta)^s u = |u|^{2_s^* - 2 - \varepsilon} u & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases} \quad (2.4)$$

where $\varepsilon \in (0, 2_s^* - 2)$. Weak solutions to (2.4) correspond to critical points of the functional

$$I_{s, \varepsilon}(u) := \frac{1}{2} \|u\|_s^2 - \frac{1}{2_s^* - \varepsilon} |u|_{2_s^* - \varepsilon}^{2_s^* - \varepsilon}.$$

The Nehari manifold and the nodal Nehari set are, respectively, defined by

$$\begin{aligned} \mathcal{N}_{s, \varepsilon}(\Omega) &:= \{u \in X_0^s(\Omega) ; I'_{s, \varepsilon}(u)[u] = 0, u \neq 0\}, \\ \mathcal{M}_{s, \varepsilon}(\Omega) &:= \{u \in X_0^s(\Omega) ; I'_{s, \varepsilon}(u)[u^\pm] = 0, u^\pm \neq 0\}. \end{aligned}$$

Since we deal with subcritical nonlinearities, by standard variational methods we know that there exists a minimizer $u_\varepsilon \in \mathcal{N}_{s, \varepsilon}(\Omega)$ of $I_{s, \varepsilon}$, and we set

$$C_{\mathcal{N}(\Omega)}(s, \varepsilon) := \inf_{v \in \mathcal{N}_{s, \varepsilon}(\Omega)} I_{s, \varepsilon}(v).$$

Moreover, the minimizer is a weak solution to (2.4) and it is of constant sign. We also remark that, equivalently, constant-sign weak solutions to (2.4) can be found as minimizers to

$$S_{s, \varepsilon} := \inf_{v \in X_0^s(\Omega) \setminus \{0\}} \frac{\|v\|_s^2}{|v|_{2_s^* - \varepsilon}^2},$$

and the following relation holds

$$C_{\mathcal{N}(\Omega)}(s, \varepsilon) = \frac{2_s^* - 2 - \varepsilon}{2(2_s^* - \varepsilon)} S_{s, \varepsilon}^{\frac{2_s^* - \varepsilon}{2_s^* - 2 - \varepsilon}}. \quad (2.5)$$

In the case of sign-changing solutions, as proved in [28], there exists a minimizer of the energy over the nodal Nehari set, and it is a weak solution to (2.4). We refer to such solutions as least energy sign-changing (or nodal) solutions and we set

$$C_{\mathcal{M}(\Omega)}(s, \varepsilon) := \inf_{v \in \mathcal{M}_{s, \varepsilon}(\Omega)} I_{s, \varepsilon}(v).$$

Let us now turn our attention to the radial case. Taking $\Omega = B_R$, where $B_R = B_R(0)$ denotes the ball in \mathbb{R}^n of radius $R > 0$ centered at the origin, we set

$$\begin{aligned} \mathcal{N}_{s, \varepsilon}^r(B_R) &:= \{u \in X_0^s(B_R) ; u \in \mathcal{N}_{s, \varepsilon}(B_R) \text{ and } u \text{ is radially symmetric}\}, \\ \mathcal{M}_{s, \varepsilon}^r(B_R) &:= \{u \in X_0^s(B_R) ; u \in \mathcal{M}_{s, \varepsilon}(B_R) \text{ and } u \text{ is radially symmetric}\}. \end{aligned}$$

As a consequence of the fractional moving plane method (see [5]), positive solutions of (2.4) in B_R are radially symmetric and radially decreasing. In particular, it holds that

$$C_{\mathcal{N}(B_R)}(s, \varepsilon) = C_{\mathcal{N}^r(B_R)}(s, \varepsilon) := \inf_{v \in \mathcal{N}_{s, \varepsilon}^r(B_R)} I_{s, \varepsilon}(v).$$

Concerning the case of nodal solutions, arguing as in [28] we obtain least energy radial sign-changing solutions as minimizers of the energy over the radial nodal Nehari set, and as before we denote

$$C_{\mathcal{M}^r(B_R)}(s, \varepsilon) := \inf_{v \in \mathcal{M}_{s, \varepsilon}^r(B_R)} I_{s, \varepsilon}(v).$$

We point out that it is not known whether or not $C_{\mathcal{M}^r(B_R)}(s, \varepsilon)$ coincide with $C_{\mathcal{M}(B_R)}(s, \varepsilon)$, but they have the same limit when $\varepsilon \rightarrow 0^+$ (see Lemma 3.3).

3. ASYMPTOTIC ANALYSIS OF THE ENERGY LEVELS AS $\varepsilon \rightarrow 0^+$

In this section we study the asymptotic behavior as $\varepsilon \rightarrow 0^+$ of the energy levels $C_{\mathcal{N}(\Omega)}(s, \varepsilon)$, $C_{\mathcal{M}(\Omega)}(s, \varepsilon)$ defined in Sect. 2. We begin with the following technical result.

Lemma 3.1. *Let $s \in (0, 1)$ and $n > 2s$. Let $\Omega \subset \mathbb{R}^n$ be a domain, let $x_0 \in \Omega$ and $\rho > 0$ be such that $B_{4\rho}(x_0) \subset \Omega$. Let $\varphi \in C_c^\infty(\Omega)$ be such that $\text{supp}(\varphi) \subset B_{2\rho}(x_0)$, $0 \leq \varphi \leq 1$ in $B_{2\rho}(x_0)$ and $\varphi \equiv 1$ in $B_\rho(x_0)$. There exists $\tau_0 > 0$ such that for every $\tau \in (0, \tau_0)$, setting*

$$u_\tau^s(x) := \varphi(x) \tau^{-\left(\frac{n-2s}{2}\right)} U_{x_0, \mu} \left(\frac{x - x_0}{\tau} + x_0 \right), \quad (3.1)$$

where $U_{x_0, \mu}$ is defined by (2.2), then the following estimates hold:

$$\begin{aligned} \|u_\tau^s\|_s^2 &\leq S_s^{\frac{n}{2s}} + C\tau^{n-2s}, \\ 0 &< S_s^{\frac{n}{2s}} - C\tau^n \leq |u_\tau^s|_{2_s^*}^2 \leq S_s^{\frac{n}{2s}}, \\ 0 &\leq |u_\tau^s|_1 \leq C\tau^{\frac{n-2s}{2}}, \end{aligned} \quad (3.2)$$

where the constants C are positive and depend only on n, s, x_0, μ and ρ . Moreover, for any $0 < \varepsilon < \frac{2s}{n-2s}$, taking $\mu = b_{n,s}^{\frac{2}{n-2s}}$, where $b_{n,s}$ is given by (2.1), we have

$$\begin{aligned} 0 &< \tau^{\left(\frac{n-2s}{2}\right)\varepsilon} \left[S_s^{\frac{n}{2s}} - C\tau^n \right] \leq |u_\tau^s|_{2_s^* - \varepsilon}^{2_s^* - \varepsilon} \leq C\tau^{\left(\frac{n-2s}{2}\right)\varepsilon}, \\ 0 &< \tau^{\left(\frac{n-2s}{2}\right)(1+\varepsilon)} \left[S_s^{\frac{n}{2s}} - C\tau^n \right] \leq |u_\tau^s|_{2_s^* - 1 - \varepsilon}^{2_s^* - 1 - \varepsilon} \leq C\tau^{\left(\frac{n-2s}{2}\right)(1+\varepsilon)}, \end{aligned} \quad (3.3)$$

where the appearing constants are positive and depend only on n, s, x_0 and ρ . Let $0 < s_0 < s_1 \leq 1$ and let $n > 2s_1$. Then, if $s \in [s_0, s_1)$ and $\varepsilon \in \left(0, \frac{2s_0}{n-2s_0}\right)$, both τ_0 and the above constants C can be taken in such a way that they depend on n, μ, ρ, s_0, s_1 , but not on s, τ and ε .

Proof. Inequalities (3.2) are proved in [26], [27] and hold true for all sufficiently small $\tau > 0$ with constants C independent on τ . Concerning the dependence of the constants C on the other parameters we refer to [8, Remark 2.2]. Let us focus on the proof of (3.3). Taking if necessary a smaller $\tau_0 > 0$ so that $\tau_0 < \min\{1, 2\rho\}$, we find that, when $q = 2_s^* - \varepsilon$ or $q = 2_s^* - 1 - \varepsilon$,

$$\int_{\mathbb{R}^n} |u_\tau^s|^q dx \leq C\tau^{n - \left(\frac{n-2s}{2}\right)q} \left(C + \int_1^{\frac{2\rho}{\tau}} r^{n - (n-2s)q - 1} dr \right),$$

where the constants $C > 0$ depend on n, s, μ , but not on τ nor on ε . Furthermore, since $0 < \varepsilon < \frac{2s}{n-2s}$ we have

$$\int_1^{\frac{2\rho}{\tau}} r^{n - (n-2s)q - 1} dr \leq C,$$

for some constant $C > 0$ independent on τ and ε . Recalling the definition of $b_{n,s}$, one can see that all the previous constants can be taken in a uniform way with respect to $s \in [s_0, s_1)$ when $n > 2s_1$ and $\varepsilon \in \left(0, \frac{2s_0}{n-2s_0}\right)$. Hence the right-hand side inequalities in (3.3) are proved.

In order to prove the left-hand side inequalities it suffice to notice that, thanks to our choice of $\mu = b_{n,s}^{\frac{2}{n-2s}}$, it follows that $|U_{\mu, x_0}^s|_\infty = 1$ and thus $|U_{\mu, x_0}^s|^q \geq |U_{\mu, x_0}^s|_{2_s^*}^{2_s^*}$ for every $x \in \mathbb{R}^n$,

where $q = 2_s^* - \varepsilon$ or $q = 2_s^* - 1 - \varepsilon$. Then, using also (3.2), we find that

$$\begin{aligned} \int_{\Omega} |u_{\tau}^s|^q dx &\geq \tau^{n - (\frac{n-2s}{2})q} \int_{B_{\rho/\tau}} |U_{\mu, x_0}^s|^q dx \\ &\geq \tau^{n - (\frac{n-2s}{2})q} \left[S_s^{\frac{n}{2s}} - \int_{\mathbb{R}^n \setminus B_{\rho/\tau}} |U_{\mu, x_0}^s|^{2_s^*} dx \right] \\ &\geq \tau^{n - (\frac{n-2s}{2})q} \left[S_s^{\frac{n}{2s}} - C\tau^n \right], \end{aligned}$$

for some constant $C > 0$ which depends only on n, s and ρ , but not on τ, ε , and which is uniform with respect to $s \in [s_0, s_1]$. The proof is complete. \square

As a consequence we obtain the following uniform asymptotic result on $C_{\mathcal{N}(\Omega)}(s, \varepsilon)$.

Lemma 3.2. *Let $s \in (0, 1)$, $n > 2s$ and let Ω be a smooth bounded domain of \mathbb{R}^n . Then, as $\varepsilon \rightarrow 0^+$, it holds*

$$C_{\mathcal{N}(\Omega)}(s, \varepsilon) \rightarrow \frac{s}{n} S_s^{\frac{n}{2s}}. \quad (3.4)$$

Moreover, if $0 < s_0 < s_1 \leq 1$ and $n > 2s_1$, for every $\varepsilon \in \left(0, \min \left\{ \tau_0, \frac{2s_0}{n-2s_0}, 1 \right\} \right)$, where τ_0 is given by Lemma 3.1, we have

$$\sup_{s \in (s_0, s_1)} \left| C_{\mathcal{N}(\Omega)}(s, \varepsilon) - \frac{s}{n} S_s^{\frac{n}{2s}} \right| \leq g_1(\varepsilon), \quad (3.5)$$

where g_1 does not depend on s and $g_1(\varepsilon) \rightarrow 0^+$ as $\varepsilon \rightarrow 0^+$.

Proof. Let $s \in (0, 1)$ and $n > 2s$. In order to prove (3.4), in view of (2.5) it is sufficient to show that $S_{s, \varepsilon} \rightarrow S_s$, as $\varepsilon \rightarrow 0^+$, where $S_{s, \varepsilon} := \inf_{u \in X_0^s(\Omega) \setminus \{0\}} \frac{\|u\|_s^2}{|u|_{2_s^* - \varepsilon}^2}$. To this end we observe that, by Hölder's inequality, for every $u \in X_0^s(\Omega)$ and any sufficiently small $\varepsilon > 0$ we have

$$|u|_{2_s^* - \varepsilon} \leq |u|_2^{\frac{2\varepsilon}{(2_s^* - \varepsilon)(2_s^* - 2)}} |u|_{2_s^*}^{\frac{2_s^*(2_s^* - 2 - \varepsilon)}{(2_s^* - \varepsilon)(2_s^* - 2)}}.$$

Then, thanks to the fractional Sobolev embedding and the variational characterization of the eigenvalues, we infer that

$$|u|_{2_s^* - \varepsilon}^2 \leq \lambda_{1, s}(\Omega)^{-\frac{2\varepsilon}{(2_s^* - \varepsilon)(2_s^* - 2)}} S_s^{-\frac{2_s^*(2_s^* - 2 - \varepsilon)}{(2_s^* - \varepsilon)(2_s^* - 2)}} \|u\|_s^2,$$

which implies, when $u \neq 0$, that

$$\lambda_{1, s}(\Omega)^{\frac{2\varepsilon}{(2_s^* - \varepsilon)(2_s^* - 2)}} S_s^{\frac{2_s^*(2_s^* - 2 - \varepsilon)}{(2_s^* - \varepsilon)(2_s^* - 2)}} \leq \frac{\|u\|_s^2}{|u|_{2_s^* - \varepsilon}^2}.$$

Taking the infimum as $u \in X_0^s(\Omega) \setminus \{0\}$ we get that

$$\lambda_{1, s}(\Omega)^{\frac{2\varepsilon}{(2_s^* - \varepsilon)(2_s^* - 2)}} S_s^{\frac{2_s^*(2_s^* - 2 - \varepsilon)}{(2_s^* - \varepsilon)(2_s^* - 2)}} \leq S_{s, \varepsilon}, \quad (3.6)$$

and thus it follows that

$$S_s \leq \liminf_{\varepsilon \rightarrow 0^+} S_{s, \varepsilon}. \quad (3.7)$$

Now, let us fix $x_0 \in \Omega$, $\rho > 0$, φ as in the statement of Lemma 3.1 and take $\mu = b_{n, s}^{\frac{2}{n-2s}}$. Let u_{τ}^s be the function defined in (3.1). Then, using both (3.2) and (3.3), for any $\tau \in (0, \tau_0)$, $\varepsilon \in (0, \frac{2s}{n-2s})$ we infer that

$$S_{s, \varepsilon} \leq \frac{\|u_{\tau}^s\|_s^2}{|u_{\tau}^s|_{2_s^* - \varepsilon}^2} \leq \frac{S_s^{\frac{n}{2s}} + C\tau^{n-2s}}{\tau^{\left(\frac{n-2s}{2_s^* - \varepsilon}\right)\varepsilon} \left[S_s^{\frac{n}{2s}} - C\tau^n \right]^{\frac{2}{2_s^* - \varepsilon}}}. \quad (3.8)$$

Hence, for $\varepsilon \in (0, \min\{\tau_0, \frac{2s}{n-2s}\})$, taking $\tau = \varepsilon$ in (3.8) we get by elementary computations that

$$\limsup_{\varepsilon \rightarrow 0^+} S_{s,\varepsilon} \leq S_s,$$

which, together with (3.7), implies (3.4). The first part of the Lemma is thus proved.

For the second part, recalling [8, (2.5), (2.8)] we have that, fixing $0 < s_0 < 1$, there exist two positive constants $\underline{\lambda}, \bar{\lambda}$ such that

$$\underline{\lambda} \leq \lambda_{1,s}(\Omega) \leq \bar{\lambda} \quad \forall s \in [s_0, 1]. \quad (3.9)$$

Hence, from (3.6), (3.9) we deduce that there exists $C > 0$ depending only on n, Ω and s_0 such that for all $s \in [s_0, 1)$

$$S_{s,\varepsilon} \geq S_s C^\varepsilon. \quad (3.10)$$

On the other hand, let us fix s_1 such that $0 < s_0 < s_1 \leq 1$ and let $s \in (s_0, s_1)$, $n > 2s_1$, $\varepsilon \in \left(0, \min\left\{\tau_0, \frac{2s_0}{n-2s_0}, 1\right\}\right)$, where τ_0 is given by Lemma 3.1. Then from (3.8), (3.9), choosing $\tau = \varepsilon$ and taking into account that $\frac{1}{2} < \varepsilon^\varepsilon \leq 1$ for any $\varepsilon \in (0, 1)$, we infer that

$$S_{s,\varepsilon} \leq S_s \left(\frac{C^\varepsilon}{(\varepsilon^\varepsilon)^\alpha [1 - C\varepsilon^n]^\beta} \right) + C\varepsilon^{n-2s}, \quad (3.11)$$

for some constants $C, \alpha, \beta > 0$ which depend only on n, s_0 and s_1 , but not on s and ε . Therefore, from (2.5), (3.10) and (3.11) we obtain

$$C^\varepsilon \frac{S}{n} S_s^{\frac{n}{2s}} \leq C_{\mathcal{N}(\Omega)}(s, \varepsilon) \leq \frac{S}{n} S_s^{\frac{n}{2s}} g(\varepsilon),$$

where g and $C > 0$ do not depend on s , and g is such that $g(\varepsilon) \rightarrow 1$ as $\varepsilon \rightarrow 0^+$. Hence, setting $g_1(\varepsilon) := \max\{|C^\varepsilon - 1|, |g(\varepsilon) - 1|\}$ we get (3.5). The proof is then complete. \square

In the next result we describe the asymptotic behavior of $C_{\mathcal{M}(\Omega)}(s, \varepsilon)$, as $\varepsilon \rightarrow 0^+$. Differently from the case of critical nonlinearities (see [8, Lemma 3.6]), there are some difficulties in proving uniform energy estimates from above which are directly related to $C_{\mathcal{N}(\Omega)}(s, \varepsilon)$. To overcome these difficulties we provide a uniform upper bound in terms of $\frac{2s}{n} S_s^{\frac{n}{2s}}$ instead, which is obtained by using as competitors for the energy superpositions of standard bubbles centered at the same point and with different concentration speeds.

Lemma 3.3. *Let $s \in (0, 1)$, $n > 2s$ and let $\Omega \subset \mathbb{R}^n$ be a smooth bounded domain. We have*

$$\lim_{\varepsilon \rightarrow 0^+} C_{\mathcal{M}(\Omega)}(s, \varepsilon) = \frac{2s}{n} S_s^{\frac{n}{2s}}. \quad (3.12)$$

Moreover, let $0 < s_0 < s_1 \leq 1$ and $n > 2s_1$. Then there exists $\hat{\varepsilon} = \hat{\varepsilon}(s_0, s_1) \in (0, 2s_0^* - 2)$ such that for every $\varepsilon \in (0, \hat{\varepsilon})$

$$\sup_{s \in (s_0, s_1)} \left| C_{\mathcal{M}(\Omega)}(s, \varepsilon) - \frac{2s}{n} S_s^{\frac{n}{2s}} \right| \leq g_2(\varepsilon), \quad (3.13)$$

where the function g_2 does not depend on s and $g_2(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0^+$. The same result holds for $\mathcal{M}_{s,\varepsilon}^r(B_R)$.

Proof. Let us fix $s \in (0, 1)$, $n > 2s$ and let $\Omega \subset \mathbb{R}^n$ be a smooth bounded domain. We claim that

$$2C_{\mathcal{N}(\Omega)}(s, \varepsilon) \leq C_{\mathcal{M}(\Omega)}(s, \varepsilon). \quad (3.14)$$

As an immediate consequence, from Lemma 3.2 we get that

$$\frac{2s}{n} S_s^{\frac{n}{2s}} \leq \liminf_{\varepsilon \rightarrow 0} C_{\mathcal{M}(\Omega)}(s, \varepsilon). \quad (3.15)$$

To prove (3.14) it suffices to notice that, given $u \in \mathcal{M}_{s,\varepsilon}(\Omega)$, then for every $\alpha, \beta > 0$ it holds

$$I_{s,\varepsilon}(\alpha u^+) + I_{s,\varepsilon}(\beta u^-) \leq I_{s,\varepsilon}(u).$$

This follows from the explicit computation of $I_{s,\varepsilon}(\alpha u^+ - \beta u^-)$, taking into account that $(u^+, u^-)_s < 0$ and that $\sup_{t \geq 0} \left(\frac{t^2}{2} - \frac{t^{2_s^* - \varepsilon}}{2_s^* - \varepsilon} \right) \leq \left(\frac{1}{2} - \frac{1}{2_s^* - \varepsilon} \right)$. Hence, choosing $u \in \mathcal{M}_{s,\varepsilon}(\Omega)$ such that $I_{s,\varepsilon}(u) = C_{\mathcal{M}(\Omega)}(s, \varepsilon)$ and α, β in such a way that $\alpha u^+, \beta u^- \in \mathcal{N}_{s,\varepsilon}(\Omega)$ (which is always possible), we obtain the desired result.

To conclude the proof of (3.12) we need to prove the limsup inequality. To this end, we consider $u_{\tau'}^s$ and $u_{\tau''}^s$ of the form (3.1), sharing all the parameters μ, φ, ρ, x_0 , apart from τ . To simplify the notation, we assume without loss of generality that $0 \in \Omega$ and we take $x_0 = 0$. Moreover, we choose ρ and μ as in Lemma 3.1 so that (3.2), (3.3) hold true whenever ε is small enough. Finally, for the concentration parameters, we take τ', τ'' of the form $\tau' = \varepsilon^{\frac{2_s}{n-2_s}}$, $\tau'' = \varepsilon^{\frac{2_s}{n-2_s}}$, where $\delta > 0$ is such that

$$\delta > \max \left\{ 1, \frac{(2_s^* - \varepsilon)(2_s^* - 1 - \varepsilon)}{2_s^* - 2\varepsilon}, 2_s^* - 1 \right\}.$$

Notice that $\delta = 2_s^* - 1$ when ε is small enough, and that it can be taken in a uniform way with respect to s when $s \in [s_0, s_1]$.

Arguing as in [8, Theorem 3.5, Step 2], we infer that

$$C_{\mathcal{M}(\Omega)}(s, \varepsilon) \leq \sup_{\alpha, \beta \geq 0} I_{s,\varepsilon}(\alpha u_{\tau'}^s - \beta u_{\tau''}^s). \quad (3.16)$$

To conclude we need to estimate the right-hand side of (3.16). The first crucial fact is that, in (3.16), it is sufficient to consider only linear combinations $\alpha u_{\tau'}^s - \beta u_{\tau''}^s$ with α, β in a compact subset of $\mathbb{R}^+ \cup \{0\}$. More precisely, we prove that there exists $\tilde{C} > 0$ independent on ε (and depending only on s_0, s_1 when $s \in [s_0, s_1]$) such that, for any $\alpha, \beta \geq 0$ satisfying $\alpha + \beta \geq \tilde{C}$, it holds

$$I_{s,\varepsilon}(\alpha u_{\tau'}^s - \beta u_{\tau''}^s) \leq 0. \quad (3.17)$$

Indeed, by a straightforward computation and using Lemma 3.1 we have

$$\|\alpha u_{\tau'}^s - \beta u_{\tau''}^s\|_s^2 \leq C(\alpha + \beta)^2, \quad (3.18)$$

for some constant C independent on both $\varepsilon, \tau', \tau''$ and s , when $s \in [s_0, s_1]$. On the other hand, arguing exactly as in [8, Lemma 3.6] and using again Lemma 3.1, we infer that for any $\theta \in (0, 1)$

$$\begin{aligned} & |\alpha u_{\tau'}^s - \beta u_{\tau''}^s|_{2_s^* - \varepsilon}^{2_s^* - \varepsilon} \\ & \geq C \alpha^{2_s^* - \varepsilon} (\tau')^{\frac{(n-2_s)}{2}} \varepsilon \left(C - \frac{(\tau')^{\frac{(n-2_s)}{2}} (2_s^* - 2\varepsilon)}{\theta^{2_s^* - 1 - \varepsilon}} - \frac{(\tau')^{\frac{(n-2_s)}{2}} \frac{2_s^*}{2_s^* - \varepsilon - 1}}{\theta^{\frac{1}{2_s^* - 1 - \varepsilon}}} \right) \\ & + \beta^{2_s^* - \varepsilon} (|u_{\tau''}^s|_{2_s^* - \varepsilon}^{2_s^* - \varepsilon} - \theta |u_{\tau''}^s|_{\infty}^{2_s^* - \varepsilon}). \end{aligned} \quad (3.19)$$

Now, thanks to our choice of μ we have

$$|u_{\tau''}^s|_{\infty} = u_{\tau''}^s(0) = (\tau'')^{-\frac{n-2_s}{2}}.$$

Hence, recalling that $\tau'' = \varepsilon^{\frac{2_s}{n-2_s}}$ and taking $\theta = C' \varepsilon^{2_s^* - \varepsilon}$, where C' will be chosen later, from Lemma 3.1 we obtain that

$$|u_{\tau''}^s|_{2_s^* - \varepsilon}^{2_s^* - \varepsilon} - \theta |u_{\tau''}^s|_{\infty}^{2_s^* - \varepsilon} \geq C - C',$$

for any $\varepsilon > 0$ small enough, where C does not depend on ε , nor on s when $s \in [s_0, s_1]$. Therefore, taking $C' = \frac{1}{2}C$ we get that

$$|u_{\tau'}^s|_{2_s^{*-2\varepsilon}}^{2_s^*-2\varepsilon} - \theta |u_{\tau''}^s|_{\infty}^{2_s^*-2\varepsilon} \geq \frac{1}{2}C > 0. \quad (3.20)$$

Thus, recalling that $\tau' = \varepsilon^{\frac{2\delta}{n-2s}}$, from (3.19), (3.20) we obtain

$$\begin{aligned} & |\alpha u_{\tau'}^s - \beta u_{\tau''}^s|_{2_s^{*-2\varepsilon}}^{2_s^*-2\varepsilon} \\ & \geq (\varepsilon^\varepsilon)^\delta C \alpha^{2_s^*-2\varepsilon} \left(C - \varepsilon^{\delta(2_s^*-2\varepsilon)-(2_s^*-\varepsilon)(2_s^*-1-\varepsilon)} - \varepsilon^{\frac{2_s^*(\delta-1)+\varepsilon}{2_s^*-1-\varepsilon}} \right) + \frac{1}{2}C \beta^{2_s^*-2\varepsilon}. \end{aligned}$$

Then, exploiting the properties of the function $t \mapsto t^t$, thanks to the definition of δ , we find that there exists $C > 0$ such that for all sufficiently small $\varepsilon > 0$

$$|\alpha u_{\tau'}^s - \beta u_{\tau''}^s|_{2_s^{*-2\varepsilon}}^{2_s^*-2\varepsilon} \geq C(\alpha^{2_s^*-2\varepsilon} + \beta^{2_s^*-2\varepsilon}) \geq C(\alpha + \beta)^{2_s^*-2\varepsilon}. \quad (3.21)$$

Finally, thanks to (3.18) and (3.21) we infer that

$$I_{s,\varepsilon}(\alpha u_{\tau'}^s - \beta u_{\tau''}^s) \leq C(\alpha + \beta)^2(1 - C(\alpha + \beta)^{2_s^*-2-2\varepsilon}),$$

which implies that there exists $\tilde{C} > 0$, not depending on ε , such that if $(\alpha + \beta) \geq \tilde{C}$ then $I_{s,\varepsilon}(\alpha u_{\tau'}^s - \beta u_{\tau''}^s) \leq 0$, as claimed. We observe that \tilde{C} can be taken in a uniform way with respect to s , when $s \in [s_0, s_1]$.

It remains to treat the case $\alpha + \beta \leq \tilde{C}$. To this end we begin with a preliminary estimate on the scalar product between two bubbles. A careful analysis of the argument carried out in [27, Proposition 21] shows that

$$|(u_{\tau'}^s, u_{\tau''}^s)_s| \leq (\tau')^{-\frac{n-2s}{2}} (\tau'')^{-\frac{n-2s}{2}} \left| \left(U_{0,\mu}^s \left(\frac{x}{\tau'} \right), U_{0,\mu}^s \left(\frac{x}{\tau''} \right) \right)_s \right| + C(\tau')^{\frac{n-2s}{2}} (\tau'')^{\frac{n-2s}{2}},$$

where the constant C does not depend on τ' nor on τ'' , and it is uniformly bounded with respect to $s \in [s_0, s_1]$. Performing a change of variables, and recalling that $U_{0,\mu}^s$ solves problem (2.3), we infer that

$$\begin{aligned} \left| \left(U_{0,\mu}^s \left(\frac{x}{\tau'} \right), U_{0,\mu}^s \left(\frac{x}{\tau''} \right) \right)_s \right| &= \left| (\tau')^{n-2s} \left(U_{0,\mu}^s, U_{0,\mu}^s \left(\frac{\tau'}{\tau''} x \right) \right)_s \right| \\ &\leq (\tau')^{n-2s} \int_{\mathbb{R}^n} |U_{0,\mu}^s|^{2_s^*-1} \left| U_{0,\mu}^s \left(\frac{\tau'}{\tau''} x \right) \right| dx \\ &\leq C(\tau')^{n-2s} \int_{\mathbb{R}^n} (\mu^2 + |x|^2)^{-\frac{n+2s}{2}} dx \leq C(\tau')^{n-2s}, \end{aligned}$$

where we used the fact that thanks to our choice of μ it holds $|U_{0,\mu}^s|_\infty = 1$, and where the constant $C > 0$ does not depend on τ' nor on τ'' and it is uniformly bounded with respect to $s \in [s_0, s_1]$. Summing up, and recalling the definition of τ' and τ'' , we infer that

$$|(u_{\tau'}^s, u_{\tau''}^s)_s| \leq C(\varepsilon^{\delta-1} + \varepsilon^{\delta+1}) \leq C\varepsilon^{\delta-1}. \quad (3.22)$$

Let us finally consider the case of $\alpha + \beta \leq \tilde{C}$. Arguing again as in [8, Lemma 3.6] and applying Lemma 3.1 and (3.22), we obtain

$$\begin{aligned}
I_{s,\varepsilon}(\alpha u_{\tau'}^s - \beta u_{\tau''}^s) &\leq \frac{\alpha^2}{2} \|u_{\tau'}^s\|_s^2 + \frac{\beta^2}{2} \|u_{\tau''}^s\|_s^2 - \frac{\alpha^{2_s^* - \varepsilon}}{2_s^* - \varepsilon} |u_{\tau'}^s|_{2_s^* - \varepsilon}^{2_s^* - \varepsilon} - \frac{\beta^{2_s^* - \varepsilon}}{2_s^* - \varepsilon} |u_{\tau''}^s|_{2_s^* - \varepsilon}^{2_s^* - \varepsilon} \\
&\quad + C \int_{\mathbb{R}^n} |u_{\tau'}^s|^{2_s^* - 1 - \varepsilon} |u_{\tau''}^s| dx + C \int_{\mathbb{R}^n} |u_{\tau''}^s|^{2_s^* - 1 - \varepsilon} |u_{\tau'}^s| dx + C\varepsilon^{\delta - 1} \\
&\leq \frac{\alpha^2}{2} S_s^{\frac{n}{2_s^*}} + \frac{\beta^2}{2} S_s^{\frac{n}{2_s^*}} - \frac{\alpha^{2_s^* - \varepsilon}}{2_s^* - \varepsilon} (\varepsilon^\varepsilon)^\delta (S_s^{\frac{n}{2_s^*}} - C\varepsilon^{2_s^* \delta}) \\
&\quad - \frac{\beta^{2_s^* - \varepsilon}}{2_s^* - \varepsilon} (\varepsilon^\varepsilon) (S_s^{\frac{n}{2_s^*}} - C\varepsilon^{2_s^*}) + C\varepsilon^{-1} \varepsilon^{\delta(1+\varepsilon)} + C\varepsilon^{-(2_s^* - 1 - \varepsilon)} \varepsilon^\delta \\
&\quad + C\varepsilon^{2\delta} + C\varepsilon^2 + C\varepsilon^{\delta - 1},
\end{aligned}$$

where we used that $|u_{\tau''}^s|_\infty = \varepsilon^{-1}$. Even in this case all the appearing constants are independent on ε , and they are uniformly bounded with respect to s when $s \in [s_0, s_1]$. Using again that $\sup_{t \geq 0} \left(\frac{t^2}{2} - \frac{t^{2_s^* - \varepsilon}}{2_s^* - \varepsilon} \right) \leq \left(\frac{1}{2} - \frac{1}{2_s^* - \varepsilon} \right)$, we get that

$$\begin{aligned}
I_{s,\varepsilon}(\alpha u_{\tau'}^s - \beta u_{\tau''}^s) &\leq \frac{2_s^* - 2 - \varepsilon}{2_s^* - \varepsilon} S_s^{\frac{n}{2_s^*}} + C(1 - (\varepsilon^\varepsilon)^\delta) + C(1 - \varepsilon^\varepsilon) + C\varepsilon^{2_s^* \delta} + C\varepsilon^{2_s^*} \\
&\quad + C\varepsilon^{\delta - 1 + \delta\varepsilon} + C\varepsilon^{\delta - (2_s^* - 1 - \varepsilon)} + C\varepsilon^{2\delta} + C\varepsilon^2 + C\varepsilon^{\delta - 1} \\
&=: \frac{2s}{n} S_s^{\frac{n}{2_s^*}} + g(\varepsilon),
\end{aligned} \tag{3.23}$$

where all the constants $C > 0$, and thus g , do not depend on s , when $s \in [s_0, s_1]$. In particular, g satisfies $g(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0^+$.

At the end, putting together (3.16), (3.23), taking into account (3.17), we obtain

$$C_{\mathcal{M}(\Omega)}(s, \varepsilon) \leq \frac{2s}{n} S_s^{\frac{n}{2_s^*}} + g(\varepsilon), \tag{3.24}$$

and thus we get that

$$\limsup_{\varepsilon \rightarrow 0^+} C_{\mathcal{M}(\Omega)}(s, \varepsilon) \leq \frac{2s}{n} S_s^{\frac{n}{2_s^*}},$$

which, together with (3.15) gives (3.12).

For the proof of the second part, fixing $0 < s_0 < s_1 \leq 1$, thanks to Lemma 3.2 and the definition of g , we deduce that inequalities (3.14) and (3.24) are uniform with respect to s when $s \in [s_0, s_1]$. At the end, arguing as in Lemma 3.2 we obtain (3.13), for some function g_2 independent on s and such that $g_2(\varepsilon) \rightarrow 0$, as $\varepsilon \rightarrow 0^+$.

In the radial case the proof is identical. Indeed, since in the construction we take standard bubbles centered at the same point, then the functions $\alpha u_{\tau'}^s - \beta u_{\tau''}^s$ are radial and thus admissible competitors. The proof is then complete. \square

4. ASYMPTOTIC ANALYSIS OF LEAST ENERGY RADIAL SIGN-CHANGING SOLUTIONS

In this section we study the asymptotic behavior of least energy radial nodal solutions to (1.2), as $\varepsilon \rightarrow 0^+$. Theorem 1.1 will be a consequence of the results contained in this section. We begin by a couple of preliminary known results.

Lemma 4.1. *Let $s \in (0, 1)$, let $n > 2s$ and let $\Omega \subset \mathbb{R}^n$ be a smooth bounded domain. Let $(u_{s,\varepsilon}) \subset \mathcal{M}_{s,\varepsilon}(\Omega)$ be a family of solutions of Problem (2.4) such that $I_{s,\varepsilon}(u_{s,\varepsilon}) = C_{\mathcal{M}(\Omega)}(s, \varepsilon)$ and set $M_{s,\varepsilon,\pm} := |u_{s,\varepsilon}^\pm|_\infty$. As $\varepsilon \rightarrow 0^+$ we have:*

- (i) $\|u_{s,\varepsilon}^\pm\|_s^2 \rightarrow S_s^{\frac{n}{2s}}$;
- (ii) $|u_{s,\varepsilon}^\pm|_{2_s^{*-2\varepsilon}}^2 \rightarrow S_s^{\frac{n}{2s}}$;
- (iii) $(u_{s,\varepsilon}^+, u_{s,\varepsilon}^-)_s \rightarrow 0$;
- (iv) $u_{s,\varepsilon} \rightarrow 0$ in $X_0^s(\Omega)$;
- (v) $M_{s,\varepsilon,\pm} \rightarrow +\infty$.

The same results hold for a family $(u_{s,\varepsilon}) \subset \mathcal{M}_{s,\varepsilon}^r(B_R)$ of radial solutions to Problem (1.2) such that $I_{s,\varepsilon}(u_{s,\varepsilon}) = C_{\mathcal{M}^r(B_R)}(s, \varepsilon)$. Moreover, for every $0 < s_0 < s_1 \leq 1$ and $n > 2s_1$, the limits (i) – (iii) are uniform with respect to $s \in [s_0, s_1]$.

Proof. It suffices to argue as in [8, Lemma 4.3], with some minor modifications. \square

The following estimate will play a central role in this paper.

Proposition 4.2. *Let $0 < s_0 < s_1 \leq 1$ and let $s \in [s_0, s_1]$, $n > 2s_1$. Let $0 < R_0 \leq R$, $g \in L^\infty(B_R)$ and v be a weak solution of*

$$\begin{cases} (-\Delta)^s v = g & \text{in } B_R, \\ v = 0 & \text{in } \mathbb{R}^n \setminus B_R, \end{cases}$$

Then $v \in C^{0,s}(\mathbb{R}^n)$ and

$$\|v\|_{C^{0,s}(\mathbb{R}^n)} \leq C|g|_{L^\infty(B_R)}$$

where the constant $C > 0$ depends only on n , s_0 , s_1 and R_0 , but neither on s nor on R .

Proof. The estimate is a consequence of results contained in [24]. Concerning the dependence on the parameters s_0 , s_1 , it can be deduced from a careful analysis of the proof in [24] (see also [8, Proposition 2.3]). As for the dependence of the constant C on the domain, it turns out that C depends only on the radii coming from the outer and inner ball conditions for B_R . Hence, it is clear that C can be chosen in a uniform way with respect to R if we assume that $R \geq R_0$, for some $R_0 > 0$. \square

From now on $u_{s,\varepsilon} \in \mathcal{M}_{s,\varepsilon}^r(B_R)$ will denote a least energy radial solution to Problem (1.2), i.e. $I_{s,\varepsilon}(u_{s,\varepsilon}) = C_{\mathcal{M}^r(B_R)}(s, \varepsilon)$. Moreover, we set $M_{s,\varepsilon} := |u_{s,\varepsilon}|_\infty$. In the next result we characterize the asymptotic behavior of the points where the blow-up occurs.

Lemma 4.3. *Let $s \in (0, 1)$, $n > 2s$. Let $x_\varepsilon \in B_R$ be such that $|u_{s,\varepsilon}(x_\varepsilon)| = O(M_{s,\varepsilon})$ as $\varepsilon \rightarrow 0^+$. Then*

$$M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| \not\rightarrow +\infty,$$

where $\beta_{s,\varepsilon} := \frac{2}{n+2s} - \frac{\varepsilon}{2s}$. In particular, we infer that $|x_\varepsilon| \rightarrow 0$.

Proof. If $s \in (\frac{1}{2}, 1)$ this is a consequence of the fractional Strauss inequality (1.3) (see [6, Proposition 1]). Indeed, suppose that eventually $x_\varepsilon \neq 0$, otherwise there is nothing to prove. Then

$$\begin{aligned} (M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon|)^{\frac{n-2s}{2}} &= M_{s,\varepsilon}^{1-\frac{\varepsilon}{2s-2}} |x_\varepsilon|^{\frac{n-2s}{2}} \\ &\leq M_{s,\varepsilon} |x_\varepsilon|^{\frac{n-2s}{2}} \leq C |u_{s,\varepsilon}(x_\varepsilon)| |x_\varepsilon|^{\frac{n-2s}{2}} \leq CK_{n,s} \|u_{s,\varepsilon}\|_s^2 \leq C, \end{aligned}$$

where we used that eventually $M_{s,\varepsilon} \geq 1$, thanks to Lemma 4.1,(v). Unfortunately, as pointed out in Sect. 1, the fractional Strauss inequality does not hold in general when $s \in (0, \frac{1}{2}]$. To overcome this difficulty we use the following argument, which is valid for any $s \in (0, 1)$.

Assume by contradiction that there exists a subsequence (still denoted by ε for simplicity), such that $M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| \rightarrow +\infty$ as $\varepsilon \rightarrow 0^+$. Let us define the rescaled functions

$$\tilde{u}_{s,\varepsilon}(x) = \frac{1}{M_{s,\varepsilon}} u_{s,\varepsilon} \left(\frac{x}{M_{s,\varepsilon}^{\beta_{s,\varepsilon}}} \right), \quad x \in \mathbb{R}^n. \quad (4.1)$$

It is immediate to see that the functions $\tilde{u}_{s,\varepsilon}$ satisfy

$$\begin{cases} (-\Delta)^s \tilde{u}_{s,\varepsilon} = |\tilde{u}_{s,\varepsilon}|^{2^*_s-2-\varepsilon} \tilde{u}_{s,\varepsilon}, & \text{in } B_{M_{s,\varepsilon}^{\beta_{s,\varepsilon}}} R, \\ \tilde{u}_{s,\varepsilon} = 0 & \text{in } \mathbb{R}^n \setminus B_{M_{s,\varepsilon}^{\beta_{s,\varepsilon}}} R. \end{cases} \quad (4.2)$$

Since by construction $|\tilde{u}_{s,\varepsilon}|_\infty \leq 1$ and $M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| \rightarrow +\infty$, then from Proposition 4.2 we deduce that

$$\|\tilde{u}_{s,\varepsilon}\|_{C^{0,s}(\mathbb{R}^n)} \leq C, \quad (4.3)$$

for some $C > 0$ independent on ε .

Now we observe that, by definition of x_ε , there exists $C_1 \in (0, 1]$ such that for all sufficiently small $\varepsilon > 0$ it holds

$$|\tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} x_\varepsilon)| = \frac{|u_{s,\varepsilon}(x_\varepsilon)|}{M_{s,\varepsilon}^{\beta_{s,\varepsilon}}} \geq C_1 + o(\varepsilon). \quad (4.4)$$

Using (4.3), (4.4) and the triangle inequality, for every $\tau > 0$ and $\xi \in \mathbb{R}^n$ such that $|\xi| \leq 1$, we infer that

$$\frac{C_1 - |\tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} x_\varepsilon + \tau\xi)| + o(\varepsilon)}{|\tau\xi|^s} \leq \frac{|\tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} x_\varepsilon) - \tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} x_\varepsilon + \tau\xi)|}{|\tau\xi|^s} \leq C.$$

In particular, we can find $\varepsilon_0 > 0$, $\tau_0 > 0$ and $C_2 > 0$ such that, for every $\varepsilon \in (0, \varepsilon_0)$, it holds

$$0 < C_2 \leq C_1 - C\tau_0^s + o(\varepsilon) \leq |\tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} x_\varepsilon + \tau_0\xi)|, \quad \forall |\xi| \leq 1.$$

Therefore, since $\tilde{u}_{s,\varepsilon}$ is radial and ξ is arbitrary we obtain that

$$0 < C_2 \leq |\tilde{u}_{s,\varepsilon}(x)|, \quad \forall x \in B_{M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| + \tau_0} \setminus B_{M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| - \tau_0}.$$

Now, since we are assuming by contradiction that $M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| \rightarrow +\infty$ and since $M_{s,\varepsilon} \rightarrow +\infty$ we get that

$$\begin{aligned} |u_{s,\varepsilon}|_{2_{s,\varepsilon}^{2^*_s-2-\varepsilon}} &\geq M_{s,\varepsilon}^{-\varepsilon \left(\frac{n-2s}{2s}\right)} |u_{s,\varepsilon}|_{2_{s,\varepsilon}^{2^*_s-2-\varepsilon}} = |\tilde{u}_{s,\varepsilon}|_{2_{s,\varepsilon}^{2^*_s-2-\varepsilon}} \\ &\geq C_2 [(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| + \tau_0)^n - (M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| - \tau_0)^n] \\ &= 2nC_2\tau_0 (M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon|)^{n-1} + \psi(\varepsilon), \end{aligned}$$

where $\psi(\varepsilon)$ is such that $\frac{\psi(\varepsilon)}{(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon|)^{n-1}} \rightarrow 0$, as $\varepsilon \rightarrow 0^+$. From this we get that $|u_{s,\varepsilon}|_{2_{s,\varepsilon}^{2^*_s-2-\varepsilon}} \rightarrow +\infty$, as $\varepsilon \rightarrow 0^+$, which contradicts Lemma 4.1, (ii). The proof is complete. \square

In the next result we study the asymptotic behavior of the rescaled solutions defined in (4.1).

Lemma 4.4. *Let $s \in (0, 1)$, $n > 2s$ and let $(\tilde{u}_{s,\varepsilon})_\varepsilon$ be the sequence of rescaled functions associated to $(u_{s,\varepsilon})_\varepsilon$, defined in (4.1). Then we have that, up to a subsequence, $\tilde{u}_{s,\varepsilon} \rightarrow \tilde{u}_s$ in $C_{loc}^{0,\alpha}(\mathbb{R}^n)$ for some $\alpha \in (0, s)$, as $\varepsilon \rightarrow 0^+$, where $\tilde{u}_s \in \mathcal{D}^s(\mathbb{R}^n)$ is a nontrivial weak solution to*

$$\begin{cases} (-\Delta)^s \tilde{u}_s = |\tilde{u}_s|^{2^*_s-2} \tilde{u}_s & \text{in } \mathbb{R}^n, \\ \tilde{u}_s > 0. \end{cases} \quad (4.5)$$

Moreover, \tilde{u}_s is radial and $|\tilde{u}_s|_\infty = |\tilde{u}_s(0)|$.

Proof. As seen in the proof of Lemma 4.3, the functions $\tilde{u}_{s,\varepsilon}$ weakly satisfy (4.2) and by construction it holds that $|\tilde{u}_{s,\varepsilon}|_\infty \leq 1$. Then, since $M_{s,\varepsilon}^{\beta_{s,\varepsilon}} R \rightarrow +\infty$, thanks to Proposition 4.2 and a standard argument, up to a subsequence, we have

$$\tilde{u}_{s,\varepsilon} \rightarrow \tilde{u}_s \text{ in } C_{loc}^{0,\alpha}(\mathbb{R}^n),$$

for some $\tilde{u}_s \in C_{loc}^{0,\alpha}(\mathbb{R}^n)$, $\alpha \in (0, s)$. We point out that $\tilde{u}_s \not\equiv 0$. Indeed, let $x_\varepsilon \in B_R$ be such that $|u_{s,\varepsilon}(x_\varepsilon)| = M_{s,\varepsilon}$. By construction we have $|\tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} x_\varepsilon)| = 1$, and thanks to Lemma 4.3 we infer that the point $M_{s,\varepsilon}^{\beta_{s,\varepsilon}} x_\varepsilon$ stays in a compact subset of \mathbb{R}^n . Therefore, from the $C_{loc}^{0,\alpha}$ -convergence of $\tilde{u}_{s,\varepsilon}$ in \mathbb{R}^n , we get that \tilde{u}_s is non trivial.

Now we show that $\tilde{u}_s \in \mathcal{D}^s(\mathbb{R}^n)$. In fact, by Lemma 4.1,(i) and since $M_{s,\varepsilon} \rightarrow +\infty$, we infer that

$$\|\tilde{u}_{s,\varepsilon}\|_s^2 = M_{s,\varepsilon}^{-\varepsilon(\frac{n-2s}{2s})} \|u_{s,\varepsilon}\|_s^2 \leq \|u_{s,\varepsilon}\|_s^2 \rightarrow 2S_s^{\frac{n}{2s}}, \quad \text{as } \varepsilon \rightarrow 0^+,$$

and in particular, up to a subsequence, $\tilde{u}_{s,\varepsilon} \rightharpoonup v$ for some $v \in \mathcal{D}^s(\mathbb{R}^n)$. Then, since $\tilde{u}_{s,\varepsilon} \rightarrow \tilde{u}_s$ in $C_{loc}^{0,\alpha}(\mathbb{R}^n)$, we get that $v = \tilde{u}_s$ and we are done. In addition, applying Fatou's Lemma we also deduce that

$$\|\tilde{u}_s\|_s^2 \leq \liminf_{\varepsilon \rightarrow 0^+} \|\tilde{u}_{s,\varepsilon}\|_s^2 \leq 2S_s^{\frac{n}{2s}}. \quad (4.6)$$

Let us prove now that \tilde{u}_s is a weak solution to (4.5). Indeed, for every $\varphi \in C_c^\infty(\mathbb{R}^n)$, since $\tilde{u}_{s,\varepsilon}$ is a weak solution to (4.2) we have

$$(\tilde{u}_{s,\varepsilon}, \varphi)_s = \int_{B_{M_{s,\varepsilon}^{\beta_{s,\varepsilon}} R}} |\tilde{u}_{s,\varepsilon}|^{2_s^* - 2 - \varepsilon} \tilde{u}_{s,\varepsilon} \varphi \, dx, \quad (4.7)$$

where ε is small enough so that $\text{supp } \varphi \subset B_{M_{s,\varepsilon}^{\beta_{s,\varepsilon}} R}$. Since $\tilde{u}_{s,\varepsilon} \rightarrow \tilde{u}_s$ for a.e. $x \in \mathbb{R}^n$, using the well known relations (see e.g. [8])

$$(u, \varphi)_s = \int_{\mathbb{R}^n} u(-\Delta)^s \varphi \, dx, \quad \forall u \in \mathcal{D}^s(\mathbb{R}^n), \quad \forall \varphi \in C_c^\infty(\mathbb{R}^n), \quad (4.8)$$

and

$$|(-\Delta)^s \varphi(x)| \leq C(\varphi) \frac{1}{(1 + |x|)^{n+2s}}, \quad \forall x \in \mathbb{R}^n, \quad \forall \varphi \in C_c^\infty(\mathbb{R}^n),$$

and thanks to Lebesgue's dominated convergence theorem, passing to the limit as $\varepsilon \rightarrow 0^+$ in (4.7) we infer that

$$\int_{\mathbb{R}^n} \tilde{u}_s(-\Delta)^s \varphi \, dx = \int_{\mathbb{R}^n} |\tilde{u}_s|^{2_s^* - 2} \tilde{u}_s \varphi \, dx.$$

Now, since $\tilde{u}_s \in \mathcal{D}^s(\mathbb{R}^n)$ we are allowed to use again (4.8), obtaining that \tilde{u}_s weakly satisfies

$$(-\Delta)^s \tilde{u}_s = |\tilde{u}_s|^{2_s^* - 2} \tilde{u}_s \quad \text{in } \mathbb{R}^n. \quad (4.9)$$

We prove now that \tilde{u}_s is of constant sign. To this end, assume by contradiction that \tilde{u}_s is sign-changing. Then, using $\tilde{u}_s^\pm \in \mathcal{D}^s(\mathbb{R}^n)$ as test functions in (4.9) and recalling that $(\tilde{u}_s^+, \tilde{u}_s^-)_s < 0$, we get that

$$\|\tilde{u}_s^\pm\|_s^2 = (\tilde{u}_s^+, \tilde{u}_s^-)_s + |\tilde{u}_s^\pm|_{2_s^*}^{2_s^*} < |\tilde{u}_s^\pm|_{2_s^*}^{2_s^*}.$$

Hence, by the Sobolev inequality we infer that

$$S_s \leq \frac{\|\tilde{u}_s^\pm\|_s^2}{|\tilde{u}_s^\pm|_{2_s^*}^2} < |\tilde{u}_s^\pm|_{2_s^*}^{2_s^* - 2},$$

and thus $2S_s^{\frac{n}{2s}} < |\tilde{u}_s^+|_{2_s^*}^{2_s^*} + |\tilde{u}_s^-|_{2_s^*}^{2_s^*} = |\tilde{u}_s|_{2_s^*}^{2_s^*}$. Finally, using \tilde{u}_s as a test function in (4.9) we have $\|\tilde{u}_s\|_s^2 = |\tilde{u}_s|_{2_s^*}^{2_s^*}$, and we obtain that $2S_s^{\frac{n}{2s}} < \|\tilde{u}_s\|_s^2$, which contradicts (4.6).

At the end we notice that, since \tilde{u}_s is a pointwise limit of radial functions, it is radial too. Moreover, since \tilde{u}_s is of constant sign, assuming without loss of generality that $\tilde{u}_s \geq 0$, we easily deduce, by the fractional strong maximum principle and the fractional moving plane method (see [5]), that \tilde{u}_s is also decreasing along the radii and thus \tilde{u}_s achieves its maximum at the origin. The proof is complete. \square

A consequence of the above result is that least energy radial sign-changing solutions to (1.2) cannot vanish at the origin.

Lemma 4.5. *Let $s \in (0, 1)$ and $n > 2s$. There exist $\bar{\varepsilon} > 0$ and $C > 0$ such that for every $\varepsilon \in (0, \bar{\varepsilon})$ it holds*

$$|u_{s,\varepsilon}(0)| \geq C. \quad (4.10)$$

Moreover, for every $0 < s_0 < s_1 \leq 1$, $n > 2s_1$ estimate (4.10) holds with $\bar{\varepsilon} > 0$, $C > 0$ independent on $s \in [s_0, s_1]$.

Proof. We prove directly the second part of the Lemma. Assume by contradiction that there exist $0 < s_0 < s_1 \leq 1$, three sequences $\varepsilon_k \rightarrow 0^+$, $C_k \rightarrow 0^+$, $(s_k)_k \in [s_0, s_1]$, and a sequence of nodal radial least energy solutions $u_k := u_{s_k, \varepsilon_k}$ such that $|u_k(0)| < C_k$. Up to a subsequence, we can always assume that $s_k \rightarrow \sigma$, with $\sigma \in [s_0, s_1]$.

Now, only two possibilities can occur: setting $M_k := |u_k|_\infty$, either $(M_k)_k$ is a bounded sequence or there exists a subsequence such that $M_k \rightarrow +\infty$.

Assume that $(M_k)_k$ is bounded. We first observe that $(M_k)_k$ is bounded away from zero, otherwise we could find a subsequence such that $M_k \rightarrow 0$, but this would contradict Lemma 4.1. Therefore, up to a subsequence we can assume that $M_k \rightarrow l$, for some real number $l > 0$. Adapting the arguments of Lemma 4.4 and using Lemma 3.3 we readily infer that, up to a subsequence, $u_k \rightharpoonup u$ in $X_0^{s_0}(B_R)$ and $u_k \rightarrow u$ in $C^{0,\alpha}(\mathbb{R}^n)$, for some $\alpha \in (0, s_0)$. Furthermore we have $u \not\equiv 0$ and it holds that

$$\int_{\mathbb{R}^n} u(-\Delta)^\sigma \varphi \, dx = \int_{\mathbb{R}^n} |u|^{2_\sigma^* - 2} u \varphi \, dx \quad \forall \varphi \in C_c^\infty(B_R).$$

Using that $u_k \rightarrow u$ in $L^2(B_R)$, thanks to the fractional Sobolev embedding and Fatou's Lemma we find

$$\|u\|_\sigma^2 = \int_{\mathbb{R}^n} |\xi|^{2\sigma} |\hat{u}(\xi)|^2 \, dx \leq \liminf_{k \rightarrow +\infty} \int_{\mathbb{R}^n} |\xi|^{2s_k} |\hat{u}_k(\xi)|^2 \, dx = \liminf_{k \rightarrow +\infty} \|u\|_s^2 \leq \frac{2\sigma}{n} S_{\sigma}^{\frac{n}{2\sigma}}, \quad (4.11)$$

where the last inequality is a consequence of the second part of Lemma 3.3, while the equalities are due to the interpretation via the Fourier transform of the fractional Laplacian (see e.g. [13]). From this discussion it follows that u is a non trivial weak solution of

$$\begin{cases} (-\Delta)^\sigma u = |u|^{2_\sigma^* - 2} u & \text{in } B_R, \\ u = 0 & \text{in } \mathbb{R}^n \setminus B_R. \end{cases} \quad (4.12)$$

This readily contradicts the Pohozaev identity when $\sigma = 1$. If $\sigma < 1$, the fractional Pohozaev identity only implies the nonexistence of constant-sign solutions to (4.12) (see [25]). In order to obtain a contradiction we show that u is of constant sign. Indeed, arguing as in the proof of Lemma 4.4, we have that any sign-changing solution u to (4.12) must satisfy $\|u\|_\sigma^2 > \frac{2\sigma}{n} S_{\sigma}^{\frac{n}{2\sigma}}$. Hence, thanks to (4.11) it follows that u is of constant sign and we get the desired contradiction.

Let us analyze the second case. Assume that $M_k \rightarrow +\infty$ and consider the rescaled functions

$$\tilde{u}_k(x) = \frac{1}{M_k} u_k \left(\frac{x}{M_k^{\beta_k}} \right), \quad x \in \mathbb{R}^n,$$

where $\beta_k = \frac{2}{n-2s_k} - \frac{\varepsilon_k}{2s_k}$. Arguing as in Lemma 4.4, and taking into account Lemma 3.3, we obtain that $\tilde{u}_k \rightarrow \tilde{u}$ in $C_{loc}^{0,\alpha}(\mathbb{R}^n)$, as $k \rightarrow +\infty$, for some $\alpha \in (0, s_0)$, where the function \tilde{u} belongs

to $\mathcal{D}^\sigma(\mathbb{R}^n) \cap C_{loc}^{0,\alpha}(\mathbb{R}^n) \setminus \{0\}$, it is radial, verifies $|\tilde{u}|_\infty = |\tilde{u}(0)|$ and weakly satisfies

$$\begin{cases} (-\Delta)^\sigma \tilde{u} = |\tilde{u}|^{2_\sigma^* - 2} \tilde{u} & \text{in } \mathbb{R}^n, \\ \tilde{u} > 0. \end{cases}$$

The only delicate point is when $\sigma = 1$. Indeed, in this case we cannot simply argue via Fatou's Lemma to show that $\tilde{u} \in \mathcal{D}^1(\mathbb{R}^n)$. Nevertheless, since

$$|\tilde{u}|_{2_1^*}^{2_1^*} \leq \liminf_{k \rightarrow +\infty} |\tilde{u}_k|_{2_{s_k}^*}^{2_{s_k}^*} \leq \liminf_{k \rightarrow +\infty} S_{s_k}^{-\frac{2_{s_k}^*}{2}} \|\tilde{u}_k\|_{s_k}^{\frac{2_{s_k}^*}{2}} \leq C,$$

we have that $\tilde{u} \in L^{2_1^*}(\mathbb{R}^n)$. Therefore we can apply [18, Theorem 2, Corollary 3], obtaining that $\tilde{u} \in \mathcal{D}^1(\mathbb{R}^n)$ and $\|\tilde{u}\|_1^2 = |\tilde{u}|_{2_1^*}^{2_1^*}$.

To conclude notice that, since we are assuming $|\tilde{u}_k(0)| = \frac{1}{M_k} |u_k(0)| \rightarrow 0$ and since we have $\tilde{u}_k \rightarrow \tilde{u}$ in $C_{loc}^{0,\alpha}(\mathbb{R}^n)$, then it follows that $|\tilde{u}(0)| = 0$, which contradicts the non triviality of \tilde{u} . The proof is then complete. \square

In the next lemma we show, independently on the number of sign-changes, that $M_{s,\varepsilon}$ is achieved in the nodal component containing the origin and blows up faster than every other extremal value achieved in the other components. Before stating the result we introduce some notation. Assuming without loss of generality that $u_{s,\varepsilon}(0) > 0$, thanks to Lemma 4.5, for all sufficiently small $\varepsilon > 0$ the following quantities are well defined:

$$\begin{aligned} r_\varepsilon^1 &:= \min\{r \in (0, R] ; u_{s,\varepsilon}(x) = 0, |x| = r\}, \\ M_{s,\varepsilon}^+ &:= \max\{u_{s,\varepsilon}(x) ; 0 \leq |x| \leq r_\varepsilon^1\}, \\ \hat{M}_{s,\varepsilon} &:= \max\{|u_{s,\varepsilon}(x)| ; r_\varepsilon^1 \leq |x| \leq R\}. \end{aligned}$$

In other words, r_ε^1 is the first nodal radius, $M_{s,\varepsilon}^+$ is the maximum of the solution in the first nodal component, while $\hat{M}_{s,\varepsilon}$ is the absolute maximum achieved in the other nodal components.

Lemma 4.6. *Let $s \in (0, 1)$ and $n > 2s$. There exists $\varepsilon' > 0$ such that for every $\varepsilon \in (0, \varepsilon')$ it holds*

$$M_{s,\varepsilon} = M_{s,\varepsilon}^+.$$

Moreover, up to a subsequence, as $\varepsilon \rightarrow 0^+$

$$(i) \quad (M_{s,\varepsilon}^+)^{\beta_{s,\varepsilon}} r_\varepsilon^1 \rightarrow +\infty, \quad (ii) \quad \frac{M_{s,\varepsilon}^+}{M_{s,\varepsilon}} \rightarrow +\infty.$$

Proof. We begin by proving that eventually $M_{s,\varepsilon} = M_{s,\varepsilon}^+$. Suppose by contradiction that there exists a sequence $\varepsilon \rightarrow 0^+$ such that $M_{s,\varepsilon} = \hat{M}_{s,\varepsilon}$. By Lemma 4.3 we get that $M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| \not\rightarrow \infty$, where x_ε is any point such that $|u_{s,\varepsilon}(x_\varepsilon)| = M_{s,\varepsilon}$. Notice that by construction we have $|x_\varepsilon| \geq r_\varepsilon^1$ and thus $M_{s,\varepsilon}^{\beta_{s,\varepsilon}} r_\varepsilon^1 \not\rightarrow +\infty$ too.

As a consequence, up to a subsequence, $M_{s,\varepsilon}^{\beta_{s,\varepsilon}} r_\varepsilon^1 \rightarrow l$ for some real number $l \geq 0$. Let $\tilde{u}_{s,\varepsilon}$ be the rescaling defined in (4.1). Then, by Lemma 4.4 we infer that $\tilde{u}_{s,\varepsilon} \rightarrow \tilde{u}_s$ in $C_{loc}^{0,\alpha}(\mathbb{R}^n)$ for some $\alpha \in (0, s)$, $\tilde{u}_s \in C_{loc}^{0,\alpha}(\mathbb{R}^n)$. On the other hand, let $(y_\varepsilon) \subset \mathbb{R}^n$ be such that $|y_\varepsilon| = r_\varepsilon^1$. Up to a further subsequence, $M_{s,\varepsilon}^{\beta_{s,\varepsilon}} y_\varepsilon \rightarrow \hat{y}$ as $\varepsilon \rightarrow 0^+$, where $|\hat{y}| = l$. Then, thanks to Proposition 4.2 and since $\tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} y_\varepsilon) = 0$, $\tilde{u}_{s,\varepsilon} \rightarrow \tilde{u}_s$ a.e., we get that

$$|\tilde{u}_s(\hat{y})| \leq |\tilde{u}_{s,\varepsilon}(\hat{y}) - \tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} y_\varepsilon)| + |\tilde{u}_{s,\varepsilon}(\hat{y}) - \tilde{u}_s(\hat{y})| \leq C|\hat{y} - M_{s,\varepsilon}^{\beta_{s,\varepsilon}} y_\varepsilon|^\alpha + o(1) = o(1).$$

From this we deduce that $\tilde{u}_s(\hat{y}) = 0$, which is a contradiction, since by Lemma 4.4 it holds that $\tilde{u}_s > 0$.

The proof of (i) is identical and we omit it. Let us prove (ii). Let $x_\varepsilon \in \mathbb{R}^n$ be such that $|u_{s,\varepsilon}(x_\varepsilon)| = \hat{M}_{s,\varepsilon}$. We claim that $\tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} x_\varepsilon) \rightarrow 0$.

Indeed, if it is not the case, up to a subsequence, we find $c \in (0, 1]$ such that $\tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} x_\varepsilon) \rightarrow c$. Thanks to Proposition 4.2 and arguing as in the proof of Lemma 4.3 we obtain that there exists a positive constant C_1 and a small positive number τ_0 , both independent on ε , such that for all sufficiently small $\varepsilon > 0$

$$|\tilde{u}_{s,\varepsilon}| \geq C_1 > 0, \quad \forall x \in B_{M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| + \tau_0} \setminus B_{M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| - \tau_0}.$$

Since $M_{s,\varepsilon}^{\beta_{s,\varepsilon}} |x_\varepsilon| > M_{s,\varepsilon}^{\beta_{s,\varepsilon}} r_\varepsilon^1 \rightarrow +\infty$, because of (i), we obtain that $|\tilde{u}_{s,\varepsilon}|_{2_s^* - \varepsilon} \rightarrow +\infty$, which contradicts Lemma 4.1, (ii). The claim is thus proved.

Now, in order to conclude the proof of (ii), we notice that

$$1 = \frac{|u_{s,\varepsilon}(x_\varepsilon)|}{\hat{M}_{s,\varepsilon}} = \frac{|\tilde{u}_{s,\varepsilon}(M_{s,\varepsilon}^{\beta_{s,\varepsilon}} x_\varepsilon)| M_{s,\varepsilon}}{\hat{M}_{s,\varepsilon}},$$

and thanks to the previous claim we obtain the desired result. \square

An immediate consequence of the previous result is the following

Lemma 4.7. *Let $s \in (0, 1)$ and $n > 2s$. The function \tilde{u}_s given by Lemma 4.4 is a standard bubble centered at the origin, i.e. \tilde{u}_s is of the form (2.2) with $x_0 = 0$.*

Proof. Since \tilde{u}_s is radial and satisfies (4.5), in order to prove the desired result we only need to show that \tilde{u}_s realizes the infimum in the fractional Sobolev inequality, that is,

$$S_s = \frac{\|\tilde{u}_s\|_s^2}{|\tilde{u}_s|_{2_s^*}^2}.$$

Now, since $\|\tilde{u}_s\|_s^2 = |\tilde{u}_s|_{2_s^*}^{2_s^*}$, by the Sobolev inequality, we readily infer that

$$S_s \leq \frac{\|\tilde{u}_s\|_s^2}{|\tilde{u}_s|_{2_s^*}^2} \leq |\tilde{u}_s|_{2_s^*}^{2_s^* - 2},$$

and to conclude it suffices to show that $|\tilde{u}_s|_{2_s^*}^{2_s^*} \leq S_s^{\frac{n}{2s}}$. To this end, we set

$$u_{s,\varepsilon}^1(x) := \begin{cases} u_{s,\varepsilon}(x) & |x| \leq r_\varepsilon^1, \\ 0 & \text{otherwise,} \end{cases}$$

and define

$$\tilde{u}_\varepsilon^1(x) := \frac{1}{M_{s,\varepsilon}^+} u_{s,\varepsilon}^1 \left(\frac{x}{(M_{s,\varepsilon}^+)^{\beta_{s,\varepsilon}}} \right), \quad x \in \mathbb{R}^n.$$

Thanks to Lemma 4.6, for all sufficiently small $\varepsilon > 0$ we have $M_{s,\varepsilon} = M_{s,\varepsilon}^+$. Then we readily infer that $|\tilde{u}_{s,\varepsilon} - \tilde{u}_{s,\varepsilon}^1|_\infty \leq \frac{\hat{M}_{s,\varepsilon}}{M_{s,\varepsilon}}$ which, again by Lemma 4.6, implies that $\tilde{u}_{s,\varepsilon} - \tilde{u}_{s,\varepsilon}^1 \rightarrow 0$ uniformly in \mathbb{R}^n , as $\varepsilon \rightarrow 0^+$. Then, by Fatou's Lemma, taking into account that $|\tilde{u}_{s,\varepsilon}^1|_{2_s^* - \varepsilon} = (M_{s,\varepsilon}^+)^{-\varepsilon \left(\frac{n-2s}{2s}\right)} |u_{s,\varepsilon}^1|_{2_s^* - \varepsilon}$, that eventually $M_{s,\varepsilon}^+ \geq 1$, and Lemma 4.1, we get that

$$|\tilde{u}_s|_{2_s^*}^{2_s^*} \leq \liminf_{\varepsilon \rightarrow 0^+} |\tilde{u}_{s,\varepsilon}^1|_{2_s^* - \varepsilon}^{2_s^* - \varepsilon} \leq \liminf_{\varepsilon \rightarrow 0^+} |u_{s,\varepsilon}^1|_{2_s^* - \varepsilon}^{2_s^* - \varepsilon} \leq \liminf_{\varepsilon \rightarrow 0^+} |u_{s,\varepsilon}^+|_{2_s^* - \varepsilon}^{2_s^* - \varepsilon} = S_s^{\frac{n}{2s}}.$$

The proof is then complete. \square

5. CHARACTERIZATION OF THE NODAL SET

In this section we study the nodal set of least energy radial sign-changing solutions to Problem (1.2). We begin with a couple of known preliminary results, which provide, respectively, an upper bound on the number of sign changes and a characterization of the nodal set.

Lemma 5.1. *Let $n > 2s$, $s \in (0, 1)$. Let $u_{s,\varepsilon}$ be a least energy radial sign-changing solution to Problem (1.2). There exists $\tilde{\varepsilon}_s \in (0, 2s^* - 2)$ such that, if $\varepsilon \in (0, \tilde{\varepsilon}_s)$, then $u_{s,\varepsilon} = u_{s,\varepsilon}(r)$ changes sign at most twice.*

Let $0 < s_0 < s_1 \leq 1$ and $n > 2s_1$. Then there exists $\tilde{\varepsilon} > 0$, independent on s , such that the same result holds for every $s \in [s_0, s_1]$ and $\varepsilon \in (0, \tilde{\varepsilon})$.

Proof. It suffices to argue as in [8, Theorem 5.1] first, and then as in [8, Theorem 5.2], taking into account Lemma 3.3 and Lemma 4.1. In particular $\hat{\varepsilon} > 0$ is given by Lemma 3.3. \square

Lemma 5.2. *Let $s \in (0, 1)$ and $n > 2s$. There exists $\check{\varepsilon}_s > 0$ such that for all $\varepsilon \in (0, \check{\varepsilon}_s)$ any least energy radial sign-changing solution $u_{s,\varepsilon}$ to (1.2) vanishes only at the nodes.*

Moreover, let $0 < s_0 < s_1 \leq 1$, $n > 2s_1$. Then the above result hold true for every $s \in [s_0, s_1]$ and $\varepsilon \in (0, \check{\varepsilon})$, for some $\check{\varepsilon} > 0$ independent on s .

Proof. It suffices to take $\check{\varepsilon}_s := \min\{\bar{\varepsilon}_s, \tilde{\varepsilon}_s\}$, where $\bar{\varepsilon}_s, \tilde{\varepsilon}_s$ are given by Lemma 4.5 and Lemma 5.1, respectively. Then, the results follows immediately by adapting the arguments of [8, Theorem 1.2]. \square

In the next Lemma we prove the upper semi-continuity of the map $s \rightarrow C_{\mathcal{M}^r(B_R)}(s, \varepsilon)$.

Lemma 5.3. *Let $0 < s_0 < s_1 \leq 1$, $n > 2s_1$ and $\varepsilon \in (0, \check{\varepsilon})$, where $\check{\varepsilon}$ is given by in Lemma 5.2. Then for every $\sigma \in [s_0, s_1]$ we have*

$$\limsup_{s \rightarrow \sigma} C_{\mathcal{M}^r(B_R)}(s, \varepsilon) \leq C_{\mathcal{M}^r(B_R)}(\sigma, \varepsilon).$$

Proof. Let us fix s_0, s_1, n and ε as in the statement. Let $(s_k)_k \subset [s_0, s_1]$ be a sequence such that $s_k \rightarrow \sigma \in [s_0, s_1]$, and consider a radial solution $u_{\sigma,\varepsilon}$ of (1.2) which realizes $C_{\mathcal{M}^r(B_R)}(\sigma, \varepsilon)$. Assume that $\sigma < 1$. We aim to construct a sequence of almost minimizers of $C_{\mathcal{M}^r}(s_k, \varepsilon)$. We proceed in three different steps. We point out that when $\sigma = 1$ the proof is identical, taking into account the conventions $(-\Delta)^1 u = -\Delta u$, $\|u\|_1^2 = |\nabla u|_2^2$, and that $(u^+, u^-)_1 \equiv 0$ for all $u \in H_0^1(B_R)$.

Step 1. There exists a sequence $(\varphi_j)_j \subset C_c^\infty(B_R) \cap \mathcal{M}_{\sigma,\varepsilon}^r(B_R)$ such that

- (1) $\text{supp}(\varphi_j^\pm) \subset \text{supp}(u_{\sigma,\varepsilon}^\pm)$,
- (2) $\varphi_j \rightarrow u_{\sigma,\varepsilon}$ in $X_0^\sigma(B_R)$, as $j \rightarrow +\infty$,
- (3) $I_{\sigma,\varepsilon}(\varphi_j) \rightarrow I_{\sigma,\varepsilon}(u_{\sigma,\varepsilon})$, as $j \rightarrow +\infty$.

We first observe that, thanks to Lemma 5.2, the boundaries of $\text{supp}(u_{\sigma,\varepsilon}^\pm)$ consist in a finite union of spheres. Therefore, adapting known density results (see e.g. [20]) we find two sequences of radial functions $(\tilde{\varphi}_j^\pm)_j \subset C_c^\infty(B_R)$ such that $\tilde{\varphi}_j^\pm \geq 0$, $\text{supp}(\tilde{\varphi}_j^\pm) \subset \text{supp}(u_{\sigma,\varepsilon}^\pm)$ for all j , and $\tilde{\varphi}_j^\pm \rightarrow u_{\sigma,\varepsilon}^\pm$ in $X_0^\sigma(B_R)$. Observe that, from the continuity of the scalar product, we have $(\tilde{\varphi}_j^+, \tilde{\varphi}_j^-)_\sigma \rightarrow (u_{\sigma,\varepsilon}^+, u_{\sigma,\varepsilon}^-)_\sigma$.

Now we recall that it is always possible to find $\alpha_j > 0$, $\beta_j > 0$ such that $\alpha_j \tilde{\varphi}_j^+ - \beta_j \tilde{\varphi}_j^- \in \mathcal{M}_{\sigma,\varepsilon}^r(B_R)$ (see e.g. [8, Remark 3.4]), which is equivalent to solve the following

$$\begin{aligned} \alpha_j^{2_\sigma^*-2-\varepsilon} |\tilde{\varphi}_j^+|_{2_\sigma^*-\varepsilon}^{2_\sigma^*-\varepsilon} + \frac{\beta_j}{\alpha_j} (\tilde{\varphi}_j^+, \tilde{\varphi}_j^-)_\sigma &= \|\tilde{\varphi}_j^+\|_\sigma^2, \\ \beta_j^{2_\sigma^*-2-\varepsilon} |\tilde{\varphi}_j^-|_{2_\sigma^*-\varepsilon}^{2_\sigma^*-\varepsilon} + \frac{\alpha_j}{\beta_j} (\tilde{\varphi}_j^+, \tilde{\varphi}_j^-)_\sigma &= \|\tilde{\varphi}_j^-\|_\sigma^2. \end{aligned} \quad (5.1)$$

We claim that, eventually, $0 < \underline{\alpha} < \alpha_j < \bar{\alpha}$ and $0 < \underline{\beta} < \beta_j < \bar{\beta}$ for some positive constants $\underline{\alpha}, \bar{\alpha}, \underline{\beta}, \bar{\beta}$. Indeed, since $\tilde{\varphi}_j^\pm \rightarrow u_{\sigma,\varepsilon}^\pm$, and $u_{\sigma,\varepsilon}^\pm$ are non trivial, then the quantities $|\tilde{\varphi}_j^\pm|_{2_\sigma^*-\varepsilon}^{2_\sigma^*-\varepsilon}$, $\|\tilde{\varphi}_j^\pm\|_\sigma^2$, $(\tilde{\varphi}_j^+, \tilde{\varphi}_j^-)_\sigma$ are uniformly bounded and uniformly away from zero. Moreover, by the definition of the scalar product we always have $(\tilde{\varphi}_j^+, \tilde{\varphi}_j^-)_\sigma < 0$. Then, treating (5.1) as an algebraic system in α_j, β_j having as coefficients $|\tilde{\varphi}_j^\pm|_{2_\sigma^*-\varepsilon}^{2_\sigma^*-\varepsilon}$, $\|\tilde{\varphi}_j^\pm\|_\sigma^2$, $(\tilde{\varphi}_j^+, \tilde{\varphi}_j^-)_\sigma$, it is easy to verify that it cannot happen that, up to a sequence, $\alpha_j \rightarrow +\infty$ or $\alpha_j \rightarrow 0^+$, and the same holds for β_j . The claim is thus proved.

Let us consider the sequence defined by $\varphi_j := \alpha_j \tilde{\varphi}_j^+ - \beta_j \tilde{\varphi}_j^-$. By construction $(\varphi_j)_j \subset C_c^\infty(B_R)$, and in view of (5.1) we have $(\varphi_j)_j \subset \mathcal{M}_{\sigma,\varepsilon}^r(B_R)$. We claim that $\varphi_j \rightarrow u_{\sigma,\varepsilon}$ in $X_0^\sigma(B_R)$.

Indeed, observe that, since $u_{\sigma,\varepsilon} \in \mathcal{M}_{\sigma,\varepsilon}^r(B_R)$, then

$$\|u_{\sigma,\varepsilon}^\pm\|_\sigma^2 = |u_{\sigma,\varepsilon}^\pm|_{2_\sigma^*-\varepsilon}^{2_\sigma^*-\varepsilon} + (u_{\sigma,\varepsilon}^+, u_{\sigma,\varepsilon}^-)_\sigma,$$

then, up to a sequence, setting $\alpha := \lim_{j \rightarrow +\infty} \alpha_j$, $\beta := \lim_{j \rightarrow +\infty} \beta_j$ and passing to the limit in (5.1) we infer that

$$\begin{aligned} \alpha(\alpha^{2_\sigma^*-2-\varepsilon} - 1) |u_{\sigma,\varepsilon}^+|_{2_\sigma^*-\varepsilon}^{2_\sigma^*-\varepsilon} &= -(\beta - \alpha)(u_{\sigma,\varepsilon}^+, u_{\sigma,\varepsilon}^-)_\sigma, \\ \beta(\beta^{2_\sigma^*-2-\varepsilon} - 1) |u_{\sigma,\varepsilon}^-|_{2_\sigma^*-\varepsilon}^{2_\sigma^*-\varepsilon} &= -\alpha(\alpha^{2_\sigma^*-2-\varepsilon} - 1) |u_{\sigma,\varepsilon}^+|_{2_\sigma^*-\varepsilon}^{2_\sigma^*-\varepsilon}. \end{aligned}$$

Recalling that $(u_{\sigma,\varepsilon}^+, u_{\sigma,\varepsilon}^-)_\sigma < 0$ it is immediate to see that both $0 < \alpha < 1$ and $\alpha > 1$ lead to a contradiction. Hence $\alpha = 1$, and as a consequence we obtain that $\beta = 1$. Finally, from this and since

$$\|u_{\sigma,\varepsilon} - \varphi_j\|_\sigma \leq |\alpha_j - 1| \|\tilde{\varphi}_j^+\|_\sigma + |\beta_j - 1| \|\tilde{\varphi}_j^-\|_\sigma + \|\tilde{\varphi}_j^+ - u_{\sigma,\varepsilon}^+\|_\sigma + \|\tilde{\varphi}_j^- - u_{\sigma,\varepsilon}^-\|_\sigma,$$

we obtain that $\varphi_j \rightarrow u_{\sigma,\varepsilon}$ in $X_0^\sigma(B_R)$, as $j \rightarrow +\infty$. At the end, the last point of Step 1 is a straightforward consequence of the strong convergence of φ_j to $u_{\sigma,\varepsilon}$, together with the fractional Sobolev embedding. The proof of Step 1 is complete.

Step 2. Let $(\varphi_j)_j \in C_c^\infty(B_R) \cap \mathcal{M}_{\sigma,\varepsilon}^r(B_R)$ be the sequence given by Step 1. Let $(s_k)_k$ be a sequence such that $s_k \rightarrow \sigma$, as $k \rightarrow +\infty$. For every j fixed, there exists a sequence $(\varphi_{j,k})_k \subset C_c^\infty(B_R)$ such that $\varphi_{j,k} \in \mathcal{M}_{s_k,\varepsilon}^r(B_R)$ for every k , and

$$\|\varphi_{j,k}\|_{s_k} \rightarrow \|\varphi_j\|_\sigma, \quad I_{s_k,\varepsilon}(\varphi_{j,k}) \rightarrow I_{\sigma,\varepsilon}(\varphi_j), \quad \text{as } k \rightarrow +\infty.$$

Let us fix j and let φ_j be as in the statement. From (4.8) and [17, Lemma 2.4], as $k \rightarrow +\infty$ we have $\|\varphi_j^\pm\|_{s_k} \rightarrow \|\varphi_j^\pm\|_\sigma$ and $\|\varphi_{j,k}\|_{s_k} \rightarrow \|\varphi_j\|_\sigma$. This easily implies that $(\varphi_{j,k}^+, \varphi_{j,k}^-)_{s_k} \rightarrow (\varphi_j^+, \varphi_j^-)_\sigma$, while by a standard computation we get that $|\varphi_{j,k}^\pm|_{2_{s_k}^*-\varepsilon}^{2_{s_k}^*-\varepsilon} \rightarrow |\varphi_j^\pm|_{2_\sigma^*-\varepsilon}^{2_\sigma^*-\varepsilon}$.

Let $\alpha_k = \alpha(j, k) > 0$, $\beta_k = \beta(j, k) > 0$ be such that $\alpha_k \varphi_{j,k}^+ - \beta_k \varphi_{j,k}^- \in \mathcal{M}_{s_k,\varepsilon}^r(B_R)$ and define $\varphi_{j,k} := \alpha_k \varphi_{j,k}^+ - \beta_k \varphi_{j,k}^-$. Arguing as in Step 1 we get that, up to a subsequence, $\alpha_k, \beta_k \rightarrow 1$ as $k \rightarrow +\infty$. This easily implies that $\|\varphi_{j,k}\|_{s_k} \rightarrow \|\varphi_j\|_\sigma$ and $|\varphi_{j,k}|_{2_{s_k}^*-\varepsilon}^{2_{s_k}^*-\varepsilon} \rightarrow |\varphi_j|_{2_\sigma^*-\varepsilon}^{2_\sigma^*-\varepsilon}$, as $k \rightarrow +\infty$. The proof of step 2 is complete.

Step 3. Conclusion.

Let $(s_k)_k \subset (0, 1)$ be a sequence such that $s_k \rightarrow \sigma$. Let us fix a small number $\tau > 0$. Thanks to Step 1, there exists a function $\varphi_\tau \in C_c^\infty(B_R) \cap \mathcal{M}_{\sigma, \varepsilon}^r(B_R)$ such that

$$|I_{\sigma, \varepsilon}(u_{\sigma, \varepsilon}) - I_{\sigma, \varepsilon}(\varphi_\tau)| < \frac{\tau}{2}.$$

On the other hand, thanks to Step 2 there exist $\hat{k} = \hat{k}(\tau) > 0$ and a sequence of functions $(\varphi_k)_k$ such that $\varphi_k \in C_c^\infty(B_R) \cap \mathcal{M}_{s_k, \varepsilon}^r(B_R)$ and

$$|I_{s_k, \varepsilon}(\varphi_k) - I_{\sigma, \varepsilon}(\varphi_\tau)| < \frac{\tau}{2}, \quad \forall k \geq \hat{k}(\tau).$$

As a consequence, we get that

$$|I_{s_k, \varepsilon}(\varphi_k) - I_{\sigma, \varepsilon}(u_{\sigma, \varepsilon})| < \tau, \quad \forall k \geq \hat{k}(\tau).$$

Therefore, since $u_{\sigma, \varepsilon}$ is a minimizer and $\varphi_k \in \mathcal{M}_{s_k, \varepsilon}^r(B_R)$, we infer that for all $k \geq \hat{k}(\tau)$

$$C_{\mathcal{M}^r(B_R)}(s_k, \varepsilon) \leq I_{s_k, \varepsilon}(\varphi_k) \leq C_{\mathcal{M}^r(B_R)}(\sigma, \varepsilon) + \tau.$$

Taking the limsup as $k \rightarrow +\infty$ we get that

$$\limsup_{k \rightarrow +\infty} C_{\mathcal{M}^r(B_R)}(s_k, \varepsilon) \leq C_{\mathcal{M}^r(B_R)}(\sigma, \varepsilon) + \tau,$$

and since $\tau > 0$ is arbitrary we obtain the desired result. The proof is then complete. \square

In the next result we prove a uniform bound with respect to s for the L^∞ -norm of the solutions.

Lemma 5.4. *Let $0 < s_0 < s_1 \leq 1$, $n > 2s_1$ and $\varepsilon \in (0, \hat{\varepsilon})$, where $\hat{\varepsilon}$ is given by Lemma 3.3. Then there exists $C > 0$, depending on ε but not on s , such that*

$$C^{-1} \leq \sup_{s \in [s_0, s_1)} |u_{s, \varepsilon}|_\infty \leq C,$$

for every least energy radial sign-changing solution $u_{s, \varepsilon} \in \mathcal{M}_{s, \varepsilon}^r(B_R)$ of (2.4).

Proof. Let us fix s_0, s_1, n and ε as in the statement. The first inequality is trivial. As for the second one, it can be proved in two different ways. Indeed, from [21, Theorem 3.2] there exists $M \in C(\mathbb{R}^+)$ such that

$$|u_{s, \varepsilon}|_\infty \leq M(|u_{s, \varepsilon}|_{2_s^*}).$$

A careful analysis of the proof shows that the function M can be chosen in such a way that M depends only on n, R, s_0, s_1 and ε , but not on s . Since $u_{s, \varepsilon} \in \mathcal{M}_{s, \varepsilon}^r(B_R) \subset \mathcal{N}_{s, \varepsilon}(B_R)$ and $u_{s, \varepsilon}$ is a least energy sign-changing solution to (1.2), we infer that

$$C_{\mathcal{M}^r(B_R)}(s, \varepsilon) = \frac{2_s^* - 2 - \varepsilon}{2(2_s^* - \varepsilon)} \|u_{s, \varepsilon}\|_s^2. \quad (5.2)$$

Thus, thanks to the fractional Sobolev embedding and Lemma 3.3 we deduce that $|u_{s, \varepsilon}|_{2_s^*} \leq C_1$, for some constant $C_1 > 0$ independent on s . Similarly, using that $2C_{\mathcal{N}(B_R)}(s, \varepsilon) \leq C_{\mathcal{M}^r(B_R)}(s, \varepsilon)$ and Lemma 3.2 we obtain that $|u_{s, \varepsilon}|_{2_s^*} \geq C_0 > 0$, where C_0 does not depend on s , and the desired result easily follows.

Alternatively, we can argue as follows: fix s_0, s_1, n and ε as in the statement. Since $u_{s, \varepsilon}$ is a least energy sign-changing solution to (1.2) with $u_{s, \varepsilon} \in \mathcal{M}_{s, \varepsilon}^r(B_R) \subset \mathcal{N}_{s, \varepsilon}(B_R)$, and since Lemma 3.3 holds, by (5.2) we get that the quantity $|u_{s, \varepsilon}|_{2_s^* - \varepsilon}$ is uniformly bounded with respect to $s \in [s_0, s_1)$. Now, suppose by contradiction that there exists a sequence $(s_k)_k \subset [s_0, s_1)$

and a sequence $(u_{s_k, \varepsilon})_k$ such that $\delta_{s_k} := |u_{s_k, \varepsilon}|_\infty \rightarrow +\infty$, as $k \rightarrow +\infty$. Up to a subsequence, $s_k \rightarrow \sigma \in [s_0, s_1]$, as $k \rightarrow +\infty$. Let us consider the rescaled functions

$$v_k(x) := \frac{1}{\delta_{s_k}} u_{s_k, \varepsilon} \left(\frac{x}{\delta_{s_k}^{\beta_{s_k}}} \right), \quad x \in \mathbb{R}^n$$

where $\beta_{s_k} := \frac{2}{n-2s_k}$. We recall that

$$\|v_k\|_{s_k}^2 = \|u_{s_k, \varepsilon}\|_{s_k}^2, \quad |v_k|_{2_{s_k}^{*-2-\varepsilon}} = (\delta_{s_k})^{-\varepsilon \left(\frac{n-2s_k}{2s_k} \right)} |u_{s_k, \varepsilon}|_{2_{s_k}^{*-2-\varepsilon}},$$

and v_k weakly satisfies

$$\begin{cases} (-\Delta)^{s_k} v_k = \frac{1}{\delta_{s_k}^\varepsilon} |v_k|_{2_{s_k}^{*-2-\varepsilon}} v_k & \text{in } B_{\delta_{s_k}^{\beta_{s_k}}}, \\ v_k = 0 & \text{in } B_{\delta_{s_k}^{\beta_{s_k}}}^c. \end{cases} \quad (5.3)$$

Arguing exactly as in Lemma 4.3 we see that $v_k \rightarrow v$ in $C_{loc}^{0, \alpha}(\mathbb{R}^n)$, for some $\alpha \in (0, s_0)$, where $v \not\equiv 0$. On the other hand, by Fatou's Lemma we have

$$|v|_{2_{\sigma}^{*-2-\varepsilon}} \leq \liminf_{k \rightarrow +\infty} |v_k|_{2_{s_k}^{*-2-\varepsilon}} = \liminf_{k \rightarrow +\infty} (\delta_{s_k})^{-\varepsilon \left(\frac{n-2s_k}{2s_k} \right)} |u_{s_k, \varepsilon}|_{2_{s_k}^{*-2-\varepsilon}} = 0,$$

because $\delta_{s_k} \rightarrow +\infty$ and $|u_{s_k, \varepsilon}|_{2_{s_k}^{*-2-\varepsilon}}$ is bounded. Hence $v \equiv 0$ and we get a contradiction. The proof is complete. \square

In the next result we study the asymptotic behavior of the solutions as s goes to some limit value.

Lemma 5.5. *Let $0 < s_0 < s_1 \leq 1$, $n > 2s_1$ and $\varepsilon \in (0, \hat{\varepsilon})$, where $\hat{\varepsilon}$ is given by Lemma 3.3. Let $s_k \rightarrow \sigma$, where $\sigma \in [s_0, s_1]$, and let $(u_{s_k, \varepsilon})_k$ be a sequence of least energy nodal radial solution to (1.2). Then*

$$u_{s_k, \varepsilon} \rightarrow u_{\sigma, \varepsilon} \text{ in } C_{loc}^{0, s_0}(\mathbb{R}^n),$$

where $u_{\sigma, \varepsilon} \in \mathcal{M}_{\sigma, \varepsilon}^r(B_R)$ weakly satisfies

$$\begin{cases} (-\Delta)^s u_{\sigma, \varepsilon} = |u_{\sigma, \varepsilon}|_{2_{\sigma}^{*-2-\varepsilon}} u_{\sigma, \varepsilon} & \text{in } B_R, \\ u_{\sigma, \varepsilon} = 0 & \text{in } \mathbb{R}^n \setminus B_R. \end{cases}$$

In addition, it holds that

$$\lim_{s_k \rightarrow \sigma} I_{s_k, \varepsilon}(u_{s_k, \varepsilon}) = I_{\sigma, \varepsilon}(u_{\sigma, \varepsilon}).$$

Proof. It suffices to argue as in [8, Theorem 6.7], taking into account Lemma 3.3 and Lemma 5.4. \square

As a corollary of the previous results we obtain the continuity of the map $s \mapsto C_{\mathcal{M}^r(B_R)}(s, \varepsilon)$.

Proposition 5.6. *Let $0 < s_0 < s_1 \leq 1$, $n > 2s_1$ and let $\varepsilon \in (0, \tilde{\varepsilon})$, where $\tilde{\varepsilon}$ is given by Lemma 5.2. Let $(s_k)_k \subset [s_0, s_1]$, $\sigma \in [s_0, s_1]$, $(u_{s_k, \varepsilon})_k$ and $u_{\sigma, \varepsilon}$ be as in Lemma 5.5. Then $u_{\sigma, \varepsilon}$ is a least energy solution, that is, $I_{\sigma, \varepsilon}(u_{\sigma, \varepsilon}) = C_{\mathcal{M}^r(B_R)}(\sigma, \varepsilon)$.*

In particular, for any $\varepsilon \in (0, \tilde{\varepsilon})$ the map from $[s_0, s_1]$ to \mathbb{R} , defined by $s \mapsto C_{\mathcal{M}^r(B_R)}(s, \varepsilon)$, is continuous.

Proof. Fixing s_0, s_1, n, ε as in the statement, applying both Lemma 5.3 and Lemma 5.5, since $0 < \tilde{\varepsilon} < \hat{\varepsilon}$, we infer that

$$C_{\mathcal{M}^r(B_R)}(\sigma, \varepsilon) \leq I_{\sigma, \varepsilon}(u_{\sigma, \varepsilon}) = \lim_{s \rightarrow \sigma} I_{s, \varepsilon}(u_{s, \varepsilon}) = \lim_{s \rightarrow \sigma} C_{\mathcal{M}^r(B_R)}(s, \varepsilon) \leq C_{\mathcal{M}^r(B_R)}(\sigma, \varepsilon),$$

which implies both stated results. \square

The following Lemma grants that every least energy nodal radial solution in a ball changes sign exactly once, when s is close to one.

Lemma 5.7. *Let $s_0 \in (0, 1)$ and $n \geq 3$. There exists $\varepsilon_0 > 0$ such that, for any $\varepsilon \in (0, \varepsilon_0)$, there exists $\bar{s} = \bar{s}(\varepsilon) \in (0, 1)$ such that for any $s \in (\bar{s}, 1)$ any least energy radial sign-changing solution $u_{s,\varepsilon}$ to (1.2) changes sign exactly once.*

Proof. We begin by recalling that, in the local case, when $n \geq 3$ there exists $\varepsilon_1 > 0$ such that, for every $\varepsilon \in (0, \varepsilon_1)$, least energy radial sign-changing solutions to

$$\begin{cases} -\Delta u = |u|^{2_1^* - 2 - \varepsilon} u & \text{in } B_R, \\ u = 0 & \text{in } \mathbb{R}^n \setminus B_R, \end{cases} \quad (5.4)$$

change sign exactly once (see e.g. [3]). Now, let us fix $s_0 \in (0, 1)$ and define $\varepsilon_0 := \min\{\check{\varepsilon}, \varepsilon_1\}$, where $\check{\varepsilon}$ is given by Lemma 5.2 for s_0 and $s_1 = 1$. Let us fix $\varepsilon \in (0, \varepsilon_0)$ and assume by contradiction that there exist $(s_k)_k \subset [s_0, 1)$ such that $s_k \rightarrow 1^-$ and a sequence $(u_{s_k,\varepsilon})_k$ of least energy radial sign-changing solutions in B_R which change sign exactly twice for any k (these functions change sign at most twice in view of Lemma 5.2). Then, by Proposition 5.6 we have that $u_{s_k,\varepsilon} \rightarrow u_{1,\varepsilon}$ in $C^{0,\alpha}(\mathbb{R}^n)$, for some $\alpha \in (0, s_0)$, and that $u_{1,\varepsilon}$ is a least energy sign-changing solution to (5.4). In particular, $u_{1,\varepsilon}$ changes sign exactly once, in view of our choice of ε .

On the other hand, arguing as in the proof of [8, Theorem 1.3] we infer that the number of sign changes is preserved when passing to the limit as $s \rightarrow 1^-$, and thus $u_{1,\varepsilon}$ has to change sign twice. This gives a contradiction and concludes the proof. \square

Finally, we can state and prove Theorem 5.8. We first recall that, when speaking of a $L^2(B_R)$ -continuous family $\mathcal{A} = \{v_{s,\varepsilon}\}_{s \in [s_0, 1)}$ of least energy nodal radial solutions to Problem (1.2), we mean a map $\Phi : [s_0, 1) \rightarrow L^2(B_R)$ such that Φ is continuous and $\Phi(s) = v_{s,\varepsilon} \in \mathcal{M}_{s,\varepsilon}^r(B_R)$ is a least energy radial sign-changing solution to Problem (1.2) for any $s \in [s_0, 1)$.

Theorem 5.8. *Let $s_0 \in (0, 1)$ and $n \geq 3$. There exists $\varepsilon_0 > 0$ such that, for any $\varepsilon \in (0, \varepsilon_0)$, if there exists a $L^2(B_R)$ -continuous family $\mathcal{A} = \{v_{s,\varepsilon}\}_{s \in [s_0, 1)}$ of least energy nodal radial solutions to Problem (1.2), then every element of the family changes sign exactly once.*

Proof. Let us fix $s_0 \in (0, 1)$, and let $\varepsilon_0 > 0$ be the number given by Lemma 5.7. Let us fix $\varepsilon \in (0, \varepsilon_0)$ and observe that, in view of Lemma 5.7, there exists $\bar{s} \in (0, 1)$ such that for any $s \in (\bar{s}, 1)$, every least energy radial sign-changing solution to (1.2) changes sign only once. Let us fix $s_1 \in (\bar{s}, 1)$, let \mathcal{A} be as in the statement, and set

$$\mathfrak{S}_\varepsilon := \{s \in [s_0, s_1] ; v_{s,\varepsilon} \text{ changes sign exactly once}\}.$$

In view of the previous discussion \mathfrak{S}_ε is not empty. We claim that \mathfrak{S}_ε is closed.

Indeed, let $(s_k)_k \subset \mathfrak{S}_\varepsilon$ be a sequence such that $s_k \rightarrow \sigma$, for some $\sigma \in [s_0, s_1]$, and consider the associated sequence $(v_{s_k,\varepsilon})_k \subset \mathcal{A}$. By Lemma 5.5 and thanks to Proposition 5.6, up to a subsequence, we have $v_{s_k,\varepsilon} \rightarrow u_\varepsilon$ in $C^{0,\alpha}(\overline{B_R})$ for some $\alpha \in (0, s_0)$, where $u_\varepsilon \in X_0^\sigma(B_R)$ is a least energy nodal radial solution of (1.2) with $s = \sigma$. In particular, $v_{s_k,\varepsilon} \rightarrow u_\varepsilon$ in $L^2(B_R)$ and, since we are assuming that \mathcal{A} is $L^2(B_R)$ -continuous, it holds that $u_\varepsilon = v_{\sigma,\varepsilon} \in \mathcal{A}$. Now, taking into account Lemma 5.2, since $v_{s_k,\varepsilon} \rightarrow u_\varepsilon$ in $C^{0,\alpha}(\overline{B_R})$ and $v_{s_k,\varepsilon}$ changes sign once for all k , we infer that the only possibility is that $v_{\sigma,\varepsilon}$ changes sign only once. Hence $\sigma \in \mathfrak{S}_\varepsilon$, and the claim is proved.

We claim that $\mathfrak{S}_{s_0,\varepsilon}$ is open. To prove the claim we show that the complementary set $\mathfrak{S}_\varepsilon^c$ is closed. By definition and thanks to Lemma 5.2 we have

$$\mathfrak{S}_\varepsilon^c = \{s \in [s_0, s_1] ; v_{s,\varepsilon} \text{ changes sign exactly twice}\}.$$

Let $(s_k)_k \subset \mathfrak{S}_\varepsilon^c$ be such that $s_k \rightarrow \sigma$ for some $\sigma \in [s_0, s_1]$, as $k \rightarrow +\infty$. Arguing as before, up to a subsequence, we get that $v_{s_k, \varepsilon} \rightarrow v_{\sigma, \varepsilon}$ in $C^{0, \alpha}(\overline{B_R})$, for $\alpha \in (0, s_0)$.

Let us denote by $0 < r'_k < r''_k < R$ the nodes of $v_{s_k, \varepsilon}(r) = v_{s_k, \varepsilon}(x)$, $|x| = r$. We observe that $r'_k \not\rightarrow 0$. Indeed, if $r'_k \rightarrow 0$, as $k \rightarrow +\infty$, as a consequence of the $C^{0, \alpha}$ -convergence we infer that $v_{\sigma, \varepsilon}(0) = 0$. But this contradicts Lemma 5.2, and we are done.

Secondly, we claim that $r'_k - r''_k \not\rightarrow 0$. Indeed, assume by contradiction that $r'_k - r''_k \rightarrow 0$. Thanks to Lemma 5.4 we get that

$$\begin{aligned} |v_{s_k, \varepsilon}^-|_{2_{s_k}^* - \varepsilon}^{2_{s_k}^* - \varepsilon} &= \int_{B_{r''_k} \setminus B_{r'_k}} |v_{s_k, \varepsilon}^-|_{2_{s_k}^* - \varepsilon}^{2_{s_k}^* - \varepsilon} dx \\ &\leq C |v_{s_k, \varepsilon}^-|_{\infty}^{2_{s_k}^* - \varepsilon} \int_{r'_k}^{r''_k} \rho^{n-1} d\rho \leq C ((r''_k)^n - (r'_k)^n) \rightarrow 0. \end{aligned} \quad (5.5)$$

On the other hand, since $v_{s_k, \varepsilon} \in \mathcal{M}_{s_k, \varepsilon}^r(B_R)$ and thanks to Lemma 3.2 we find a constant $C > 0$ independent on k such that

$$C^\varepsilon S_{s_k} \leq S_{s_k, \varepsilon} \leq \frac{\|v_{s_k, \varepsilon}^-\|_{s_k}^2}{|v_{s_k, \varepsilon}^-|_{2_{s_k}^* - \varepsilon}^{2_{s_k}^* - 2 - \varepsilon}} < |v_{s_k, \varepsilon}^-|_{2_{s_k}^* - \varepsilon}^{2_{s_k}^* - 2 - \varepsilon}. \quad (5.6)$$

Hence, $|v_{s_k, \varepsilon}^-|_{2_{s_k}^* - \varepsilon}^{2_{s_k}^* - 2 - \varepsilon}$ is bounded away from zero and this contradicts (5.5).

It remains to prove that $r''_k \not\rightarrow R$. To this end, we first point out that, thanks [24, Theorem 1.2], it holds

$$\left\| \frac{v_{s_k, \varepsilon}}{\delta^{s_k}} \right\|_{C^{0, \alpha}(\overline{B_R})} \leq C |v_{s_k, \varepsilon}|_{\infty}^{2_{s_k}^* - 1 - \varepsilon},$$

where $0 < \alpha < \min\{s, 1 - s\}$,

$$\delta^s(x) := d(x, \partial B_R)^s = (R - |x|)^s \quad \text{and} \quad \frac{v_{s_k, \varepsilon}}{\delta^{s_k}}(R) := \lim_{\tau \rightarrow 0} \frac{v_{s_k, \varepsilon}(R - \tau)}{\delta^{s_k}(R - \tau)}.$$

A careful analysis of the proof shows that the constant $C > 0$ is uniform for $s \in [s_0, s_1]$ because s_1 is strictly less than one. Moreover we can fix α by choosing $0 < \alpha < \min\{s_0, 1 - s_1\}$.

Assume now by contradiction that $R - r''_k \rightarrow 0$ as $k \rightarrow +\infty$. Using the previous estimate, Lemma 5.4 and since $v_{s_k, \varepsilon}(r''_k) = 0$, we have

$$\left| \frac{v_{s_k, \varepsilon}}{\delta^{s_k}}(R) \right| = \left| \frac{v_{s_k, \varepsilon}}{\delta^{s_k}}(R) - \frac{v_{s_k, \varepsilon}(r''_k)}{\delta^{s_k}(r''_k)} \right| \leq C |R - r''_k|^\alpha \rightarrow 0. \quad (5.7)$$

On the other hand, applying the fractional Pohozaev identity (see [25]) to (1.2) we get that

$$\frac{2n - (n - 2s_k)(2_{s_k}^* - \varepsilon)}{2_{s_k}^* - \varepsilon} |v_{s_k, \varepsilon}|_{2_{s_k}^* - \varepsilon}^{2_{s_k}^* - \varepsilon} = \Gamma(1 + s_k)^2 R |\partial B_R| \left| \frac{v_{s_k, \varepsilon}}{\delta^{s_k}}(R) \right|^2$$

which, together with (5.7), implies that $|v_{s_k, \varepsilon}|_{2_{s_k}^* - \varepsilon}^{2_{s_k}^* - \varepsilon} \rightarrow 0$ as $s_k \rightarrow \sigma$, thus contradicting (5.6).

From this discussion it follows that r'_k and r''_k definitely stay in the interior of the domain, away from the origin and their distance does not tend to zero. Thanks to the $C^{0, \alpha}$ -convergence and by Lemma 5.2 we infer that also $v_{\sigma, \varepsilon}$ changes sign exactly twice. Hence $\sigma \in \mathfrak{S}_\varepsilon^c$, and thus $\mathfrak{S}_\varepsilon^c$ is a closed set. At the end, \mathfrak{S}_ε is not empty and both open and closed, thus it coincides with the whole interval $[s_0, s_1]$. Since by construction $s_1 > \bar{s}$ we conclude that every element of \mathcal{A} changes sign exactly once. The proof is complete. \square

REFERENCES

- [1] F. V. Atkinson, H. Brezis and L. A. Peletier, Solutions d'équations elliptiques avec exposant de Sobolev critique qui changent de signe, *C. R. Acad. Sci. Paris Sér. I Math.*, **306** (1988), no.16, 711–714.
- [2] M. Ben Ayed, K. El Mehdi and F. Pacella, Blow-up and symmetry of sign-changing solutions to some critical elliptic equations, *Journal of Diff. Eq.*, **230** (2006), 771–795.
- [3] M. Ben Ayed, K. El Mehdi and F. Pacella, Classification of low energy sign-changing solutions of an almost critical problem, *Journal of Funct. Anal.*, **250** (2007), 347–373.
- [4] X. Cabré and Y. Sire, Nonlinear equations for fractional Laplacians, I: Regularity, maximum principles, and Hamiltonian estimates, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **31** (2014), 2353.
- [5] W. Chen, C. Li and Y. Li, A direct method of moving planes for the fractional Laplacian, *Adv. in Math.*, **308** (2017), 404–437.
- [6] Y. Cho and T. Ozawa, Sobolev inequalities with symmetry, *Commun. Contemp. Math.*, **11** (2009), 355–365.
- [7] W. Choi, S. Kim and K.-A. Lee, Asymptotic behavior for solutions for nonlinear elliptic problems with the fractional Laplacian, *J. Funct. Anal.*, **266** (2014), 6531–6598.
- [8] G. Cora and A. Iacopetti, On the structure of the nodal set and asymptotics of least energy sign-changing radial solutions of the fractional Brezis-Nirenberg problem, *Nonlinear Analysis*, **176** (2018), 226–271.
- [9] A. Cotsiolis and N. Tavoularis, Best constants for Sobolev inequalities for higher order fractional derivatives, *J. Math. Anal. Appl.*, **295** (2004), 225–236.
- [10] R. L. Frank and E. Lenzmann, Uniqueness of non-linear ground states for fractional Laplacians in \mathbb{R} , *Acta Math.*, **210** (2013), 261–318.
- [11] R. L. Frank, E. Lenzmann, and L. Silvestre, Uniqueness of radial solutions for the fractional Laplacian, *Comm. Pure Appl. Math.*, **69** (2016), 1671–1726.
- [12] Z.-C. Han, Asymptotic approach to singular solutions for nonlinear elliptic equations involving critical Sobolev exponent, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **8** (1991), no.2, 159–174.
- [13] E. Di Nezza, G. Palatucci and E. Valdinoci, Hitchhiker's guide to the fractional Sobolev spaces, *Bull. Sci. Math.*, **136** (2012), 521–573.
- [14] A. Iacopetti and F. Pacella, A nonexistence result for sign-changing solutions of the Brezis-Nirenberg problem in low dimensions, *Journal of Diff. Eq.*, **258** (2015), no.12, 4180–4208.
- [15] A. Iacopetti and G. Vaira, Sign-changing tower of bubbles for the Brezis-Nirenberg problem, *Commun. Contemp. Math.*, **18** (2016), no.1, 1550036.
- [16] A. Iacopetti and G. Vaira, Sign-changing blowing-up solutions for the Brezis-Nirenberg problem in dimensions four and five, *Ann. Sc. Norm. Sup. di Pisa*, Vol. XVIII (2018), 1–38.
- [17] M.M. Fall and E. Valdinoci, Uniqueness and Nondegeneracy of Positive Solutions of $(-\Delta)^s u + u = u^p$ in \mathbb{R}^N when s is Close to 1, *Commun. Math. Phys.*, **329** (2014), 383–404.
- [18] A. Farina, On the classification of solutions of the Lane-Emden equation on unbounded domains of \mathbb{R}^n , *J. Math. Pures Appl.*, **87** (2007), 537–561.
- [19] M. M. Fall and V. Felli, Unique Continuation Property and Local Asymptotics of Solutions to Fractional Elliptic Equations, *Comm. in PDE*, **39** (2014), no.2, 354–397.
- [20] A. Fiscella, R. Servadei and E. Valdinoci, Density properties for fractional Sobolev spaces, *Annales Academiæ Scientiarum Fennicæ Mathematica*, **40** (2015), 235–253.
- [21] A. Iannizzotto, S. Mosconi and M. Squassina, H^s versus C^0 -weighted minimizers, *Nonlinear Differ. Equ. Appl.*, **22** (2015), no.3, 477–497.
- [22] A. Pistoia and T. Weth, Sign-changing bubble tower solutions in a slightly subcritical semilinear Dirichlet problem, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **24** (2007), no.2, 325–340.
- [23] O. Rey, Proof of two conjectures of H. Brezis and L. A. Peletier, *Manuscripta Math.*, **65** (1989), 19–37.
- [24] X. Ros-Oton and J. Serra, The Dirichlet problem for the fractional Laplacian: regularity up to the boundary, *J. Math. Pures et Appliq.*, **101** (2014), 275–302.
- [25] X. Ros-Oton and J. Serra, The Pohozaev identity for the fractional Laplacian, *Arch. for Rat. Mech. and Anal.*, **213** (2014), no.2, 587 – 628.
- [26] R. Servadei, The Yamabe equation in a non-local setting, *Adv. Nonlinear Anal.*, **2** (2013), 235–270.
- [27] R. Servadei and E. Valdinoci, The Brezis-Nirenberg result for the fractional Laplacian, *Trans. Amer. Math. Soc.*, **367** (2015), 67–102.
- [28] K. Teng, K. Wang and R. Wang, A sign-changing solution for nonlinear problems involving the fractional Laplacian, *Electronic Journal of Differential Equations*, **109** (2015), 1–12.

E-mail address: gabriele.cora@unito.it

E-mail address: iacopetti@mat.uniroma1.it