



Characterization of Protective Immune Responses Induced by Pneumococcal Surface Protein A in Fusion with Pneumolysin Derivatives

Cibelly Goulart, Thais Raquel da Silva, Dunia Rodriguez, Walter Rodrigo Politano, Luciana C. C. Leite, Michelle Darrieux

Published: March 22, 2013 • <https://doi.org/10.1371/journal.pone.0059605>

Abstract

Pneumococcal surface protein A (PspA) and Pneumolysin derivatives (Pds) are important vaccine candidates, which can confer protection in different models of pneumococcal infection. Furthermore, the combination of these two proteins was able to increase protection against pneumococcal sepsis in mice. The present study investigated the potential of hybrid proteins generated by genetic fusion of PspA fragments to Pds to increase cross-protection against fatal pneumococcal infection. Pneumolysoids were fused to the N-terminus of clade 1 or clade 2 *pspA* gene fragments. Mouse immunization with the fusion proteins induced high levels of antibodies against PspA and Pds, able to bind to intact pneumococci expressing a homologous PspA with the same intensity as antibodies to rPspA alone or the co-administered proteins. However, when antibody binding to pneumococci with heterologous PspAs was examined, antisera to the PspA-Pds fusion molecules showed stronger antibody binding and C3 deposition than antisera to co-administered proteins. In agreement with these results, antisera against the hybrid proteins were more effective in promoting the phagocytosis of bacteria bearing heterologous PspAs *in vitro*, leading to a significant reduction in the number of bacteria when compared to co-administered proteins. The respective antisera were also capable of neutralizing the lytic activity of Pneumolysin on sheep red blood cells. Finally, mice immunized with fusion proteins were protected against fatal challenge with pneumococcal strains expressing heterologous PspAs. Taken together, the results suggest that PspA-Pd fusion proteins comprise a promising vaccine strategy, able to increase the immune response mediated by cross-reactive antibodies and complement deposition to heterologous strains, and to confer protection against fatal challenge.

Citation: Goulart C, Silva TRd, Rodriguez D, Politano WR, Leite LCC, Darrieux M (2013) Characterization of Protective Immune Responses Induced by Pneumococcal Surface Protein A in Fusion with Pneumolysin Derivatives. PLoS ONE 8(3): e59605. <https://doi.org/10.1371/journal.pone.0059605>

Editor: Bernard Beall, Centers for Disease Control & Prevention, United States of America

Received: August 15, 2012; **Accepted:** February 15, 2013; **Published:** March 22, 2013

Copyright: © 2013 Goulart et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) and Fundação Butantan. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript

Competing interests: The authors have declared that no competing interests exist.

Introduction

Streptococcus pneumoniae is a major human pathogen, accounting for over 10% of total deaths in children under the age of five [1]. Despite the well established efficacy of conjugate vaccines against invasive disease, the high production costs involved in the conjugation processes limit their implementation in lower income countries, in which the burden of pneumococcal diseases is highest. Also, due to the limited number of polysaccharides included in the formulations, the extent of vaccine coverage tends to decrease as less prevalent serotypes emerge. In fact, serotype replacement has been observed after the introduction of PCV7 in different populations [2], [3]. Finally, serotype replacement is associated with the emergence of antibiotic resistant clones [4], reinforcing the need for cost-effective strategies that confer broad protection, such as protein-based vaccines.

PspA and Pneumolysin (Ply) are among the most well studied pneumococcal proteins; their contribution to virulence has been demonstrated with mutant strains lacking either one or both proteins, which have shown reduced fitness in different models of colonization, lung infection and bacteremia [5]. Mutant strains were cleared more rapidly from the lungs and blood of mice when compared to wild type counterparts [5], [6] and deposited more C3 *in vitro* [7]. Furthermore, the combination of both mutations had an additive effect on C3 deposition and pneumococcal clearance [6], suggesting that these proteins contribute synergistically to bacterial evasion of innate immune responses [6], [7].

Recombinant forms of PspA and Pneumolysin derivatives (Pds) have been investigated as potential vaccine candidates in different animal models, with promising results. The N-terminal region of PspA, which is responsible for inhibiting complement deposition on the bacterial surface [8], [9] and contains most of the immunogenic epitopes of the molecule [10], confers protection against invasive infection [11]–[13], lobar pneumonia [14] and colonization [15], [16]. Furthermore, it has been recently demonstrated that maternal immunization with PspA protects the offspring against pneumococcal infection [17]. The N-terminus of PspA, however, exhibits structural and serological variability [18]. Based on the observation that different PspA molecules induce antibodies with distinct degrees of cross-reactivity [19], [20] and cross-protection [11], [21], it has been suggested that PspA-based anti-

pneumococcal vaccines should include more than one molecule in order to extend coverage. The potential of PspA as a vaccine candidate has been further supported by human clinical trials, which have demonstrated the induction of antibodies with high cross-reactivity against heterologous molecules [21], which can passively protect mice against fatal pneumococcal infection [21].

Pneumolysin (Ply) is a cholesterol dependent cytolysin with several biological effects, such as activation of classical complement pathway [22], induction of apoptosis in numerous cells types [23], [24], impairment of ciliary function in the lungs and induction of oxidative burst by neutrophils [22]. In fact, the instillation of purified Ply in the lungs is sufficient to reproduce many aspects of pneumococcal pneumonia in rats (reviewed in [22]). Furthermore, Ply has been shown to interact with TLR-4 [25] and to induce TLR-4 independent activation of the NLRP3 inflammasome, contributing to host protection against pneumococcal pneumonia [26] and lethal infection [25].

Since Ply is toxic in its native form, several detoxified forms – named pneumolysoids (Pds) – have been produced, by site-directed mutagenesis or chemical detoxification, and evaluated for their immunogenicity and protective effect in different animal models, with variable results, including protection in rhesus macaques [14], [27]–[33]. Of those toxoids, the best characterized are PdB, carrying a Trp-Phe substitution at position 433 [30], and PdT, a triple mutant containing Asp-385 to Asn, Cys-428 to Gly and Trp-433 to Phe substitutions [34]. While PdT alone or co-administered with other pneumococcal antigens, did not induce significant protection against lethal intraperitoneal challenge [29], PdB has been shown to elicit protection against nasal challenge with some pneumococcal strains, which was enhanced by co-administration of other pneumococcal proteins, such as PspA and PhTB [14], [35]. The combination of PspA and PdB elicited the highest protection levels in mouse models of sepsis and focal pneumonia, suggesting a complementary role for these two antigens [14], [36].

On a whole, the results indicate that effective protein-based anti-pneumococcal vaccines tend to require the combination of different proteins in order to extend protection. In the present work, we investigated the ability of fusion proteins including the N-terminal region of family 1 PspAs and detoxified derivatives of Pneumolysin to induce protective immune responses in a mouse model of fatal pneumococcal challenge.

Materials and Methods

Pneumococcal Strains

All pneumococcal strains used in this study are shown in Table 1. Pneumococci were maintained as frozen stocks (−80°C) in Todd-Hewitt broth supplemented with 0.5% yeast extract (THY), with 10% glycerol. In each experiment, the isolates were plated on blood agar prior to growth in THY.

Strain	Serotype	PspA Clades	Source	Reference
P69	10A	1	UFG	20
94/01	18A	2	IAL	20
245/00	14	1	IAL	20
491/00	6B	1	IAL	20
472/96	6B	1	IAL	30
A66.1	3	2	UAB	42
D39	2	2	UAB	5

IAL: Instituto Adolfo Lutz, São Paulo, Brazil.
 UFG: Universidade Federal de Goiás, Goiânia, Brazil.
 UAB: University of Alabama at Birmingham.
 doi:10.1371/journal.pone.0059605.t001

Table 1. Pneumococcal strains used in this study.

<https://doi.org/10.1371/journal.pone.0059605.t001>

Cloning of pspAs, pds and Hybrid Genes

Gene fragments encoding the N-terminal region of *pspA* were amplified from pneumococcal strains 245/00 (PspA1) or 94/01 (PspA2) by PCR (Figure 1). The mutant detoxified Pneumolysin gene *pdT* was obtained by PCR from the pQE-30-pdT, kindly provided by Drs. Richard Malley and James Paton. Two more Ply mutants were obtained by site-directed mutagenesis, using the protocol described by Withers-Martinez *et al.* (1999). *Pd_{H367}* or *pid1* was amplified from pneumococcal strain D39 and contains one mutation on the His 367 residue, which was substituted by Arg. This mutant was first described by Berry *et al.*, 1995, and retains 0.02% of the hemolytic activity of the native protein. The same mutation was inserted in the *ply* gene from strain 472/96, which contains a natural Asp-380 to Asn substitution, in a region described as involved in complement activation (Mitchell *et al.*, 1991). This second mutant, *pd_{H367R380}* or *pid2*, contains, therefore, two mutations. The primers used to obtain the mutants are listed in Table S1. The *pspA* fragments and *ply* mutant genes were inserted into pGEM-T easy vector (Promega) and fused through complementary cohesive ends added to the primers, generating three chimeric genes: *pspA1-pid1*, *pspA1-pid2*, and *pspA2-pdT*. *pspA2-pdT* was digested with the appropriate restriction endonucleases and ligated to the linearized pQE30 (QIAGEN) expression vector; *pspA1-pid1* and *pspA1-pid2* were excised from pGEM-T easy and subcloned into linearized pAE-6xHis expression vector [37].

Hemolysis Inhibition Assay

The recombinant Ply was expressed using *E. coli* M15 - RM 86 clone, kindly provided by Drs. Richard Malley and James Paton, and purified by affinity chromatography. The hemolysis inhibition assay was performed in 96 wells plates. 200 μ L of sheep whole blood were washed 3 times with PBS and resuspended in 10 mL of PBS. The antisera produced against PspA2, PdT, PspA2+PdT or PspA2-PdT were incubated with 8 HU (hemolytic units) of Ply at 1:40 dilution at 37°C for 30 min. The antisera generated by immunization with PspA1, PID1, PID2, co-administered proteins or fused proteins PspA1-PID1 and PspA1-PID2 were incubated with 4 HU of Ply at 1:10 dilution at 37°C for 30 min. Serum from mice that received saline and Al(OH)₃ was used as control. 50 μ L aliquots of 2% red blood cells were added at a final concentration of 1%, followed by incubation at 37°C for 30 min. The plates were harvested by centrifugation at 1000 \times g for 10 min, and the supernatant absorbance was determined at 540 nm.

Challenge

Two weeks after the last immunization the animals were challenged with 4×10^3 CFUs of *S. pneumoniae* strain A66.1 or 5×10^6 CFUs of strain 491/00 injected by the intravenous route. The mice were monitored for 15 days and the differences between the survival rates in each group were analyzed by Mann–Whitney U test. Moribund mice or animals that developed paralysis were euthanized by CO₂ narcosis. At the endpoint, all surviving animals were euthanized.

Results

Expression and Purification of rPspA-Pd Hybrid Proteins

The gene fragments encoding rPspAs and rPds were fused with restriction enzymes and expressed in *E. coli* using vectors that add a Histidine tag to the beginning of the amino acid sequence. Both *pspA* gene fragments contain the N-terminal region including the proline rich region and the non-proline block; the scheme of proteins and hybrids is shown in Figure 1. The rPspA1-PID1 and rPspA1-PID2 hybrids were expressed in inclusion bodies, denatured with urea and refolded. rPspA2-PdT was expressed in soluble form. All proteins were purified through Ni²⁺-affinity chromatography and analyzed by immunoblotting using anti-rPspA1 or anti-rPspA2 and anti-rPly antibodies (Figure 2). All hybrid proteins were expressed and purified integrally and were recognized by antibodies against PspA or Ply individually.




Figure 2. Recognition of hybrid proteins by antibodies against PspA and Pds.

Recombinant proteins were separated by SDS-PAGE and transferred to PVDF membrane - A) rPspA2 and rPspA2-PdT; B) rPspA1 and rPspA1-PID1; C) rPspA1 and rPspA1-PID2; D) rPdT and rPspA2-PdT; E) rPID1 and rPspA1-PID1; and F) rPID2 and rPspA1-PID2. The membranes were incubated with anti-rPspA2 (A) anti-rPspA1 (B and C) or anti-rPly (D, E and F) followed by incubation with anti-mouse IgG conjugated with HRP. Detection was performed with an ECL kit (GE Healthcare). Molecular mass markers (kDa) are indicated on the left.

<https://doi.org/10.1371/journal.pone.0059605.g002>

Mouse Immunization with Hybrid Proteins Induce Levels of Antibodies Comparable to Proteins Administered Individually

Sera obtained by BALB/c immunization with rPspAs, rPds, and rPspA-Pds were quantified by ELISA against each recombinant protein. Immunization with the hybrid proteins induced antibody levels comparable to those obtained in the groups immunized with each antigen alone (Table 2).

Coating antigen	Antisera	Antibodies concentration ($\mu\text{g/mL}$)	P value
PspA 2	PspA2	13,350	<0,01
	PspA2-PdT	13,740	<0,01
PdT	PdT	91	<0,05
	PspA2-PdT	126	<0,01
PspA1	PspA1	10,140	<0,001
	PspA1-PID1	12,530	<0,001
	PspA1-PID2	7,000	<0,01
PID1	PID1	245	<0,05
	PspA1-PID1	313	<0,01
PID2	PID2	167	<0,05
	PspA1-PID2	241	<0,01

*p values were calculated in comparison with the control group.
doi:10.1371/journal.pone.0059605.t002

Table 2. Antibody levels in mice immunized with the recombinant proteins.
<https://doi.org/10.1371/journal.pone.0059605.t002>

Antibodies Generated by Mouse Immunization with Hybrid Proteins Bind to the Surface of Pneumococci Bearing Different PspAs

Sera from mice immunized with the recombinant proteins and hybrids were tested for their ability to bind onto the pneumococcal surface. Pneumococci bearing family 1 PspAs were incubated with the antisera followed by incubation with anti-mouse IgG-FITC. Antibodies generated against rPspA2-PdT were able to bind to the surface of pneumococci bearing homologous PspAs (clade 2), similarly to anti-rPspA2 or anti-rPspA2+PdT antisera (Figure 3–A and B). Interestingly, this same antiserum was able to bind with significantly higher affinity to pneumococcal strains bearing heterologous clade 1 PspAs when compared with antiserum generated against the co-administered proteins (Figure 3 G and H). Antisera generated against the other hybrids, rPspA1-PID1 and rPspA1-PID2, also revealed strong binding to pneumococcal strains, usually comparable to that observed with antisera against the co-administered proteins (Figures 3 C, E, D, I, K and L), with few exceptions (Figure 3 F and J). However, in some cases, the hybrid proteins showed a binding capacity lower than that observed for anti-PspA antisera alone (Figures 3 D, J and L). No binding was observed when anti-Pds antisera were used.




Figure 3. Antibody binding onto pneumococcal surface.

Pneumococcal strains containing PspA2 or PspA1 were incubated with antisera from mice immunized with rPspAs, rPds, co-administered proteins or hybrids, followed by incubation with anti-IgG mouse conjugated with FITC and analyzed by FACS. Serum from mice that received saline/ $\text{Al}(\text{OH})_3$ was used as a control. The percentage of fluorescent bacteria (>10 fluorescence intensity units) was calculated for each sample. Statistical analysis was performed by one-way ANOVA with a Tukey's Multiple Comparison Test : * $p<0.05$; ** $p<0.01$; *** $p<0.001$ for treated versus control or between immunized groups, as indicated.

<https://doi.org/10.1371/journal.pone.0059605.g003>

Antibodies Induced against rPspA-Pd Hybrids Induced an Increased C3 Deposition on Pneumococci Bearing Heterologous PspAs

The antisera were evaluated as to their ability to increase complement deposition on the surface of pneumococci. *S. pneumoniae* strains expressing PspA clade 1 or clade 2 were incubated with the antisera generated against the rPspA-Pd hybrids and respective controls, in the presence of a complement source, followed by incubation with anti-C3 conjugated with FITC, and these were analyzed by FACS. When we used a bacterium containing a homologous PspA, we observed that antibodies from mice immunized with co-administered proteins showed increased complement deposition when compared with anti-serum from the group immunized with rPspA alone (Figure 4–94/00 strain). However, comparable levels of C3 deposition were observed when the same bacteria was incubated with serum from the group immunized with rPspA2-PdT or the antisera induced against the co-administered proteins, which is in agreement with the results of antibody binding. On the other hand, the anti-rPspA2-PdT antibodies induced significantly higher amounts of C3 deposition on the bacterial surface, as compared to antibodies induced

against the co-administered proteins for all the other strains used in this study, including those bearing PspA2 or PspA1 (Figure 4 A66.1, 245/00, P69 strains). In relation to the antisera induced against the rPspA1-PID1 and rPspA1-PID2 hybrids, no significant differences were observed in C3 deposition on pneumococci when compared with antisera generated against rPspA1 alone or the co-administered proteins, also in accordance with antibody binding (Figure S1).




Figure 4. Complement deposition on pneumococcal surface in the presence of specific antibodies.

Pneumococcal strains were incubated with antisera from mice immunized with rPspAs, rPds, co-administered proteins or hybrids and NMS as complement source. After incubation with anti-C3 mouse conjugated with FITC, the samples were analyzed by FACS. Serum from mice that received saline/Al(OH)₃ was used as a control. The median of fluorescence intensity (MFI) was calculated for each sample. Statistical analysis was performed by one-way ANOVA with a Tukey's Multiple Comparison Test: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ for treated versus control or between immunized groups, as indicated. <https://doi.org/10.1371/journal.pone.0059605.g004>

In-vitro Opsonophagocytosis Mediated by Anti-hybrid Antisera is More Efficient than Antisera against Co-administered Proteins in Heterologous Strains

Since the immunization with the rPspA2-PdT hybrid induces antibodies that exhibited both a stronger binding capacity and a more pronounced C3 deposition on the bacterial surface when compared to the other groups, these antibodies were investigated for their ability to mediate the opsonophagocytosis and killing of pneumococci *in vitro*. Bacterial strains bearing PspA clade 1 or clade 2 were incubated with antisera against rPspA2, rPdT, rPspA2+rPdT and rPspA2-PdT and a complement source, followed by incubation with murine peritoneal phagocytic cells. The samples were plated and the number of CFUs recovered after 18 h were counted. When we used the pneumococcal strain 94/00, expressing the homologous PspA clade 2, antibodies generated against the hybrid, rPspA2-PdT, promoted a significant reduction in the number of CFU recovered when compared to control or anti-rPdT anti-serum (Figure 5). This reduction was similar to that observed when incubating with anti-rPspA2 or the sera from the co-administered antigens, anti-rPspA2+rPdT (Figure 5–94/00 strain). However, when a pneumococcal strain bearing a heterologous PspA molecule (clade 1) was used, the anti-rPspA2-PdT anti-serum was more efficient in promoting the opsonophagocytic killing of the bacterium than antisera against the co-administered proteins (Figure 5–245/00 strain).




Figure 5. Pneumococcal phagocytosis mediated by specific antibodies in the presence of complement.

Pneumococcal strains bearing PspA2 (94/01 strain) or PspA1 (245/00 strain) were incubated with antisera from mice immunized with rPspA2, rPdT, rPspA2+rPdT or rPspA2-PdT and NMS as complement source, followed by incubation with mouse peritoneal phagocytes and plated on blood agar plates. CFU recovered were counted after 18 h. Statistical analysis was performed by one-way ANOVA with a Tukey's Multiple Comparison Test. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ for treated versus control or between immunized groups, as indicated. <https://doi.org/10.1371/journal.pone.0059605.g005>

Inhibition of Ply Cytolytic Activity by Anti-pneumolysoids Antibodies

The ability of antibodies generated against PspA1, PspA2, pneumolysoids, co-administered proteins or fused proteins to inhibit the lytic effects of Pneumolysin was tested by incubation with sheep red blood cells in the presence of recombinant Ply. All antisera from formulations including pneumolysoids significantly reduced the hemolysis of red blood cells by Pneumolysin, with the exception of anti-PID1 (Figure 6). As expected, antibodies to PspA1 and 2 did not induce a significant inhibition of hemolysis (Figure 6).




Figure 6. Inhibition of hemolytic activity of Ply on red blood cells by antisera generated against recombinant and hybrids proteins. Ply was incubated with antisera and sheep red blood cells, and the supernatant absorbance was measured at 540 nm. Results are shown as percentages of the hemolytic activity in presence of sera from mice receiving saline and Al(OH)₃ (control). Statistical analysis was performed by one-way ANOVA with a Tukey's Multiple Comparison Test.; **p<0.01; ***p<0.001.
<https://doi.org/10.1371/journal.pone.0059605.g006>

Immunization with PspA-Pds Leads to an Increased Survival against Fatal Pneumococcal Challenge

The protective effect of rPspA-Pd fusions was evaluated in comparison with the isolated rPspA, rPds or the co-administered proteins, by intravenous lethal challenge with the virulent pneumococcal strains St 491/00 (Figure 7 - A) or A66.1 (Figure 7 - B and C), expressing PspA clades 1 or 2, respectively. Figure 6 shows the survival of mice up to 15 days after challenge, when the experiment was terminated. All rPspA-Pd hybrids induced higher survival rates in comparison with the control group or Pds. Furthermore, the animals immunized with the hybrids showed an increased, but not significant, survival when compared to those immunized with rPspA alone. In fact, two of the hybrids, rPspA1-PID1 and rPspA1-PID2 induced 100% protection against pneumococcal challenge. No differences in protection were observed between mice receiving the hybrids or co-administered proteins. Immunization with rPds alone, on the other hand, did not induce protection in this model of systemic infection.




Figure 7. Mouse immunization with rPspA-Pds confers protection against pneumococcal sepsis with strains bearing heterologous PspAs. BALB/c mice immunized with 3 doses of rPspAs, rPds, co-administered or hybrid proteins were challenged with lethal doses of pneumococcal strains bearing heterologous PspA: A – Mice were challenged with 491/00 strain (PspA1); B and C – Mice were challenged with A66.1 strain (PspA2). The mice were monitored for 15 days and differences between the survival rates in each group were analyzed by Mann–Whitney U test, *p<0.05; **p<0.01; ***p<0.001 for treated versus control groups.
<https://doi.org/10.1371/journal.pone.0059605.g007>

Discussion

PspA and Pds have been studied as vaccine candidates against pneumococcal infections for decades, with striking success in different animal models and, in the case of rPspA, induction of antibodies with high cross-reactivity [21] and protective potential in humans [21].

Since rPspA exhibits structural and serological variability [18], it has been suggested that the inclusion of two or more fragments would be necessary in order to increase vaccine coverage. Fusion of PspA fragments of families 1 and 2 have been demonstrated to increase protection against invasive pneumococcal infection [13], [39], as well as rPspA fusion or co-administration with adjuvant molecules [40], [41]. Nevertheless, the width of protection could be further extended by including a more conserved protein in the formulation. Several formulations containing more than one pneumococcal protein have been studied [14], [33], [42]–[44]. Furthermore, it has been suggested that vaccines including the same components as mixtures or fused formulations can differ in the levels of protection that they induce. A study from Lu *et al.* [42] demonstrated protection against fatal pneumococcal infection in mice immunized with trivalent vaccine containing fusions of rPdT, rPsaA and cell wall polysaccharide, but not with the antigens mixture. The same was observed using rPspA fused or mixed with flagellin, as an adjuvant molecule [40], suggesting that fused antigens could be more effective than co-administered formulations.

The present work investigated the potential of rPspA-Pd fusions to protect mice against invasive pneumococcal infections. Three fusions were produced, including two rPspA fragments and three different pneumolysoids. These fusion proteins were recognized by antibodies made against both PspA and the Pneumolysoids by Western blotting, showing that both proteins in the constructs were expressed and purified in the integral form, allowing for specific antibody recognition. The antibody levels were measured by ELISA against each recombinant protein. Although PspA was more immunogenic than the Pds (inducing antibody levels around a thousand times higher), immunization with the hybrids induced antibody levels comparable to those produced in mice immunized with each antigen alone, indicating that no antigenic competition occurred in the fusion.

Antibodies against the recombinant proteins and hybrids were evaluated for the ability to recognize and bind to intact pneumococci by FACS. Sera from mice immunized with the fusions revealed a strong binding capacity to all pneumococci tested, comparable – and in some cases superior – to that of sera against the co-administered proteins. Particularly, sera from mice immunized with rPspA2-PdT showed a stronger binding capacity to a PspA clade 1 bearing strain, when compared to the co-administered proteins. The correlation between sequence identity among PspAs and antibody binding has been demonstrated in other studies, with variable results [11], [13], [19], [20]. In general, a strong association between PspA type and the ability of induced antibodies to recognize the PspA molecules in the bacterial surface has been observed. Therefore, the significant increase in recognition of a PspA clade 1 strain by antibodies against a clade 2-containing hybrid suggests that the genetic fusion of PspA and Pd, may have a positive effect on the immune response induced against PspA.

Recent studies using cell fractionation and Western-blotting have demonstrated that Ply localizes in the pneumococcal cell wall compartment [45], [46]. However, the absence of antibody recognition observed with antibodies to Pds alone suggests that this protein is not accessible to antibodies in intact pneumococci. These results were in accordance with previous studies, which suggest that Ply is not displayed on the pneumococcal surface [47], [48]. Since antibodies to Pds alone do not interact with the bacterial surface, the increased binding capacity of anti-hybrid antiserum onto pneumococci bearing heterologous PspAs when compared with anti-rPspA antiserum could be explained by a possible modification in the PspA structure caused by PdT fusion, that promoted the presentation of more conserved epitopes or by an adjuvant effect of Ply [49].

Complement deposition is the key for opsonization and phagocytosis of pneumococci. Therefore, since PspA and Ply have both been shown to interact with complement components, we investigated the ability of the induced antibodies to promote C3 deposition onto the bacterial surface. In agreement with the binding results, antibodies to the rPspA2-PdT fusion protein mediated a

significant enhancement in the levels of C3 deposited on the bacterial surface in relation to antibodies generated against co-administered proteins.

In order to investigate whether the produced antibodies were able to mediate the opsonophagocytosis and killing of pneumococci, bacteria were incubated in the presence of sera induced against either the recombinant proteins alone, co-administered, or the fusion proteins, and mouse peritoneal phagocytes. Corroborating with the C3 complement deposition results, serum induced against the rPspA2-PdT fusion protein was able to promote the opsonophagocytosis and killing of pneumococcal strains. Furthermore, when compared with antisera induced against the co-administered proteins, the anti-hybrid antiserum showed a significantly increased ability to reduce the number of CFU recovered in a strain containing a heterologous PspA.

In accordance with the in vitro assays, which revealed a strong ability of antibodies against the hybrids to mediate complement deposition and phagocytic killing of bacteria, immunization with rPspA-Pd fusions protected mice against fatal challenge with pneumococcal strains bearing heterologous PspA molecules. The results also provide an insight on the mechanism responsible for protection in this model, with the induction of antibodies capable of enhancing C3 deposition on the bacterial surface, which in turn become more susceptible to phagocytic killing.

Quin et al (2007), using mutant Ply-negative pneumococci, observed that deletion of Ply did not affect blood clearance in comparison with a wild type strain. Therefore, we did not expect the Pds alone to be protective against an intravenous challenge. In fact, none of the Pds tested confer protection in this model of infection. However, anti-Pd antibodies could neutralize the lytic effects of Ply. The ability of such antibodies to mediate protective responses was evaluated through a hemolysis inhibition assay using sheep red blood cells. All formulations including Pds induced antibodies able to inhibit hemolysis by Ply, except PID1. This result indicates that antibodies against Pds may play a role in protection from *Streptococcus pneumoniae* infections by inhibiting Ply's lytic effects. The correlation between the capacity to inhibit Ply induced hemolysis and protection conferred by Pds has been confirmed by Salha et al (2012), using a detoxified mutant PlyD1 [32].

Taken together, the results suggest that PspA-Pd fusion proteins comprise a promising vaccine strategy, able to increase the immune response mediated by cross-reactive antibodies and complement deposition to heterologous strains, and to confer protection against fatal challenge.

Supporting Information

Figure S1.

Complement deposition on pneumococcal surface in the presence of specific antibodies. Pneumococcal strain D39 was incubated with antisera from mice immunized with rPspA1, rPID1, co-administered proteins or PspA1-PID1 hybrid (A), or rPspA1, PID2, co-administered proteins or PspA1-PID2 hybrid (B) and NMS as complement source. After incubation with anti-C3 mouse conjugated with FITC, the samples were analyzed by FACS. Serum from mice that received saline/Al(OH)₃ was used as a control. The median of fluorescence intensity (MFI) is shown for each sample.

<https://doi.org/10.1371/journal.pone.0059605.s001>
(TIF)

Table S1.

Oligonucleotides used in this study: sequence of the primers used for amplification of the PspA fragments and for insertion of the mutation on the Ply gene is shown.

<https://doi.org/10.1371/journal.pone.0059605.s002>
(DOCX)

Author Contributions

Conceived and designed the experiments: CG DR LCCL MD. Performed the experiments: CG TRS DR WRP MD. Analyzed the data: CG DR LCCL MD. Contributed reagents/materials/analysis tools: LCCL MD. Wrote the paper: CG DR LCCL MD.

References

- O'Brien KL, Wolfson LJ, Watt JP, Henkle E, Deloria-Knoll M, et al. (2009) Burden of disease caused by *Streptococcus pneumoniae* in children younger than 5 years: global estimates. *Lancet* 374: 893–902.
View Article • PubMed/NCBI • Google Scholar
- Singleton RJ, Hennessy TW, Bulkow LR, Hammitt LL, Zulz T, et al. (2007) Invasive pneumococcal disease caused by nonvaccine serotypes among alaska native children with high levels of 7-valent pneumococcal conjugate vaccine coverage. *JAMA* 297: 1784–1792.
View Article • PubMed/NCBI • Google Scholar
- Hsu HE, Shutt KA, Moore MR, Beall BW, Bennett NM, et al. (2009) Effect of pneumococcal conjugate vaccine on pneumococcal meningitis. *N Engl J Med* 360: 244–256.
View Article • PubMed/NCBI • Google Scholar
- Song JH, Dagan R, Klugman KP, Fritzell B (2012) The relationship between pneumococcal serotypes and antibiotic resistance. *Vaccine* 30: 2728–2737.
View Article • PubMed/NCBI • Google Scholar
- Ogunniyi AD, LeMessurier KS, Graham RM, Watt JM, Briles DE, et al. (2007) Contributions of pneumolysin, pneumococcal surface protein A (PspA), and PspC to pathogenicity of *Streptococcus pneumoniae* D39 in a mouse model. *Infect Immun* 75: 1843–1851.
View Article • PubMed/NCBI • Google Scholar
- Berry AM, Paton JC (2000) Additive attenuation of virulence of *Streptococcus pneumoniae* by mutation of the genes encoding pneumolysin and other putative pneumococcal virulence proteins. *Infect Immun* 68: 133–140.
View Article • PubMed/NCBI • Google Scholar

7. Quin LR, Moore QC 3rd, McDaniel LS (2007) Pneumolysin, PspA, and PspC contribute to pneumococcal evasion of early innate immune responses during bacteremia in mice. *Infect Immun* 75: 2067–2070.
View Article • PubMed/NCBI • Google Scholar
8. Ren B, Szalai AJ, Hollingshead SK, Briles DE (2004) Effects of PspA and antibodies to PspA on activation and deposition of complement on the pneumococcal surface. *Infect Immun* 72: 114–122.
View Article • PubMed/NCBI • Google Scholar
9. Tu AH, Fulgham RL, McCrory MA, Briles DE, Szalai AJ (1999) Pneumococcal surface protein A inhibits complement activation by *Streptococcus pneumoniae*. *Infect Immun* 67: 4720–4724.
View Article • PubMed/NCBI • Google Scholar
10. McDaniel LS, Ralph BA, McDaniel DO, Briles DE (1994) Localization of protection-eliciting epitopes on PspA of *Streptococcus pneumoniae* between amino acid residues 192 and 260. *Microb Pathog* 17: 323–337.
View Article • PubMed/NCBI • Google Scholar
11. Moreno AT, Oliveira ML, Ferreira DM, Ho PL, Darrieux M, et al. (2010) Immunization of mice with single PspA fragments induces antibodies capable of mediating complement deposition on different pneumococcal strains and cross-protection. *Clinical and vaccine immunology: CVI* 17: 439–446.
View Article • PubMed/NCBI • Google Scholar
12. Roche H, Hakansson A, Hollingshead SK, Briles DE (2003) Regions of PspA/EF3296 best able to elicit protection against *Streptococcus pneumoniae* in a murine infection model. *Infect Immun* 71: 1033–1041.
View Article • PubMed/NCBI • Google Scholar
13. Darrieux M, Miyaji EN, Ferreira DM, Lopes LM, Lopes AP, et al. (2007) Fusion proteins containing family 1 and family 2 PspA fragments elicit protection against *Streptococcus pneumoniae* that correlates with antibody-mediated enhancement of complement deposition. *Infect Immun* 75: 5930–5938.
View Article • PubMed/NCBI • Google Scholar
14. Briles DE, Hollingshead SK, Paton JC, Ades EW, Novak L, et al. (2003) Immunizations with pneumococcal surface protein A and pneumolysin are protective against pneumonia in a murine model of pulmonary infection with *Streptococcus pneumoniae*. *J Infect Dis* 188: 339–348.
View Article • PubMed/NCBI • Google Scholar
15. Ferreira DM, Oliveira ML, Moreno AT, Ho PL, Briles DE, et al. Protection against nasal colonization with *Streptococcus pneumoniae* by parenteral immunization with a DNA vaccine encoding PspA (Pneumococcal surface protein A). *Microb Pathog* 48: 205–213.
View Article • PubMed/NCBI • Google Scholar
16. Arulanandam BP, Lynch JM, Briles DE, Hollingshead S, Metzger DW (2001) Intranasal vaccination with pneumococcal surface protein A and interleukin-12 augments antibody-mediated opsonization and protective immunity against *Streptococcus pneumoniae* infection. *Infect Immun* 69: 6718–6724.
View Article • PubMed/NCBI • Google Scholar
17. Kono M, Hotomi M, Hollingshead SK, Briles DE, Yamanaka N (2011) Maternal immunization with pneumococcal surface protein A protects against pneumococcal infections among derived offspring. *PLoS One* 6: e27102.
View Article • PubMed/NCBI • Google Scholar
18. Hollingshead SK, Becker R, Briles DE (2000) Diversity of PspA: mosaic genes and evidence for past recombination in *Streptococcus pneumoniae*. *Infect Immun* 68: 5889–5900.
View Article • PubMed/NCBI • Google Scholar
19. Csordas FC, Perciani CT, Darrieux M, Goncalves VM, Cabrera-Crespo J, et al. (2008) Protection induced by pneumococcal surface protein A (PspA) is enhanced by conjugation to a *Streptococcus pneumoniae* capsular polysaccharide. *Vaccine* 26: 2925–2929.
View Article • PubMed/NCBI • Google Scholar
20. Goulart C, Darrieux M, Rodriguez D, Pimenta FC, Brandileone MC, et al. (2011) Selection of family 1 PspA molecules capable of inducing broad-ranging cross-reactivity by complement deposition and opsonophagocytosis by murine peritoneal cells. *Vaccine* 29: 1634–1642.
View Article • PubMed/NCBI • Google Scholar
21. Briles DE, Hollingshead SK, King J, Swift A, Braun PA, et al. (2000) Immunization of humans with recombinant pneumococcal surface protein A (rPspA) elicits antibodies that passively protect mice from fatal infection with *Streptococcus pneumoniae* bearing heterologous PspA. *The Journal of infectious diseases* 182: 1694–1701.
View Article • PubMed/NCBI • Google Scholar
22. Marriott HM, Mitchell TJ, Dockrell DH (2008) Pneumolysin: a double-edged sword during the host-pathogen interaction. *Curr Mol Med* 8: 497–509.
View Article • PubMed/NCBI • Google Scholar
23. Marriott HM, Dockrell DH (2006) *Streptococcus pneumoniae*: the role of apoptosis in host defense and pathogenesis. *Int J Biochem Cell Biol* 38: 1848–1854.
View Article • PubMed/NCBI • Google Scholar
24. Srivastava A, Henneke P, Visintin A, Morse SC, Martin V, et al. (2005) The apoptotic response to pneumolysin is Toll-like receptor 4 dependent and protects against pneumococcal disease. *Infect Immun* 73: 6479–6487.

[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

25. Malley R, Henneke P, Morse SC, Cieslewicz MJ, Lipsitch M, et al. (2003) Recognition of pneumolysin by Toll-like receptor 4 confers resistance to pneumococcal infection. *Proc Natl Acad Sci U S A* 100: 1966–1971.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
26. Witznath M, Pache F, Lorenz D, Koppe U, Gutbier B, et al. (2011) The NLRP3 inflammasome is differentially activated by pneumolysin variants and contributes to host defense in pneumococcal pneumonia. *J Immunol* 187: 434–440.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
27. Alexander JE, Lock RA, Peeters CC, Poolman JT, Andrew PW, et al. (1994) Immunization of mice with pneumolysin toxoid confers a significant degree of protection against at least nine serotypes of *Streptococcus pneumoniae*. *Infect Immun* 62: 5683–5688.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
28. Kirkham LA, Kerr AR, Douce GR, Paterson GK, Dilts DA, et al. (2006) Construction and immunological characterization of a novel nontoxic protective pneumolysin mutant for use in future pneumococcal vaccines. *Infect Immun* 74: 586–593.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
29. Ferreira DM, Areas AP, Darrieux M, Leite LC, Miyaji EN (2006) DNA vaccines based on genetically detoxified derivatives of pneumolysin fail to protect mice against challenge with *Streptococcus pneumoniae*. *FEMS Immunol Med Microbiol* 46: 291–297.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
30. Paton JC, Lock RA, Lee CJ, Li JP, Berry AM, et al. (1991) Purification and immunogenicity of genetically obtained pneumolysin toxoids and their conjugation to *Streptococcus pneumoniae* type 19F polysaccharide. *Infect Immun* 59: 2297–2304.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
31. Wu K, Zhang X, Shi J, Li N, Li D, et al. (2010) Immunization with a combination of three pneumococcal proteins confers additive and broad protection against *Streptococcus pneumoniae* infections in mice. *Infect Immun* 78: 1276–1283.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
32. Salha D, Szeto J, Myers L, Claus C, Sheung A, et al. (2012) Neutralizing antibodies elicited by a novel detoxified pneumolysin derivative, PlyD1, provide protection against both pneumococcal infection and lung injury. *Infect Immun* 80: 2212–2220.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
33. Denoel P, Philipp MT, Doyle L, Martin D, Carletti G, et al. (2011) A protein-based pneumococcal vaccine protects rhesus macaques from pneumonia after experimental infection with *Streptococcus pneumoniae*. *Vaccine* 29: 5495–5501.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
34. Berry AM, Alexander JE, Mitchell TJ, Andrew PW, Hansman D, et al. (1995) Effect of defined point mutations in the pneumolysin gene on the virulence of *Streptococcus pneumoniae*. *Infect Immun* 63: 1969–1974.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
35. Ogunniyi AD, Grabowicz M, Briles DE, Cook J, Paton JC (2007) Development of a vaccine against invasive pneumococcal disease based on combinations of virulence proteins of *Streptococcus pneumoniae*. *Infect Immun* 75: 350–357.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
36. Ogunniyi AD, Folland RL, Briles DE, Hollingshead SK, Paton JC (2000) Immunization of mice with combinations of pneumococcal virulence proteins elicits enhanced protection against challenge with *Streptococcus pneumoniae*. *Infect Immun* 68: 3028–3033.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
37. Ramos CR, Abreu PA, Nascimento AL, Ho PL (2004) A high-copy T7 *Escherichia coli* expression vector for the production of recombinant proteins with a minimal N-terminal His-tagged fusion peptide. *Braz J Med Biol Res* 37: 1103–1109.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
38. Rodrigues DC, Cavada BS, Abreu-De-Oliveira JT, Moreira RA, Russo M (1992) Differences in Macrophage stimulation and leukocyte accumulation in response to intraperitoneal administration of Glucose/Mannose-Binding Plant Lectins. *Brazilian J Med Biol Res* 25: 823–826.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
39. Xin W, Li Y, Mo H, Roland KL, Curtiss R 3rd (2009) PspA family fusion proteins delivered by attenuated *Salmonella enterica* serovar Typhimurium extend and enhance protection against *Streptococcus pneumoniae*. *Infect Immun* 77: 4518–4528.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
40. Nguyen CT, Kim SY, Kim MS, Lee SE, Rhee JH (2011) Intranasal immunization with recombinant PspA fused with a flagellin enhances cross-protective immunity against *Streptococcus pneumoniae* infection in mice. *Vaccine* 29: 5731–5739.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
41. Oliveira ML, Miyaji EN, Ferreira DM, Moreno AT, Ferreira PC, et al. (2010) Combination of pneumococcal surface protein A (PspA) with whole cell pertussis vaccine increases protection against pneumococcal challenge in mice. *PLoS One* 5: e10863.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

42. Lu YJ, Forte S, Thompson CM, Anderson PW, Malley R (2009) Protection against Pneumococcal colonization and fatal pneumonia by a trivalent conjugate of a fusion protein with the cell wall polysaccharide. *Infect Immun* 77: 2076–2083.
View Article • PubMed/NCBI • Google Scholar

43. Ljutic B, Ochs M, Messham B, Ming M, Dookie A, et al. (2012) Formulation, stability and immunogenicity of a trivalent pneumococcal protein vaccine formulated with aluminum salt adjuvants. *Vaccine* 30: 2981–2988.
View Article • PubMed/NCBI • Google Scholar

44. Ogunniyi AD, Folland RL, Briles DB, Hollingshead SK, Paton JC (2000) Immunization of mice with combinations of pneumococcal virulence proteins elicits enhanced protection against challenge with *Streptococcus pneumoniae*. *Infect Immun* 68: 3028–3033.
View Article • PubMed/NCBI • Google Scholar

45. Price KE, Camilli A (2009) Pneumolysin localizes to the cell wall of *Streptococcus pneumoniae*. *J Bacteriol* 191: 2163–2168.
View Article • PubMed/NCBI • Google Scholar

46. Price KE, Greene NG, Camilli A (2012) Export Requirements of Pneumolysin in *Streptococcus pneumoniae*. *J Bacteriol* 194: 3651–3660.
View Article • PubMed/NCBI • Google Scholar

47. Berry AM, Lock RA, Hansman D, Paton JC (1989) Contribution of autolysin to virulence of *Streptococcus pneumoniae*. *Infect Immun* 57: 2324–2330.
View Article • PubMed/NCBI • Google Scholar

48. Balachandran P, Hollingshead SK, Paton JC, Briles DE (2001) The autolytic enzyme LytA of *Streptococcus pneumoniae* is not responsible for releasing pneumolysin. *J Bacteriol* 183: 3108–3116.
View Article • PubMed/NCBI • Google Scholar

49. Malley R, Henneke P, Morse SC, Cieslewicz MJ, Lipsitch M, et al. (2003) Recognition of pneumolysin by Toll-like receptor 4 confers resistance to pneumococcal infection. *Proc Natl Acad Sci U S A* 100: 1966–1971.
View Article • PubMed/NCBI • Google Scholar