

Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Preventive Medicine

journal homepage: www.elsevier.com/locate/ypmed

Q1 Abdominal obesity as a mediator of the influence of physical activity on 2 insulin resistance in Spanish adults[☆]

Q2 Antonio García-Hermoso^a, Vicente Martínez-Vizcaíno^{b,*}, Jose I. Recio Rodriguez^c, Ana Díez-Fernández^d,
4 Manuel A. Gómez Marcos^c, Luis García Ortiz^c, for the EVIDENT Group^e

5 ^a Laboratorio de Ciencias de la Actividad Física, el Deporte y la Salud, Facultad de Ciencias Médicas, Universidad de Santiago de Chile, USACH, Chile

6 ^b Universidad de Castilla–La Mancha, Health and Social Research Center, Cuenca, Spain

7 ^c The Alamedilla Health Center, Castilla y León Health Service, SACYL, USAL, IBSAL, Salamanca, Spain

8 ^d Faculty of Occupational Therapy, Speech Therapy and Nursing, University of Castilla–La Mancha, Talavera de la Reina, Toledo, Spain

9 ^e EVIDENT Group, RedIAPP: Red de Investigación en Actividades Preventivas y Promoción de la Salud (Research Network on Preventive Activities and Health Promotion), Spain

10

1 1 A R T I C L E I N F O

12 Available online xxxx

34 Keywords:

35 Diabetes

36 Moderate-to-vigorous physical activity

37 Obesity

38 Waist circumference

39 Obesity

40 Cardiometabolic risk factors

A B S T R A C T

Objective. The aim of the study was to analyze the relationship between moderate-to-vigorous physical activity (MVPA) and insulin resistance (IR) in Spanish adults and to examine whether this relationship is mediated by abdominal obesity (waist circumference – WC).

Methods. The cross-sectional study included 1162 healthy subjects belonging to the EVIDENT study (mean age 55.0 ± 13.3 years; 61.8% women) from six different Spanish provinces. Moderate-to-vigorous physical activity (MVPA) was measured objectively over 7 days using Actigraph accelerometers, collecting data in 60-second epochs, and retaining respondents with ≥4 valid days for the analysis. The homeostasis model of assessment (HOMA-IR) was used to determine IR, and its individual components – fasting glucose and insulin – were determined using standard protocols. Linear regression models were fitted according to Baron and Kenny's procedures for mediation analysis.

Results. Fasting insulin and HOMA-IR levels were significantly worse in adults who spent fewer minutes in MVPA (first quartile ≤ 30.1 and 22.7 min/day in men and women, respectively) after adjusting for age, sex, smoking habits, drinking habits, accelerometer wear time, sedentary time, and Mediterranean diet adherence. However, when WC was added to the ANCOVA models as a covariate, the effects disappeared. Mediation analysis reported that WC acts as a full mediator in the relationship between MVPA and IR (HOMA-IR and fasting insulin).

Conclusion. These findings show that WC plays a pivotal role in the relationship between MVPA and IR, and therefore highlights that decreasing abdominal obesity might be considered as an intermediate outcome for evaluating interventions aimed at preventing diabetes mellitus.

© 2015 Published by Elsevier Inc.

41

42

43

45 Introduction

46 Physical activity is considered a protector factor against type 2 diabetes and cardiovascular risks (Jeon et al., 2007), and its benefits depend on the intensity, duration, and frequency of the activity performed (United States. Department of Health, 1996). International recommendations for adults include at least 150 min/week of moderate-to-vigorous physical activity (MVPA) or 75 min of vigorous-intensity aerobic activity for at least 20 min 3 days a week (Haskell et al., 2007). Previous studies have shown an inverse association between MVPA with insulin resistance (IR) (Ekelund et al., 2009; Nelson et al., 2013) and abdominal obesity assessed by waist circumference (WC),

(Henson et al., 2013; Scheers et al., 2013; Tremblay et al., 1990; Yang et al., 2006), which has shown to be an appropriate proxy for abdominal obesity (WHO, 2008). These are both related to cardiovascular risk (CVR) (Haffner et al., 1998; Muniyappa et al., 2008), metabolic syndrome (Zimmet et al., 2005) and type 2 diabetes (Wareham et al., 1999; Wilson et al., 2005). Therefore, increased MVPA levels both decrease cardiovascular risk factors and improve the cardiometabolic profile in the adult population.

A recent study with obese sedentary women has shown that WC combined with cardiorespiratory fitness mediate the association between MVPA and IR (Shalev-Goldman et al., 2013), while other researchers have reported that this association remains unaffected when adjusting for WC (Balkau et al., 2008; Ekelund et al., 2009). Therefore, it seems necessary to clarify whether physical activity plays an independent role as a protective factor of IR, or whether its influence is mediated by changes in WC, given the close physiological link between central obesity and IR (Carey et al., 1996). Mediation analysis

[☆] Trial registration: ClinicalTrials.gov identifier: NCT01083082.

* Corresponding author at: Universidad de Castilla–La Mancha, Edificio Melchor Cano, Centro de Estudios Socio-Sanitarios, Santa Teresa Jornet s/n, 16071 Cuenca, Spain.

E-mail address: Vicente.Martinez@uclm.es (V. Martínez-Vizcaíno).

is a statistical procedure that can be used to clarify the process underlying the relationship between two variables and the extent to which this relationship can be modified, mediated, or confounded by a third variable (Baron and Kenny, 1986). Thus, the aim of this study was twofold: i) to analyze the relationship between objectively measured MVPA and IR in Spanish adults and ii) to examine whether this relationship is mediated by abdominal obesity.

Subjects and methods

Study design

We conducted a cross-sectional analysis of the EVIDENT baseline study. Our study evaluated the association of lifestyles with the circadian pattern of blood pressure, arterial stiffness, and endothelial function in subjects with different levels of physical activity. The protocol of the EVIDENT study (NCT01083082) was published elsewhere (García-Ortiz et al., 2010).

Subjects

Subjects aged 20–80 years were selected through random sampling from the offices of general practitioners in six primary care centers from six different Spanish provinces (response rate ranged from 65% to 85%, depending on the center). The exclusion criteria have been published elsewhere (García-Ortiz et al., 2010). From the 1553 subjects included in the EVIDENT study, 391 were excluded due to lack of HOMA-IR (237) or accelerometer data (154), and therefore, 1162 patients were ultimately included in the analysis.

The Research Ethics Committee of Salamanca University Hospital (Spain) and the other centers involved approved this study. All participants gave written informed consent according to the general recommendations of the Declaration of Helsinki (World Medical Association, 2013).

Measurements

A detailed description was published elsewhere detailing how the measurements were collected (García-Ortiz et al., 2010). In short, we measured the following variables:

Anthropometric measurements

Body weight was determined on two occasions using a homologated electronic scale (Seca 770) (precision ± 0.1 kg). Height, in turn, was measured with a portable system (Seca 222). Body mass index (BMI) and WC were also measured. WC was measured as follows: the upper border of the iliac crest was located, and the tape was wrapped around above this point, parallel to the floor, ensuring that it was adjusted without compressing the skin. The reading was taken at the end of a normal breath according to the recommendations of the 2007 SEEDO Conference (Salas-Salvado et al., 2007). In order to ensure the quality and reliability of the anthropometric measurements, the nurses responsible for the measurements were trained at the start of the project. We considered WC to be normal when the value was below 94 cm in men and 80 cm in women, according to the WHO cut-off points (WHO, 2008).

Biochemical determinations

Venous blood sampling was performed between 8:00 and 9:00 AM after the individuals had fasted (avoided smoking and the consumption of alcohol and caffeinated beverages for the previous 12 h). Blood samples were collected in the respective health centers and were analyzed at the hospital of the city participating in external quality assurance programs of the Spanish Society of Clinical Chemistry and Molecular Pathology. Fasting plasma glucose was measured using standard enzymatic automated methods. The blood concentration of insulin was determined using a chemiluminescent microparticle immunoassay.

The insulin sensitivity was determined using the Homeostasis Model Assessment Insulin Resistance (HOMA-IR) index with the following formula: fasting glucose (mmol/l) \times fasting insulin (mU/ml)/22.5.

Physical activity and sedentary behavior

ActiGraph GT3X accelerometers (ActiGraph, Shalimar, FL, USA) were used to assess physical activity and sedentary behavior. This previously validated accelerometer (Matthews et al., 2013; Melanson and Freedson, 1995) measures acceleration in three individual orthogonal planes (vertical, anteroposterior, and medio-lateral) and provides activity counts as a composite vector magnitude of these three axes.

Participants were verbally instructed on how to use the accelerometer, which was worn fastened with an elastic band to the right side of the waist for 7 consecutive days. All subjects were instructed to wear the accelerometer throughout the day from the time they woke up in the morning until they went to bed at night, except for bathing and performing activities in the water. Wear time was determined by subtracting non-wear time from 24 h. Non-wear time was defined as an interval of at least 60 consecutive minutes of zero activity counts, with allowance of up to 2 min of counts between 0 and 100. The intensity of PA was categorized according to the cut-off points Troiano et al. (2008) proposed: sedentary (<100 counts/min), light (100–2019 counts/min), moderate (2020–5998 counts/min), and vigorous (>5999 counts/min). MVPA time was calculated as the mean daily minutes ≥ 2020 counts per minute from all valid days.

Mediterranean diet adherence screener (MEDAS)

Adherence to the Mediterranean diet was assessed using the validated 14-point MEDAS (Schroder et al., 2011), an adaptation of a previously validated nine-item index. The MEDAS was developed by the PREDIMED study group (Martinez-Gonzalez et al., 2004). The questionnaire allows for easily estimating Mediterranean diet adherence and may be useful in clinical settings. The 14-item version includes 12 questions about food consumption frequency and two questions about food intake habit characteristics of the Spanish Mediterranean diet. Further information about the diet assessment is available elsewhere (Patino-Alonso et al., 2014).

Smoking and alcohol consumption

Smoking and alcohol consumption history was assessed through questions on smoking status (current smokers or nonsmokers) and alcohol consumption status (current consume alcohol or not consume).

Statistical analysis

The continuous variables were expressed as the mean \pm SD and frequency distribution for categorical data. Statistical normality was tested using both graphical (normal probability plot) and statistical (Kolmogorov–Smirnov test) procedures. Because of their skewed distribution, glucose, insulin and HOMA-IR were log-transformed (natural logarithm) before being included in the models. To aid with interpretation, the data were back-transformed from the log scale for presentation in the results.

ANCOVA models were estimated to test the differences in the mean of fasting plasma glucose, insulin and HOMA-IR by categories of MVPA (low = quartile 1; medium = quartiles 2–3; high = quartile 4), controlling for age, smoking habits, drinking habits, accelerometer wear time, total sedentary time, and Mediterranean diet adherence in the first step (model 1), and including WC in the second step (model 2), by sex. Pairwise post hoc comparisons were examined using the Bonferroni test.

To examine whether the association between physical activity and IR parameters was mediated by WC, linear regression models were fitted using bootstrapped mediation procedures included in the PROCESS SPSS macro (Preacher and Hayes, 2008). The first equation regressed the mediator (WC) on the independent variable (MVPA entered as a continuous variable, min/day). The second equation regressed

the dependent variable (fasting glucose, fasting plasma insulin and HOMA-IR) on the independent variable. The third equation regressed the dependent variable on both the independent and the mediator variables.

The following criteria were used to establish mediation: (1) the independent variable must be significantly related to the mediator, (2) the independent variable must be significantly related to the dependent variable, (3) the mediator must be significantly related to the dependent variable, and (4) the association between the independent and dependent variables must be attenuated when the mediator is included in the regression model (Baron and Kenny, 1986). In addition, we tested mediation using the steps Sobel (1982) outlined. First, we estimated the attenuation or indirect influence (i.e. the influence of the independent variable on the mediator from the first regression model multiplied by the influence of the mediator on the dependent variable obtained from the third regression model). Second, we divided the indirect influence by its standard error and performed a Z test under the null hypothesis that the indirect influence is equal to zero. This analysis was adjusted for age, sex, smoking habits, drinking habits and accelerometer wear time.

Statistical analyses were performed using IBM SPSS statistics software and the level of significance was set at $\alpha = 0.05$.

Results

Data were obtained from 1162 subjects (mean age 55.0 ± 13.3 years), 718 (61.8%) of whom were women. The anthropometric, IR parameters and accelerometer descriptive data of the sample are shown in Table 1. Most subjects (925; 80.0%) were non-smokers and did not consume alcohol (741; 63.8%), and on average, the men and women were 4.5 cm (for a total of 98.5 cm) and 9.2 cm above WC cut-off point (for a total of 89.2 cm), respectively. Mean fasting plasma glucose was higher in men than in women and the accelerometry data showed that men did more MVPA than women.

Mean differences in IR parameters according to MVPA categories are shown in Table 2. Men and women in the high MVPA categories had lower fasting plasma insulin and HOMA-IR values compared to those

in the low categories after controlling for age, smoking habits, drinking habits, accelerometer wear time, and sedentary time. However, when we included additional adjustment for WC (model 2), the differences disappeared.

In both sexes, WC acted as a full mediator of the relationship between MVPA and fasting plasma insulin (Fig. 1A). Similar results were obtained when HOMA-IR was the dependent variable in the mediation model (Fig. 1B). The estimated percentage of total influence mediated by WC was 12.3% in men ($z = -2.03, p = 0.042$) and 20.6% in women ($z = -3.07, p = 0.002$) for fasting plasma insulin, and 14.1% ($z = -2.06; p = 0.039$) in men and 23.2% ($z = -3.13, p = 0.002$) in women for HOMA-IR. Finally, the criteria necessary for mediation analysis were not achieved for fasting glucose levels (data not reported).

Discussion

Although a growing number of studies have examined the association between physical activity and IR, none of them have evaluated the mediation role of excess weight in this relationship in healthy adults. Our results show that MVPA was negatively associated with IR parameters, even after adjusting for sociodemographic and lifestyle potential confounders, but this association disappeared when additional adjustments for WC were included. Moreover, the mediation analysis disclosed that the influence of MVPA on IR was mediated by abdominal obesity, as measured by WC.

There is consistent evidence regarding the beneficial influence of MVPA on insulin sensitivity and glycemic control in adults, independent of weight change (Ekelund et al., 2007; Ivy, 1997), as well as on abdominal obesity, as measured by WC (Staiano et al., 2012; Stamatakis et al., 2009). Previous studies have shown that subjects with higher values of WC presented worse IR profiles, and those who were physically active presented lower IR values in the same WC category compared to their sedentary peers (Ekelund et al., 2007; Healy et al., 2006; Nelson et al., 2013; Weinstein et al., 2004).

Some other studies have described that there is a relationship between PA and IR, independent of WC (Balkau et al., 2008; Ekelund et al., 2009), while others have reported that the amount of physical

Table 1
Demographic and clinical characteristics of patients, by sex.

	Total (n = 1162)	Men ^a (n = 444)	Women ^a (n = 718)	p
Age (years)	55.0 (13.3)	57.3 (14.5)	53.5 (16.0)	<0.001
Smoking status, n (%)				
Yes	237 (20.0)	83 (18.6)	154 (21.4)	0.431
No or past	925 (80.0)	361 (81.4)	564 (78.6)	
Alcohol status, n (%)				
Yes	421 (36.2)	116 (26.1)	305 (42.5)	<0.001
No or past	741 (63.8)	328 (73.9)	413 (57.5)	
Weight (kg)	72.8 (17.0)	81.8 (12.7)	67.2 (13.4)	<0.001
Height (cm)	163.2 (10.2)	170.8 (8.4)	158.6 (8.1)	<0.001
Body mass index (kg/m ²)	27.3 (3.4)	28.0 (4.2)	26.8 (5.4)	<0.001
Waist circumference (cm)	92.8 (13.6)	98.5 (10.5)	89.2 (13.4)	<0.001
Obesity ^b n (%)				
Yes	863 (74.3)	291 (65.5)	572 (79.7)	<0.001
No	299 (25.7)	153 (34.5)	146 (20.3)	
Fasting plasma glucose (mmol/L)	5.08 (1.4)	5.26 (1.3)	4.96 (1.1)	<0.001
Fasting plasma insulin (µmol/L)	7.48 (6.5)	7.78 (5.9)	7.30 (6.7)	0.228
HOMA-IR	1.75 (1.7)	1.88 (1.7)	1.68 (1.9)	0.068
Accelerometer wear time (min/day)	931.2 (218.0)	941.6 (231.0)	924.8 (210.6)	0.239
MVPA (min/day)	46.5 (28.6)	53.8 (31.3)	42.0 (25.9)	<0.001
Meet recommendations for MVPA, ^c n (%)	527 (45.3)	233 (52.5)	294 (41.0)	<0.001
VPA (min/day)	1.22 (5.4)	2.0 (7.8)	0.7 (3.0)	<0.001
Sedentary time (min/day)	580.3 (175.6)	601.7 (186.6)	567.2 (167.4)	0.003
Healthy diet score (0–14)	7.4 (1.8)	7.4 (1.9)	7.5 (1.7)	0.386
Mediterranean diet adherence, ^b n (%)	337 (29.0)	120 (27.1)	217 (30.2)	0.304

MVPA, moderate-vigorous physical activity; VPA, vigorous physical activity.

^a Values are means \pm SD, except for categorical data (n (%)).

^b Cut-off for obesity were waist circumference greater than 94 cm for men and 80 cm for women (WHO, 2008).

^c 150 min/week of MVPA (Haskell et al., 2007).

Table 2
ANCOVA models testing differences in fasting plasma glucose, insulin, and HOMA-IR by MVPA categories, adjusting for potential confounders, by sex.

		Fasting plasma glucose ^a (mmol/L)		Fasting plasma insulin ^a (μmol/L)		HOMA-IR ^a	
		Men	Women	Men	Women	Men	Women
	Low (L)	5.44 ± 1.18	5.20 ± 1.44	9.62 ± 6.30	8.97 ± 6.86	2.46 ± 2.14	2.21 ± 2.24
	Medium (M)	5.41 ± 1.48	4.92 ± 0.79	8.41 ± 4.09	7.16 ± 5.14	2.06 ± 1.09	1.58 ± 1.13
	High (H)	5.15 ± 0.99	4.87 ± 0.76	7.24 ± 4.47	6.64 ± 5.13	1.69 ± 1.13	1.47 ± 1.18
<i>Model 1</i>	<i>p</i>	0.139	0.223	0.030	0.001	0.004	0.002
	Post-hoc tests ^b	ns	ns	L > H	L > H	L > H	L > H
<i>Model 2</i>	<i>p</i>	0.134	0.719	0.213	0.114	0.201	0.191
	Post-hoc tests ^b	ns	ns	ns	ns	ns	ns

Values are means ± SD.
Abbreviations: HOMA-IR, homeostatic model assessment of insulin resistance; MVPA, moderate-vigorous physical activity; ns, non-significant.
Model 1: adjusted for age, smoking habit, drinking habit, accelerometer wear time, sedentary time, and Mediterranean diet adherence. Model 2: adjusted for the same covariates as Model 1 and waist circumference.
^a Log transformed for comparing means.
^b Bonferroni pairwise comparisons.

activity was not associated with IR after controlling for WC (Shalev-Goldman et al., 2013). Our data show that participants in the low MVPA category had higher fasting plasma insulin and HOMA-IR levels than did those in the higher category, but after adjusting for WC, the relationship between MVPA and IR disappeared. Likewise, Healy et al. (2008) found that the inverse association of MVPA with cardiometabolic risk factors did not remain significant when WC was included as a confounder, except for triglyceride levels.

Discrepancies between studies can be explained by differences in the populations studied (e.g. health risk status, weight status, ethnicity and age), in the methods for physical activity assessment (e.g. calibrated heart rate monitoring or accelerometers), and in the control of confounders (e.g. sedentary time, light intensity physical activity). The main difference between our results and those presented by Shalev-Goldman et al. (2013) is that we calculated the magnitude of WC as a single mediator, finding that it affects individuals in the causal pathway between MVPA and IR.

This evidence allows us to conclude that active people are more likely to present a healthy IR and cardiometabolic profile, and thus being physically active may help to prevent metabolic syndrome and type 2 diabetes. It also confirms the potential mediation role of abdominal obesity, as other authors have suggested (Alberti et al., 2005). Therefore, our cross-sectional study suggests that the influence of

physical activity on improving IR is substantially greater when accompanied by a decrease in WC.

Furthermore, our non-significant findings for fasting plasma glucose are in agreement with the results from most previous studies (Henson et al., 2013; Scheers et al., 2013; Shalev-Goldman et al., 2013), and emphasizing that fasting plasma glucose might not be sensitive to changes in physical activity, and two-hour plasma glucose appears to be a more appropriate biomarker for use as an outcome measure in interventions that are based on physical activity (Healy et al., 2008).

Mediation analysis

Our data, by using mediation analysis, reveal that abdominal obesity has a powerful influence on the relationship between MVPA and both IR and fasting plasma insulin in adults. Visceral or central obesity increases the flux of free fatty acids to the liver, and an important role in the genesis of IR of these free fatty acids has been described (Petersen et al., 2004). Inflammation has also been reported as an alternative candidate for the common link between central obesity and IR (Ghanim et al., 2004). Finally, increased abdominal obesity has been associated not only with impaired insulin sensitivity but also with reduced glycogen synthesis (non-oxidative glucose disposal) and a reduced responsiveness of muscle




Fig. 1. WC mediation model of the relationship between MVPA (min/day), fasting insulin (1A) and HOMA-IR (1B) controlling for potential confounders (age, smoking habits, drinking habits, accelerometer wear time, sedentary time, and Mediterranean diet adherence), by sex. Data in roman type refer to men. Data in italics refer to women. **p* ≤ 0.05; ***p* ≤ 0.001.

glycogen synthase (Carey et al., 1996), all of which are related to central obesity and an increased risk of IR.

These physiological pathways support the hypothesis that programs aimed at increasing physical activity could diminish IR through reductions of abdominal obesity, thus you can expect a weak impact on the insulin metabolism of those physical activity programs that, while improving other cardiometabolic factors, fail to diminish adiposity. Besides, the relationships analyzed here are probably related to more than a single mediator variable (e.g. BMI, accumulated daily activity, sedentary time); future studies using structural equations or complex mediator model procedures might be useful in clarifying more specifically the potential mediator or confounder role of each factor.

Finally, it might be also interesting to address how these results may affect those subjects defined as metabolically healthy obese (MHO) vs. metabolically unhealthy obese (MUHO) or non-obese. Previous studies have found that MHO subjects had intermediate levels of WC (Després, 2012), a more favorable fat distribution (lower visceral fat and greater thigh subcutaneous fat), normal levels of C-reactive protein (Phillips, 2013) and higher PA levels (Wildman et al., 2008) compared to MUHO and non-obese subjects. A recent study in a Spanish population (Martínez-Larrad et al., 2014) reported that the prevalence of MHO subjects was 4.78% according to Wildman modified criteria (Ascaso et al., 2001). Thus, because of this relatively low prevalence, our coefficients' estimations will probably remain very similar if MHO individuals are removed from the analyses, but our mediation results should be applied with caution in these subjects, since indirect coefficients through WC might be substantially different in this MHO phenotype.

This study has several limitations. First, the cross-sectional design prevents us from establishing a causal relationship. For example, we cannot establish that abdominal obesity predicts greater MVPA, since it seems reasonable to think that many adults begin physical activity due to an awareness of excess weight. Second, we did not use an intravenous glucose tolerance test or hyperinsulinemic–euglycemic clamp technique, both considered better measures of IR (DeFronzo et al., 1979), but instead, we used HOMA-IR, which is probably the most appropriate method for measuring IR in large epidemiological studies. Third, PA was assessed using an accelerometer during the same week, and the fact that the subjects were evaluated with the device might have influenced their activity levels that week. However, without any activity goals communicated, the possible effect seems to be negligible. Fourth, we studied subjects that were randomly selected from outpatient clinics in different regions of Spain; however, the subjects may not be representative of the general Spanish population, but rather, of people that attend primary care centers. We must also take into consideration the limited generalizability of the study because the participants were people who met the rigorous inclusion criteria of the parent trial. The generalizability is also limited because of the impossibility of adjusting for all potential confounders, and this causes the residual confounding to tease out independent associations. Finally, because of the limited amount of time spent in vigorous physical activity in our sample (8 out of 10 participants accumulated no vigorous physical activity), statistics that are similar to those reported by other authors (Ekelund et al., 2009), it was not feasible to examine the relative importance of vigorous physical activity on IR. However, we found a relationship between insulin, IR and vigorous physical activity in men, independent of WC, suggesting that obesity mechanisms may not be so relevant (Janssen and Ross, 2012) in this population.

Conclusion

Our findings are important from a clinical and public health point-of-view because they disclose that abdominal obesity plays a critical role in the relationship between MVPA and IR, and they alert us that physical activity only produces benefits in IR if the activity is not accompanied by compensatory behaviors that might mitigate its influence on WC (e.g. less sedentary time or nutritional changes).

References

- Alberti, K.G.M., Zimmet, P., Shaw, J., 2005. The metabolic syndrome—a new worldwide definition. *Lancet* 366, 1059–1062.
- Ascaso, J.F., Romero, P., Real, J.T., Priego, A., Valdecabres, C., Carmena, R., 2001. Insulin resistance quantification by fasting insulin plasma values and HOMA index in a non-diabetic population. *Med. Clin.* 117, 530–533.
- Balkau, B., Mhamdi, L., Opper, J.-M., Nolan, J., Golay, A., Porcellati, F., Laakso, M., Ferrannini, E., 2008. Physical activity and insulin sensitivity the RISC study. *Diabetes* 57, 2613–2618.
- Baron, R.M., Kenny, D.A., 1986. The moderator–mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J. Pers. Soc. Psychol.* 51, 1173.
- Carey, D.G., Jenkins, A.B., Campbell, L.V., Freund, J., Chisholm, D.J., 1996. Abdominal fat and insulin resistance in normal and overweight women: direct measurements reveal a strong relationship in subjects at both low and high risk of NIDDM. *Diabetes* 45, 633–638.
- DeFronzo, R.A., Tobin, J.D., Andres, R., 1979. Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am. J. Physiol. Gastrointest. Liver Physiol.* 237, G214–G223.
- Després, J.-P., 2012. What is “metabolically healthy obesity”? from epidemiology to pathophysiological insights. *J. Clin. Endocrinol. Metab.* 97, 2283–2285.
- Ekelund, U., Franks, P.W., Sharp, S., Wareham, N.J., 2007. Increase in physical activity energy expenditure is associated with reduced metabolic risk independent of change in fitness and fitness. *Diabetes Care* 30, 2101–2106.
- Ekelund, U., Brage, S., Griffin, S.J., Wareham, N.J., 2009. Objectively measured moderate- and vigorous-intensity physical activity but not sedentary time predicts insulin resistance in high-risk individuals. *Diabetes Care* 32, 1081–1086.
- García-Ortiz, L., Recio-Rodríguez, J.L., Martín-Cantera, C., Cabrejas-Sánchez, A., Gómez-Arranz, A., González-Viejo, N., Nicolás, E.I., Patino-Alonso, M.C., Gómez-Marcos, M.A., 2010. Physical exercise, fitness and dietary pattern and their relationship with circadian blood pressure pattern, augmentation index and endothelial dysfunction biological markers: EVIDENT study protocol. *BMC Public Health* 10, 233.
- Ghanim, H., Aljada, A., Hofmeyer, D., Syed, T., Mohanty, P., Dandona, P., 2004. Circulating mononuclear cells in the obese are in a proinflammatory state. *Circulation* 110, 1564–1571.
- Haffner, S.M., Lehto, S., Rönnemaa, T., Pyörälä, K., Laakso, M., 1998. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N. Engl. J. Med.* 339, 229–234.
- Haskell, W.L., Lee, I.M., Pate, R.R., Powell, K.E., Blair, S.N., Franklin, B.A., Macera, C.A., Heath, G.W., Thompson, P.D., et al., 2007. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med. Sci. Sports Exerc.* 39, 1423–1434.
- Healy, G.N., Dunstan, D.W., Shaw, J.E., Zimmet, P.Z., Owen, N., 2006. Beneficial associations of physical activity with 2-h but not fasting blood glucose in Australian adults: the AusDiab study. *Diabetes Care* 29, 2598–2604.
- Healy, G.N., Wijndaele, K., Dunstan, D.W., Shaw, J.E., Salmon, J., Zimmet, P.Z., Owen, N., 2008. Objectively measured sedentary time, physical activity, and metabolic risk the Australian Diabetes, Obesity and Lifestyle study (AusDiab). *Diabetes Care* 31, 369–371.
- Henson, J., Yates, T., Biddle, S., Edwardson, C., Khunti, K., Wilmot, E., Gray, L., Gorely, T., Nimmo, M., et al., 2013. Associations of objectively measured sedentary behaviour and physical activity with markers of cardiometabolic health. *Diabetologia* 1–9.
- Ivy, J.L., 1997. Role of exercise training in the prevention and treatment of insulin resistance and non-insulin-dependent diabetes mellitus. *Sports Med.* 24, 321–336.
- Janssen, I., Ross, R., 2012. Vigorous intensity physical activity is related to the metabolic syndrome independent of the physical activity dose. *Int. J. Epidemiol.* 41, 1132–1140.
- Jeon, C.Y., Lokken, R.P., Hu, F.B., Van Dam, R.M., 2007. Physical activity of moderate intensity and risk of type 2 diabetes: a systematic review. *Diabetes Care* 30, 744–752.
- Martínez-González, M.A., Fernández-Jarne, E., Serrano-Martínez, M., Wright, M., Gómez-Gracia, E., 2004. Development of a short dietary intake questionnaire for the quantitative estimation of adherence to a cardioprotective Mediterranean diet. *Eur. J. Clin. Nutr.* 58, 1550–1552.
- Martínez-Larrad, M.T., Anchuelo, A.C., Del Prado, N., Rueda, J.M.I., Gabriel, R., Serrano-Ríos, M., 2014. Profile of individuals who are metabolically healthy obese using different definition criteria. A population-based analysis in the Spanish population. *PLoS One* 9, e106641.
- Matthews, C.E., Keadle, S.K., Sampson, J., Lyden, K., Bowles, H.R., Moore, S.C., Libertine, A., Freedson, P.S., Fowke, J.H., 2013. Validation of a previous-day recall measure of active and sedentary behaviors. *Med. Sci. Sports Exerc.* 45 (8), 1629–1638.
- Melanson, E., Freedson, P.S., 1995. Validity of the Computer Science and Applications, Inc. (CSA) activity monitor. *Med. Sci. Sports Exerc.* 27, 934–940.
- Muniyappa, R., Lee, S., Chen, H., Quon, M.J., 2008. Current approaches for assessing insulin sensitivity and resistance in vivo: advantages, limitations, and appropriate usage. *Am. J. Physiol. Endocrinol. Metab.* 294, E15–E26.
- Nelson, R.K., Horowitz, J.F., Holleman, R.G., Swartz, A.M., Strath, S.J., Kriska, A.M., Richardson, C.R., 2013. Daily physical activity predicts degree of insulin resistance: a cross-sectional observational study using the 2003–2004 National Health and Nutrition Examination Survey. *Int. J. Behav. Nutr. Phys. Act.* 10, 10.1186.
- Patino-Alonso, M.C., Recio-Rodríguez, J.L., Belio, J.F.M., Colominas-Garrido, R., Lema-Bartolomé, J., Arranz, A.G., Agudo-Conde, C., Gómez-Marcos, M.A., García-Ortiz, L., 2014. Factors associated with adherence to the Mediterranean diet in the adult population. *J. Acad. Nutr. Diet.* 114, 583–589.
- Petersen, K.F., Dufour, S., Befroy, D., Garcia, R., Shulman, G.I., 2004. Impaired mitochondrial activity in the insulin-resistant offspring of patients with type 2 diabetes. *N. Engl. J. Med.* 350, 664–671.

- 454 Phillips, C.M., 2013. Metabolically healthy obesity: definitions, determinants and clinical 483
455 implications. *Rev. Endocr. Metab. Disord.* 14, 219–227. 484
- 456 Preacher, K.J., Hayes, A.F., 2008. Asymptotic and resampling strategies for assessing and 485
457 comparing indirect effects in multiple mediator models. *Behav. Res. Methods* 40, 486
458 879–891. 487
- 459 Salas-Salvado, J., Rubio, M.A., Barbany, M., Moreno, B., Grupo, C.d.I.S., 2007. SEEDO 2007 488
460 Consensus for the evaluation of overweight and obesity and the establishment of 489
461 therapeutic intervention criteria. *Med. Clín.* 128, 184. 490
- 462 Scheers, T., Philippaerts, R., Lefevre, J., 2013. SenseWear-determined physical activity and 491
463 sedentary behavior and metabolic syndrome. *Med. Sci. Sports Exerc.* 45, 481–489. 492
- 464 Schroder, H., Fito, M., Estruch, R., Martinez-Gonzalez, M.A., Corella, D., Salas-Salvado, J., 493
465 Lamuela-Raventos, R., Ros, E., Salaverria, I., Fiol, M., Lapetra, J., Vinyoles, E., Gómez- 494
466 Gracia, E., Lahoz, C., Serra-Majem, L., Pintó, X., Ruiz-Gutierrez, V., Covas, M.I., 2011. 495
467 A short screener is valid for assessing Mediterranean diet adherence among older 496
468 Spanish men and women. *J. Nutr.* 141, 1140–1145. 497
- 469 Shalev-Goldman, E., McGuire, K.A., Ross, R., 2013. Waist circumference and cardiorespiratory 498
470 fitness are independently associated with glucose tolerance and insulin 499
471 resistance in obese women. *Appl. Physiol. Nutr. Metab.* 39, 358–362. 500
- 472 Sobel, M.E., 1982. Asymptotic confidence intervals for indirect effects in structural 501
473 equation models. *Sociol. Methodol.* 13, 290–312. 502
- 474 Staiano, A.E., Reeder, B.A., Elliott, S., Joffres, M.R., Pahwa, P., Kirkland, S.A., Paradis, G., 503
475 Katzmarzyk, P.T., 2012. Physical activity level, waist circumference, and mortality. 504
476 *Appl. Physiol. Nutr. Metab.* 37, 1008–1013. 505
- 477 Stamatakis, E., Hirani, V., Rennie, K., 2009. Moderate-to-vigorous physical activity and 506
478 sedentary behaviours in relation to body mass index-defined and waist 507
479 circumference-defined obesity. *BJN* 101, 765–773. 508
- 480 Tremblay, A., Després, J.-P., Leblanc, C., Craig, C.L., Ferris, B., Stephens, T., Bouchard, C., 509
481 1990. Effect of intensity of physical activity on body fatness and fat distribution. 510
482 *Am. J. Clin. Nutr.* 51, 153–157. 511
- Troiano, R.P., Berrigan, D., Dodd, K.W., Mâsse, L.C., Tilert, T., McDowell, M., 2008. Physical 483
484 activity in the United States measured by accelerometer. *Med. Sci. Sports Exerc.* 40, 485
486 181–188. 487
- United States, Department of Health, 1996. *Physical Activity and Health: A Report of the* 488
489 *Surgeon General.* DIANE Publishing. 490
- Wareham, N.J., Byrne, C.D., Williams, R., Day, N.E., Hales, C.N., 1999. Fasting proinsulin 491
492 concentrations predict the development of type 2 diabetes. *Diabetes Care* 22, 493
494 262–270. 495
- Weinstein, A.R., Sesso, H.D., Lee, I.M., Cook, N.R., Manson, J.E., Buring, J.E., Gaziano, J.M., 496
497 2004. Relationship of physical activity vs body mass index with type 2 diabetes in 498
499 women. *JAMA* 292, 1188–1194. 500
- Wildman, R.P., Muntner, P., Reynolds, K., McGinn, A.P., Rajpathak, S., Wylie-Rosett, J., 501
502 Sowers, M.R., 2008. The obese without cardiometabolic risk factor clustering and 503
504 the normal weight with cardiometabolic risk factor clustering: prevalence and corre- 505
506 lates of 2 phenotypes among the US population (NHANES 1999–2004). *Arch. Intern. 507
508 Med.* 168, 1617–1624. 509
- Wilson, P.W., D'Agostino, R.B., Parise, H., Sullivan, L., Meigs, J.B., 2005. Metabolic syndrome 510
511 as a precursor of cardiovascular disease and type 2 diabetes mellitus. *Circulation* 112, 512
513 3066–3072. 514
- World Health Organization, 2008. *Waist Circumference and Waist–Hip Ratio Report of a* 515
516 *WHO Expert Consultation (Geneva, 8–11 December 2008).* 517
- World Medical Association, 2013. *World Medical Association Declaration of Helsinki: eth-* 518
519 *ical principles for medical research involving human subjects.* *JAMA* 310, 2191–2194. 520
- Yang, X., Telama, R., Viikari, J., Raitakari, O.T., 2006. Risk of obesity in relation to physical 521
522 activity tracking from youth to adulthood. *Med. Sci. Sports Exerc.* 38, 919–925. 523
- Zimmet, P., Magliano, D., Matsuzawa, Y., Alberti, G., Shaw, J., 2005. The metabolic 524
525 syndrome: a global public health problem and a new definition. *J. Atheroscler. 526
527 Thromb.* 12 (6), 295–300. 528