

FACTA UNIVERSITATIS (NIŠ)  
 SER. MATH. INFORM. Vol. 35, No 3 (2020), 647-672  
<https://doi.org/10.22190/FUMI2003647M>

## EXISTENCE AND BLOW UP FOR A NONLINEAR VISCOELASTIC HYPERBOLIC PROBLEM WITH VARIABLE EXPONENTS \*

Fatima Z. Mahdi and Ali Hakem

© by University of Niš, Serbia | Creative Commons Licence: CC BY-NC-ND  
**Abstract.** Our aim in this paper is to establish the weak existence theorem and find under suitable assumptions sufficient conditions on  $m, p$  and the initial data for which the blow up takes place for the following boundary value problem:

$$|u_t|^p u_{tt} - \Delta u - \Delta u_{tt} + \int_0^t g(t-s) \Delta u(s) ds + |u_t|^{m(x)-2} u_t = |u|^{p(x)-2} u.$$

This paper extends some of the results obtained by the authors and it is focused on new results which are consequence of the presence of variable exponents.

**Keywords:** Variable exponents; weak solutions; blow up.

### 1. Introduction

Let  $\Omega \subset \mathbb{R}^n (n \geq 2)$  be a bounded Lipschitz domain and  $0 < T < \infty$ . We consider the following initial boundary value problem:

$$(1.1) \quad \begin{cases} |u_t|^p u_{tt} - \Delta u - \Delta u_{tt} + \int_0^t g(t-s) \Delta u(s) ds \\ + |u_t|^{m(x)-2} u_t = |u|^{p(x)-2} u, & (x, t) \in Q_T, \\ u(x, t) = 0, & (x, t) \in S_T, \\ u(x, 0) = u_0(x), u_t(x, 0) = u_1(x), & x \in \Omega, \end{cases}$$

where  $Q_T = \Omega \times (0, T]$  and  $S_T$  denote the lateral boundary of the cylinder  $Q_T$ . It is assumed throughout the paper that the exponents  $m(x)$  and  $p(x)$  are continuous in  $\Omega$  with logarithmic module of continuity:

Received May 27, 2019; accepted May 6, 2020.

2010 *Mathematics Subject Classification.* Primary 35B40; Secondary: 35L70, 35L45

\*The authors were supported in part by PRFU/C00L03UN220120200002/ALGERIA

$$(1.2) \quad 1 < m^- = \operatorname{ess\,inf}_{x \in \Omega} m(x) \leq m(x) \leq m^+ = \operatorname{ess\,sup}_{x \in \Omega} m(x) < \infty,$$

$$(1.3) \quad 1 < p^- = \operatorname{ess\,inf}_{x \in \Omega} p(x) \leq p(x) \leq p^+ = \operatorname{ess\,sup}_{x \in \Omega} p(x) < \infty,$$

$$(1.4) \quad \forall z, \xi \in \Omega, |z - \xi| < 1, |m(x) - m(\xi)| + |p(z) - p(\xi)| \leq \omega(|z - \xi|),$$

where

$$(1.5) \quad \lim_{\tau \rightarrow 0^+} \sup \omega(\tau) \ln \frac{1}{\tau} = C < +\infty.$$

**Remark 1.1.** We use the standard Lebesgue space  $L^p(\Omega)$  and the Sobolev space  $H_0^1(\Omega)$  with their usual scalar product and norms. We will use the embedding  $H_0^1(\Omega) \hookrightarrow L^s(\Omega)$  for  $2 \leq s \leq 2n/(n-2)$  if  $n \geq 3$  or  $s \geq 2$  if  $n = 1, 2$ . The generic embedding constant, denoted by  $C_*$  is given by

$$(1.6) \quad \|u\|_s \leq C_* \|\nabla u\|_2.$$

And we also assume that

$(H_1)$  :  $\rho$  is a constant that satisfies

$$0 < \rho \leq \frac{2}{n-2} \text{ if } n \geq 3 \quad \text{and} \quad 0 < \rho \quad \text{if } n = 1, 2.$$

$(H_2)$  :  $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  is bounded  $\mathcal{C}^1$  function satisfying

$$g(0) > 0, \quad 1 - \int_0^\infty g(s) ds = l > 0.$$

$(H_3)$  : There exists  $\xi > 0$  such that

$$g'(t) < -\xi(t)g(t), \quad t \geq 0.$$

If  $m, p$  are constants, there have been many results about the existence and blow-up properties of the solutions, we refer the readers to the bibliography given in [5]-[25]. In recent years, a great attention has been focused on the study of mathematical models of electro-rheological fluids. These models include hyperbolic, parabolic or elliptic equations which are nonlinear with respect to gradient of the thought solution and with variable exponents of nonlinearity see ([3]-[12]-[15]-[23]-[24]) and the references therein. It should be mentioned that questions of existence, uniqueness and regularity of weak solutions for parabolic and elliptic equations have been studied by many authors under various conditions on the data and by different methods- (see [[1],[2]] and the further references therein).

To the best of our knowledge, there are only a few works about viscoelastic hyperbolic equations with variable exponents of nonlinearity. In [4] the authors investigated the finite time blow-up of solutions for viscoelastic hyperbolic equations, and in [5] the authors discussed only the viscoelastic hyperbolic problem with constant exponents. Motivated by the works of [[5],[4]], we shall study the existence and energy decay of the solutions to problem (1.1) and state some properties to the

solutions.

The present paper is organized as follows. In Section 2, we introduce the function spaces of Orlicz-Sobolev type and a brief description of their main properties, give the definition of the weak solution to the problem and prove the existence of weak solutions for problem (1.1) with Galerkin’s method. In the last sections, we finally prove the desired results.

### 2. Existence of weak solutions

In this section, the existence of weak solutions is studied. Firstly, we introduce some Banach spaces

$$L^{p(x)}(\Omega) = \left\{ u(x) : u \text{ is measurable in } \Omega, A_{p(\cdot)}(u) = \int_{\Omega} |u(x)|^{p(x)} dx < \infty \right\},$$

with the following Luxembourg-type norm

$$\|u\|_{p(\cdot)} = \inf \{ \lambda > 0, A_{p(\cdot)}(u/\lambda) \leq 1 \}.$$

We, next, define the variable-exponent Lebesgue Sobolev space  $W^{1,p(\cdot)}(\Omega)$  as follows:

$$W^{1,p(\cdot)}(\Omega) = \left\{ u \in L^{p(\cdot)}(\Omega) \text{ such that } \nabla u \text{ exists and } |\nabla u| \in L^{p(\cdot)}(\Omega) \right\}.$$

This space is a Banach space with respect to the norm  $\|u\|_{W^{1,p(\cdot)}(\Omega)} = \|u\|_{p(\cdot)} + \|\nabla u\|_{p(\cdot)}$ . Furthermore, we set  $W_0^{1,p(\cdot)}(\Omega)$  to be the closure of  $C_0^\infty(\Omega)$  in  $W^{1,p(\cdot)}(\Omega)$ . Here we note that the space  $W_0^{1,p(\cdot)}(\Omega)$  is usually defined in a different way for the variable exponent case. However, both definitions are equivalent (see [10]). The dual of  $W_0^{1,p(\cdot)}(\Omega)$  is defined as  $W^{-1,p'(\cdot)}(\Omega)$ ; in the same way as the classical Sobolev spaces, where  $\frac{1}{p(\cdot)} + \frac{1}{p'(\cdot)} = 1$ .

**Lemma 2.1.** ([3]) For  $u \in L^{p(x)}(\Omega)$ , the following relations hold:

1.  $\|u\|_{p(\cdot)} < 1 (= 1; > 1) \Leftrightarrow A_{p(\cdot)}(u) < 1 (= 1; > 1)$ ;
2.  $\|u\|_{p(\cdot)} < 1 \Rightarrow \|u\|_{p(\cdot)}^{p^+} \leq A_{p(\cdot)}(u) \leq \|u\|_{p(\cdot)}^{p^-}$ ;  
 $\|u\|_{p(\cdot)} > 1 \Rightarrow \|u\|_{p(\cdot)}^{p^+} \geq A_{p(\cdot)}(u) \geq \|u\|_{p(\cdot)}^{p^-}$ ;
3.  $\|u\|_{p(\cdot)} \rightarrow 0 \Leftrightarrow A_{p(\cdot)}(u) \rightarrow 0; \|u\|_{p(\cdot)} \rightarrow \infty \Leftrightarrow A_{p(\cdot)}(u) \rightarrow \infty$ .

**Lemma 2.2.** ([26]) For  $u \in W_0^{1,p(\cdot)}(\Omega)$ , if  $p$  satisfies condition (1.2), the  $p(\cdot)$ -Poincaré’s inequality

$$\|u\|_{p(x)} \leq C \|\nabla u\|_{p(x)},$$

holds, where the positive constant  $C$  depends on  $p$  and  $\Omega$ .

**Remark 2.1.** Note that the following inequality

$$\int_{\Omega} |u|^{p(x)} dx \leq C \int_{\Omega} |\nabla u|^{p(x)} dx,$$

does not in general hold.

**Lemma 2.3.** ([10]). Let  $\Omega$  be an open domain (that may be unbounded) in  $\mathbb{R}^n$  with cone property. If  $p(x) : \bar{\Omega} \rightarrow \mathbb{R}$  is Lipschitz continuous function satisfying  $1 < p^- \leq p^+ < \frac{n}{k}$  and  $r(x) : \bar{\Omega} \rightarrow \mathbb{R}$  is measurable and satisfies

$$p(x) \leq r(x) \leq p^*(x) = \frac{np(x)}{n - kp(x)} \text{ a.e } x \in \bar{\Omega},$$

then there is a continuous embedding  $W^{k,p(x)}(\Omega) \hookrightarrow L^{r(x)}(\Omega)$ .

The main theorem in this section is the following:

**Theorem 2.1.** Let  $u_0, u_1 \in H_0^1(\Omega)$  be given. Assume that the exponents  $m(x)$  and  $p(x)$  satisfy conditions (1.2)-(1.4). Then the problem (1.1) has at least one weak solution  $u : \Omega \times (0, \infty) \rightarrow \mathbb{R}$  in the class

$$u \in L^\infty(0, \infty; H_0^1(\Omega)), \quad u' \in L^\infty(0, \infty; H_0^1(\Omega)), \quad u'' \in L^\infty(0, \infty; H_0^1(\Omega)).$$

And one of the following conditions holds:

$$(A_1) \quad 2 < p^- < p^+ < \max \left\{ n, \frac{np^-}{n - p^-} \right\}, \quad 2 < m^- < m^+ < p^-;$$

$$(A_2) \quad \max \left\{ 1, \frac{2n}{n+2} \right\} < p^- < p^+ < 2, \quad 1 < m^- < m^+ < \frac{3p^- - 2}{p^-}.$$

*Proof.* Let us take for  $\{w_j\}_{j=1}^\infty$  the orthogonal basis of  $H_0^1(\Omega)$  such that

$$-\Delta w_j = \lambda_j w_j, \quad x \in \Omega, \quad w_j = 0, \quad x \in \partial\Omega.$$

We denote by  $V_k = \text{span} \{w_1, \dots, w_k\}$  the subspace generated by the first  $k$  vectors of the basis  $\{w_j\}_{j=1}^\infty$ . By normalization, we have  $\|w_j\|_2 = 1$ . Let us define the operator:

$$\begin{aligned} \langle Lu, \phi \rangle &= \int_{\Omega} [|u_t|^p u_{tt} \phi + \nabla u \nabla \phi + \nabla u_{tt} \nabla \phi - \int_0^t g(t-s) \nabla u \nabla \phi ds \\ &\quad + |u_t|^{m(x)-2} u_t \phi - \alpha |u|^{p(x)-2} u \phi] dx, \quad \phi \in V_k. \end{aligned}$$

For any given integer  $k$ , we consider the approximate solution  $u_k = \sum_{i=1}^k c_i^k(t)w_i$ ,

which satisfies

$$(2.1) \quad \begin{cases} \langle Lu_k, w_i \rangle = 0 & i = 1, 2, \dots, k, \\ u_k(0) = u_{0k}, & u_{kt}(0) = u_{1k}, \end{cases}$$

where

$$u_{0k} = \sum_{i=1}^k (u_0, w_i)w_i, u_{1k} = \sum_{i=1}^k (u_1, w_i)w_i \text{ and } u_{0k} \rightarrow u_0, u_{1k} \rightarrow u_1 \text{ in } H_0^1(\Omega).$$

Here we denote by  $(\cdot, \cdot)$  the inner product in  $\mathbb{L}^2(\Omega)$ .

Problem (1.1) generates the system of  $k$  ordinary differential equations

$$(2.2) \quad \begin{cases} \left| \sum_{i=1}^k (c_i^k(t))', w_i \right|^\rho \left( \sum_{i=1}^k c_i^k(t), w_i \right)'' = -\lambda_i c_i^k(t) + \lambda_i \int_0^t g(t-s)c_i^k(s)ds \\ + \left| \sum_{i=1}^k (c_i^k(t))', w_i \right|^{m(x)-2} \sum_{i=1}^k (c_i^k(t))', w_i \\ -\alpha \left| \sum_{i=1}^k c_i^k(t), w_i \right|^{p(x)-2} \sum_{i=1}^k c_i^k(t), w_i, \\ c_i^k(0) = (u_0, w_i), (c_i^k(0))' = (u_1, w_i), \quad i = 1, 2, \dots, k. \end{cases}$$

By the standard theory of the ODE system, we infer that the problem (2.2) admits a unique solution  $c_i^k(t)$  in  $[0, t_k]$ , where  $t_k > 0$ . Then we can obtain an approximate solution  $u_k(t)$  for (1.1) in  $V_k$ , over  $[0, t_k]$ . This solution can be extended to  $[0, T]$ , for any given  $T > 0$ , by the estimate below. Multiplying (2.1) by  $(c_i^k(t))'$  and summing with respect to  $i$  we arrive at the relation

$$(2.3) \quad \begin{aligned} 0 &= \frac{d}{dt} \left( \frac{1}{\rho+2} \|u_k'\|_{\rho+2}^{\rho+2} + \frac{1}{2} \|\nabla u_k\|_2^2 + \frac{1}{2} \|\nabla u_k'\|_2^2 \right) + \int_{\Omega} |u_k'|^{m(x)} dx \\ &- \frac{d}{dt} \left( \int_0^t g(t-s) \int_{\Omega} (\nabla u_k(s) \nabla u_k'(t)) dx ds \right) - \alpha \frac{d}{dt} \left( \int_{\Omega} \frac{1}{p(x)} |u_k|^{p(x)} dx \right). \end{aligned}$$

Multiplying (2.1) by  $(c_i^k(t))'$ , integrating over  $Q_T$ , using integration by part and Green formula, one obtains

$$(2.4) \quad \begin{aligned} &- \int_0^t g(t-s) \int_{\Omega} (\nabla u_k(s), \nabla u_k'(t)) dx ds = \frac{1}{2} \frac{d}{dt} (g \diamond \nabla u_k)(t) \\ &- \frac{1}{2} (g' \diamond \nabla u_k)(t) - \frac{1}{2} \frac{d}{dt} \int_0^t g(s) ds \|\nabla u_k\|_2^2 + \frac{1}{2} g(t) \|\nabla u_k\|_2^2, \end{aligned}$$

here

$$(\varphi \diamond \nabla \psi)(t) = \int_0^t \varphi(t-s) \|\nabla \psi(t) - \nabla \psi(s)\|_2^2 ds.$$

Combining (2.3)-(2.4) and  $(H_2) - (H_3)$ , we get

$$(2.5) \quad \begin{aligned} & \frac{d}{dt} \left( \frac{1}{\rho+2} \|u'_k\|_{\rho+2}^{\rho+2} + \frac{1}{2} \|\nabla u'_k\|_2^2 + \frac{1}{2} (g \diamond \nabla u_k)(t) \right. \\ & \left. + \frac{1}{2} \left( 1 - \int_0^t g(s) ds \right) \|\nabla u_k\|_2^2 - \alpha \int_{\Omega} \frac{1}{p(x)} |u_k|^{p(x)} dx \right) \\ & = \frac{1}{2} (g \diamond \nabla u_k)(t) - \frac{1}{2} g(t) \|\nabla u_k\|_2^2 - \int_{\Omega} |u'_k|^{m(x)} dx. \end{aligned}$$

Integrating (2.5) over  $(0, t)$ , and using the assumptions (1.2)-(1.4), it is easy to verify that

$$\begin{aligned} & \frac{1}{\rho+2} \|u'_k\|_{\rho+2}^{\rho+2} + \frac{1}{2} \|\nabla u'_k\|_2^2 + \frac{1}{2} \left( 1 - \int_0^t g(s) ds \right) \|\nabla u_k\|_2^2 + \frac{1}{2} (g \diamond \nabla u_k)(t) \\ & - \alpha \frac{1}{p(x)} |u_k|^{p(x)} \leq C_1, \end{aligned}$$

where  $C_1$  is a positive constant depending only on  $\|u_0\|_{H_0^1}, \|u_1\|_{H_0^1}$ .

According to the Lemma 2.1, we also have

$$(2.6) \quad \begin{aligned} & \frac{1}{\rho+2} \|u'_k\|_{\rho+2}^{\rho+2} + \frac{1}{2} \|\nabla u'_k\|_2^2 + \frac{1}{2} \left( 1 - \int_0^t g(s) ds \right) \|\nabla u_k\|_2^2 + \frac{1}{2} (g \diamond \nabla u_k)(t) \\ & - \max \left\{ \alpha \frac{1}{p^-} \|u_k\|_{p(x)}^-, \alpha \frac{1}{p^+} \|u_k\|_{p(x)}^+ \right\} \leq C_1. \end{aligned}$$

In view of  $(H_1) - (H_2) - (H_3)$  and  $(A_1) - (A_2)$ , we get

$$(2.7) \quad \|u'_k\|_{\rho+2}^{\rho+2} + \|\nabla u'_k\|_2^2 + (g \diamond \nabla u_k)(t) \leq C_2,$$

where  $C_2$  is positive constant depending only on  $\|u_0\|_{H_0^1}, \|u_1\|_{H_0^1}, l, p^-, p^+$ . It follows from (2.7) that

$$(2.8) \quad u_k \text{ is uniformly bounded in } \mathbb{L}^\infty(0, T; H_0^1(\Omega)).$$

$$(2.9) \quad u'_k \text{ is uniformly bounded in } \mathbb{L}^\infty(0, T; H_0^1(\Omega)).$$

Next, multiplying (1.1) by  $(c_i^k(t))''$  and then summing with respect to  $i$ , we get the following

$$(2.10) \quad \begin{aligned} & \int_{\Omega} |u'_k|^\rho |u''_k|_2^2 dx + \|\nabla u''_k\|_2^2 + \frac{d}{dt} \left( \frac{1}{m(x)} |u'_k|^{m(x)} \right) = - \int_{\Omega} \nabla u_k \nabla u''_k dx \\ & + \int_{\Omega} g(t-s) \int_{\Omega} \nabla u_k(s) \nabla u''_k dx ds + \alpha \int_{\Omega} |u_k|^{p(x)-2} u_k u''_k dx. \end{aligned}$$

Note that we have the estimates for  $\varepsilon > 0$

$$(2.11) \quad \int_{\Omega} |u'_k|^\rho |u''_k|_2^2 dx \leq C_\varepsilon \| |u'_k|^\rho \|_2^2 + \frac{1}{4\varepsilon} \|u''_k\|_2^2,$$

$$(2.12) \quad \left| - \int_{\Omega} \nabla u_k \nabla u_k'' dx \right| \leq \varepsilon \|\nabla u_k''\|_2^2 + \frac{1}{4\varepsilon} \|\nabla u_k\|_2^2,$$

$$(2.13) \quad \begin{aligned} & \left| - \int_0^t g(t-s) \int_{\Omega} \nabla u_k(s) \nabla u_k''(t) dx ds \right| \\ & \leq \frac{1}{4\varepsilon} \int_{\Omega} \left( \int_0^t g(t-s) \nabla u_k(s) ds \right)^2 dx + \varepsilon \|\nabla u_k''\|_2^2 \\ & \leq \varepsilon \|\nabla u_k''\|_2^2 + \frac{1}{4\varepsilon} \int_0^t g(s) ds \int_0^t g(t-s) \int_{\Omega} |\nabla u_k(s)|^2 dx ds \\ & \leq \varepsilon \|\nabla u_k''\|_2^2 + \frac{(1-l)g(0)}{4\varepsilon} \int_0^t \|\nabla u_k(s)\|_2^2 ds, \end{aligned}$$

and

$$(2.14) \quad \begin{aligned} \alpha \|\|u_k|^{p(x)-2} u_k u_k''\| & \leq \alpha \varepsilon \|\nabla u_k''\|_2^2 + \frac{\alpha}{4\varepsilon} \|\|u_k|^{p(x)-2} u_k\|_2^2 \\ & \leq \alpha \varepsilon \|\nabla u_k''\|_2^2 + \frac{\alpha}{4\varepsilon} \int_{\Omega} (\|u_k|^{p(x)-2} u_k)^2 dx. \end{aligned}$$

From Lemma 2.2, we have

$$(2.15) \quad \|u_k''\|_2^2 \leq C^2 \|\nabla u_k''\|_2^2,$$

and

$$(2.16) \quad \begin{aligned} & \int_{\Omega} (\|u_k'|^{p(x)-2} u_k')^2 dx = \int_{\Omega} |u_k|^{2(p(x)-1)} u_k dx \\ & \leq \max \left\{ \int_{\Omega} |u_k|^{2(p^- - 1)} dx, \int_{\Omega} |u_k|^{2(p^+ - 1)} dx \right\} \\ & \leq \max \left\{ C^{* \frac{1}{2(p^- - 1)}} \|\nabla u_k\|_{2(p^- - 1)}, C^{* \frac{1}{2(p^+ - 1)}} \|\nabla u_k'\|_{2(p^+ - 1)} \right\}, \end{aligned}$$

where  $C, C^*$  are embedding constants. Taking into account (2.10)-(2.16), we obtain

$$(2.17) \quad \begin{aligned} & C_{\varepsilon} \int_{\Omega} |\nabla u_t|^{2\rho} dx + \frac{1}{4\varepsilon} \int_{\Omega} |u_k''|^2 dx + (1 - 2\varepsilon - \alpha\varepsilon C) \|\nabla u_k''\|_2^2 \\ & + \frac{d}{dt} \left( \frac{1}{m(x)} |u_k'|^{m(x)} \right) \leq \frac{1}{4\varepsilon} \|\nabla u_k\|_2^2 + \frac{(1-l)g(0)}{4\varepsilon} \int_0^t \|\nabla u_k(s)\|_2^2 ds \\ & + \max \left\{ C^{* \frac{1}{2(p^- - 1)}} \|\nabla u_k\|_{\frac{1}{p^- - 1}}, C^{* \frac{1}{2(p^+ - 1)}} \|\nabla u_k\|_{\frac{1}{p^+ - 1}} \right\}. \end{aligned}$$

Integrating (2.17) over  $(0, t)$  and using (2.7), Lemma 2.3 we get

$$(2.18) \quad \begin{aligned} & C_{\varepsilon} T C_2^{2\rho} + \frac{1}{4\varepsilon} \int_0^t \|u_k''\|^2 dx + (1 - 2\varepsilon - \alpha\varepsilon C) \int_{\Omega} \|\nabla u_k''\|_2^2 ds \\ & + \int_{\Omega} \frac{1}{m(x)} |u_k'|^{m(x)} dx \leq \frac{1}{4\varepsilon} (C_3 + (1-l)g(0)T) + C_4, \end{aligned}$$

where  $C_4$  is a positive constant depending only on  $\|u_1\|_{H_0^1}$ . Taking  $\alpha, \varepsilon$  small enough in (2.18), we obtain the estimate

$$(2.19) \quad \frac{1}{4\varepsilon} \int_0^t \|u_k''\|^2 ds + \int_{\Omega} \frac{1}{m(x)} |u_k'|^{m(x)} dx \leq C_5.$$

Hence according to the Lemma 2.1, we have that

$$(2.20) \quad \frac{1}{4\varepsilon} \int_0^t \|u_k''\|^2 ds + \min \left\{ \frac{1}{m^+} \|u_k'\|_{m(x)}^-, \frac{1}{m^+} \|u_k'\|_{m(x)}^+ \right\} \leq C_5,$$

where  $C_5$  is a positive constant depending only on  $\|u_0\|_{H_0^1}$ ,  $\|u_1\|_{H_0^1}$ ,  $l$ ,  $g(0)$ ,  $T$ . From estimate (2.20), we get

$$(2.21) \quad u_k'' \text{ is uniformly bounded in } \mathbb{L}^2(0, T; H_0^1(\Omega)).$$

By (2.7)-(2.9) and (2.25), we infer that there exists a subsequence  $u_i$  of  $u_k$  and function  $u$  such that

$$(2.22) \quad u_i \rightharpoonup u \text{ weakly star in } L^\infty(0, T; H_0^1(\Omega)),$$

by (2.7)-(2.9) and (2.25), we infer that there exists a subsequence  $u_i$  of  $u_k$  and a function  $u$  such that

$$(2.23) \quad u_i \rightharpoonup u \text{ weakly star in } \mathbb{L}^\infty(0, T; H_0^1(\Omega)),$$

$$(2.24) \quad u_i \rightharpoonup u \text{ weakly in } \mathbb{L}^{p^-}(0, T; W^{1,p(x)}(\Omega)),$$

where  $C_5$  is a positive constant depending only on  $\|u_0\|_{H_0^1}$ ,  $\|u_1\|_{H_0^1}$ ,  $l$ ,  $g(0)$ ,  $T$ . From estimate (2.20), we get

$$(2.25) \quad u_k'' \text{ is uniformly bounded in } \mathbb{L}^2(0, T; H_0^1(\Omega)).$$

By (2.7)-(2.9) and (2.25), we infer that there exists a subsequence  $u_i$  of  $u_k$  and function  $u$  such that

$$(2.26) \quad u_i \rightharpoonup u \text{ weakly star in } \mathbb{L}^\infty(0, T; H_0^1(\Omega)),$$

by (2.7)-(2.9) and (2.25), we infer that there exists a subsequence  $u_i$  of  $u_k$  and function  $u$  such that

$$(2.27) \quad u_i \rightharpoonup u \text{ weakly star in } \mathbb{L}^\infty(0, T; H_0^1(\Omega)),$$

$$(2.28) \quad u_i \rightharpoonup u \text{ weakly in } \mathbb{L}^{p^-}(0, T; W^{1,p(x)}(\Omega)),$$

$$(2.29) \quad u_i' \rightharpoonup u' \text{ weakly star in } \mathbb{L}^\infty(0, T; H_0^1(\Omega)),$$

$$(2.30) \quad u_i'' \rightharpoonup u'' \text{ weakly in } \mathbb{L}^2(0, T; H_0^1(\Omega)).$$

Next, we will deal with the nonlinear term. From the Aubin-Lions theorem, see ([20], pp.57-58], it follows from (2.29) and (2.30) that there exists a subsequence of  $u_i$ , still represented by the same notation, such that



$u'_i \rightarrow u'$  strongly in  $\mathbb{L}^2(0, T; \mathbb{L}^2(\Omega))$ , which implies that  $u'_i \rightarrow u'$  almost everywhere in  $\Omega \times (0, T)$ . Hence, by (2.27) – (2.30), we have

$$(2.31) \quad |u'_i|^\rho u''_i \rightharpoonup |u'|^\rho u'' \quad \text{weakly in } \Omega \times (0, T),$$

$$(2.32) \quad |u_i|^{p(x)-2} u_i \rightharpoonup |u|^{p(x)-2} u \quad \text{weakly in } \Omega \times (0, T),$$

$$(2.33) \quad |u'_i|^{m(x)-2} u'_i \rightarrow |u'|^{m(x)-2} u' \quad \text{almost everywhere in } \Omega \times (0, T).$$

Multiplying (2.2) by  $\phi(t) \in C(0, T)$  (which  $C(0, T)$  is space of  $C^\infty$  function with compact support in  $(0, T)$ ) and integrating the obtained result over  $(0, T)$ , we obtain that

$$(2.34) \quad \langle Lu_k, w_i \phi(t) \rangle = 0, \quad i = 1, 2, \dots, k.$$

Note that  $\{w_i\}_{i=1}^\infty$  is basis of  $H_0^1(\Omega)$ . Convergence (2.27)-(2.33) is sufficient to pass to the limit in (2.34) in order to get

$$|u_t|^p u_{tt} - \Delta u - \Delta u_{tt} + \int_0^t g(t-s) \Delta u(s) ds + |u_t|^{m(x)-2} u_t = |u|^{p(x)-2} u \quad \text{in } \mathbb{L}^2(0, T; H_0^1(\Omega)),$$

for arbitrary  $T > 0$ . In view of (2.27) – (2.30) and Lemma 3.3.17 in [?], we derive that

$$u_k(0) \rightharpoonup u(0) \text{ weakly in } H_0^1(\Omega), \quad u'_k(0) \rightharpoonup u'(0) \text{ weakly in } H_0^1(\Omega).$$

Hence, we get  $u(0) = u_0, u_1(0) = u_1$ . Then, we conclude the proof of the Theorem 2.1.  $\square$

### 3. Blow up

In this section, we shall prove our main result concerning the blow-up of solutions to Theorem 2.1. For this task, we define

$$(3.1) \quad \begin{aligned} E(t) &= \frac{1}{\rho+2} \|u_t\|_{\rho+2}^{\rho+2} + \frac{1}{2} \|\nabla u_t\|_2^2 + \frac{1}{2} \left( 1 - \int_0^t g(s) ds \right) \|\nabla u\|_2^2 \\ &+ \frac{1}{2} (g \circ \nabla u)(t) - \frac{1}{p} \|u\|_p^p, \end{aligned}$$

where

$$(3.2) \quad (g \circ v)(t) = \int_0^\infty g(s) ds < \frac{\frac{p}{2} - 1}{\frac{p}{2} - 1 + \frac{1}{2p}}$$

**Lemma 3.1.** ([22]) *The modified energy functional satisfies the solution of (1.1)*

$$(3.3) \quad E'(t) \leq \frac{1}{2} (g' \circ \nabla u)(t) - \frac{1}{2} g(t) \|\nabla u(t)\|_2^2 - \|u_t\|_m^m \leq \frac{1}{2} (g' \circ \nabla u)(t).$$

**Theorem 3.1.** *Suppose that*

$$(3.4) \quad \max\{m, p\} \leq \frac{2(n-1)}{n-2}, \quad n \geq 3,$$

holds. Assume further that  $u_0, u_1 \in H_0^1(\Omega)$  and  $E(0) < 0$ . Then the solution of theorem 2.1 blows up in finite time

$$T^* \leq \frac{C(1-\alpha)}{\epsilon\gamma\alpha L^{\frac{\alpha}{1-\alpha}}(0)}$$

**Lemma 3.2.** *Suppose that (3.4) holds. Then there exists a positive constant  $C > 1$  depending on  $\Omega$  only such that for any  $u \in H_0^1(\Omega)$  and  $2 \leq s \leq p$ , we have*

$$(3.5) \quad \|u\|_p^s \leq C (\|\nabla u\|_2^2 + \|u\|_p^p).$$

*Proof.* 1. If  $\|u\|_p \leq 1$  then  $\|u\|_p^s \leq \|u\|_p^2 \leq C\|\nabla u\|_2^2$  by Sobolev embedding theorems.

2. If  $\|u\|_p > 1$  then  $\|u\|_p^s \leq \|u\|_p^p$ . Therefore (3.5) follows.

□

We set

$$H(t) = -E(t).$$

We use, throughout this paper,  $C$  to denote a generic positive constant depending on  $\Omega$  only. As a result of (3.1) and (3.5) we have

**Corollary 3.1.** *Let the assumptions of the lemma 3.2 hold. Then we have the following for all  $t \in [0, T)$ ,*

$$(3.6) \quad \|u\|_p^s \leq C \left( -H(t) - \|u_t\|_{\rho+2}^{\rho+2} + \|\nabla u_t\|_2^2 - \|\nabla u\|_2^2 - (g \circ \nabla u)(t) + \|u\|_p^p \right).$$

*Proof.* (Theorem 3.1) By multiplying equation (1.1) by  $-u_t$  and integrating over  $\Omega$  we obtain

$$(3.7) \quad \frac{d}{dt} \left\{ -\frac{1}{\rho+2} \int_{\Omega} |u_t|^{\rho+2} dx - \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx - \frac{1}{2} \int_{\Omega} |\nabla u_t|^2 dx + \frac{1}{p} \int_{\Omega} |u|^p dx \right\} + \int_0^t g(t-s) \int_{\Omega} \nabla u_t(t) \cdot \nabla u(s) dx ds = \int_{\Omega} |u_t|^m dx,$$

for any regular solution. This result can be extended to weak solutions by density

argument. But

$$\begin{aligned}
 & \int_0^t g(t-s) \int_{\Omega} \nabla u_t(t) \cdot \nabla u(\tau) dx ds \\
 &= \int_0^t \int_{\Omega} \nabla u_t(t) \cdot |\nabla u(s) - \nabla u(t)| dx ds + \int_0^t g(t-s) \int_{\Omega} \nabla u_t(t) \cdot \nabla u(t) dx ds \\
 &= -\frac{1}{2} \int_0^t g(t-s) \frac{d}{dt} \int_{\Omega} |\nabla u(s) - \nabla u(t)|^2 dx ds \\
 (3.8) \quad &+ \int_0^t g(s) \left( \frac{d}{dt} \frac{1}{2} \int_{\Omega} |\nabla u(t)|^2 dx \right) ds \\
 &= -\frac{1}{2} \frac{d}{dt} \left[ \int_0^t g(t-s) \int_{\Omega} |\nabla u(s) - \nabla u(t)|^2 dx ds \right] \\
 &+ \frac{1}{2} \frac{d}{dt} \left[ \int_0^t g(s) \int_{\Omega} |\nabla u(t)|^2 dx ds \right] \\
 &+ \frac{1}{2} \int_0^t g'(t-s) \int_{\Omega} |\nabla u(s) - \nabla u(t)|^2 dx ds - \frac{1}{2} g(t) \int_{\Omega} |\nabla u(t)|^2 dx ds.
 \end{aligned}$$

We then insert (3.8) in (3.7) to get

$$\begin{aligned}
 & \frac{d}{dt} \left\{ -\frac{1}{\rho+2} \int_{\Omega} |u_t|^{\rho+2} dx - \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx - \frac{1}{2} \int_{\Omega} |\nabla u_t|^2 dx + \frac{1}{p} \int_{\Omega} |u|^p dx \right\} \\
 (3.9) \quad & - \frac{1}{2} \frac{d}{dt} \left[ \int_0^t g(t-s) \int_{\Omega} |\nabla u(s) - \nabla u(t)|^2 dx d\tau \right] \\
 & + \frac{1}{2} \frac{d}{dt} \left[ \int_0^t g(s) \int_{\Omega} |\nabla u(t)|^2 dx ds \right] \\
 & = \int_{\Omega} |u_t|^m dx - \frac{1}{2} \int_0^t g'(t-s) \int_{\Omega} |\nabla u(s) - \nabla u(t)|^2 dx ds + \frac{1}{2} g(t) \|\nabla u(t)\|^2.
 \end{aligned}$$

By using the definition of  $H(t)$  the estimate (3.9) becomes

$$\begin{aligned}
 (3.10) \quad H'(t) &= \int_{\Omega} |u_t|^m dx - \frac{1}{2} \int_0^t g'(t-s) \int_{\Omega} |\nabla u(s) - \nabla u(t)|^2 dx ds \\
 &+ \frac{1}{2} g(t) \|\nabla u(t)\|^2 \geq 0.
 \end{aligned}$$

Consequently, we have

$$(3.11) \quad 0 < H(0) \leq H(t) \leq \frac{1}{p} \|u\|_p^p.$$

We define

$$(3.12) \quad L(t) = H^{1-\alpha}(t) + \frac{\epsilon}{\rho+1} \int_{\Omega} |u_t|^\rho u_t u dx + \epsilon \int_{\Omega} \nabla u_t \cdot \nabla u dx,$$

where  $\epsilon$  small to be chosen later and

$$0 < \alpha \leq \frac{p-m}{m-1}$$

By taking derivative of (3.12) and using (1.1) we obtain

$$\begin{aligned}
 (3.13) \quad L'(t) &= -\frac{1}{2}(1-\alpha)H^{-\alpha}(t) \int_0^t g'(t-s) \int_{\Omega} |\nabla u(s) - \nabla u(t)|^2 dx ds \\
 &+ (1-\alpha)H^{-\alpha}(t) \left\{ \int_{\Omega} |u_t|^m dx + \frac{1}{2}g(t)\|\nabla u(t)\|^2 \right\} \\
 &+ \frac{\epsilon}{\rho+1}\|u_t\|_{\rho+2}^{\rho+2} - \epsilon\|\nabla u\|_2^2 + \epsilon\|\nabla u_t\|_2^2 + \epsilon \int_0^t g(t-s) \int_{\Omega} \nabla u(s) \cdot \nabla u(t) ds \\
 &- \epsilon \int_{\Omega} |u_t|^{m-2} u_t u dx + \epsilon \|u\|_p^p.
 \end{aligned}$$

We then exploit Young's inequality, and use (3.1) to substitute for  $\int_{\Omega} |u(x,t)|^p dx$  hence (3.13) becomes

$$\begin{aligned}
 (3.14) \quad L'(t) &\geq (1-\alpha)H^{1-\alpha}(t)\|u_t\|_m^m + \frac{\epsilon}{\rho+1}\|u_t\|_{\rho+2}^{\rho+2} \\
 &- \epsilon \left( 1 - \int_0^t g(s) ds \right) \|\nabla u\|_2^2 + \epsilon\|\nabla u_t\|_2^2 - \epsilon\eta(g \circ \nabla u) \\
 &- \frac{\epsilon}{4\eta} \int_0^t g(s) ds \|\nabla u\|_2^2 - \epsilon \int_{\Omega} |u_t|^{m-2} u_t u dx + \epsilon \frac{p}{2}(g \circ \nabla u) \\
 &+ \epsilon \left( pH(t) + \frac{p}{\rho+2}\|u_t\|_{\rho+2}^{\rho+2} + \frac{p}{2}\|\nabla u_t\|_2^2 + \frac{p}{2} \left( 1 - \int_0^t g(s) ds \right) \|\nabla u\|_2^2 \right) \\
 &\geq (1-\alpha)H^{1-\alpha}(t)\|u_t\|_m^m + \epsilon \left( \frac{1}{\rho+1} + \frac{p}{\rho+2} \right) \|u_t\|_{\rho+2}^{\rho+2} + \epsilon pH(t) \\
 &+ \epsilon \left( \left( \frac{p}{2} - 1 \right) - \left( \frac{p}{2} - 1 + \frac{1}{4\eta} \right) \int_0^t g(s) ds \right) \|\nabla u\|_2^2 \\
 &+ \epsilon \left( \frac{p}{2} - \eta \right) (g \circ \nabla u) + \epsilon \left( \frac{p}{2} + 1 \right) \|\nabla u_t\|_2^2,
 \end{aligned}$$

for some number  $\eta$  with  $0 < \eta < \frac{p}{2}$ . By recalling (3.2), the estimate (3.14) is reduced to

$$\begin{aligned}
 (3.15) \quad L'(t) &\geq (1-\alpha)H^{-\alpha}(t)\|u_t\|_m^m + \epsilon \left( \frac{1}{\rho+1} + \frac{p}{\rho+2} \right) \|u_t\|_{\rho+2}^{\rho+2} \\
 &+ \epsilon pH(t) + \epsilon a_1(g \circ \nabla u) + \epsilon a_2\|\nabla u\|_2^2 + \epsilon a_3\|\nabla u_t\|_2^2 - \epsilon \int_{\Omega} |u_t|^{m-2} u_t u dx,
 \end{aligned}$$

where

$$a_1 = \frac{p}{2} - \eta > 0, \quad a_2 = \left( \frac{p}{2} - 1 \right) - \left( \frac{p}{2} - 1 + \frac{1}{4\eta} \right) \int_0^t g(s) ds > 0, \quad a_3 = \frac{p}{2} + 1 > 0.$$

To estimate the last term of (3.15), we use again Young's inequality

$$XY \leq \frac{\delta^r}{r} X^r + \frac{\delta^{-q}}{q} Y^q, \quad X, Y \geq 0, \quad \text{for all } \delta > 0, \quad \frac{1}{r} + \frac{1}{q} = 1,$$

with  $r = m$  and  $q = \frac{m}{(m-1)}$ . So we have

$$\int_{\Omega} |u_t|^{m-1} |u| dx \geq \frac{\delta^m}{m} \|u\|_m^m + \frac{m-1}{m} \delta^{\frac{-m}{(m-1)}} \|u_t\|_m^m,$$

which yields, by substitution in (3.15), for all  $\delta > 0$

$$(3.16) \quad \begin{aligned} L'(t) \geq & \left[ (1-\alpha)H^{-\alpha}(t) - \frac{m-1}{m} \delta^{\frac{-m}{(m-1)}} \right] \|u_t\|_m^m + \epsilon a_3 \|\nabla u_t\|_2^2 - \epsilon \frac{\delta^m}{m} \|u\|_m^m \\ & + \epsilon \left( \frac{1}{\rho+1} + \frac{p}{\rho+2} \right) \|u_t\|_{\rho+2}^{\rho+2} + \epsilon p H(t) + \epsilon a_1 (g \circ \nabla u) + \epsilon a_2 \|\nabla u\|_2^2. \end{aligned}$$

The inequality (3.16) remains valid even if  $\delta$  is time dependant since the integral is taken over the  $x$  variable. Therefore by taking  $\delta$  so that  $\delta^{\frac{-m}{(m-1)}} = kH^{-\alpha}(t)$ , for large  $k$  to be specified later, and substituting in (3.16) we arrive at

$$(3.17) \quad \begin{aligned} L'(t) \geq & \left[ (1-\alpha) - \frac{m-1}{m} \epsilon k \right] H^{-\alpha}(t) \|u_t\|_m^m + \epsilon \left( \frac{1}{\rho+1} \right) \|u_t\|_{\rho+2}^{\rho+2} \\ & + \epsilon a_1 (g \circ \nabla u) + \epsilon a_2 \|\nabla u\|_2^2 + \epsilon a_3 \|\nabla u_t\|_2^2 + \epsilon \left( \frac{p}{\rho+2} \right) \|u_t\|_{\rho+2}^{\rho+2} \\ & + \epsilon \left[ p H(t) - \frac{k^{1-m}}{m} H^{\alpha(m-1)}(t) \|u\|_m^m \right]. \end{aligned}$$

By exploiting (3.11) and inequality  $\|u\|_m^m \leq C \|u\|_p^m$ , we obtain

$$H^{\alpha(m-1)}(t) \|u\|_m^m \leq \left( \frac{1}{p} \right)^{\alpha(m-1)} C \|u\|_p^{m+\alpha p(m-1)},$$

therefore, from (3.17), one obtains

$$(3.18) \quad \begin{aligned} L'(t) \geq & \left[ (1-\alpha) - \frac{m-1}{m} \epsilon k \right] H^{-\alpha}(t) \|u_t\|_m^m \\ & + \epsilon \left( \frac{1}{\rho+1} + \frac{p}{\rho+2} \right) \|u_t\|_{\rho+2}^{\rho+2} + \epsilon a_1 (g \circ \nabla u) + \epsilon a_2 \|\nabla u\|_2^2 + \epsilon a_3 \|\nabla u_t\|_2^2 \\ & + \epsilon \left[ p H(t) - \frac{k^{1-m}}{m} \left( \frac{1}{p} \right)^{\alpha(m-1)} C \|u\|_p^{m+\alpha p(m-1)} \right]. \end{aligned}$$

At this stage, we use Corollary 3.1 for  $s = m + \alpha(m - 1) \leq p$ , to deduce from (3.18)

$$\begin{aligned}
 L'(t) &\geq \left[ (1 - \alpha) - \frac{m - 1}{m} \epsilon k \right] H^{-\alpha}(t) \|u_t\|_m^m \\
 &+ \epsilon \left( \frac{1}{\rho + 1} + \frac{p}{\rho + 2} \right) \|u_t\|_{\rho+2}^{\rho+2} + \epsilon a_1 (g \circ \nabla u) + \epsilon a_2 \|\nabla u\|_2^2 + \epsilon a_3 \|\nabla u_t\|_2^2 \\
 &+ \epsilon \left[ p H(t) - C_1 k^{1-m} \left\{ -H(t) - \|u_t\|_{\rho+2}^{\rho+2} + \|\nabla u_t\|_2^2 - \|\nabla u\|_2^2 \right. \right. \\
 (3.19) \quad &\left. \left. - (g \circ \nabla u)(t) + \|u\|_p^p \right\} \right] \\
 &\geq \left[ (1 - \alpha) - \frac{m - 1}{m} \epsilon k \right] H^{-\alpha}(t) \|u_t\|_m^m \\
 &+ \epsilon \left( \frac{1}{\rho + 1} + \frac{p}{\rho + 2} + C_1 k^{1-m} \right) \|u_t\|_{\rho+2}^{\rho+2} \\
 &+ \epsilon (a_1 + C_1 k^{1-m}) (g \circ \nabla u) + \epsilon (a_2 + C_1 k^{1-m}) \|\nabla u\|_2^2 \\
 &+ \epsilon (a_3 - C_1 k^{1-m}) \|\nabla u_t\|_2^2 + \epsilon (p + C_1 k^{1-m}) H(t) - \epsilon C_1 k^{1-m} \|u\|_p^p,
 \end{aligned}$$

where  $C_1 = \left(\frac{1}{p}\right)^{\alpha(m-1)} C/m$ . By noting that

$$H(t) \geq -\frac{1}{\rho + 2} \|u_t\|_{\rho+2}^{\rho+2} - \frac{1}{2} \|\nabla u_t\|_2^2 - \frac{1}{2} \|\nabla u\|_2^2 - \frac{1}{2} (g \circ \nabla u) + \frac{1}{p} \|u\|_p^p,$$

and writing  $p = 2a_4 + (p - 2a_4)$ , where  $a_4 = \min\{a_1, a_2, a_3\}$ , the estimate (3.19) yields

$$\begin{aligned}
 L'(t) &\geq \left[ (1 - \alpha) - \frac{m - 1}{m} \epsilon k \right] H^{-\alpha}(t) \|u_t\|_m^m \\
 (3.20) \quad &+ \epsilon \left( \frac{1}{\rho + 1} + \frac{p}{\rho + 2} + C_1 k^{1-m} - a_4 \right) \|u_t\|_{\rho+2}^{\rho+2} + \epsilon \left( \frac{2a_4}{p} - C_1 k^{1-m} \right) \|u\|_p^p \\
 &+ \epsilon (a_1 + C_1 k^{1-m} - a_4) (g \circ \nabla u) + \epsilon (a_2 + C_1 k^{1-m} - a_4) \|\nabla u\|_2^2 \\
 &+ \epsilon (a_3 - C_1 k^{1-m} - a_4) \|\nabla u_t\|_2^2 + \epsilon (p + C_1 k^{1-m} - 2a_4) H(t).
 \end{aligned}$$

We choose  $k$  large enough so that (3.20) becomes

$$\begin{aligned}
 (3.21) \quad L'(t) &\geq \left[ (1 - \alpha) - \frac{m - 1}{m} \epsilon k \right] H^{-\alpha}(t) \|u_t\|_m^m \\
 &+ \epsilon \gamma \left[ H(t) + \|u_t\|_{\rho+2}^{\rho+2} + \|\nabla u_t\|_2^2 + \|\nabla u\|_2^2 + (g \circ \nabla u)(t) + \|u\|_p^p \right],
 \end{aligned}$$

where  $\gamma > 0$  is the minimum of the coefficients of  $H(t)$ ,  $\|u_t\|_2^2$ ,  $\|u\|_p^p$ , and  $(g \circ \nabla u)(t)$  in (3.21). Once  $k$  is fixed (hence  $\gamma$ ), we pick  $\epsilon$  small enough so that

$$(1 - \alpha) - \frac{\epsilon k(m - 1)}{m} \geq 0,$$

and

$$L(0) = H^{1-\alpha}(0) + \frac{\epsilon}{\rho + 1} \int_{\Omega} |u_t|^\rho u_1 u_0 dx + \epsilon \int_{\Omega} \nabla u_1 \cdot \nabla u_0 dx > 0.$$

Therefore (3.21) takes the form

$$(3.22) \quad L'(t) \geq \epsilon\gamma \left[ H(t) + \|u_t\|_{\rho^2}^{\rho+2} + \|\nabla u_t\|_2^2 + \|\nabla u\|_2^2 + (g \circ \nabla u)(t) + \|u\|_p^p \right].$$

Consequently, we have

$$L(t) \geq L(0) > 0, \quad \text{for all } t \geq 0.$$

We now estimate

$$\left| \int_{\Omega} |u_t|^\rho u_t u dx \right| \leq \|u_t\|_{\rho+2}^{\rho+1} \|u\|_{\rho+2} \leq C \|u_t\|_{\rho+2}^{\rho+2} \|u\|_p,$$

we have

$$\left| \int_{\Omega} |u_t|^\rho u_t u dx \right|^{\frac{1}{1-\alpha}} \leq C \|u_t\|_{\rho+2}^{\frac{\rho+1}{1-\alpha}} \|u\|_p^{\frac{1}{1-\alpha}} \leq C \left( \|u_t\|_{\rho+2}^{\frac{\rho+1}{1-\alpha}\mu} + \|u\|_p^{\frac{\theta}{1-\alpha}} \right).$$

Where  $\frac{1}{\mu} + \frac{1}{\theta} = 1$ . Choose  $\mu = \frac{(1-\alpha)(\rho+2)}{\rho+1} (> 1)$ , then

$$\frac{\theta}{1-\alpha} = \frac{\rho+2}{(1-\alpha)(\rho+2) - (\rho+1)} < p.$$

Using Corollary 3.1, we obtain for all  $t \geq 0$

$$\left| \int_{\Omega} |u_t|^\rho u_t u dx \right|^{\frac{1}{1-\alpha}} \leq C \left[ -H(t) + \|u_t\|_{\rho+2}^{\rho+2} + \|\nabla u_t\|_2^2 - \|\nabla u\|_2^2 - (g \circ \nabla u)(t) + \|u\|_p^p \right].$$

Therefore,

$$(3.23) \quad \begin{aligned} L^{\frac{1}{1-\alpha}}(t) &= \left( H^{1-\alpha}(t) + \frac{\epsilon}{\rho+1} \int_{\Omega} |u_t|^\rho u_t u dx + \epsilon \int_{\Omega} \nabla u_t \cdot \nabla u dx \right)^{\frac{1}{1-\alpha}} \\ &\leq C \left[ \|u_t\|_{\rho+2}^{\rho+2} + H(t) + \|\nabla u_t\|_2^2 + \|\nabla u\|_2^{\frac{2}{1-2\alpha}} + \|u\|_p^p \right], \quad \forall t \geq 0. \end{aligned}$$

Noting that

$$(3.24) \quad \|\nabla u\|_2^{\frac{2}{1-2\alpha}} \leq C^{\frac{1}{1-2\alpha}} \leq \frac{C^{\frac{1}{1-2\alpha}}}{H(0)} H(t),$$

it follows from (3.23) and (3.24) that

$$(3.25) \quad L^{\frac{1}{1-\alpha}}(t) \leq C \left[ \|u_t\|_{\rho+2}^{\rho+2} + \|\nabla u_t\|_2^2 + \|u\|_p^p \right], \quad \forall t \geq 0.$$

Combining (3.22) and (3.25), we arrive at

$$(3.26) \quad L'(t) \geq \frac{\epsilon\gamma}{C} L^{\frac{1}{1-\alpha}}(t), \quad \forall t \geq 0.$$

A simple integration of (3.26) over  $(0, t)$  yields

$$(3.27) \quad L^{\frac{\alpha}{1-\alpha}}(t) \geq \frac{1}{L^{\frac{\alpha}{1-\alpha}}(0) - \epsilon\gamma t\alpha/|C(1-\alpha)|}$$

This shows that  $L(t)$  blows up in finite time.

$$(3.28) \quad T^* \leq \frac{C(1-\alpha)}{\epsilon\gamma\alpha L^{\frac{\alpha}{1-\alpha}}(0)}$$

Summarizing, the proof is completed.  $\square$

#### 4. Asymptotic Behavior

In this section, we investigate the asymptotic behavior of the problem (1.1). We define

$$(4.1) \quad G(t) = ME(t) + \epsilon\psi(t) + \chi(t),$$

where  $\epsilon$  and  $M$  are positive constants which shall be determined later, and

$$(4.2) \quad \psi(t) = \frac{1}{\rho+1}\xi(t) \int_{\Omega} |u_t|^\rho u_t u dx + \xi(t) \int_{\Omega} \nabla u_t \cdot \nabla u dx,$$

$$(4.3) \quad \chi(t) = \xi(t) \int_{\Omega} \left( \Delta u_t - \frac{|u_t|^\rho u_t}{\rho+1} \right) \int_0^t g(t-s) [u(t) - u(s)] ds dx.$$

**Theorem 4.1.** *Let  $(u_0, u_1) \in H_0^1(\Omega) \times H_0^1(\Omega)$  be given. Assume that  $(H_1) - (H_3)$  and (3.4) hold. Then for each  $t_0 > 0$ , there exists two positive constants  $K$  and  $\kappa$  such that the solution of (1.1) satisfies*

$$(4.4) \quad E(t) \leq K e^{-\kappa \int_0^t \xi(s) ds}, \quad t \geq t_0.$$

For our purposes, we need:

**Theorem 4.2.** *([22]) Suppose that  $(H_1) - (H_3)$  and (3.4) hold. If  $u_0, u_1 \in H_0^1(\Omega)$  and*

$$(4.5) \quad \frac{C_*^p}{l} \left( \frac{2p}{(p-2)l} E(0) \right)^{\frac{p-2}{2}} < 1,$$

where  $C_*$  is the best Poincaré's constant. Then the solution of the problem (1.1) is global in time and satisfies

$$(4.6) \quad l \|\nabla u(t)\| + \|\nabla u_t(t)\| \leq \frac{2p}{p-2} E(0).$$

The proof of the theorem 4.2 is detailed in [22].



**Lemma 4.1.** *Let  $u \in L^\infty(0, T; H_0^1(\Omega))$  be the solution of (1.1), then we have*

$$(4.7) \int_{\Omega} \left( \int_0^t g(t-s) [u(t) - u(s)] ds \right)^{\rho+2} dx \leq C_*^{\rho+2} (1-l)^{\rho+1} \left( \frac{4pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} \times (g \circ \nabla u)(t).$$

*Proof.* Here, we point out that

$$\int_0^t g(t-s) [u(t) - u(s)] ds = \int_0^t [g(t-s)]^{\frac{\rho+1}{\rho+2}} [g(t-s)]^{\frac{1}{\rho+2}} [u(t) - u(s)] ds,$$

then by using Hölder’s inequality, we get

$$\begin{aligned} & \int_{\Omega} \left( \int_0^t g(t-s)(u(t) - u(s)) ds \right)^{\rho+2} dx \leq \\ & \leq \left( \int_0^\infty g(s) ds \right)^{\rho+1} \int_0^t g(t-s) \int_{\Omega} |u(t) - u(s)|^{\rho+2} dx ds \\ & \leq C_*^{\rho+2} (1-l)^{\rho+1} \int_0^t g(t-s) \|\nabla u(t) - \nabla u(s)\|_2^{\rho+2} ds \\ & \leq C_*^{\rho+2} (1-l)^{\rho+1} \left( \frac{4pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} (g \circ \nabla u)(t). \end{aligned}$$

This ends the proof.  $\square$

**Lemma 4.2.** *For  $\epsilon > 0$  small enough while  $M > 0$  is large enough, the relation*

$$\alpha_1 G(t) \leq E(t) \leq \alpha_2 G(t),$$

*holds for two positive constants  $\alpha_1$  and  $\alpha_2$ .*

*Proof.* By using Young’s inequality, the Sobolev embedding theorem, (1.6),(4.6) and Lemma 4.1, we can derive that

$$\begin{aligned} \left| \frac{1}{\rho+1} \int_{\Omega} |u_t|^\rho u_t u dx \right| & \leq \frac{1}{\rho+2} \|u_t\|_{\rho+2}^{\rho+2} + \frac{1}{(\rho+1)(\rho+2)} \|u\|_{\rho+2}^{\rho+2} \\ & \leq \frac{1}{\rho+2} \|u_t\|_{\rho+2}^{\rho+2} + \frac{C_*^{\rho+2}}{(\rho+1)(\rho+2)} \left( \frac{2pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} \|\nabla u\|_2^2, \end{aligned}$$

and

$$\begin{aligned} & \left| \frac{1}{\rho+1} \int_{\Omega} |u_t|^\rho u_t \int_0^t g(t-s) [u(t) - u(s)] ds dx \right| \\ & \leq \frac{1}{\rho+1} \|u_t\|_{\rho+2}^{\rho+2} + \frac{1}{(\rho+1)(\rho+2)} \int_{\Omega} \left( \int_0^t g(t-s) [u(t) - u(s)] ds \right)^{\rho+2} dx \\ & \leq \frac{1}{\rho+1} \|u_t\|_{\rho+2}^{\rho+2} + \frac{C_*^{\rho+2}}{(\rho+1)(\rho+2)} (1-l)^{\rho+1} \left( \frac{4pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} (g \circ \nabla u)(t). \end{aligned}$$

It follows that

$$\begin{aligned}
G(t) &\leq ME(t) + \left( \frac{1}{\rho+1} + \frac{\epsilon}{\rho+2} \right) \xi(t) \|u_t\|_{\rho+2}^{\rho+2} \\
&+ \epsilon \left[ \frac{C_*^{\rho+2}}{(\rho+1)(\rho+2)} \left( \frac{2pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} + \frac{1}{2} \right] \xi(t) \|\nabla u\|_2^2 + \frac{\epsilon+1}{2} \xi(t) \|\nabla u_t\|_2^2 \\
&+ \left[ \frac{1-l}{2} + \frac{C_*^{\rho+2}}{(\rho+1)(\rho+2)} (1-l)^{\rho+1} \left( \frac{4pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} \right] \xi(t) (g \circ \nabla u)(t) \\
&\leq ME(t) + \left( \frac{1}{\rho+1} + \frac{\epsilon}{\rho+2} \right) N \|u_t\|_{\rho+2}^{\rho+2} \\
&+ \epsilon \left[ \frac{C_*^{\rho+2}}{(\rho+1)(\rho+2)} \left( \frac{2pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} + \frac{1}{2} \right] N \|\nabla u\|_2^2 + \frac{\epsilon+1}{2} N \|\nabla u_t\|_2^2 \\
&+ \left[ \frac{1-l}{2} + \frac{C_*^{\rho+2}}{(\rho+1)(\rho+2)} (1-l)^{\rho+1} \left( \frac{4pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} \right] N (g \circ \nabla u)(t) \leq \frac{1}{\alpha_1} E(t),
\end{aligned}$$

and

$$\begin{aligned}
G(t) &\geq ME(t) - \left( \frac{1}{\rho+1} + \frac{\epsilon}{\rho+2} \right) \xi(t) \|u_t\|_{\rho+2}^{\rho+2} \\
&- \epsilon \left[ \frac{C_*^{\rho+2}}{(\rho+1)(\rho+2)} \left( \frac{2pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} + \frac{1}{2} \right] \xi(t) \|\nabla u\|_2^2 - \frac{\epsilon+1}{2} \xi(t) \|\nabla u_t\|_2^2 \\
&- \left[ \frac{1-l}{2} + \frac{C_*^{\rho+2}}{(\rho+1)(\rho+2)} (1-l)^{\rho+1} \left( \frac{4pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} \right] \xi(t) (g \circ \nabla u)(t) \\
&\geq \left[ \frac{M}{\rho+2} - \left( \frac{1}{\rho+1} + \frac{\epsilon}{\rho+2} \right) N \right] \|u_t\|_{\rho+2}^{\rho+2} + \left( \frac{M}{2} - \frac{\epsilon+1}{2} N \right) \|\nabla u_t\|_2^2 \\
&+ \left\{ \frac{M}{2} l - \epsilon \left[ \frac{C_*^{\rho+2}}{(\rho+1)(\rho+2)} \left( \frac{2pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} + \frac{1}{2} \right] N \right\} - \frac{M}{p} \|u\|_p^p \\
&+ \left\{ \frac{M}{2} - \left[ \frac{1-l}{2} + \frac{C_*^{\rho+2}}{(\rho+1)(\rho+2)} (1-l)^{\rho+1} \left( \frac{4pE(0)}{(p-2)l} \right)^{\frac{\rho}{2}} \right] N \right\} (g \circ \nabla u)(t) \geq \frac{1}{\alpha_2} E(t).
\end{aligned}$$

For  $\epsilon > 0$  small enough while  $M > 0$  is large enough. This completes the proof.  $\square$

**Lemma 4.3.** Under the assumptions  $(H_1) - (H_3)$  and (4.6), the functional

$$\psi(t) = \frac{1}{\rho+1} \xi(t) \int_{\Omega} |u_t|^{\rho} u_t u dx + \xi(t) \int_{\Omega} \nabla u_t \cdot \nabla u dx,$$

satisfies the solutions of (1.1),

$$\begin{aligned}
(4.8) \quad \psi'(t) &\leq - \left[ \frac{l}{2} - \delta C_*^2 - k\delta \left( 1 + \frac{C_*^2}{\rho+1} \right) \right] \xi(t) \|\nabla u\|_2^2 \\
&+ \left\{ 1 + \frac{A}{4\delta} + \frac{k}{4\delta} \left[ 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{(p-2)l} \right)^{\rho} \right] \right\} \xi(t) \|\nabla u_t\|_2^2 \\
&+ \frac{1-l}{2l} \xi(t) (g \circ \nabla u)(t) + \frac{1}{\rho+1} \xi(t) \|u_t\|_{\rho+2}^{\rho+2} + \xi(t) \|u\|_p^p.
\end{aligned}$$

*Proof.* By using the equation of (1.1), we easily see that

$$\begin{aligned}
 \psi'(t) &= \frac{1}{\rho+1} \xi(t) \|u_t\|_{\rho+2}^{\rho+2} + \xi(t) \int_{\Omega} |u_t|^\rho u_{tt} u dx + \xi(t) \|\nabla u_t\|_2^2 \\
 &+ \xi(t) \int_{\Omega} \nabla u \cdot \nabla u_{tt} dx + \frac{1}{\rho+1} \xi'(t) \int_{\Omega} |u_t|^\rho u_t u dx + \xi'(t) \int_{\Omega} \nabla u_t \cdot \nabla u dx \\
 (4.9) \quad &= \frac{1}{\rho+1} \xi(t) \|u_t\|_{\rho+2}^{\rho+2} + \xi(t) \|\nabla u_t\|_2^2 - \xi(t) \|\nabla u\|_2^2 \\
 &+ \xi(t) \int_{\Omega} \nabla u \cdot \int_0^t g(t-s) \nabla u(s) ds dx - \xi(t) \int_{\Omega} |u_t|^{m-2} u_t u dx + \xi(t) \|u\|_p^p \\
 &+ \frac{1}{\rho+1} \xi'(t) \int_{\Omega} |u_t|^\rho u_t u dx + \xi'(t) \int_{\Omega} \nabla u_t \cdot \nabla u dx.
 \end{aligned}$$

Now we estimate

$$\begin{aligned}
 (4.10) \quad &\int_{\Omega} \nabla u(t) \cdot \int_0^t g(t-s) \nabla u(s) ds dx \leq \frac{1}{2} \|\nabla u\|_2^2 \\
 &+ \frac{1}{2} \int_{\Omega} \left( \int_0^t g(t-s) (|\nabla u(s) - \nabla u(t)| + |\nabla u(t)|) ds \right)^2 dx.
 \end{aligned}$$

We use Young's inequality and the fact that

$$\int_0^t g(s) ds \leq \int_0^\infty g(s) ds = 1 - l,$$

it follows from (4.10) for  $\eta = \frac{l}{1-l} > 0$  that

$$\begin{aligned}
 &\int_{\Omega} \nabla u(t) \cdot \int_0^t g(t-s) \nabla u(s) ds dx \leq \frac{1}{2} (1 + \eta) \int_{\Omega} \left( \int_0^t g(t-s) |\nabla u(t)| ds \right)^2 dx \\
 &+ \frac{1}{2} \left( 1 + \frac{1}{\eta} \right) \int_{\Omega} \left( \int_0^t g(t-s) |\nabla u(s) - \nabla u(t)| ds \right)^2 dx + \frac{1}{2} \|\nabla u\|_2^2 \\
 &\leq \frac{1}{2} \|\nabla u\|_2^2 + \frac{1}{2} (1 + \eta) (1 - l)^2 \|\nabla u(t)\|_2^2 + \frac{1}{2} \left( 1 + \frac{1}{\eta} \right) (1 - l) (g \circ \nabla u)(t) \\
 &\leq \frac{2-l}{2} \|\nabla u(t)\|_2^2 + \frac{1}{2l} (1 - l) (g \circ \nabla u)(t),
 \end{aligned}$$

and

$$(4.11) \quad \int_{\Omega} |u_t|^\rho u_t u dx \leq \frac{1}{4\delta} \|u_t\|_{2(\rho+1)}^{2(\rho+1)} + \delta C_*^2 \|\nabla u\|_2^2,$$

for any  $\delta > 0$ . In view of (4.6) and the Sobolev embedding

$$H_0^1(\Omega) \hookrightarrow L^{2(\rho+1)}(\Omega), \quad \text{for } 0 < \rho \leq \frac{2}{n-2} \text{ if } n \geq 3 \text{ and } \rho > 0 \text{ if } n = 1, 2,$$

we get

$$(4.12) \quad \|u_t\|_{2(\rho+1)}^{2(\rho+1)} \leq C_*^{2(\rho+1)} \left( \frac{2pE(0)}{p-2} \right)^\rho \|\nabla u_t\|_2^2.$$

It follows from (4.11) and (4.12) that

$$(4.13) \quad \frac{1}{\rho+1} \int_{\Omega} |u_t|^\rho u_t u dx \leq \frac{1}{4\delta} \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho \|\nabla u_t\|_2^2 + \frac{1}{\rho+1} \delta C_*^2 \|\nabla u\|_2^2,$$

and

$$(4.14) \quad \left| \int_{\Omega} |u_t|^{m-2} u_t u dx \right| \leq \delta \int_{\Omega} |u|^2 dx + \frac{1}{4\delta} \int_{\Omega} |u_t|^{2m-2} dx \leq \delta C_*^2 \|\nabla u\|_2^2 + \frac{1}{4\delta} C_*^{2m-2} \|\nabla u_t\|_{2m-2}^{2m-2} \leq \delta C_*^2 \|\nabla u\|_2^2 + \frac{1}{4\delta} C_*^{2m-2} \left( \frac{2pE(0)}{p-2} \right)^{\frac{m-2}{2}} \|\nabla u_t\|_2^2 \leq \delta C_*^2 \|\nabla u\|_2^2 + \frac{A}{4\delta} \|\nabla u_t\|_2^2.$$

Also

$$(4.15) \quad \int_{\Omega} \nabla u_t \cdot \nabla u dx \leq \frac{1}{4\delta} \|\nabla u_t\|_2^2 + \delta \|\nabla u\|_2^2.$$

By combining (4.9), (4.10), (4.13), (4.14) and (4.15), we deduce easily the estimate (4.8). This completes our proof.  $\square$

**Lemma 4.4.** *Under the assumptions  $(H_1) - (H_3)$ , the functional*

$$\chi(t) = \xi(t) \int_{\Omega} \left( \Delta u_t - \frac{|u_t|^\rho u_t}{\rho+1} \right) \int_0^t g(t-s) [u(t) - u(s)] ds dx,$$

satisfies, along solutions of (1.1) and for  $\delta > 0$

$$(4.16) \quad \begin{aligned} \chi'(t) &\leq \delta_1 \left[ 1 + 2(1-l)^2 + C_*^{2p-2} \left( \frac{2pE(0)}{(p-2)l} \right)^{p-2} \right] \xi(t) \|\nabla u\|_2^2 \\ &\quad \left[ \frac{(\rho+2)k}{4\delta_1(\rho+1)} + 2\delta_1 + \frac{1}{2\delta_1} + \frac{C_*^2}{2\delta_1} \right] (1-l)\xi(t) (g \circ \nabla u)(t) \\ &\quad - \frac{g(0)}{4\delta_1} \left( 1 + \frac{C_*^2}{\rho+1} \right) \xi(t) (g' \circ \nabla u)(t) - \frac{1}{\rho+1} \xi(t) \left( \int_0^t g(s) ds \right) \|u_t\|_{\rho+2}^{\rho+2} \\ &\quad + \left\{ (k+1)\delta_1 \left[ 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho \right] + \delta_1 \tilde{A} - \int_0^t g(s) ds \right\} \xi(t) \|\nabla u_t\|_2^2. \end{aligned}$$

*Proof.* Applying (1.1), the computation yields

$$\begin{aligned} \chi'(t) &= \xi(t) \int_{\Omega} (\Delta u_{tt} - |u_t|^\rho u_{tt}) \int_0^t g(t-s) [u(t) - u(s)] ds dx \\ &\quad + \xi(t) \int_{\Omega} \left( \Delta u_t - \frac{|u_t|^\rho u_t}{\rho+1} \right) \int_0^t g'(t-s) [u(t) - u(s)] ds dx \\ &\quad + \xi(t) \int_0^t g(s) ds \int_{\Omega} \left( \Delta u_t - \frac{|u_t|^\rho u_t}{\rho+1} \right) u_t dx \\ &\quad + \xi'(t) \int_{\Omega} \Delta u_t \int_0^t g(t-s) [u(t) - u(s)] ds dx \\ &\quad - \frac{1}{\rho+1} \xi'(t) \int_{\Omega} |u_t|^\rho u_t \int_0^t g(t-s) [u(t) - u(s)] ds dx. \end{aligned}$$

By integrating the parts, it follows that

$$\begin{aligned}
 \chi'(t) &= \xi(t) \int_{\Omega} \nabla u(t) \cdot \int_0^t g(t-s)[\nabla u(t) - \nabla u(s)] ds dx \\
 &\quad - \xi(t) \int_{\Omega_t} \int_0^t g(t-s) \nabla u(s) ds \cdot \int_0^t g(t-s)[\nabla u(t) - \nabla u(s)] ds dx \\
 &\quad - \xi(t) \int_0^t g(s) ds \|\nabla u_t\|_2^2 - \xi(t) \int_{\Omega} \nabla u_t \cdot \int_0^t g'(t-s)[\nabla u(t) - \nabla u(s)] ds dx \\
 &\quad - \frac{1}{\rho+1} \xi(t) \int_{\Omega} |u_t|^\rho u_t \int_0^t g'(t-s)[u(t) - u(s)] ds dx \\
 (4.17) \quad &\quad - \frac{1}{\rho+1} \xi(t) \|u_t\|_{\rho+2}^{\rho+2} \int_0^t g(s) ds \\
 &\quad + \xi(t) \int_{\Omega} |u_t|^{m-2} u_t \int_0^t g(t-s)[u(t) - u(s)] ds dx \\
 &\quad - \xi(t) \int_{\Omega} |u|^{p-2} u \int_0^t g(t-s)[u(t) - u(s)] ds dx \\
 &\quad - \xi'(t) \int_{\Omega} \nabla u_t \cdot \int_0^t g(t-s)[\nabla u(t) - u(s)] ds dx \\
 &\quad - \frac{1}{\rho+1} \xi'(t) \int_{\Omega} |u_t|^\rho u_t \int_0^t g(t-s)[u(t) - u(s)] ds dx.
 \end{aligned}$$

In fact, by exploiting Young's inequality, we get that for any  $\delta_1 > 0$

$$\begin{aligned}
 (4.18) \quad \int_{\Omega} \nabla u(t) \cdot \left( \int_0^t g(t-s)[\nabla u(t) - \nabla u(s)] ds \right) dx &\leq \delta_1 \|\nabla u\|_2^2 \\
 &\quad + \frac{1}{4\delta_1} (1-l)(g \circ \nabla u)(t),
 \end{aligned}$$

$$\begin{aligned}
 &\int_{\Omega} \int_0^t g(t-s) \nabla u(s) ds \cdot \int_0^t g(t-s)[\nabla u(t) - \nabla u(s)] ds dx \\
 &\leq \delta_1 \int_{\Omega} \left( \int_0^t g(t-s) |\nabla u(s) - \nabla u(t)| + |\nabla u(t)| ds \right)^2 dx \\
 (4.19) \quad &+ \frac{1}{4\delta_1} \int_{\Omega} \left( \int_0^t g(t-s) |\nabla u(t) - \nabla u(s)| ds \right)^2 dx \\
 &\leq \left( 2\delta_1 + \frac{1}{4\delta_1} \right) \int_{\Omega} \left( \int_0^t g(t-s) |\nabla u(t) - \nabla u(s)| ds \right)^2 dx \\
 &\leq \left( 2\delta_1 + \frac{1}{4\delta_1} \right) (1-l)(g \circ \nabla u)(t) + 2\delta_1 (1-l)^2 \|\nabla u\|_2^2 + 2\delta_1 (1-l)^2 \|\nabla u\|_2^2,
 \end{aligned}$$

and

$$\begin{aligned}
 (4.20) \quad \int_{\Omega} \nabla u_t \cdot \int_0^t g'(t-s)[\nabla u(t) - \nabla u(s)] ds dx &\leq \delta_1 \|\nabla u_t\|_2^2 \\
 &\quad + \frac{g(0)}{4\delta_1} (-g' \circ \nabla u)(t).
 \end{aligned}$$

Also

$$\begin{aligned}
 (4.21) \quad & \frac{1}{\rho + 1} \int_{\Omega} |u_t|^\rho u_t \int_0^t g'(t - s)[u(t) - u(s)] ds dx \\
 & \leq \delta_1 \frac{C_*^{2(\rho+1)}}{\rho + 1} \left( \frac{2pE(0)}{p - 2} \right)^\rho \|\nabla u_t\|_2^2 + \frac{g(0)}{4\delta_1(\rho + 1)} C_*^2(-g' \circ \nabla u)(t).
 \end{aligned}$$

Similarly, we get

$$\begin{aligned}
 (4.22) \quad & \left| \int_{\Omega} |u_t|^{m-2} u_t \int_0^t g(t - s)(u(t) - u(s)) ds dx \right| \\
 & \leq \delta_1 \|u_t\|_{2m-2}^{2m-2} + \frac{1}{4\delta_1} \int_{\Omega} \left( \int_0^t g(t - s)|u(t) - u(s)| ds \right)^2 dx \\
 & \leq \delta_1 C_*^{2m-2} \|\nabla u_t\|_{2m-2}^{2m-2} + \frac{1}{4\delta_1} \int_0^t g(s) ds \int_0^t g(t - s) \int_{\Omega} |u(t) - u(s)|^2 ds dx \\
 & \leq \delta_1 \tilde{A} \|\nabla u_t\|_2^2 + \frac{1}{4\delta_1} (1 - l) C_*^2(g \circ \nabla u)(t),
 \end{aligned}$$

$$\begin{aligned}
 (4.23) \quad & - \int_{\Omega} |u|^{p-2} u \int_0^t g(t - s)[u(t) - u(s)] ds dx \\
 & \leq \frac{1}{4\delta_1} \int_{\Omega} \left( \int_0^t g(t - s)[u(t) - u(s)] ds \right)^2 dx \\
 & + \delta_1 \|u\|_{2p-2}^{2p-2} \leq \delta_1 \|u\|_{2p-2}^{2p-2} + \frac{C_*^2(1 - l)}{4\delta_1} (g \circ \nabla u)(t) \\
 & \leq \delta_1 C_*^{2p-2} \left( \frac{2pE(0)}{(p - 2)l} \right)^{p-2} \|\nabla u\|_2^2 + \frac{C_*^2(1 - l)}{4\delta_1} (g \circ \nabla u)(t),
 \end{aligned}$$

and

$$\begin{aligned}
 (4.24) \quad & \int_{\Omega} \nabla u_t \cdot \int_0^t g(t - s)[\nabla u(t) - \nabla u(s)] ds dx \leq \delta_1 \|\nabla u_t\|_2^2 \\
 & + \frac{1}{4\delta_1} (1 - l)(g \circ \nabla u)(t).
 \end{aligned}$$

We estimate

$$\begin{aligned}
 (4.25) \quad & \frac{1}{\rho + 1} \int_{\Omega} |u_t|^\rho u_t \int_0^t g(t - s)[u(t) - u(s)] ds dx \\
 & \leq \frac{1}{\rho + 1} \delta_1 \|u_t\|_{2(\rho+1)}^{2(\rho+1)} + \frac{1}{4\delta_1(\rho + 1)} \int_{\Omega} \left( \int_0^t g(t - s)[u(t) - u(s)] ds \right)^2 dx \\
 & \leq \delta_1 \frac{C_*^{2(\rho+1)}}{\rho + 1} \left( \frac{2pE(0)}{p - 2} \right)^\rho \|\nabla u_t\|_2^2 + \frac{1 - l}{4\delta_1(\rho + 1)} (g \circ \nabla u)(t).
 \end{aligned}$$

Combining the estimates (4.18)-(4.25) and (4.17), the assertion of the lemma 4.4 is established.  $\square$

*Proof.* (Theorem 4.1). Since  $g$  is positive, we have that, for any  $t_0 > 0$ ,

$$\int_0^t g(s)ds \geq \int_0^{t_0} g(s)ds = g_0 > 0, \quad t \geq t_0.$$

By using (4.1),(4.8),(4.16) and Lemma 4.1, a series of computations yields, for  $t \geq t_0$ ,

$$\begin{aligned} G'(t) &\leq \frac{M}{2}(g' \circ \nabla u)(t) - \epsilon \left[ \frac{l}{2} - \delta C_*^2 - k\delta \left( 1 + \frac{C_*^2}{\rho + 1} \right) \right] \xi(t) \|\nabla u\|_2^2 \\ &+ \epsilon \frac{1-l}{2l} \xi(t)(g \circ \nabla u)(t) + \epsilon \frac{1}{\rho+1} \xi(t) \|u_t\|_{\rho+2}^{\rho+2} + \epsilon \xi(t) \|u\|_p^p \\ &+ \epsilon \left[ 1 + \frac{A}{4\delta} + \frac{k}{4\delta} \left[ 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho \right] \right] \xi(t) \|\nabla u_t\|_2^2 \\ &+ \delta_1 \left[ 1 + 2(1-l)^2 + C_*^{2p-2} \left( \frac{2pE(0)}{(p-2)l} \right)^{p-2} \right] \xi(t) \|\nabla u\|_2^2 \\ &+ \left[ \frac{(\rho+2)k}{4\delta_1(\rho+1)} + 2\delta_1 + \frac{1}{2\delta_1} + \frac{C_*^2}{2\delta_1} \right] (1-l)\xi(t)(g \circ \nabla u)(t) \\ &- \frac{g(0)}{4\delta_1} \left( 1 + \frac{C_*^2}{\rho+1} \right) \xi(t)(g' \circ \nabla u)(t) - \frac{1}{\rho+1} \xi(t) \left( \int_0^t g(s)ds \right) \|u_t\|_{\rho+2}^{\rho+2} \\ (4.26) \quad &+ \left\{ (k+1)\delta_1 \left[ 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho \right] + \delta_1 \tilde{A} - \int_0^t g(s)ds \right\} \xi(t) \|\nabla u_t\|_2^2 \\ &\leq - \left\{ \left[ g_0 - \epsilon \left( 1 + \frac{A}{4\delta} + \frac{k}{4\delta} \left( 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho \right) \right) \right] \right. \\ &\quad \left. - (k+1)\delta_1 \left[ 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho + \delta_1 \tilde{A} \right] \right\} \xi(t) \|u_t\|_2^2 \\ &- \left\{ \epsilon \left[ \frac{l}{2} - \delta C_*^2 - k\delta \left( 1 + \frac{C_*^2}{\rho+1} \right) \right] \right. \\ &\quad \left. - \delta_1 \left[ 1 + 2(1-l)^2 + C_*^{2p-2} \left( \frac{2pE(0)}{(p-2)l} \right)^{p-2} \right] \right\} \xi(t) \|\nabla u\|_2^2 + \epsilon \xi(t) \|u\|_p^p \\ &- \left[ \frac{M}{2} - \frac{g(0)}{4\delta_1} \left( 1 + \frac{C_*^2}{\rho+1} \right) N \right] (-g' \circ \nabla u)(t) - (g_0 - \epsilon) \frac{1}{\rho+1} \xi(t) \|u_t\|_{\rho+2}^{\rho+2} \\ &+ \left[ \frac{\epsilon}{2l} + \frac{(\rho+2)k}{4\delta_1(\rho+1)} + 2\delta_1 + \frac{1}{2\delta_1} + \frac{C_*^2}{2\delta_1} \right] (1-l)\xi(t)(g \circ \nabla u)(t). \end{aligned}$$

At this point, we choose  $\delta > 0$  so small that

$$\frac{l}{2} - \delta C_*^2 - k\delta \left( 1 + \frac{C_*^2}{\rho+1} \right) > \frac{l}{4}.$$

Hence  $\delta$  is fixed, we choose  $\epsilon > 0$  small enough so that Lemma 4.2 holds and that

$$\epsilon < \frac{g_0}{2 \left[ 1 + \frac{A}{4\delta} + \frac{k}{4\delta} \left( 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho \right) \right]}$$

Once  $\delta$  and  $\epsilon$  are fixed, we choose a positive constant  $\delta_1$  satisfying

$$\delta_1 < \min \{ \xi_1, \xi_2 \},$$

where

$$\xi_1 = \frac{g_0}{2(k+1) \left[ 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho + \delta_1 \tilde{A} \right]},$$

and

$$\xi_2 = \frac{g_0}{2(k+1) \left[ 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho + \delta_1 \tilde{A} \right]}.$$

and be such that

$$g_0 - \epsilon \left[ 1 + \frac{A}{4\delta} + \frac{k}{4\delta} \left( 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho \right) \right] - (k+1)\delta_1 \left[ 1 + \frac{C_*^{2(\rho+1)}}{\rho+1} \left( \frac{2pE(0)}{p-2} \right)^\rho + \delta_1 \tilde{A} \right] > 0,$$

also

$$\epsilon \left[ \frac{l}{2} - \delta C_*^2 - k\delta \left( 1 + \frac{C_*^2}{\rho+1} \right) \right] - \delta_1 \left[ 1 + 2(1-l)^2 + C_*^{2p-2} \left( \frac{2pE(0)}{(p-2)l} \right)^{p-2} \right] > 0.$$

We then pick  $M$  sufficiently large so that Lemma 4.2 holds and that

$$\left[ \frac{M}{2} - \frac{g(0)}{4\delta_1} \left( 1 + \frac{C_*^2}{\rho+1} \right) N \right] - \left[ \frac{\epsilon}{2l} + \frac{(\rho+2)k}{4\delta_1(\rho+1)} + 2\delta_1 + \frac{1}{2\delta_1} + \frac{C_*^2}{2\delta_1} \right] (1-l) > 0.$$

Hence, using  $(H_3)$ , we get

$$\begin{aligned} & \left[ \frac{M}{2} - \frac{g(0)}{4\delta_1} \left( 1 + \frac{C_*^2}{\rho+1} \right) N \right] (-g' \circ \nabla u)(t) \\ & - \left[ \frac{\epsilon}{2l} + \frac{(\rho+2)k}{4\delta_1(\rho+1)} + 2\delta_1 + \frac{1}{2\delta_1} + \frac{C_*^2}{2\delta_1} \right] (1-l)\xi(t)(g \circ \nabla u)(t) \geq k_3 \xi(t)(g \circ \nabla u)(t). \end{aligned}$$

By using Lemma 4.2 and (4.26), we arrive  $\forall t \geq t_0$  at

$$(4.27) \quad G'(t) \leq -\beta_1 \xi(t)E(t) \leq \alpha_1 \beta_1 \xi(t)G(t),$$

for some positive constant  $\beta_1$ . A simple integration of (4.27) leads to

$$(4.28) \quad G(t) \leq G(t_0)e^{-\alpha_1 \beta_1 \int_{t_0}^t \xi(s)ds}, \quad \forall t \geq t_0.$$

Thus, from Lemma 4.2 and (4.28), we get

$$(4.29) \quad E(t) \leq \alpha_2 G(t_0)e^{-\alpha_1 \beta_1 \int_{t_0}^t \xi(s)ds} = Ke^{-\kappa \int_{t_0}^t \xi(s)ds}, \quad \forall t \geq t_0.$$

This completes our proof.  $\square$



## REFERENCES

1. R. ABOULAICHA, D. MESKINEA and A. SOUISSIA: *New diffusion models in image processing*. Comput. Math. Appl. **56** (2008), 874–882.
2. F. ANDREU-VAILLO, V. CASELLES and JM. MAZN: *Parabolic Quasilinear Equations Minimizing Linear Growth Functions*. Progress in Mathematics, Birkhuser, Basel, 2004.
3. SN. ANTONTSEV and V. ZHIKOV: *Higher integrability for parabolic equations of  $p(x, t)$ -Laplacian type*. Adv. Differ. Equ. **10** (2005), 1053–1080.
4. SN. ANTONTSEV: *Wave equation with  $p(x, t)$ -Laplacian and damping term: blow-up of solutions. Existence and blow-up*. Differ. Equ. Appl. **3(4)** (2011), 503–525.
5. M M. CAVALCANTI, VN. D. CAVALCANTI and J. FERREIRA: *Existence and uniform decay for nonlinear viscoelastic equation with strong damping*. Math. Methods Appl. Sci. **24** (2001), 1043–1053.
6. M M. CAVALCANTI, VN. D. CAVALCANTI and JA. SORIANO: *Exponential decay for the solution of semilinear viscoelastic wave equations with localized damping*. Electron. J. Differ. Equ. **44** (2002), 1–14.
7. M M. CAVALCANTI and HP. OQUENDO: *Frictional versus viscoelastic damping in a semilinear wave equation*. SIAM J. Control Optim. **42(4)** (2003), 1310–1324.
8. Y. CHEN, S. LEVINE and M. RAO: *Variable exponent, linear growth functions in image restoration*. SIAM J. Appl. Math. **66** (2006), 1383–1406.
9. D. EDMUNDS and J. RAKOSNIK: *Sobolev embeddings with variable exponent*. Mathematische Nachrichten. **246(1)** (2002), 53–67.
10. X. FAN, J. SHEN and D. ZHAO: *Sobolev embedding theorems for spaces  $W^{k,p(x)}(\Omega)$* . J. Math. Anal. Appl. **262** (2001), 749–760.
11. X. FAN and D. ZHAO: *On the spaces  $L^{p(x)}$  and  $L^{m,p(x)}$* . J. Math. Anal. Appl. **263** (2001), 424–446.
12. Y. GAO, B. GUO and W. GAO: *Weak solutions for a high-order pseudo-parabolic equation with variable exponents*. Appl. Anal. (2013). doi:10.1080/00036811.2013.772138.
13. V.A. GALAKTIONOV and S.I. POHOZAEV: *Blow-up and critical exponents for nonlinear hyperbolic equations*. Nonlinear Analysis: Theory, Methods and Applications. **53(3)** (2003), 453–466.
14. V. GEORGIEV and G. TODOROVA: *Existence of solutions of the wave equation with nonlinear damping and source terms*. J. Diff. Eqns. **109 (2)** (1994), 295–308.
15. C. GOODRICH and M.A. RAGUSA: *Hölder continuity of weak solutions of  $p$ -Laplacian PDEs with VMO coefficients*. Nonlinear Analysis. **185** (2019), 336–355.
16. A. HARAUX and E. ZUAZUA: *Decay estimates for some semilinear damped hyperbolic problems*. Arch. Rational Mech. Anal. **150** (1988), 191–206.
17. A. M. KBIRI, S. A. MESSAOUDI and H. B. KHENOUS: *A blow-up result for nonlinear generalized heat equation*. Computers and Mathematics with Applications. **68(12)** (2014), 1723–1732.
18. O. KOVCIK and J. RAKOSNIK: *On spaces  $L^{p(x)}$  and  $W^{1,p(x)}$* . Czechoslov. Math. J. **41(116)** (1991), 592–618.

19. SZ. LIAN, WJ. GAO, CL. CAO and HJ. YUAN: *Study of the solutions to a model porous medium equation with variable exponents of nonlinearity*. J. Math. Anal. Appl. **342** (2008), 27–38.
20. JL. LIONS: *Quelques méthodes De résolution des Problèmes aux limites non linéaires*. Dunod, Paris, 1969.
21. W. LIU: *General decay and blow-up of solution for a quasilinear viscoelastic problem with nonlinear source*. Nonlinear Analysis. **73** (2010), 1890–1904.
22. S. A. MESSAOUDI and N. TATAR: *Global existence and uniform stability of solutions for a quasilinear viscoelastic problem*. Math. Meth. Appl. Sci. **30** (2007), 665–680.
23. M.A. RAGUSA and A. TACHIKAWA: *On continuity of minimizers for certain quadratic growth functionals*: Journal of the Mathematical Society of Japan. **57** (3) (2005) , 691–700.
24. M.A. RAGUSA and A. TACHIKAWA: *Regularity for minimizers for functionals of double phase with variable exponents*. Advances in Nonlinear Analysis. **9** (2020), 710–728.
25. E. VITILLARO: *Global nonexistence theorems for a class of evolution equations with dissipation*. Archive for Rational Mechanics and Analysis. **149**(2) (1999), 155–182.
26. JN. ZHAO: *Existence and nonexistence of solutions for  $u_t = \operatorname{div}(|\nabla u|^{p-2}\nabla u) + f(\nabla u, u, x, t)$* . J. Math. Anal. Appl. **172** (1993), 130–146.
27. SM. ZHENG: *Nonlinear Evolution Equation*. CRC Press, Boca Raton, 2004.

Fatima Zohra Mahdi  
 Laboratory ACEDP  
 Djillali Liabes university  
 22000 Sidi Bel Abbes, Algeria  
 mahdifatimazohra@yahoo.fr

Ali Hakem  
 Laboratory ACEDP  
 Djillali Liabes university  
 P.O. Box 73  
 22000 Sidi Bel Abbes, Algeria  
 hakemali@yahoo.com