

Type A Aortic Dissection, 5-year Operative Outcome Study

Thomas Perillo

Kaitlin Kelly

Caitlin Hoeing

James K. Wu MD

Follow this and additional works at: <https://scholarlyworks.lvhn.org/research-scholars-posters>



Part of the [Cardiology Commons](#)

This Poster is brought to you for free and open access by LVHN Scholarly Works. It has been accepted for inclusion in LVHN Scholarly Works by an authorized administrator. For more information, please contact LibraryServices@lvhn.org.

Type A Aortic Dissection, 5-year Operative Outcome Study

Thomas Perillo, Kaitlin Kelly, Caitlin Hoeing, James Wu, MD
 Division of Cardiothoracic Surgery
 Lehigh Valley Health Network, Allentown, Pennsylvania


Background

- An aortic dissection is a medical emergency, where a tear of the inner layer of the aorta causes blood accumulation between the inner/outer layers of the aorta
 - Type A dissection: occurrence in ascending aorta
 - Most common among older males
 - Common risk factors include hypertension, atherosclerosis, smoking, and structural heart problems
- Previous studies have found that rates of post operative complications and mortality are significantly lower when compared to those of dissections treated with medication alone

Objective

- Evaluate outcomes of type A aortic dissection repair via median sternotomy method

Methods



Results

Type A Dissection Patients N = 23		
History	N	%
Hypertension	6	26.09
Smoking (Tabacco)	8	34.78
Hyperlipidemia	6	26.09
Obesity	8	34.78
Family History	4	17.39
Atrial Fibrillation	8	34.78
Anemia	4	17.39
Operative Cannulation	N	%
Right Axillary Artery	14	60.97
Right Femoral Artery	6	26.09
Aorta	1	4.35
Right Atrium	2	8.69
Mortality	N	%
<30 Days Post Op	3	13.04
<90 Days Post Op	2	8.69
>90 Days Post Op	0	0

Table 1: Patient data for Type A aortic dissection patients and general procedural outcomes

- Median age at time of surgery: 64 years
- Percentage of male patients: 78.26%
- Mean BMI: 26.04 kg/m² [overweight]
- Overall mortality: 21.74% (N=5)

Outcomes




Figure 1: Adjusted Kaplan-Meier curve of Type A aortic dissection repair operations. A comparison to ascending aortic aneurysm repair is made, signifying the higher mortality rate of dissections

Conclusion and Future Direction

- Aortic dissection is a medical emergency, and repair surgery has a higher mortality rate compared to other aortic surgeries
- Instances of death occurred all within the first three months post operation
- More research into advanced surgical repair and preventative medicine would better improve outcomes of those with type A aortic dissection

References:

Cabasa, Alduz, and Alberto Pochettino. "Surgical Management and Outcomes of Type A Dissection—the Mayo Clinic Experience." *Annals of Cardiothoracic Surgery*, vol. 5, no. 4, 2016, pp. 296–309., doi:10.21037/acs.2016.06.01.

Harris, Christopher et al. "Type A aortic dissection." *Annals of cardiothoracic surgery* vol. 5,3 (2016): 256. doi:10.21037/acs.2016.05.04

Tsai, T. T. "Long-Term Survival in Patients Presenting With Type A Acute Aortic Dissection: Insights from the International Registry of Acute Aortic Dissection (IRAD)." *Circulation*, vol. 114, no. 1_suppl, 2006, doi:10.1161/circulationaha.105.000497.