

## Some Vector $FK$ Sequence Spaces Generated by Modulus Function

E. Herawati<sup>1\*</sup>, N. Irsyad<sup>2</sup> and E. Rosmaini<sup>1</sup>

<sup>1</sup>Department of Mathematics, Universitas Sumatera Utara, Medan, 20155, Indonesia

<sup>2</sup>School of Mathematical Sciences, Universiti Sains Malaysia, Penang, 11800, Malaysia

**Abstract.** In this paper, some vector valued sequence spaces  $\Gamma_f(X)$  and  $\Lambda_f(X)$  using modulus function are presented. Furthermore, we examined some topological properties of these sequence spaces equipped with a paranorm.

**Keywords:** Modulus function, Paranorm, Vector valued sequence space.

**Abstrak.** Pada paper ini, diperkenalkan beberapa ruang barisan bernilai vektor  $\Gamma_f(X)$  dan  $\Lambda_f(X)$  menggunakan fungsi modulus. Lebih lanjut, dipelajari beberapa sifat-sifat topologi dari ruang-ruang barisan ini dikenakan suatu paranorma tertentu.

**Kata Kunci:** Fungsi modulus, Paranorma, Ruang barisan bernilai vektor.

Received 06 August 2020 | Revised 01 September 2020 | Accepted 30 September 2020

### 1 Introduction

Let  $X$  be a vector space and  $\mathbb{R}$  be the set of real numbers. A function  $f : \mathbb{R}^+ \cup \{0\} \rightarrow \mathbb{R}^+ \cup \{0\}$  is called modulus function if following condition of  $f$  satisfying:

1.  $f$  is vanishing at zero
2.  $f$  satisfies triangle inequality
3.  $f$  is an increasing function i.e.  $f(\cdot) \uparrow$
4.  $f$  is a continuous function from the right at 0 [1]

The function  $f$  must be continuous for every element  $x$  in  $(0, \infty)$ . The space of all real number sequences  $(x_n)$  such that the infinite series of absolute modulus function is finite denoted by  $\ell(f)$  [2]

$$\sum_{n=1}^{\infty} f(|x_n|) < \infty.$$

The space  $\ell(f)$  becomes a  $FK$ -space under the  $F$ -norm

$$p(x) = \sum_{n=1}^{\infty} f(|x_n|) < \infty.$$

\*Corresponding author at: Department of Mathematics, Universitas Sumatera Utara, Medan, 20155, Indonesia

E-mail address: elvina@usu.ac.id

Adnan [3] examined the  $FK$ -space properties of an analytic and entire real sequence space using modulus function. He showed the characterization to matrix transformation of Ruckle's space  $\ell(f)$  into analytic  $FK$ -space. For the theory of  $FK$ -space we refer to Banas and Mursaleen [4].

Through the article  $\Omega(X)$ ,  $\Gamma_f(X)$ ,  $\Lambda_f(X)$  denoted by the space of vector value sequences, entire vector value sequence space and analytic vector value sequence space. The vector value sequence space studied by some authors [5, 6, 7, 8, 9, 10, 11, 12, 13]. Further, the concept of sequence space using modulus function was investigated by [14, 15, 16, 17, 18].

Recently, Herawati [5] studied the geometric of the vector value sequence spaces defined by order- $\varphi$  function under Lattice norm. Further, Gultom [6] studied some topologies properties of a finite arithmetic mean vector value sequence space denoted by  $W_f(X)$  for  $X$  is a linear space and  $f$  is a  $\varphi$ -function.

A functional is called paranormed if satisfies the properties  $p : X \rightarrow \mathbb{R}$  that satisfies the properties  $p(\theta) = 0$ , with  $\theta$  is the zero vector in  $X$ , non-negative,  $p$  satisfies triangle inequalities, even and every real sequence  $(\lambda_n)$  with  $|\lambda_n - \lambda| \rightarrow 0$ . The space  $X$  with paranorm  $p$  is called *paranormed space*, written as  $X = (X, p)$  [1, 19].

In this work, we define the space of vector value sequences  $\Gamma_f(X)$  and  $\Lambda_f(X)$  called entire and analytic vector valued sequence spaces generated by modulus function and study the topological properties of the sets equipped with paranorm.

## 2 Main Results

In this main result section, firstly, we introduce paranorm on this space and examine some topological properties such as complete properties. Let  $X$  be a Banach space and  $f$  be a modulus function. Let  $y(n) = f(\|x(n)\|_X) \in \mathbb{R}$  for all natural numbers  $n$ , then we get a sequence  $y = (y(n))$ . We define the sets

$$\begin{aligned}\Gamma_f(X) &= \left\{ x = (x(n))_{n \in \mathbb{N}} : x(n) \in X \text{ and } (y(n))^{\frac{1}{n}} \rightarrow 0, n \rightarrow \infty \right\} \\ \Lambda_f(X) &= \left\{ x = (x(n))_{n \in \mathbb{N}} : x(n) \in X \text{ and } \sup_{n \in \mathbb{N}} \{(y(n))^{\frac{1}{n}}\} < \infty \right\}\end{aligned}$$

### Theorem 1.

The sets  $\Gamma_f(X)$  and  $\Lambda_f(X)$  are vector spaces.

#### Proof.

Let  $x, z$  be any elements in  $\Gamma_f(X)$ , then

$$\lim_{n \rightarrow \infty} (y(n))^{\frac{1}{n}} = 0 \text{ and } \lim_{n \rightarrow \infty} (w(n))^{\frac{1}{n}} = 0$$

for  $n \rightarrow \infty$ , with  $y(n) = f(x(n))$  and  $w(n) = f(z(n))$  for each natural number  $n$ . We will apply the following inequality : if  $a_n, b_n \in \mathbb{R}$  and  $0 \leq q_n \leq \sup q_n = H$  for each natural number  $n$ , then

$$|a_n + b_n|^{q_n} \leq M(|a_n|^{q_n}) + |b_n|^{q_n}$$

where  $M = \max\{1, 2^{H-1}\}$ . Therefore,

$$(y(n) + w(n))^{\frac{1}{n}} \leq (y(n))^{\frac{1}{n}} + (w(n))^{\frac{1}{n}}$$

Since  $(q_n) = (\frac{1}{n})$ , then  $H = \sup \frac{1}{n} = 1$ . Thus

$$(y(n) + w(n))^{\frac{1}{n}} \leq (y(n))^{\frac{1}{n}} + (w(n))^{\frac{1}{n}}$$

Since  $(y(n))^{\frac{1}{n}} \rightarrow 0$  and  $(w(n))^{\frac{1}{n}}$  for  $n \rightarrow \infty$ , then  $(y(n) + w(n))^{\frac{1}{n}} \rightarrow 0$  for  $n \rightarrow \infty$ . Therefore, we obtain  $x + y \in \Gamma_f(X)$ . Further, for element  $x \in \Gamma_f(X)$  and  $\alpha \in \mathbb{R}$ , then

$$(y(n))^{\frac{1}{n}} \rightarrow 0, n \rightarrow \infty$$

Because of an increasing function  $f$  and the positivity of  $|\alpha|$ , then from the Archimedean properties, there exists natural number  $n_0$  with

$$f(|\alpha||x(n)|) \leq f(2^{n_0}|x(n)|)$$

Since  $f$  satisfies  $\Delta_2$ -condition, we get

$$(f(2^{n_0}|x(n)|))^{\frac{1}{n}} = K^{\frac{n_0}{n}} (f(|x(n)|))^{\frac{1}{n}} \rightarrow 0$$

for each natural number  $n$ . It shows that  $\alpha x \in \Gamma_f(X)$ . Because  $x + z \in \Gamma_f(X)$  and  $\alpha x \in \Gamma_f(X)$  for each  $x, y \in \Gamma_f(X)$  and each  $\alpha \in \mathbb{R}$ , we get  $\Gamma_f(X)$  is a vector or linear space and the proof of the theorem is finished. In the same way, it can be shown that  $\Lambda_f(X)$  is a vector space. ■

### Theorem 2.

A functional  $p : \Gamma_f(X) \rightarrow \mathbb{R}$  defined by

$$p(x) = \sup_{n \geq 1} \left\{ (y(n))^{\frac{1}{n}} \right\}$$

is a paranorm.

#### Proof.

Let  $x$  be an element in  $\Gamma_f(X)$ . It is clear that the functional  $p$  is non-negative,  $p(\theta) = 0$ , with  $\theta$  is the zero vector in  $X$  and even, for each  $x \in \Gamma_f(X)$ . Now, we will show that  $p$  satisfies the triangle inequality. To do that, take any  $x, z \in \Gamma_f(X)$ , then

$$\lim_{n \rightarrow \infty} (y(n))^{\frac{1}{n}} = 0 \text{ and } \lim_{n \rightarrow \infty} (w(n))^{\frac{1}{n}} = 0$$

for  $n \rightarrow \infty$ , with  $y(n) = f(x(n))$  and  $w(n) = f(z(n))$  for each  $n \in \mathbb{N}$ . we obtain

$$\sup \left\{ (y(n) + w(n))^{\frac{1}{n}} \right\} \leq \sup \left\{ (y(n))^{\frac{1}{n}} + (w(n))^{\frac{1}{n}} \right\}$$

Therefore, there's vector sequences of  $x, y \in \Gamma_f(X)$ , we get  $p$  satisfies the triangle inequality. Next, we will show that  $p$  satisfies the continuity of scalar multiplication. To do that, take any real

sequence  $(\lambda_n)$  and  $(x(n)) \in \Gamma_f(X)$  with  $|\lambda_n - \lambda| \rightarrow 0$  for  $n \in \infty$ . We have

$$\begin{aligned} (f(\|x(n)\|_X))^{\frac{1}{n}} &= (f(\|\lambda_n x(n) - \lambda x(n)\|))^{\frac{1}{n}} \\ &= (f(\|(\lambda_n - \lambda)x(n)\| + \|\lambda(x(n) - x)\|))^{\frac{1}{n}} \\ &\leq ((f|\lambda_n - \lambda|\|x(n)\| + |\lambda|\|(x(n) - x)\|))^{\frac{1}{n}} \end{aligned}$$

and

$$\begin{aligned} p(\lambda_n x(n) - \lambda x(n)) &= \sup \{ (f(\|\lambda_n x(n) - \lambda x(n)\|))^{\frac{1}{n}} \} \\ &\leq |\lambda_n - \lambda| p(x(n)) + |\lambda| p(x(n) - x) \rightarrow 0 \end{aligned}$$

Hence,  $p(\lambda_n x(n) - \lambda x(n)) \rightarrow 0$ . The proof of the theorem is finished.  $\blacksquare$

### Theorem 3.

The vector spaces of  $\Gamma_f(X)$  and  $\Lambda_f(X)$  are complete paranormed sequence space under the paranorm defined in Theorem 2.

#### Proof.

Take any Cauchy sequence  $(x^i)$  in  $\Gamma_f(X)$  with  $x^i = (x^i(n)) = (x^i(1), x^i(2), \dots)$ . Therefore, for any positive real number  $\varepsilon$ , there exists  $i_0 \in \mathbb{N}$ , for all  $j \geq i \geq i_0$ , we get

$$p(x^j - x^i) = \sup \left\{ (f(\|x^j(n) - x^i(n)\|))^{\frac{1}{n}} \right\} < \varepsilon$$

Since  $\sup (f(\|x^j(n) - x^i(n)\|))^{\frac{1}{n}} < \varepsilon$ , we have  $(f(\|x^j(n) - x^i(n)\|))^{\frac{1}{n}} < \varepsilon$  for  $\varepsilon > 0$ . Since  $f$  is a modulus function, then  $\|x^j(n) - x^i(n)\| = 0$  for each natural number  $n$ . In other words,  $\|x^j(n) - x^i(n)\| < \varepsilon$ . It shows that for each natural number  $n$  of the sequence  $(x^j(n))$  is a Cauchy. Since  $X$  is a complete normed space, then  $(x^j(n))$  converges to  $x(n) \in X$ . Hence,  $\lim_{j \rightarrow \infty} x^j(n) = x(n)$  for all  $n$ . Therefore, there's sequence  $x = (x(n)) = (x(1), x(2), \dots)$  such that

$$\begin{aligned} \sup \left\{ (f(\|x - x^i\|))^{\frac{1}{n}} \right\} &= \sup \left\{ (f(\|\lim_{i \rightarrow \infty} x - x^i\|))^{\frac{1}{n}} \right\} \\ &= \sup \left\{ \lim_{i \rightarrow \infty} (f(\|x - x^i\|))^{\frac{1}{n}} \right\} \\ &= \lim_{i \rightarrow \infty} \sup \left\{ (f(\|x - x^i\|))^{\frac{1}{n}} \right\} \end{aligned}$$

for every  $i \geq i_0$ . By using the definition of paranorm, we get

$$p(x - x^i) = \sup \left\{ (f(\|x - x^i\|))^{\frac{1}{n}} \right\} < \varepsilon$$

It shows that  $x^i \rightarrow x$  for  $i \rightarrow \infty$ . Then it will be shown that  $x \in \Gamma_f(X)$ . Using the continuous

property of  $f$ , we get

$$\begin{aligned} (f(\|x\|))^{\frac{1}{n}} &= (f(\|\lim_{i \rightarrow \infty} x^i\|))^{\frac{1}{n}} \\ &= \lim_{i \rightarrow \infty} (f(\|x^i\|))^{\frac{1}{n}} \rightarrow 0 \end{aligned}$$

for  $i \rightarrow \infty$ . Hence,  $x \in \Gamma_f(X)$ . The proof of this theorem is finished.  $\blacksquare$

### 3 Conclusions

According to the main results, it can be concluded  $\Gamma_f(X)$  and  $\Lambda_f(X)$  are complete paranormed sequence space under the paranorm.

### REFERENCES

---

- [1] H. Nakano, "Modulared sequence spaces," *Proc. Japan Acad.*, vol. 27, no. 9, pp. 508–512, 1951.
- [2] W. H. Ruckle, "Fk spaces in which the sequence of coordinate vectors is bounded," *Canadian Journal of Mathematics*, vol. 25, no. 5, pp. 973–978, 1973.
- [3] A. Alhomaidan, "Some fk spaces defined by a modulus function," *International Journal of Pure and Applied Mathematics*, vol. 30, no. 1, pp. 43–48, 2006.
- [4] J. Banas and M. Mursaleen, *Sequence Spaces and Measures of Noncompactness with Applications to Differential and Integral Equations*. Springer India, 2014.
- [5] E. Herawati, Supama, and M. Mursaleen, "Local structure of riesz valued sequence spaces defined by an order  $\phi$ -function," *Linear and Multilinear Algebra*, vol. 65, no. 3, pp. 545–554, 2017.
- [6] S. N. R. Gultom, R. Siregar, and E. Herawati, "Köthe-töeplitz duals of vector valued sequence spaces defined by  $\phi$ -function," *Journal of Physics: Conference Series*, vol. 1116, no. 2, pp. 1–6, 2018.
- [7] E. Kolk, "Topologies in generalized orlicz sequence spaces," *Filomat*, vol. 25, no. 4, pp. 191–211, 2011.
- [8] I. E. Leonard, "Banach sequence spaces," *Journal of Mathematical Analysis and Applications*, vol. 54, no. 1, pp. 245–265, 1976.
- [9] N. R. Das and A. Choudhury, "Matrix transformation of vector valued sequence space," *Bulletin of the Calcutta Mathematical Society*, vol. 84, no. 1, pp. 47–54, 1992.
- [10] M. Et, "Spaces of cesáro difference sequences of order  $r$  defined by a modulus function in a locally convex space," *Taiwanese Journal of Mathematics*, vol. 10, no. 4, pp. 865–879, 2006.
- [11] M. Et, A. Gökhan, and H. Altinok, "On statistical convergence of vector-valued sequences associated with multiplier sequences," *Ukrainian Mathematical Journal*, vol. 58, no. 139, 2006.

- [12] B. C. Tripathy and M. Sen, "Vector valued paranormed bounded and null sequence spaces associated with multiplier sequences," *Soochow Journal of Mathematics*, vol. 29, no. 3, pp. 313–326, 2003.
- [13] B. C. Tripathy and S. Mahanta, "On a class of vector-valued sequences associated with multiplier sequences," *Acta Mathematicae Applicatae Sinica*, vol. 20, no. 4, pp. 487–494, 2004.
- [14] T. Bilgin, "The sequence space  $l(p, f, q, s)$  on seminormed spaces," *Bull. Calcutta Math. Soc.*, vol. 86, no. 4, pp. 295–304, 1994.
- [15] S. Pehlivan and B. Fisher, "On some sequence spaces," *Indian Journal of Pure and Applied Mathematics*, vol. 25, no. 10, pp. 1067–1071, 1994.
- [16] A. Waszak, "On the strong convergence in some sequence spaces," *Fasciculi Mathematici*, vol. Nr 33, pp. 125–137, 2002.
- [17] V. K. Bhardwaj, "A generalization of a sequence space of ruckle," *Bull. Calcutta Math. Soc.*, vol. 95, no. 5, pp. 411–420, 2003.
- [18] Y. Altin, "Properties of some sets of sequences defined by a modulus function," *Acta Mathematica Scientia Series B English Edition*, vol. 29, no. 2, pp. 427–434, 2009.
- [19] S. Simons, "The sequence spaces  $l(p_v)$  and  $m(p_v)$ ," *Proceedings of the London Mathematical Society*, vol. s3-15, no. 1, pp. 422–436, 1965.