



ELSEVIER

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

The Journal of Arthroplasty

journal homepage: www.arthroplastyjournal.org

Complications - Infection

Tranexamic Acid Is Associated With Reduced Periprosthetic Joint Infection After Primary Total Joint Arthroplasty



Hamidreza Yazdi, MD ^{a, b}, Mitchell R. Klement, MD ^a, Mohammed Hammad ^a,
Daisuke Inoue, MD ^{a, c}, Chi Xu, MD ^{a, d}, Karan Goswami, MD ^a,
Javad Parvizi, MD, FRCS ^{a, *}

^a Rothman Orthopaedic Institute at Thomas Jefferson University, Philadelphia, PA^b Department of Orthopaedic Surgery, Iran University of Medical Sciences, Tehran, Iran^c Department of Orthopaedic Surgery, Graduate School of Medical Science, Kanazawa University, Kanazawa, Ishikawa, Japan^d Department of Orthopaedic Surgery, General Hospital of People's Liberation Army, Beijing, People's Republic of China

ARTICLE INFO

Article history:

Received 9 July 2019

Received in revised form

16 October 2019

Accepted 16 October 2019

Available online 22 October 2019

Keywords:

tranexamic acid

periprosthetic joint infection

anemia

primary joint arthroplasty

complications

ABSTRACT

Background: Previous studies have demonstrated preoperative anemia to be a strong risk factor for periprosthetic joint infection (PJI) in total joint arthroplasty (TJA). Allogeneic blood transfusion can be associated with increased risk of PJI after primary and revision TJA. Tranexamic acid (TXA) is known to reduce blood loss and the need for allogeneic blood transfusion after TJA. The hypothesis of this study is that administration of intravenous TXA would result in a reduction in PJI after TJA.

Methods: An institutional database was utilized to identify 6340 patients undergoing primary TJA between January 1, 2013 and June 31, 2017 with a minimum of 1-year follow-up. Patients were divided into 2 groups based on whether they received intravenous TXA prior to TJA or not. Patients who developed PJI were identified. All PJI patients met the 2018 International Consensus Meeting definition for PJI. A multivariate regression analysis was performed to identify variables independently associated with PJI.

Results: Of the patients included, 3683 (58.1%) received TXA and 2657 (41.9%) did not. The overall incidence of preoperative anemia was 16%, postoperative blood transfusion 1.8%, and PJI 2.4%. Bivariate analysis showed that patients who received TXA were significantly at lower odds of infection. After adjusting for all confounding variables, multivariate regression analysis showed that TXA is associated with reduced PJI after primary TJA.

Conclusion: TXA can help reduce the rate of PJI after primary TJA. This protective effect is likely interlinked to reduction in blood loss, lower need for allogeneic blood transfusion, and issues related to immunomodulation associated with blood transfusion.

© 2019 Elsevier Inc. All rights reserved.

Periprosthetic joint infection (PJI) is a rare, yet a devastating, complication of total joint arthroplasty (TJA). It occurs in 1%–2% of TJA procedures [1–3] and is one of the most common reasons for TJA revision [4–10].

By 2030, primary TJA has been projected to grow by 174% to 572,000 for primary total hip arthroplasty (THA) and by 673% to 3.48 million for primary total knee arthroplasty (TKA) [11]. Failures

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may include receipt of payment, either direct or indirect, institutional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2019.10.029>.

* Reprint requests: Javad Parvizi, MD, FRCS, Rothman Orthopaedic Institute, 125 S 9th St, Ste 1000, Philadelphia, PA 19107.

<https://doi.org/10.1016/j.arth.2019.10.029>

0883-5403/© 2019 Elsevier Inc. All rights reserved.

of TJA because of complications such as infection, loosening, periprosthetic fracture, and postoperative pain are indications for revision surgery [12]. These revisions will become a tremendous burden to the health systems, with a projected estimate cost of \$13 billion in 2030 [12]. Given the increased demands for arthroplasty, patients should be perioperatively optimized to minimize the incidence of PJI, and other complications, following TJA.

Preoperative anemia and blood loss during primary TJA leading to allogeneic blood transfusion have been identified as risk factors for surgical site infection (SSI) and PJI in TJA [13–16]. This has led to a growing increase in the use of antifibrinolytic drugs to reduce blood loss and blood transfusion post TJA [17]. The results from previous studies and extensive meta-analysis have established that tranexamic acid (TXA) is a safe and effective drug for reducing

blood loss and the need for allogeneic blood transfusion post-TJA [17–19].

Although efficacy of TXA in reducing blood loss and the need for allogeneic blood transfusion has been well established, the potential role of TXA in reducing PJI following TJA has not been elucidated. This study was conceived to examine the hypothesis that administration of intravenous (IV) TXA can lead to a reduction in PJI after primary TJA.

Materials and Methods

Following Institutional Review Board approval, a prospectively maintained, single-institution database was queried to identify 7725 patients undergoing primary TJA between January 1, 2013 and June 31, 2017. All patients who had primary THA or TKA were included. For patients who had multiple joints replacement, each joint was considered as a separate case. Patients undergoing revision arthroplasty, primary arthroplasty in tumor cases, and those with inadequate follow-up (1012) were excluded (Fig. 1).

Overall, 6340 cases were included in the study with minimum 1-year follow-up. Of these, 2927 (46.2%) were male. The mean age was 64.72 ± 10.7 years (range 12–98) and the mean body mass index (BMI) was 29.8 ± 5.3 kg/m² (range 14–58.3) (Table 1). The mean follow-up was 26.3 months (range 12–67.3).

The cohort was divided into 2 groups: patients who received IV TXA prior to arthroplasty (3683 patients) and those who did not receive TXA (2657 patients). Patients in the TXA group were all administered a single dose of TXA (15 mg/kg) intravenously, 20–30 minutes prior to incision or inflation of tourniquet. The antibiotic and postoperative care was the same for both patient groups. The majority of patients (91.8%) in the study received aspirin (325 mg/d or 81 mg bid) for 4 weeks postoperatively. Using the World

Table 1
Patient Demographics.

Gender (%)	
Male	2927 (46.2)
Female	3413 (53.8)
Age (y), mean \pm SD (range)	64.72 \pm 10.7 (12–98)
Race (%)	
Caucasian	4753 (75)
African American	1048 (16.5)
Hispanics	77 (1.2)
Asians	35 (1.2)
Others	387 (6.1)
BMI (kg/m ²), mean \pm SD (range)	29.8 \pm 5.3 (14–58.3)
Type of surgery (%)	
TKA	3028 (47.8)
THA	3312 (52.2)
TXA (%)	
Yes	3683 (58.1)
No	2657 (41.9)
Anemic (%)	
Yes	971 (16)
No	5090 (84)

SD, standard deviation; BMI, body mass index; TKA, total knee arthroplasty; THA, total hip arthroplasty; TXA, tranexamic acid.

Health Organization definition, preoperative anemia was defined as hemoglobin (Hgb) <13 g/dL in males and Hgb <12 g/dL in females. Patient demographics, comorbidities, operative details, and perioperative variables were recorded and compared across the 2 cohorts.

Patients who developed PJI following primary TJA were identified using the 2018 International Consensus Meeting definition [20] and subsequently confirmed by manual chart review.

Statistical Analyses

The 2 groups (TXA vs no TXA) were compared on categorical variables with Fisher's exact tests and on continuous variables with *t*-tests for independent groups or Mann-Whitney *U* for nonparametric variants. Preliminary bivariate logistic regression analyses were conducted to determine whether various preoperative and patient variables were significantly related to PJI outcomes. Variables which were related to PJI in the bivariate analyses ($P < .25$ as recommended by Hosmer and Lemeshow, 2000) were considered for logistic regression model. All analyses were conducted with the statistical packages SPSS (IBM Corp, 2013, version 23, Armonk, NY).

Results

Overall, 3683 patients (58.1%) received IV preoperative TXA. The overall incidence of preoperative anemia was 16%, postoperative blood transfusion was 1.8%, and PJI rate was 2.4%. Most of TJA surgeries were done under neuraxial anesthesia (91.7%) and nearly all (96.6%) of total knee replacements were done under tourniquet.

The preliminary analyses showed that patients receiving TXA were younger, more likely to be female, had a lower BMI and comorbidities, and undergoing THA. This group was also more likely to receive aspirin (for deep vein thrombosis [DVT] prophylaxis) and had less anemia (all $P < .05$; Table 2). The outcome measurement showed that the TXA group has a smaller postoperative Hgb decrease, less need for allogeneic blood transfusion, shorter length of hospitalization, less wound complications, and lower PJI rate (all $P < .05$; Table 3). Intraoperative blood loss, 90-day readmission, and postoperative thromboembolic events were similar in both groups.

With regard to PJI rate, bivariate analyses (Table 4) showed that rheumatoid arthritis (odd ratio [OR] 5.07, 95% confidence interval




Fig. 1. Flow diagram detailing patient inclusion, exclusion, and grouping. TJA, total joint arthroplasty; TXA, tranexamic acid.

Table 2
Univariate Analysis.

Variables	Total (n = 6340)	Non-TXA (n = 2657)	TXA (n = 3683)	P-Value
Age (y), mean ± SD	64.7 ± 10.7	66.9 ± 10.5	63.1 ± 10.5	<.001
Gender, female (%)	3413 (53.8)	1306 (49.2)	2107 (57.2)	<.001
BMI (kg/m ²), mean ± SD	29.8 ± 5.3	30.0 ± 5.3	29.7 ± 5.3	.012
Race, Caucasian (%)	4753 (75.0)	1912 (72.0)	2841 (77.1%)	<.001
Joint, knee (%)	3028 (47.8)	1369 (51.5)	1659 (45.0)	<.001
CCI, mean ± SD	0.5 ± 1.0	0.8 ± 1.2	0.4 ± 0.8	<.001
CCI ≥ 1 (%)	2034 (32.1)	1111 (41.8)	923 (25.1)	<.001
Diabetes (%)	379 (6.0)	190 (7.2)	189 (5.1)	<.001
Malignancy (%)	68 (1.1)	37 (1.4)	31 (0.8)	.036
Renal disease (%)	188 (3.0)	140 (5.3)	48 (1.3)	<.001
Liver disease (%)	49 (0.8)	24 (0.9)	25 (0.7)	.314
Rheumatic arthritis (%)	284 (4.5)	140 (5.3)	144 (3.9)	.01
Preoperative hemoglobin (g/L), mean ± SD	13.3 ± 1.6	13.1 ± 1.8	13.5 ± 1.5	<.001
Anemia (%)	971 (16.0)	510 (20.9)	461 (12.7)	<.001
Coagulopathy (%)	106 (1.7)	81 (3.1)	25 (0.7)	<.001
Simultaneous bilateral TJA (%)	336 (5.3)	80 (3.0)	256 (7.0)	<.001
Neuraxial anesthesia (%)	5812 (91.7)	2365 (89.2)	3447 (93.6)	<.001
Tourniquet (%)	2925 (96.6)	1322 (96.6)	1603 (96.6)	.93
Anticoagulation				<.001
Aspirin (%)	5813 (91.8)	2304 (87.0)	3509 (95.3)	
Warfarin (%)	330 (5.2)	255 (9.6)	75 (2.0)	
Others	188 (3.0)	89 (3.4)	99 (2.7)	
Intraoperative antibiotics				.601
Cephalosporin (%)	5224 (82.5)	2182 (82.5)	3042 (82.6)	
Vancomycin (%)	1082 (17.1)	452 (17.1)	630 (17.1)	
Others (%)	23 (0.4)	12 (0.5)	11 (0.3)	
Operative time (min), mean ± SD	101.5 ± 39.8	102.5 ± 40.5	100.9 ± 39.3	.111

TXA, tranexamic acid; SD, standard deviation; BMI, body mass index; CCI, Charlson comorbidity index; TJA, total joint arthroplasty.

[CI] 3.28–7.83, $P = .000$), higher BMI (OR 1.06, 95% CI 1.03–1.09, $P = .000$), higher Charlson Comorbidity Index (OR 1.27, 95% CI 1.13–1.43, $P = .000$), male gender (OR 1.58, 95% CI 1.14–2.20, $P = .005$), renal disease (OR 3.86, 95% CI 2.22–6.73, $P = .000$), liver disease (OR 3.74, 95% CI 1.33–10.54, $P = .01$), coagulopathies (OR 2.54, 95% CI 1.09–5.88, $P = .03$), longer length of hospital stay (OR 1.18, 95% CI 1.13–1.23, $P = .000$), longer operation time (OR 1.01, 95% CI 1.01–1.02, $P = .000$), and anemia (OR 4.63, 95% CI 3.31–6.46, $P = .000$) were associated with higher odds of PJI. Administration of TXA (OR 0.47, 95% CI 0.34–0.66, $P = .000$), neuraxial anesthesia (OR 0.32, 95% CI 0.21–0.48, $P = .000$), and simultaneous bilateral surgeries (OR 0.24, 95% CI 0.06–0.96, $P = .04$) were associated with lower odds of infection.

After controlling for confounding variables, multivariate analysis showed that administration of TXA (OR 0.68, 95% CI 0.46–0.99,

Table 3
Outcome Measurements.

Variables	Non-TXA (n = 2657)	TXA (n = 3683)	P-Value
IOBL (mL), mean ± SD	147.8 ± 240.3	147.7 ± 159.5	.99
Hemoglobin drop (g/L), mean ± SD	2.3 ± 1.8	2.0 ± 1.6	.000
Postoperative blood transfusion (%)	75 (2.8)	41 (1.1)	.000
Length of stay (d), mean ± SD	2.4 ± 2.5	1.8 ± 1.4	.000
Wound complication (%)	35 (1.5)	15 (0.4)	.000
DVT (%)	0 (0.0)	4 (0.1)	.16
PE (%)	5 (0.2)	2 (0.1)	.120
PJI (%)	90 (3.4)	60 (1.6)	.000
Type of PJI (%)			.68
Acute	27 (30.0)	16 (26.7)	
Chronic	62 (68.9)	42 (70.0)	
Hematogenous	1 (1.1)	2 (3.3)	
90-d readmission (%)	120 (4.5)	155 (4.2)	.55

TXA, tranexamic acid; IOBL, intraoperative blood loss; SD, standard deviation; DVT, deep vein thrombosis; PE, pulmonary embolism; PJI, periprosthetic joint infection.

$P = .04$) was associated with lower rate of PJI. TXA also was more effective in nonanemic patients (OR 0.52, 95% CI 0.32–0.84, $P = .008$) and patients undergoing hip arthroplasty (OR 0.50, 95% CI 0.31–0.81, $P = .005$) than anemic patients and those undergoing knee arthroplasty (Table 5).

Discussion

The association between preoperative anemia and SSI or PJI has been demonstrated in numerous prior studies [13–16,21–24]. Patients with preoperative anemia have higher likelihood of needing

Table 4
Bivariate Predictors of PJI via Logistic Regression.

Predictor	OR (95% CI)	P-Value
TXA	0.47 (0.34–0.66)	.000
Age (y)	1.00 (0.99–1.02)	.95
Male	1.58 (1.14–2.20)	.005
BMI (kg/m ²)	1.06 (1.03–1.09)	.000
Race, not Caucasians	0.71 (0.47–1.07)	.10
Knee	0.77 (0.55–1.06)	.11
CCI scores	1.27 (1.13–1.43)	.000
Diabetes	1.00 (0.51–1.99)	.99
Malignancy	1.25 (0.30–5.17)	.75
Renal disease	3.86 (2.22–6.73)	.000
Liver disease	3.74 (1.33–10.54)	.01
Rheumatic arthritis	5.07 (3.28–7.83)	.000
Anemia	4.63 (3.31–6.46)	.000
Coagulopathy	2.54 (1.09–5.88)	.03
Simultaneous bilateral TJA	0.24 (0.06–0.96)	.04
Neuraxial anesthesia	0.32 (0.21–0.48)	.000
Warfarin	2.20 (1.54–3.15)	.000
Vancomycin	2.20 (1.54–3.15)	.000
Operative time (min)	1.01 (1.01–1.02)	.000
Length of stay (d)	1.18 (1.13–1.23)	.000

PJI, periprosthetic joint infection; OR, odds ratio; CI, confidence interval; TXA, tranexamic acid; BMI, body mass index; CCI, Charlson comorbidity index; TJA, total joint arthroplasty.

Table 5
Multivariate Analysis for Association Between TXA and PJI.

Variables	PJI Rate	Nonadjusted OR (95% CI)	P-Value	Adjust OR (95% CI)	P-Value
Overall					
Without TXA	3.4%	Reference		Reference	
With TXA	1.6%	0.47 (0.34–0.66)	<.001	0.68 (0.46–0.99)	.045
With anemia					
Without TXA	7.8%	Reference		Reference	
With TXA	5.6%	0.70 (0.42–1.17)	.175	1.14 (0.60–2.17)	.683
No anemia					
Without TXA	2.3%	Reference		Reference	
With TXA	1.1%	0.46 (0.29–0.71)	.001	0.52 (0.32–0.84)	.008
Hip					
Without TXA	4.2%	Reference		Reference	
With TXA	1.7%	0.39 (0.25–0.60)	<.001	0.50 (0.31–0.81)	.005
Knee					
Without TXA	2.6%	Reference		Reference	
With TXA	1.6%	0.59 (0.35–0.98)	.042	1.18 (0.63–2.23)	.600

TXA, tranexamic acid; PJI, periprosthetic joint infection; OR, odds ratio; CI, confidence interval.

allogeneic blood transfusions and are at higher risk of developing PJI [13–16]. Thus, strategies to optimize anemic patients preoperatively and minimize blood loss during surgery have been implemented to reduce the need for allogeneic blood transfusion [25]. One such strategy has been the administration of TXA during TJA with proven efficacy to reduce blood loss and the need for allogeneic blood transfusion [17–19,26–30]. The current study corroborates the findings of prior studies by demonstrating an excellent efficacy for TXA. Patients in the current study who received TXA had less Hgb drop postoperatively and were less likely to require blood transfusion and had lower rate of postoperative wound complications. This group also had shorter length of hospital stay and lower rate of PJI.

A recent study demonstrated additional benefits to TXA when administered to patients undergoing TKA [31]. Patients receiving TXA had less hidden blood loss, a lower ratio of postoperative knee swelling, less postoperative knee pain, lower levels of inflammatory biomarkers, better early knee function, and better satisfaction than those treated with tourniquet alone [32]. Administration of TXA is also safe with no real concerns that can be gleaned from the literature. TXA has also been exonerated from causing venous thromboembolism (VTE). A recently published meta-analysis concluded that administration of topical, IV, and oral TXA was not associated with an increased risk of VTE [33]. We also found no differences between both groups in terms of DVT, pulmonary embolism, and 90-day readmissions.

Most of the patients who received TXA were also placed on aspirin for VTE prophylaxis. Heller et al [34] have demonstrated that TXA can be used along with aspirin safely in arthroplasty patients without increasing the risk of subsequent VTE. Our results also showed that this combination did not affect the rate of DVT, pulmonary embolism, and readmissions. In addition, a study by Kim et al [35] detected that administration of intra-articular TXA resulted in fewer wound complications compared to patients who did not receive TXA. We also observed a similar finding.

The confounding variable that influences blood loss during TKA is the issue of tourniquet. Alcelik et al [36] in a systemic review concluded that the total and intraoperative blood loss was less when a tourniquet was used. On the other hand, Yi et al [37] published a meta-analysis demonstrating that the use of tourniquet during TKA resulted in a higher rate of PJI. In our cohort, most of the knee surgeries (96.6%) were done with the use of tourniquet and both groups were similar in terms of tourniquet use. We are not able to study the influence of tourniquet on blood loss and examine its potential confounding effect. The other confounding variable relates to the type of anesthesia used during TJA. Studies comparing neuraxial anesthesia with general anesthesia during TJA have reported

conflicting observations. But the overwhelming majority of studies demonstrate that the use of hypotensive neuraxial anesthesia leads to a lower blood loss, lower rate of SSI, wound complications, and fewer overall complications than those receiving general anesthesia [38–40]. Our results also confirmed previous studies.

Bivariate analysis has shown that in addition to TXA, other variables such as male gender, BMI, Charlson Comorbidity Index, rheumatoid arthritis, renal disease, liver disease, coagulopathy, anemia, operation time, and length of stay affect the PJI rate after primary TJA which is supported by the literature [2,3,13,15,21,23,41–44]. After adjusting for all variables related to outcomes, TXA was an independent risk factor protecting against PJI. The protective effect of TXA was more in hip joints and nonanemic patients.

The unadjusted and adjusted models predicting PJI may be affected by variations in patients with or without TXA. Patients who did not receive TXA in the current study tended to be male, older, sicker, anemic, higher BMI, and less likely to receive aspirin for DVT prophylaxis or neuraxial anesthesia. These reasons for not receiving TXA were due to previous traditional contraindications of TXA in our hospital. These variables are also predisposing factors for increased PJI. Based on recent studies on the safety of TXA, the indications for TXA are increasing even in sicker patients [33,35,45,46]. In an effort to remove the influence of these variables, we did perform a regression analysis to identify the influence of TXA administration alone on the incidence of PJI. The regression analysis revealed that TXA is associated with reduced PJI after primary TJA.

This study does have several strengths. First, we extracted data from the clinical charts of all these patients that resulted in obtaining more accurate data than relying on administrative data. Second, all patients at our institution are followed closely for the first 90 days by nurse navigators. Thus, it is unlikely that any complications or readmissions in these patients, even when presented to an outside facility, were missed. Third, we attempted to study the influence of various risk factors on PJI and adjust for these confounding variables in the statistical analyses. Finally, we used a strict definition for PJI removing subjective evaluations to define PJI.

This study also has some limitations. First, it is a retrospective study with all possible biases that such study design introduces. In our hospital, TXA has been used since 2012. Initially, it was administered in small groups of healthier patients. In 2013–2014, the indications for TXA were expanded gradually and we had more patients who received TXA. Recently, almost all TJAs are done with the administration of TXA. To have more patients with enough follow-ups and less time-dependent confounding factors, we selected all primary TJAs from January 1, 2013 to June 31, 2017. This restricted the number of cases and probably affected our results in the multivariate model for anemic patients. Second, this study was not performed

using matching criteria. We attempted to perform matching based on strict criteria, but even in case of one-by-one matching, a high attrition was observed. Third, this study was completed at an urban, tertiary referral center that does a high volume of primary TJA, and results may not be generalizable to all centers.

Conclusion

Based on the findings of this study, TXA appears to be associated with reduced PJI following primary TJA. Although the exact reason for such finding is unknown, we speculate that the beneficial effect on TXA may relate to its efficacy in reducing blood loss and the need for allogeneic blood transfusion, both of which are known to influence SSI after all surgical procedures including TJA.

References

- [1] Kurtz SM, Lau E, Watson H, Schmier JK, Parvizi J. Economic burden of periprosthetic joint infection in the United States. *J Arthroplasty* 2012;27:61–65e1.
- [2] Kurtz SM, Ong KL, Lau E, Bozic KJ, Berry D, Parvizi J. Prosthetic joint infection risk after TKA in the Medicare population. *Clin Orthop Relat Res* 2010;468:52–6.
- [3] Ong KL, Kurtz SM, Lau E, Bozic KJ, Berry DJ, Parvizi J. Prosthetic joint infection risk after total hip arthroplasty in the Medicare population. *J Arthroplasty* 2009;24:105–9.
- [4] Kurtz SM, Medel FJ, MacDonald DW, Parvizi J, Kraay MJ, Rimnac CM. Reasons for revision of first-generation highly cross-linked polyethylenes. *J Arthroplasty* 2010;25:67–74.
- [5] Graves S, Davidson D, de Steiger R, Tomkins A. Annual report 2010. Australian National Joint Replacement Registry. Adelaide: Australian Orthopaedic Association; 2010.
- [6] Bozic KJ, Kurtz SM, Lau E, Ong K, Chiu V, Vail TP, et al. The epidemiology of revision total knee arthroplasty in the United States. *Clin Orthop Relat Res* 2010;468:45–51.
- [7] Bozic KJ, Kurtz SM, Lau E, Ong K, Vail TP, Berry DJ. The epidemiology of revision total hip arthroplasty in the United States. *J Bone Joint Surg Am* 2009;91:128–33.
- [8] 2010 Annual report. The Norwegian Arthroplasty Register. Bergen: Centre of Excellence of Joint Replacements; 2010.
- [9] Garellick G, Karrholm J, Rogmark C, Herbergs P. Annual report 2008. Swedish Hip Arthroplasty Register. Göteborg: Department of Orthopaedics, Sahlgrenska University Hospital; 2009. <https://registercentrum.blob.core.windows.net/shpr/r/Annual-report-2008-HkLEFUTje.pdf>. [Accessed 17 March 2011].
- [10] Lidgjen L, Sundberg M, Dahl AW, Robertsson O. Annual report 2010. The Swedish Knee Arthroplasty Register. Lund: Dept. of Orthopedics, Lund University Hospital; 2010.
- [11] Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *J Bone Joint Surg Am* 2007;89:780–5.
- [12] Bhandari M, Smith J, Miller LE, Block JE. Clinical and economic burden of revision knee arthroplasty. *Clin Med Insights Arthritis Musculoskelet Disord* 2012;5:89–94.
- [13] Akonjom M, Battenberg A, Beverland D, Choi JH, Fillingham Y, Gallagher N, et al. General assembly, prevention, blood conservation: proceedings of international Consensus on Orthopedic Infections. *J Arthroplasty* 2019;34:S147–55.
- [14] Greenky M, Gandhi K, Pulido L, Restrepo C, Parvizi J. Preoperative anemia in total joint arthroplasty: is it associated with periprosthetic joint infection? *Clin Orthop Relat Res* 2012;470:2695–701.
- [15] Viola J, Gomez MM, Restrepo C, Maltenfort MG, Parvizi J. Preoperative anemia increases postoperative complications and mortality following total joint arthroplasty. *J Arthroplasty* 2015;30:846–8.
- [16] Friedman R, Homering M, Holberg G, Berkowitz SD. Allogeneic blood transfusions and postoperative infections after total hip or knee arthroplasty. *J Bone Joint Surg Am* 2014;96:272–8.
- [17] Benoni G, Carlsson A, Petersson C, Fredin H. Does tranexamic acid reduce blood loss in knee arthroplasty? *Am J Knee Surg* 1995;8:88–92.
- [18] Sukeik M, Alshryda S, Haddad FS, Mason JM. Systematic review and meta-analysis of the use of tranexamic acid in total hip replacement. *J Bone Joint Surg Br* 2011;93 B:39–46.
- [19] Yang ZG, Chen WP, Wu LD. Effectiveness and safety of tranexamic acid in reducing blood loss in total knee arthroplasty: a meta-analysis. *J Bone Joint Surg Am* 2012;94:1153–9.
- [20] Shohat N, Bauer T, Buttaro M, Budhiparama N, Cashman J, Della Valle CJ, et al. Hip and knee section, what is the definition of a periprosthetic joint infection (PJI) of the knee and the hip? Can the same criteria be used for both joints? Proceedings of International Consensus on Orthopedic Infections. *J Arthroplasty* 2018;34:S325–7. <https://doi.org/10.1016/j.arth.2018.09.045>.
- [21] Greenky M, Gandhi K, Pulido L, Restrepo C, Parvizi J. Preoperative anemia in total joint arthroplasty: is it associated with periprosthetic joint infection? *Clin Orthop Relat Res* 2012;470:2695–701.
- [22] Bozic KJ, Ong K, Lau E, Berry DJ, Vail TP, Kurtz SM, et al. Estimating risk in Medicare patients with THA: an electronic risk calculator for periprosthetic joint infection and mortality. *Clin Orthop Relat Res* 2013;471:574–83.
- [23] Rasouli MR, Restrepo C, Maltenfort MG, Purtill JJ, Parvizi J. Risk factors for surgical site infection following total joint arthroplasty. *J Bone Joint Surg Am* 2014;96:e158.
- [24] Eka A, Chen AF. Patient-related medical risk factors for periprosthetic joint infection of the hip and knee. *Ann Transl Med* 2015;3:233.
- [25] Alshryda S, Sukeik M, Sarda P, Blenkinsopp J, Haddad FS, Mason JM. A systematic review and meta-analysis of the topical administration of tranexamic acid in total hip and knee replacement. *Bone Joint J* 2014;96-B:1005–15.
- [26] Keating EM, Meding JB. Perioperative blood management practices in elective orthopedic surgery. *J Am Acad Orthop Surg* 2002;10:393–400.
- [27] Tornero E, Pereira A, Bravo J, Angulo S, Basora M, Marcos M, et al. Transfusion of packed red blood cells stored >14 days was associated with a higher risk of infection after hip revision arthroplasty. *Hip Int* 2016;26:132–7.
- [28] Newman ET, Watters TS, Lewis JS, Jennings JM, Wellman SS, Attarian DE, et al. Impact of perioperative allogeneic and autologous blood transfusion on acute wound infection following total knee and total hip arthroplasty. *J Bone Joint Surg Am* 2014;96:279–84.
- [29] Han X, Gong G, Han N, Liu M. Efficacy and safety of oral compared with intravenous tranexamic acid in reducing blood loss after primary total knee and hip arthroplasty: a meta-analysis. *BMC Musculoskelet Disord* 2018;19:430.
- [30] Frisch NB, Wessell NM, Charters MA, Yu S, Jeffries JJ, Silverton CD. Predictors and complications of blood transfusion in total hip and knee arthroplasty. *J Arthroplasty* 2014;29:189–92.
- [31] Huerfano E, Huerfano M, Shanaghan KA, Gonzalez Della Valle A. Topical tranexamic acid in revision total knee arthroplasty reduces transfusion rates and may be associated with earlier recovery. *J Arthroplasty* 2018;34:S249–55.
- [32] Huang Z, Xie X, Li L, Huang Q, Ma J, Shen B, et al. Intravenous and topical tranexamic acid alone are superior to tourniquet use for primary total knee arthroplasty: a prospective, randomized controlled trial. *J Bone Joint Surg Am* 2017;99:2053–61.
- [33] Fillingham YA, Ramkumar DB, Jevsevar DS, et al. The safety of tranexamic acid in total joint arthroplasty: a direct meta-analysis. *J Arthroplasty* 2018;33:3070–3082.e1.
- [34] Heller S, Secrist E, Shahi A, Chen AF, Parvizi J. Tranexamic acid can be administered to arthroplasty patients who receive aspirin for venous thromboembolic prophylaxis. *J Arthroplasty* 2016;31:1437–41.
- [35] Kim YT, Kang MW, Lee JK, Lee YM, Kim JI. Combined use of topical intra-articular tranexamic acid and rivaroxaban in total knee arthroplasty safely reduces blood loss, transfusion rates, and wound complications without increasing the risk of thrombosis. *BMC Musculoskelet Disord* 2018;19:227.
- [36] Alcelik I, Pollock RD, Sukeik M, Bettany-Saltikov J, Armstrong PM, Fisman P. A comparison of outcomes with and without a tourniquet in total knee arthroplasty: a systematic review and meta-analysis of randomized controlled trials. *J Arthroplasty* 2012;27:331–40. <https://doi.org/10.1016/j.arth.2011.04.046>.
- [37] Yi S, Tan J, Chen C, Chen H, Huang W. The use of pneumatic tourniquet in total knee arthroplasty: a meta-analysis. *Arch Orthop Trauma Surg* 2014;134:1469–76.
- [38] Helwani MA, Avidan MS, Ben Abdallah A, Kaiser DJ, Clohisey JC, Hall BL, et al. Effects of regional versus general anesthesia on outcomes after total hip arthroplasty: a retrospective propensity-matched cohort study. *J Bone Joint Surg Am* 2015;97:186–93.
- [39] Pugely AJ, Martin CT, Gao Y, Schweizer ML, Callaghan JJ. The incidence of and risk factors for 30-day surgical site infections following primary and revision total joint arthroplasty. *J Arthroplasty* 2015;30:47–50.
- [40] Liu J, Ma C, Elkassabany N, Fleisher LA, Neuman MD. Neuraxial anesthesia decreases postoperative systemic infection risk compared with general anesthesia in knee arthroplasty. *Anesth Analg* 2013;117:1010–6.
- [41] Jansen E, Varonen M, Huhtala H, Lehto MU, Lumio J, Konttinen YT, et al. Incidence of prosthetic joint infections after primary knee arthroplasty. *J Arthroplasty* 2010;25:87–92.
- [42] Bozic KJ, Lau E, Kurtz S, Ong K, Berry DJ. Patient related risk factors for postoperative mortality and periprosthetic joint infection in Medicare patients undergoing TKA. *Clin Orthop Relat Res* 2012;470:130–7.
- [43] Poultsides LA, Triantafyllopoulos GK, Sakellariou VI, Memtsoudis SG, Sculco TP. Infection risk assessment in patients undergoing primary total knee arthroplasty. *Int Orthop* 2018;42:87–94.
- [44] Resende VAC, Neto AC, Nunes C, Andrade R, Espregueira-Mendes J, Lopes S. Higher age, female gender, osteoarthritis and blood transfusion protect against periprosthetic joint infection in total hip or knee arthroplasties: a systematic review and meta-analysis. *Knee Surg Sports Traumatol Arthrosc* 2018;26:1–13.
- [45] Gillette BP, DeSimone LJ, Trousdale RT, Pagnano MW, Sierra RJ. Low risk of thromboembolic complications with tranexamic acid after primary total hip and knee arthroplasty. *Clin Orthop Relat Res* 2013;471:150–4.
- [46] Sabbag OD, Abdel MP, Amundson AW, Larson DR, Pagnano MW. Tranexamic acid was safe in arthroplasty patients with a history of venous thromboembolism: a matched outcome study. *J Arthroplasty* 2017;32:S246–50.