

XXIII Congress of Chemists and Technologists of Macedonia

BF

**INFLUENCE OF YEAST STRAINS
ON PHENOLIC COMPOSITION AND
ANTIOXIDANT ACTIVITY OF VRANEC WINES**

Violeta Ivanova-Petropulos^{1*}, Arianna Ricci², Dusko Nedelkovski³, Violeta Dimovska¹,
Giuseppina P. Parpinello², Andrea Versari²

*e-mail: violeta.ivanova@ugd.edu.mk

1-Faculty of Agriculture, University “Goce Delčev”, Krste Misirkov bb, Štip, Republic of
Macedonia

2-Department of Agricultural and Food Sciences, University of Bologna, Piazza Goidanich 60,
Cesena (FC) 47521, Italy

3-Institute of Agriculture, “Ss. Cyril and Methodius” University, Aleksandar Makedonski bb, 1000
Skopje, Republic of Macedonia

The phenolic composition of ten autochthonous monovarietal Vranec red wines produced with different yeast strains was determined by HPLC-DAD analysis. Vranec wines were fermented with the following *Saccharomyces cerevisiae* yeast strains: Clos, RC212, D254, BDX (from Lallemand), and six Vinalco yeasts isolated from the Tikveš wine region (Republic of Macedonia). A total of 19 phenolic compounds were identified and quantified in wines and among them the malvidin-3-glucoside and its derivatives were the major compounds. Wines fermented with Vinalco yeast presented higher amount of anthocyanins as well as phenolic acids (hydroxycinnamic and hydroxybenzoic) compared to the wines fermented with Lallemand yeasts. The content of malvidin-3-glucoside in Vranec wines fermented with Vinalco yeasts ranged between 235–887 mg/L, whereas Vranec wines fermented with Lallemand yeasts contained 335 to 355 mg/L. The caftaric acid, the main cinnamic acid derivative in all wines, was found in concentration between 176–507 mg/L, followed by coutaric and fertaric acids (14–89 and 11–49 mg/L, respectively). Wines showed relatively high value of the antioxidant activity that ranged between 99–117 mg/L Trolox equivalents, regardless the yeast strain used for fermentation.

Key words: anthocyanins; phenolic acids; HPLC; Vinalco yeast; Vranec wine.

Acknowledgement: Financial support provided by the Research Fund of the University “Goce Delčev” – Štip for the project titled “Polyphenolic and aroma profile of Vranec wines fermented with isolated yeasts from Tikveš wine area” is gratefully acknowledged. Additionally, this work was financially supported by the ERASMUS Mundus Action 2 Project covering the study stay of Violeta Ivanova-Petropulos at the University of Bologna, Italy.